The impact of punctuated infectiousness on infectious disease dynamics

Author One[1], Author Two[1,2], Author Three[2]

[1]Department of Computer Science, University of Colorado Boulder, Boulder CO USA
[2]BioFrontiers Institute, University of Colorado Boulder, Boulder CO USA
[3]Department of Epidemiology, Colorado School of Public Health, Aurora CO USA

Abstract

1 Introduction

The trajectory of an epidemic is governed by two key factors: the basic reproduction number (R_0), which is the expected number of secondary infections that an infectious person will produce in an otherwise susceptible population; and the generation interval distribution, $g(\tau)$, which is the distribution of times at which those secondary infections are expected to occur relative to the index case's infection ($\tau = 0$) [Wallinga and Lipsitch, 2007].

Both the reproduction number and the generation interval distribution are expectations taken over a large population. At the individual level, the individual reproduction number (ν_i) is the expected number of infections that a particular infectious person i will produce, such that $R_0 = E_i[\nu_i]$ [Lloyd-Smith et al., 2005]. Similarly, the "individual infectiousness kernel" ($a_i(\tau)$) describes the time distribution of new infections caused by a particular index case i, such that $g(\tau) = E_i[a_i(\tau)]$ [Champredon and Dushoff, 2015, Svensson, 2007].

For many pathogens, the shape of the individual infectiousness kernel is poorly understood. Contact tracing to count infections is hard enough; measuring precisely the times of those infections is even harder. When it is done, it is usually averaged over many individuals, so that the "infectiousness period" assigned to a person in models reflects some average over multiple individuals—either the generation interval itself, or some intermediate between the individual infectiousness kernel and the generation interval distribution. Some evidence suggests that the infectious period for SARS-CoV-2 and influenza, at the individual level, may be highly punctuated, such that opportunities for superspreading only exist over the course of a few hours.

At the individual level, communicating about the likely timing of a person's infectiousness is important: it is good to know if you will be very infectious for a short time, or mildly infectious for a long time, so that you can shape your behavior accordingly. Additionally, a punctuated individual infectiousness kernel could impact epidemic dynamics. Even if the epidemic has the same mean-field behavior, it could impact the stochastic dynamics via highly synchronized transmission events, especially when an outbreak is small. It could interplay with control measures, especially those designed to intervene on infectiousness at the individual level like test- or symptom-based screening and isolation. Indeed, the impact of the punctuatedness of the individual infectiousness kernel has been examined as a sensitivity analysis in various epidemiological studies. **Dan; others?** Stochastic effects are known to cascade through macroscopic features of an epidemic **Morris2025**. However, we lack a comprehensive understanding of how the sharpness of the individual infectiousness kernel, independent of other features like the individual reproduction number, basic reproduction number, and generation interval distribution, impacts epidemic dynamics under both uncontrolled and controlled settings. Furthermore, we lack an understanding of how to measure the shape of the individual infectiousness kernel reliably, along with the settings when the kernel's shape is statistically identifiable.

Here, we address these gaps.

2 Expressing the Hudson–Kermack–McKendrick model in terms of individual infectiousness

2.1 The individual infectiousness profile

Early work by Hudson, and later by Kermack and McKendrick [Kermack and McKendrick, 1927], modeled disease transmission in a population using an integral equation. In modern notation [Breda et al., 2012], the force of infection is

$$F(t) = \int_0^\infty F(t-\tau)\, S(t-\tau)\, A(\tau)\, d\tau \tag{1}$$

where $S(t)$ is the density of susceptible individuals in the population at time t, and $A(\tau)$ is the population-level "infectiousness profile", a curve describing the expected contribution to the force of infection from an individual who was infected τ time units ago. The incidence of disease is simply the product of the force of infection and the susceptible density, $F(t) \cdot S(t)$; thus, the infectiousness profile $A(\tau)$ is the fundamental object that determines how an epidemic unfolds.

The infectiousness profile is related to the basic reproduction number, R_0, and the generation interval, $g(\tau)$ [Breda et al., 2012]:

$$R_0 = \int_0^\infty A(\tau)\, d\tau \tag{2}$$

$$g(\tau) = \frac{A(\tau)}{R_0} \tag{3}$$

Furthermore, particular choices of $A(\tau)$ yield disease transmission dynamics that are equivalent to more familiar ordinary differential equation-based models [Breda et al., 2012]. For example, the SEIR model

$$\frac{dS}{dt} = -\beta S I \tag{4}$$

$$\frac{dE}{dt} = \beta S I - \gamma E \tag{5}$$

$$\frac{dI}{dt} = \gamma E - \alpha I \tag{6}$$

$$\frac{dR}{dt} = \alpha I \tag{7}$$

is obtained when

$$A(\tau) = \beta \frac{\gamma}{\gamma - \alpha} \left(e^{-\alpha\tau} - e^{-\gamma\tau} \right) \tag{8}$$

An individual person's infectiousness kernel $a_i(\tau)$ may differ substantially from $A(\tau)$, and multiple different $a_i(\tau)$ may yield the same $A(\tau)$. For example, the SEIR model is often conceptually motivated by considering an individual who undergoes a latent period of length $X \sim \text{Exp}(\gamma)$ during which they are not infectious, followed by an infectious period of length $Y \sim \text{Exp}(\alpha)$ during which

they are infectious at level β. Mathematically,

$$a_i(\tau) = \begin{cases} \beta & \tau \in [\tau_i^{\text{on}}, \tau_i^{\text{on}} + \tau_i^{\text{off}}] \\ 0 & \text{otherwise} \end{cases} \quad \text{where } \tau_i^{\text{on}} \sim \text{Exp}(\gamma) \text{ and } \tau_i^{\text{off}} \sim \text{Exp}(\alpha) \tag{9}$$

It can be shown that $E_i[a_i(\tau)] = \lim_{n\to\infty} \frac{1}{n} \sum_{i=1}^{n} a_i(\tau) = A(\tau)$ [Breda et al., 2012]. However, other $a_i(\tau)$ also yield the same $A(\tau)$, including

$$a_i(\tau) = \beta \frac{\gamma}{\gamma - \alpha} \left(e^{-\alpha\tau} - e^{-\gamma\tau}\right) \tag{10}$$

where each person's infectiousness profile is a smooth curve that is identical to the population-level infectiousness profile, and

$$a_i(\tau) = \frac{\beta}{\gamma} \cdot \delta_{t_i}(\tau) \qquad \text{where } t_i \text{ is distributed according to the generation interval density, } A(\tau)/R_0 \tag{11}$$

where each person's infectiousness profile is a delta function, such that all their infectiousness is concentrated at a single moment (**Figure XX**).

Like $A(\tau)$, $a_i(\tau)$ is related to the individual reproduction number, ν_i, and the individual generation interval distribution, $\xi_i(\tau)$:

$$\nu_i = \int_0^\infty a_i(\tau)\, d\tau \tag{12}$$

$$\xi_i(\tau) = a_i(\tau)/\nu_i \tag{13}$$

Figure 1: The individual infectiousness profile impacts stochastic epidemic trajectories. (A-C) Schematic diagrms of three individual infectionsless profiles that yield equivalent mean-field dynamics: the smooth (A), stepwise (B), and spike (C) profiles (Eq xx, xx, xx). (D-I) Deterministic epidemic trajectories (blue) are numerical solutions of the mean-field renewal equation model, equivalent to a standard SEIR model with mean latent period XX and mean infectious period XX. For each individual infectiousness profile type, 1,000 stochastic simulations were generated (grey), depicting the cumulative infections (D-F) and daily new infection counts (G-I) for each profile type.

2.2 A Gamma convolutional model for the individual infectiousness profile

The "stepwise" expression for $a_i(\tau)$ in Eq. (9) differs from the "smooth" and "spike" expressions in Eqs. (10)–(11) in a fundamental way: For Eq. (9), ν_i differs across individuals ($\nu_i \sim$ XX), while for Eqs. (10)–(11), ν_i is uniform across individuals ($\nu_i = R_0$). As a result, the probability of epidemic extinction differs for the stepwise profile relative to both the smooth and spike profiles – but this is due to an inherent difference in the distribution of the individual reproduction number, not to the timing of the infections. Because of this fundamental dissimilarity—and because the impact of variation in ν_i on epidemic dynamics has been discussed extensively elsewhere [**?**]—we will focus most of the manuscript on individual infectiousness profiles of the type in Eqs. (10)–(11), where the individual infectiousness ν_i is uniform across the population. That way, the impact of the timing of infectiousness can be examined in isolation.

To achieve this, we seek a family of functions that allow us to model individual infectiousness profiles with the following characteristics:

- All have uniform total infectiousness: $\int_0^\infty a_i(\tau)\, d\tau = R_0$
- Smooth, single-peaked distributions
- All have identical shape, except for a time shift (no mixtures of broad smooth distributions with narrow spiky distributions)

A candidate set of functions that obeys this is the set of Gamma convolutional curves. Let $A(\tau)$ be Gamma distributed (up to a scaling constant):

$$A(\tau) = R_0\, f_A(\tau) \qquad \text{where } f_A(\tau) \sim \text{Gamma}(\alpha, r), \quad r = \alpha/\mu \tag{14}$$

where μ is the mean generation interval and α controls the shape of the population-level infectiousness profile.

Now, let

$$a_i(\tau) = R_0\, f_a(\tau - s_i) \qquad \text{where } f_a(\tau) \sim \text{Gamma}(\kappa, r) \quad \text{and} \quad s_i \sim \text{Gamma}(\alpha - \kappa, r) \tag{15}$$

Here, $\kappa \in (0, \alpha)$ is a parameter governing the punctuatedness of $a_i(\tau)$, with smaller κ yielding a more concentrated infectiousness profile (**Figure XX**). Since the sum of two Gamma-distributed random variables with the same rate is also Gamma distributed (with the same rate and with shape equal to the sum of the two component shapes), this formulation is guaranteed to converge to $A(\tau)$ in expectation (**Supplementary Methods**). Furthermore, the κ-values may differ across individuals; regardless of the distribution they are drawn from, the Gamma sum property ensures that $E_i[a_i(\tau)] = A(\tau)$ (**Supplementary Methods**, **Figure XX**).

2.3 Splitting the infectiousness profile into biological infectiousness and contacts

From Eqs. (12)–(13), the individual infectiousness profile can be expressed in terms of the individual generation interval distribution and the individual reproduction number:

$$a_i(\tau) = \xi_i(\tau)\,\nu_i \tag{16}$$

A different and more general decomposition is also possible, into factors relating to a person's "biological infectiousness", $b_i(\tau)$, and "contact potential", $c_i(t)$:

$$a_i(\tau) = b_i(\tau)\,c_i(t_i + \tau) \tag{17}$$

The biological infectiousness, $b_i(\tau)$, is a nonnegative density function that integrates to 1 ($\int_0^\infty b_i(\tau)\,d\tau = 1$) that describes how a person's potential infectiousness is spread over time. The contact potential, $c_i(t)$, is a nonnegative time-varying function that describes the expected number of individuals person i would infect if all their infectiousness were concentrated at the single moment t (i.e., if $b_i(\tau)$ were a delta function). Thus, $c_i(t)$ incorporates both a person's contact rate and the probability of transmission given contact. It can immediately be seen that if $c_i(t)$ is constant, then we recover Eq. (16) where $b_i(\tau)$ is the individual generation interval distribution and $c_i(t) = \nu_i$. For time-varying $c_i(t)$, this form shows that the individual reproduction number and the individual generation interval distribution depend not just on the time since infection τ, but also on clock time t. For infection time t^*, we obtain

$$\nu_i(t^*) = \int_0^\infty b_i(\tau)\,c_i(t^* + \tau)\,d\tau \tag{18}$$

$$\xi_i(t^*, \tau) = \frac{b_i(\tau)\,c_i(t^* + \tau)}{\nu_i(t^*)} \tag{19}$$

This decomposition feeds back into the population-level model. If we assume that b_i and c_i are independent (i.e., an individual's biological infectiousness is independent of their contact pattern), then

$$A(\tau) = E[a_i(\tau)] = B(\tau) \cdot C(t_i + \tau) \tag{20}$$

where $B(\tau) = E_i[b_i(\tau)]$ and $C(t_i+\tau) = E_i[c_i(t+\tau)]$. Likewise, if C is constant in time, this reduces to the familiar

$$A(\tau) = g(\tau)\,R_0 \tag{21}$$

Splitting $a_i(\tau)$ into biological infectiousness b_i and contact potential c_i allows us to separately consider how the shape of a person's infectiousness, unconditional on their contact patterns, impacts transmission dynamics under various contact patterns.

For the Gamma convolutional model, we have

$$b_i(\tau) = f_a(\tau - s_i) \tag{22}$$

and

$$c_i(t) = R_0 \quad \text{for all } i, t \tag{23}$$



Figure 2: The additive Gamma model for individual infectiousness profiles, with a split into biological and behavioral components.

2.4 Stochastic simulations

To simulate the infections produced by a single person, we generate the number of infections using a Poisson(ν_i) distribution and distribute them according to $a_i(\tau)/\nu_i$. For the Gamma convolutional model, the timing distribution has closed form and is easy to sample from using built-in packages. For more arbitrary individual infectiousness profiles (e.g., obtained by convolving a Gamma-distributed $b_i(\tau)$ with a periodic contact function $c_i(t)$, we use Poisson thinning:

3 The impact of the individual infectiousness profile on uncontrolled epidemics

3.1 Punctuated individual infectiousness profiles generate later and more variably-timed epidemics.

Even in large, well-mixed populations, stochasticity can durably impact the trajectory of an epidemic. One way this stochasticity enters is via a stochastic time shift, where epidemics take a variable amount of time to reach the exponential growth phase [Morris et al., 2024]. Here, we address the question of whether the shape of the individual infectiousness profile impacts this time shift distribution, holding all else equal.

Let $I_b(t)$ be the (stochastic) number of infections at time t for a single epidemic realization, where the subscript "b" marks that it is a branching process. Let $I_d(t)$ be its mean-field deterministic counterpart. Then, we can approximate $I_b(t)$ as $I_d(t+\varsigma)$, *i.e.,* the branching process is approximately equal to the deterministic process with a random time shift.

The time shift is

$$\varsigma = \frac{\log W - \log E[W]}{r}$$

where r is the epidemic's exponential growth rate and $W = \lim_{t\to\infty} W(t)$ is the limiting value of the rescaled branching process $W(t) = e^{-rtI_b(t)}$. The process $W(t)$ is well studied, and knowing its distribution fully determines the distribution of the time shift ς.

The expectation of W is equal to the initial condition ($E[W] = I_0$). For simplicity, we will assume $I_0 = 1$ throughout this section, though the findings can be easily extended to general initial conditions. Here, we assess the impact of the individual infectiousness profile's breadth, ϕ, on (a) the variance of W and (b) the mode of a Gamma approximation to W obtained via moment matching.

It can be shown that

$$\text{Var}(W) = \frac{c^\alpha + \sigma^{(1-\psi)\alpha} - 1}{1 - c^\alpha}$$

where c depends on R_0 and α but not ψ. Thus, broad profiles ($\psi \to 1$) minimize Var(W) and narrow profiles ($\psi \to 0$) maximize it.

We can approximate the distribution of W using moment matching to a generalized Gamma distribution. With the Gamma-shaped individual infectiousness profiles, the moments have closed-form expressions. Using two-moment matching to a Gamma distribution, we find that

$$\text{mode}_\varsigma = \frac{1}{r}\log(1 - \text{CV}^2_{\text{cond}})]$$

3.2 Narrower individual infectiousness profiles yield greater uncertainty in epidemic growth rate estimates.

Based on simulations.

3.3 Narrower biological infectiousness profiles interact with periodic contact patterns to generate overdispersion in secondary infections.

Closed-form solution:

$$k = \frac{2}{\bar{c}^2} A^2 \left(1 + \frac{\omega^2}{r^2}\right)^{\psi\alpha} = \frac{2}{\bar{c}^2} A^2 \left(1 + (\frac{z^*_{per}}{z_{per}})^2\right)^{\psi\alpha}$$

Figure 3: Impact of the individual infectiousness profile on epidemic timing. Survival curves for time-to-100. growth rate estimate box-and-whiskers. Overdispersion/k vs psi heatmap.

4 The impact of the individual infectiousness profile on epidemic control

4.1 Detection and isolation

We first consider the detect-and-isolate interventions. Detection may occur through various mechanisms, e.g., the onset of symptoms or a positive diagnostic test.

We can express the effectiveness of a detect-and-isolate intervention in terms of the *test effectiveness* [Middleton and Larremore, 2024].

If the detection mechanism is independent of the individual infectiousness profile (e.g., detection is anchored to the infection time but does not depend on what happens thereafter), the imapct of the individual infectiousness profile washes out (**Supplementary Text**). However, if detectability is anchored somehow to the infectiousness profile – which is likely to be the case in general – the individual profile begins to matter.

Here, we consider detection strategies that are anchored to the mode of the individual infectiousness profile (i.e., the time of peak infectiousness). Let $D(\tau)$ be the cumulative probability that a person has been detected by time τ.

To begin, we consider a fixed, deterministic detection time relative to the mean - this might correspond to symptoms arriving deterministically some amount of time prior to (or after) peak infectiousness.

Maybe: just show a TE profile for a spike, smooth, and $\psi = 0.5$ profile for a deterministic offset (size of offset on the horizontal axis, TE on the vertical); and then show the analogous curves for when the detection profile is random (say, normal, where now the horizontal axis captures the mean of the distribution, and we pick some reasonable standard deviation). This should illustrate how the timing of detection (with variation) interplays with the punctuation of the infectiousness profile. That might be enough here.

Then we can focus on the implications for the generation interval distribution and secondary infection distribution.

I want to try my hand at making this figure: what do we need?

- The general detection distribution framework
- Narrower infectiousness profiles transfer detection into an all-or-nothing proposition
- Detection and isolation shorten the generation interval distribution, moreso for smoooth profiles, which means that epidemics may grow (and shrink) faster than a naive application of TE would suggest...
- while detection and isolation increases overdispersion in secondary infections, moreso for spike profiles, so that epidemics are less likely to take off (but more explosive) than would be expected with a naive application of TE.
- The takehome: detection and isolation reduce R_{eff} via TE, but a simple application of TE won't tell us everything we need to know: instead, we need to know something about the

individual infectiousness profile to properly anticipate how detect-and-isolate interventions will impact epidemic dynamics.

- Furthermore: narrower infectiousness profiles make detect-and-isolate interventions sensitive to small shifts, especially near the infectiousnesss profile itself. So, if symptoms move from slightly after to slighlty before infectiousness on average, this will have an outsize impact on narrow profiles vs. smooth ones. Similarly, if a small increase in testing cadence could have a huge impact for spike profiles, if we shift the mean of D from slightly after to slightly before peak infectiousness; but in other areas, it'll hardly have any impact on spike profiles, whereas the change in effectiveness for smooth profiles should be more gradual.

- This last point is relevant for when detection is anchored to the mode (or otherwise anchored to the individual infectiousness profile); otherwise, using the population average I think should be fine (e.g. if detection depends only on time of infection but not subsequent infectiousness).

- The takeaway from this: if we want to know the impact of shifting biology (changes in the symptom timing) or shifts in recommendations (changes in the cadence of testing), we should know something about the individual infectiousness profile (and how it relates to detectability).

- Adding in delays/turnaround time will also impact all this.

4.2 Gathering size restrictions

5 Identifiability and inference of the individual infectiousness profile's shape

While the concentration of the individual infectiousness profile can impact how an epidemic unfolds, we lack frameworks for estimating it. Many approaches exist for estimating the generation interval, but there has not been a systematic approach developed to assess how individual infectiousness varies and to specify where on the continuum we sit—i.e., how much of the variation in the generation interval distribution is attributable to variation between individuals, and how much to variation in infection timing within individuals.

We begin by assessing under which conditions we can hope to infer the shape of the individual infectiousness profile. We restrict to gamma-distributed timings, and consider cases where we assume that κ is shared across individuals and where κ comes from a beta distribution governed by hyperparameters α and β. Then, when κ is identifiable, we introduce a hierarchical Bayesian inference strategy to infer the shape of the individual infectiousness profile given contact tracing data.

For this section: let's imagine that there's a single psi value shaed among everyone. We can use a Beta prior. Two things to consider: how do we estimate ψ given a list of infection times (assume we know them exactly)? And how do we estimate ψ given infection times and knowledge of $c_i(t)$? My hope is that this will emphasize that knowing the "negatives" – i.e., the contacts that "should" have led to transmission but did not – improves our inferential power substantially. Then, I want to think about how to achieve a given power in each scenario across R0, assuming some fixed generation interval distribution. That should be enough for now. We should probably include some amount of uncertainty in infection times for this base analysis, and we should post code for running the analysis given a line list of infection times and contacts. In the Discussion, we need to address the fact that there could be different ψ_i values across individuals, themselves drawn from a distribution, so we'd want some kind of hierarhcical Bayesian model for inference here; also, we don't consider differences in the height (i.e. inherent biological differences in ν_i / b_i), which is another key thing to include in the inference.

6 Discussion

References

D. Breda, O. Diekmann, W. F. De Graaf, A. Pugliese, and R. Vermiglio. On the formulation of epidemic models (an appraisal of Kermack and McKendrick). *Journal of Biological Dynamics*, 6(sup2):103–117, September 2012. ISSN 1751-3758, 1751-3766. doi: 10.1080/17513758.2012.716454. URL http://www.tandfonline.com/doi/abs/10.1080/17513758.2012.716454.

David Champredon and Jonathan Dushoff. Intrinsic and realized generation intervals in infectious-disease transmission. *Proceedings of the Royal Society B: Biological Sciences*, 282(1821): 20152026, December 2015. ISSN 0962-8452, 1471-2954. doi: 10.1098/rspb.2015.2026. URL https://royalsocietypublishing.org/doi/10.1098/rspb.2015.2026.

William Ogilvy Kermack and A. G. McKendrick. A contribution to the mathematical theory of epidemics. *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, 115(772):700–721, August 1927. ISSN 0950-1207, 2053-9150. doi: 10.1098/rspa.1927.0118. URL https://royalsocietypublishing.org/rspa/article/115/772/700/2165/A-contribution-to-the-mathematical-theory-of.

J. O. Lloyd-Smith, S. J. Schreiber, P. E. Kopp, and W. M. Getz. Superspreading and the effect of individual variation on disease emergence. *Nature*, 438(7066):355–359, November 2005. ISSN 0028-0836, 1476-4687. doi: 10.1038/nature04153. URL https://www.nature.com/articles/nature04153.

Casey Middleton and Daniel B. Larremore. Modeling the transmission mitigation impact of testing for infectious diseases. *Science Advances*, 10(24):eadk5108, June 2024. ISSN 2375-2548. doi: 10.1126/sciadv.adk5108. URL https://www.science.org/doi/10.1126/sciadv.adk5108.

Dylan Morris, John Maclean, and Andrew J. Black. Computation of random time-shift distributions for stochastic population models. *Journal of Mathematical Biology*, 89(3):33, September 2024. ISSN 0303-6812, 1432-1416. doi: 10.1007/s00285-024-02132-6. URL https://link.springer.com/10.1007/s00285-024-02132-6.

Åke Svensson. A note on generation times in epidemic models. *Mathematical Biosciences*, 208 (1):300–311, July 2007. ISSN 00255564. doi: 10.1016/j.mbs.2006.10.010. URL https://linkinghub.elsevier.com/retrieve/pii/S0025556406002094.

J Wallinga and M Lipsitch. How generation intervals shape the relationship between growth rates and reproductive numbers. *Proceedings of the Royal Society B: Biological Sciences*, 274(1609): 599–604, February 2007. ISSN 0962-8452, 1471-2954. doi: 10.1098/rspb.2006.3754. URL https://royalsocietypublishing.org/doi/10.1098/rspb.2006.3754.

Acknowledgements

Funding

Author contributions

Supplementary Information

Supplementary Methods

Supplementary Figures

Supplementary Tables