



UNSW
A U S T R A L I A



UNIVERSITY OF NEW SOUTH WALES

SCHOOL OF MATHEMATICS AND STATISTICS

Assignment 2

Graph Theory

Author:
Ed McDonald

Student Number:
z3375335

SCHOOL OF MATHEMATICS & STATISTICS

ASSIGNMENT COVER SHEET

Student last name: _____

Student given name: _____

Student ID: _____

Course no & Name: _____

Lecturer/Tutor: _____

Class no: _____ Day: _____

Class time: _____ Due date: _____

Assessment/ Assignment Topic: _____

Student Academic Misconduct Declaration

www.lc.unsw.edu.au/plagiarism

I declare that this assessment item is my own work, except where acknowledged, and has not been submitted for academic credit elsewhere, and acknowledge that the assessor of this item may, for the purpose of assessing this item:

- Reproduce this assessment item and provide a copy to another member of the University; and or,
- Communicate a copy of this assessment item to a plagiarism checking service (which may then retain a copy of the assessment item on its database for the purpose of future plagiarism checking).

I certify that I have read and understand the University rules in respect of Student Academic Misconduct.

(By signing this declaration you are agreeing to the conditions above)

Signature: _____

Date: _____

School date stamp:

Question 1

For graphs G and H , we let consider the graph product $G \times H$ as having the vertex set $V(G) \times V(H)$, and we have an edge $(v, w)(v', w') \in E(G \times H)$ if and only if $v = v'$ and $ww' \in E(H)$ or $vv' \in E(G)$ and $w = w'$.

For a graph G , $\alpha(G)$ denotes the size of the largest independent set in G .

Lemma 1 (Part (a)). *Let G be a graph on $n \geq 2$ vertices, and let $r \geq 1$. Then*

$$\alpha(G \times K_r) \leq n.$$

Proof. It is sufficient to show that any set of $n + 1$ vertices in $G \times K_r$ has an adjacent pair. Let $(v_1, w_1), (v_2, w_2), \dots, (v_{n+1}, w_{n+1}) \in V(G \times K_r)$, where each $v_k \in G$ and each $w_k \in K_r$.

By the pigeonhole principle, not all the v_k can be distinct, so we must have $v_k = v_j$ for some $j \neq k$. Now since K_r is complete, $w_k w_j \in E(K_r)$. Hence, $(v_k, w_k)(v_j, w_j) \in E(G \times K_r)$, so any set of $n + 1$ vertices must have an adjacent pair.

Hence, $\alpha(G \times K_r) \leq n$. □

Lemma 2 (Part (b)). *For a graph G on $n \geq 2$ vertices, and $r \geq 1$, we have $\alpha(G \times K_r) = n$ if and only if $r = \chi(G)$.*

Proof. Let $r = \chi(G)$. We must show that $\alpha(G \times K_r)$ has an independent set of size n . Choose an r -colouring for G , $c : G \rightarrow \{1, 2, \dots, r\}$. Let $V_j = c^{-1}(\{j\})$ be the set of vertices of G with colour j .

Label the vertices of K_r as w_1, w_2, \dots, w_r .

Consider the set of vertices $S = \{(v, w_j) \mid v \in V_j, 0 \leq j \leq r\}$. We shall show that S is an independent set. Let $(v, w_j), (v', w_k) \in S$. Then the vertices $(v, w_j), (v', w_k)$ are adjacent if and only if one of two possibilities holds:

Firstly, if $w_j = w_k$, then $(v, w_j), (v', w_k)$ are adjacent if and only if $vv' \in E(G)$. But this is impossible since $v, v' \in V_j$. So v and v' both have colour j so cannot be adjacent.

The second case is that $j \neq k$, but we have $v = v'$. But then v has colour j , and v' has colour k . But this is impossible since $v = v'$.

Thus, the vertices $(v, w_j), (v', w_k)$ are never adjacent.

Now, the size of S is just $|S| = \sum_{j=1}^r |\{(v, w_j) \mid v \in V_j\}| = \sum_{j=1}^r |V_j| = |V(G)|$, since the colour classes form a partition of G .

Hence, $|S| = n$, so we have an independent set of size n . □

Corollary 1 (Part (c)). *For a graph G on $n \geq 2$ vertices, we have:*

$$\chi(G) = \min\{r > 0 : \alpha(G \times K_r) = n\}.$$

Proof. By lemma 2, we know that $\alpha(G \times K_r) = n$ when $r = \chi(G)$. We must show that $\alpha(G \times K_r) < n$ when $r < \chi(G)$.

We will show this by demonstrating how an independent set of size n in $G \times K_r$ induces a colouring of G with at most r colours. Since this is possible only when $r \geq \chi(G)$, this will prove that there are no independent sets of size n in $G \times K_r$ when $r < \chi(G)$.

Let $r \geq 1$, and suppose that S is an independent set of size n in $G \times K_r$, and label the vertices of K_r as w_1, w_2, \dots, w_r .

First we show that for all $v \in V$, there is some j so that $(v, w_j) \in S$. If this is not the case, then by the pigeonhole principle there must be some $v \in V$ such that $(v, w), (v, u) \in S$ for distinct $w, u \in K_r$. But then $wu \in E(K_r)$, so (v, w) and (v, u) are adjacent, which is impossible since S is independent.

Now we can define a function $c : V(G) \rightarrow \{1, 2, \dots, r\}$ as follows.

We define $c(v) = j$ if $(v, w_j) \in S$. This is well defined, since if $(v, w_j), (v, w_k) \in S$ are distinct, we must have that $w_j w_k \in E(K_r)$. Hence $c(v)$ is defined uniquely for at least some vertices in G .

In fact c is a colouring of G . To see this, let $vv' \in E(G)$. If $c(v) = c(v') = j$, then $(v, w_j), (v', w_j) \in S$. But then since v and v' are adjacent, we must have that (v, w_j) and (v', w_j) are adjacent. Since S is independent, we must then have $c(v) \neq c(v')$.

Hence c is a colouring of v with at most r colours, and we are done. \square

Question 2

For this question, G is a 3-connected graph and $xy \in E(G)$. We use the notation G/xy to denote the graph obtained from G by contracting xy .

Proposition 1 (Part (a)). *If G/xy is 3-connected, then $G - \{x, y\}$ is 2-connected.*

Proof. To show that $G - \{x, y\}$ is 2-connected, at the very least it must have at least three vertices. To show this, we note that by assumption G/xy is 3-connected, and $|V(G/xy)| = |V(G)| - 1$. Since G/xy is itself 3-connected, we have $|V(G/xy)| > 3$. Hence, $|V(G)| > 4$. Thus, $|V(G - \{x, y\})| > 2$.

Now we need to show that $G - \{x, y\}$ is connected, and for any vertex $v \in V(G - \{x, y\})$, $G - \{x, y, v\}$ is connected.

Now since G is by assumption 3-connected, automatically we have that $G - \{x, y\}$ is connected. Hence it is only required to show that $G - \{x, y, v\}$ is connected for all vertices $v \in V(G - \{x, y\})$.

So let $v \in V(G - \{x, y\})$. Hence $v \in V(G/xy)$. Let w be the vertex in G/xy formed from merging x and y .

Thus, $G/xy - \{v, w\}$ is connected since G/xy is 3-connected by assumption.

Let $p, q \in V(G - \{x, y, v\})$. Since $G/xy - \{v, w\}$ is connected, there is a path P joining p and q in G/xy which avoids v and w .

Hence every edge of P consists of edges of $G/xy - \{v, w\}$. Since every edge of $G/xy - \{v, w\}$ is an edge of G , P can be considered as a path in G . Since it avoids v and w , it must avoid x, y and v in G .

Thus $G - \{x, y, v\}$ is connected, and so $G - \{x, y\}$ is connected. \square

Proposition 2 (Part (b)). *Now if G/xy is not 3-connected, then $G - \{x, y\}$ is not 2-connected.*

Proof. We shall prove the contrapositive. Suppose that $G - \{x, y\}$ is 2-connected. In particular, this means that $G - \{x, y\}$ has at least three vertices, so G has at least five vertices. Hence, G/xy has at least four vertices.

Now to complete the proof we need to show that removing any 2 element subset of G/xy does not disconnect G/xy .

Now by assumption, $G - \{x, y\}$ is 2-connected. In particular this means that it is connected, and for any $v \in V(G) - \{x, y\}$ we have that $G - \{x, y, v\}$ is connected.

Now, let $u, w \in V(G/xy)$. Then there are two possibilities:

1. One of u or v is formed by contracting x and y .

2. Neither of u and v is formed by contracting x and y .

In the first case, we may assume without loss of generality that u is formed from contracting x and y . We now consider $G/xy - \{u, w\}$. This is simply $G - \{x, y, w\}$. But by assumption this is connected, hence $G/xy - \{u, w\}$ is connected.

In the second case, we have $u, v \in V(G - \{x, y\})$. Now, let $H = G/xy - \{u, v\}$.

Now, let $p, q \in V(H)$. We must show that there is a path joining p and q in H . Since neither u nor v is formed from contracting x and y , it is sufficient to show that there is a path joining p and q in G which avoids u and v . Hence, we search for a path in $G - \{u, v\}$. However, by assumption G is 3-connected, so such a path exists. Hence, H is connected.

Thus, G/xy is 3-connected, as required. \square

Question 3

For this question, G is a graph and $P_G(k)$ denotes the number of k -colourings of G . It is known that $P_G(k)$ is polynomial in k , and is hence called the chromatic polynomial of G .

From the problem sheets, we know that if G has n vertices then $P_G(k)$ is a monic polynomial of degree n , and if G has m edges then the coefficient of k^{n-1} is $-m$.

Since this is not proved in the notes, we give a short proof here. Let $x, y \in V(G)$, and let $G' = G - xy$ and G'' be the graph obtained by contracting x and y . Now a k -colouring of G is exactly a k -colouring of G' which assigns different colours to x and y . The k -colourings of G which assign the same colours to x and y are exactly the k -colourings of G'' . Hence, $P_G(k) = P_{G'}(k) - P_{G''}(k)$. Hence if we do induction on the number of edges of G , we can prove the required properties of $P_G(k)$.

Lemma 3 (Part (a)). *There is no graph G with chromatic polynomial*

$$P_G(k) = k^4 - 4k^3 + 8k^2 - 5k.$$

Proof. Let G be such a graph. Then G must have 4 vertices and 4 edges. Hence, by the handshaking lemma the average degree of G is 2. Hence G is 2-regular or has a vertex of degree 3.

If G is 2-regular, then $G = C_4$ is a cyclic graph on 4 vertices. Hence, in this case, the number of 2-colourings of G is 2.

However $P_G(2) = 16 - 32 + 32 - 10 = 6$, so $G \neq C_4$.

The only other case is that G has a vertex of degree 3. Since $|V(G)| = 4$, this means that G has a vertex v , connected to every other vertex. Hence if we have a 2-colouring of G , $c : V(G) \rightarrow \{1, 2\}$ we either have $c(v) = 1$ and all other vertices have colour 2, or $c(v) = 2$ and all other vertices have colour 1. However this cannot be a true colouring since there are four edges: so one edge must connect two vertices of colour 2. Hence there is no 2-colouring of G . Thus, $p_G(k) = 0 \neq 6$.

Thus we have exhausted all possible graphs with four vertices and four edges. Hence the given polynomial cannot be a chromatic polynomial. \square

Lemma 4 (Part (b)). *Let G be a graph, with $v \in V(G)$. Then the number of k -colourings of G which assign colour 1 to v is given by $P_G(k)/k$.*

Proof. Let C_i be the number of colourings which assign colour i to v . Then,

$$P_G(k) = C_1 + C_2 + \cdots + C_k.$$

Now by symmetry, we have $C_1 = C_i$ for all i , hence,

$$P_G(k) = kC_1$$

as required. \square

Lemma 5 (Part (c)). *Let G be a graph with connected components G_1, G_2, \dots, G_r . Then*

$$P_G(k) = \prod_{j=1}^r P_{G_j}(k).$$

Proof. Let $k \geq 1$. Then for each $1 \leq j \leq r$, a k -colouring of G induces a k -colouring of G_j . Thus each k -colouring of G is produced by an individual k colouring of each G_j . Hence, the number of k -colourings of G is given by a k -colouring of G_1 , times the number of k -colourings of G_2 , times the number of k -colourings of G_2 , and etc., until we have a k -colouring for G . \square

Lemma 6 (Part (d)). *Let $r > 0$ and let G have r connected components. If:*

$$P_G(k) = c_0 + c_1k + c_2k^2 + \cdots + c_nk^n$$

we have $c_0 = c_1 = \cdots = c_{r-1} = 0$ and $c_r \neq 0$.

In other words, if G has r connected components, then r is the minimal positive integer such that the coefficient of k^r in $P_G(k)$ is nonzero.


Alternatively, k divides $P_G(k)$ at least r times.

Proof. Let the connected components of G be $\{G_i\}_{i=1}^r$. Then by lemma 5, we have:

$$P_G(k) = \prod_{i=1}^r P_{G_i}(k).$$

By lemma 4, k divides $P_{G_i}(k)$. Hence, k divides $P_G(k)$ at least r times. \square

Now for Part (e), the following graph has chromatic polynomial $P_G(k) = k^2(k-1)^3(k-2)$:



Question 4

For this question, $k \geq 1$ and $G = (V, E)$ is a k -edge connected graph. We let $F_1, \dots, F_m \subseteq E$ be distinct sets of k edges which disconnected G . We let $\mathcal{F} = F_1 \cup F_2 \dots \cup F_m$, and let C_1, \dots, C_t be the connected components of $G - \mathcal{F}$.

Lemma 7 (Part (a)). *For each $i = 1, \dots, m$, we have:*

$$|\{e \in \mathcal{F} : e \cap C_i \neq \emptyset\}| \geq k.$$

Proof. Since G is k -connected, it suffices to show that $S_i = \{e \in \mathcal{F} : e \cap C_i \neq \emptyset\}$ disconnects G .

Consider $G - S_i$. Suppose that e is an edge joining C_i to $G - C_i$. Then since C_i is a connected component of $G - \mathcal{F}$, we must have $e \in \mathcal{F}$, and in particular $e \in S_i$. Hence $e \notin G - S_i$, so $G - S_i$ contains no edges joining C_i to $G - C_i$. Thus $G - S_i$ is disconnected, and hence $|S_i| \geq k$. \square

Corollary 2 (Part (b)). *We have $|\mathcal{F}| \geq kt/2$.*

Proof. Now, each $e \in \mathcal{F}$ is incident with at least one C_i , and at most two C_i . Let S_i as in lemma 7, and we have:

$$\mathcal{F} = S_1 \cup S_2 \cup \dots \cup S_t.$$

However, each element of \mathcal{F} occurs in at most two of the S_i . Thus the sum

$$\sum_{i=1}^t |S_i|$$

counts each element of \mathcal{F} at most twice. Hence,

$$\sum_{i=1}^t |S_i| \leq 2|\mathcal{F}|.$$

Hence, since for each i we have $|S_i| \geq k$ from lemma 7,

$$kt \leq 2|\mathcal{F}|.$$

So the result follows. □

Corollary 3 (Part (c)). *We have $|\mathcal{F}| \leq mk$, and hence $t \leq 2m$.*

Proof. Since $\mathcal{F} = F_1 \cup F_2 \cup \dots \cup F_m$, hence $|\mathcal{F}| \leq |F_1| + |F_2| + \dots + |F_m|$. Since each $|F_i| = k$, we then have $|\mathcal{F}| \leq mk$. Thus,

$$kt/2 \leq |\mathcal{F}| \leq mk.$$

and hence $t \leq 2m$. □

Question 5

For this question, H denotes a 3-uniform hypergraph on n vertices with n element vertex set V and m element hyperedge set E , and $m \geq n/3$.

Lemma 8 (Part (a)). *Let $U \subset V$ be a random vertex set, with $v \in U$ with probability p . Then $\mathbb{E}(|U|) = np$.*

Proof. Let $X : G \rightarrow \{0, 1\}$ be the indicator function of the set U . Then by the linearity of expectation,

$$\mathbb{E}(|U|) = \sum_{v \in V} \mathbb{E}(X(v)).$$

Now, $X(v) = 1$ with probability p and 0 with probability $1-p$. Hence $\mathbb{E}(X(v)) = p$, so $\mathbb{E}(|U|) = np$. \square

Lemma 9 (Part (b)). *Define Y to be the number of hyperedges contained in U , randomly chosen as in lemma 8, that is,*

$$Y = |\{e \in E : e \subseteq U\}|. \quad (1)$$

Then $\mathbb{E}(Y) = mp^3$.

Proof. Let $X : E \rightarrow \{0, 1\}$ be the indicator function of the set of hyperedges in U . Then by the linearity of expectation,

$$\mathbb{E}(Y) = \sum_{e \in E} \mathbb{E}(X(e)).$$

So we need to compute $\mathbb{E}(X(e))$ for a given hyperedge e .

Since $X(e) = 1$ if $e \subseteq U$, and 0 otherwise, we have that $\mathbb{E}(X(e))$ equal the probability that $e \subseteq U$. In otherwords, $\mathbb{E}(X(e))$ is the probability that U contains e .

Suppose that $e = \{u, v, w\}$. Then $\mathbb{E}(X(e)) = \mathbb{P}(u, v, w \in U)$. Since each vertex is chosen to be in U with probability p independently, we have that $\mathbb{E}(X(e)) = p^3$. Thus,

$$\begin{aligned} \mathbb{E}(Y) &= \sum_{e \in E} p^3 \\ &= mp^3. \end{aligned}$$

\square

Corollary 4 (Part (c)). *Hence, H contains an independent set with at least $\frac{2n^{3/2}}{3\sqrt{3m}}$ vertices.*

Proof. We will use the probabilistic method: we will define a random independent subset of H and show that the expected value of its size is at least $\frac{2n^{3/2}}{3\sqrt{3m}}$.

Let $U \subseteq H$ be chosen as in lemma 8, and let Y be the number of edges in U as in lemma 9.

If we remove one vertex from every edge in U , then we create an independent set. Hence we can remove Y vertices from H to form an independent set of size $|U| - Y$. By the linearity of expectation, and lemmas 8 and 9:

$$\begin{aligned}\mathbb{E}(|U| - Y) &= \mathbb{E}(|U|) - \mathbb{E}(Y) \\ &= np - mp^3.\end{aligned}$$

Hence there exists an independent set of size at least $np - mp^3$.

Since this holds for all $p \in [0, 1]$, we can choose p freely in this range. Let $p = \sqrt{n/3m}$. This is a true probability since by assumption $m \geq n/3$. Then,

$$\begin{aligned}np - mp^3 &= \sqrt{\frac{n}{3m}} \left(n - m \frac{n}{3m} \right) \\ &= \frac{2n^{3/2}}{3\sqrt{3m}}.\end{aligned}$$

Hence there is an independent set of at least this size. □

Remark 1. *Our choice for p in the proof of corollary 4 in fact maximises $np - mp^3$ for $p \in [0, 1]$, but it is not necessary to show this since we only require a lower bound.*