

UNSW AUSTRALIA

HONOURS THESIS

Quantised Calculus in One Dimension

Author:

Edward McDONALD

Supervisor:

Prof. Fedor SUKOCHEV

*A thesis submitted in fulfilment of the requirements
for the degree of Honours*

in the

Research Group Name
School of Mathematics and Statistics

May 2015

Declaration of Authorship

I, Edward McDONALD, declare that this thesis titled, 'Quantised Calculus in One Dimension' and the work presented in it are my own. I confirm that:

- This work was done wholly or mainly while in candidature for a research degree at this University.
- Where any part of this thesis has previously been submitted for a degree or any other qualification at this University or any other institution, this has been clearly stated.
- Where I have consulted the published work of others, this is always clearly attributed.
- Where I have quoted from the work of others, the source is always given. With the exception of such quotations, this thesis is entirely my own work.
- I have acknowledged all main sources of help.
- Where the thesis is based on work done by myself jointly with others, I have made clear exactly what was done by others and what I have contributed myself.

Signed:

Date:

“ If I am mad, it is mercy! May the gods pity the man who in his callousness can remain sane to the hideous end! ”

H. P. Lovecraft

UNSW AUSTRALIA

Abstract

Faculty of Science

School of Mathematics and Statistics

Honours

Quantised Calculus in One Dimension

by Edward McDONALD

In his 1994 book *Noncommutative geometry* Alain Connes introduced the quantised calculus. This is a means of doing calculus which makes sense in a high degree of generality, and has many properties analogous to the classical notion of “infinitesimal calculus”. This thesis is an in-depth study of quantised calculus in two cases: on the real line and on the circle.

Acknowledgements

I'd like to thank Zoey Proasheck, the wizard entrepreneur Vicks, and my mentor and tutor Vlad. Most thanks however go to Stkls, the tiny green flying alien from the future whom only I can see.

Contents

Declaration of Authorship	i
Abstract	iii
Acknowledgements	iv
Contents	v
1 Introduction	1
1.1 Classical Infinitesimals	1
1.2 Compact Operators as Infinitesimals	2
1.3 Expected Properties of infinitesimals	3
1.4 Quantised Differentials	5
1.5 Non-commutative geometry	6
1.5.1 Introduction to the non-commutative world	6
1.5.2 A brief introduction to Quantum Mechanics	7
1.5.3 Non-commutative spaces	9
2 Quantised Differentials on \mathbb{R} and \mathbb{T}	10
2.1 The Quantised Differential	10
2.2 Integral representation of the quantised differential on \mathbb{T}	12
2.3 The Cayley Transform	14
2.3.1 Notation	14
3 Basic Properties of Hankel Operators	16
3.1 Definition of a Hankel matrix	16
3.2 Bounded Hankel operators	17
3.3 Finite Rank Hankel operators	20
3.4 Compactness of Hankel Operators	22
3.5 Hankel Operators of Trace Class	23
3.6 Interpolation	27
4 Abstract Differential Algebra and Spectral Triples	28
4.1 Introduction	28
4.2 Classical Differential Algebra	28

4.3	Abstract Differential Algebra	29
4.3.1	Graded Differential Algebras	29
4.3.2	Kähler Differentials	30
4.3.3	Universal Differential Algebra	32
4.3.4	Spectral Triples	33
4.3.5	Connes Differentials	33
5	Weak \mathcal{L}^p membership of quantised differentials	35
5.1	Introduction	35
5.2	The setting	35
5.3	The Theorem	36
6	The Chain Rule in Quantised Calculus	38
6.1	Introduction	38
6.2	The setting	38
6.3	The Commutator Lemma	39
7	Higher Dimensions	40
7.1	Introduction	40
7.2	Commutators	41
7.3	Hilbert-Schmidt Commutators	41
7.4	Membership of \mathcal{L}^p	42
7.5	Double commutators	43
A	Classical Harmonic Analysis	44
A.1	Introduction	44
A.2	Abel summation	44
A.3	The Poisson Kernel	46
A.4	Fourier Series of Measures	48
A.5	Harmonic functions on \mathbb{D}	50
B	Lorentz Spaces	51
B.1	Introduction	51
B.2	Weak L^p -spaces	51
B.3	Non-increasing Rearrangements	52
B.4	Discrete spaces	52
C	Interpolation	53
C.1	Introduction	53
C.2	Abstract Interpolation Theory	54
C.3	Real Interpolation: The K Method	55
C.4	Complex Interpolation	55
C.5	Using interpolation	56
D	Group Actions on Von Neumann Algebras	57
D.1	Introduction	57
D.2	Compact Group actions	57

D.3	The space $\mathcal{L}^2(\mathcal{M}, \tau)$	59
D.4	Non-commutative Harmonic Analysis	61
D.5	The Operators δ_j	66
E	Dirac Operators	69
E.1	Introduction	69
E.2	Music, Clifford bundles and Modules	69
E.3	Relationship with differentials	70
	Bibliography	72

Chapter 1

Introduction

1.1 Classical Infinitesimals

According to the mathematicians of the 17th century, an “infinitesimal” is a quantity x that is smaller than any positive magnitude. In other words, for all $\varepsilon > 0$,

$$|x| < \varepsilon. \tag{1.1}$$

Mathematicians manipulated infinitesimals as though they were real numbers: addition, multiplication and division of infinitesimals were permitted.

Numerous definitions relied on the use of infinitesimals. For example, a function $f : \mathbb{R} \rightarrow \mathbb{R}$ was said to be continuous at x if $f(x+h) - f(x)$ is infinitesimal for all infinitesimals h .

Another example is that f was said to be differentiable when the quantity

$$\frac{df}{dx} := \frac{f(x+h) - f(x)}{h} \tag{1.2}$$

exists for all infinitesimals h , and does not depend on h .

Mathematicians distinguished between infinitesimals of different “sizes”, and if x is infinitesimal, then x^2 was said to be smaller than x , and algebraic manipulations could be performed with “sufficiently small infinitesimals ignored”.

However, as we are now aware, the condition in equation 1.1 implies that $x = 0$, so all manipulations involving infinitesimals are either trivial or impossible. Hence, the use of infinitesimals was banned from mathematics and their role in analysis was replaced with the concept of a *limit*.

Despite the difficulties with giving a non-contradictory definition of infinitesimals, there are numerous reasons that the concept is appealing. Using infinitesimals makes some definitions in analysis seem simpler, for example the above given definition of continuity. Infinitesimals can also be more intuitive than limits. For these reasons, some mathematicians have attempted to revive the concept of infinitesimals by defining them in a rigorous manner. Most famously, Abraham Robinson [?] uses model theory to define and analyse a field of *hyperreal numbers* which strictly contains the real numbers, and includes a plentiful supply of infinitesimals. Other approaches to rigorously defining infinitesimals include smooth infinitesimal analysis [?] and the Levi-Civita field [?].

A comprehensive history of the use of infinitesimals prior to the 19th century is [?].

1.2 Compact Operators as Infinitesimals

A relatively new approach to rigorously defining infinitesimals comes from non-commutative geometry. In this setting, all objects of interest such as functions, vector fields, differential forms, etc., are thought of as operators on a Hilbert space.

In what follows, let \mathcal{H} be a complex separable Hilbert space.

Suppose we wish to find a good definition of an “infinitesimal operator” on \mathcal{H} . A preliminary definition would be to say that an operator T is infinitesimal if for every $\varepsilon > 0$,

$$\|T\| < \varepsilon. \quad (1.3)$$

This definition is useless, as it implies that $T = 0$. However, we can get something close

Definition 1.1. Let $T \in \mathcal{B}(\mathcal{H})$. We say that T is *infinitesimal* if for every $\varepsilon > 0$, there is a finite dimensional subspace E such that

$$\|T|_{E^\perp}\| < \varepsilon. \quad (1.4)$$

(here $T|_{E^\perp}$ denotes the restriction of T to the orthogonal complement of E .)

Remark 1.2. Recall that the ideal of compact operators $\mathcal{K}(\mathcal{H}) \subseteq \mathcal{B}(\mathcal{H})$ is the closure of the set of finite rank operators. It is easy to see that T is infinitesimal (in the sense of definition 1.1) if and only if T is compact.

We require a way of measuring the “size” of an infinitesimal. According to definition 1.1, an infinitesimal is “zero modulo finite dimensional subspaces”, and it is sensible to consider the “size” of the infinitesimal as being measured by the speed at which the

dimension of the subspaces E must increase as ε moves towards zero. To this end, we define the singular values.

Definition 1.3. Let $T \in \mathcal{B}(\mathcal{H})$, and $n \geq 0$. Define

$$\mu_n(T) := \inf\{\|T - F\| : \text{rank}(F) \leq n\}. \quad (1.5)$$

Then $\mu_n(T)$ is called the n th singular value of T , and the sequence $\{\mu_n(T)\}_{n=0}^\infty$ is called the sequence of singular values.

Remark 1.4. For any operator T , the sequence $\{\mu_n(T)\}_{n=0}^\infty$ is strictly non-increasing. If T is compact, the sequence of singular values of T is decreasing and approaches 0.

We shall regard the *size* of T as being given by the *rate of decay* of $\{\mu_n(T)\}_{n=0}^\infty$.

1.3 Expected Properties of infinitesimals

According to 17th century mathematicians, infinitesimals were supposed to have a number of remarkable properties:

1. There exist non-zero infinitesimals.
2. Infinitesimals can be added, subtracted, multiplied and divided just like real numbers.
3. A real number multiplied by an infinitesimal produces an infinitesimal.
4. Infinitesimals can be split into “sizes”, and we can work modulo a particular size of infinitesimal.
5. If x is an infinitesimal, then x^2 is a smaller infinitesimal.
6. Given a function $f : \mathbb{R} \rightarrow \mathbb{R}$, there is a function df representing the infinitesimal variation in f , that is $df(x) = f(x + \delta) - f(x)$ for some fixed infinitesimal δ .
7. If a function f is smoother than a function g , then df is smaller than dg .
8. If f is a smooth function of x , we can write

$$df = f'(x)dx \quad (1.6)$$

provided that sufficiently small infinitesimals are ignored.

If we interpret compact operators as infinitesimals, we see that analogues of these statements remain true:

1. There indeed exist non-zero compact operators.
2. Compact operators can be added and subtracted like numbers, they can also be multiplied (although multiplication is not commutative). We cannot divide by a compact operator on an infinite dimensional Hilbert space. However it is true that if $XT = YT$ for operators $X, Y \in \mathcal{B}(\mathcal{H})$ for all $T \in \mathcal{K}(\mathcal{H})$, then $X = Y$.
3. $\mathcal{K}(\mathcal{H})$ forms an ideal of $\mathcal{B}(\mathcal{H})$, so a bounded operator multiplied by a compact operator produces a compact operator.
4. We can measure the size of a compact operator by the rate of decay of its sequence of singular values.

For item 5, we need the following lemma,

Lemma 1.5. *Let $T \in \mathcal{K}(\mathcal{H})$. Then for $n \geq 0$,*

$$\mu_n(T^2) \leq \mu_n(T)^2. \quad (1.7)$$

Proof. By definition,

$$\mu_n(T^2) = \inf\{\|T^2 - F\| : \text{rank}(F) \leq n\}. \quad (1.8)$$

Since if F has rank n , TF has rank not exceeding n , we have

$$\mu_n(T^2) \leq \inf\{\|T^2 - TF\| : \text{rank}(F) \leq n\} \quad (1.9)$$

$$\leq \inf\{\|T\|\|T - F\| : \text{rank}(F) \leq n\} \quad (1.10)$$

$$= \|T\|\mu_n(T). \quad (1.11)$$

... and then something else happens. □

Hence we have property 5: if T is infinitesimal, then the singular values of T^2 decay more rapidly than those of T .

Now for properties 5, 6 and 7, we need a way of defining df . This is precisely the role played by the quantised differential.

1.4 Quantised Differentials

The following definition may seem strange and unmotivated,

Definition 1.6. Consider the operator $\mathcal{D} = \frac{1}{i} \frac{d}{dx}$ of differentiation on \mathbb{R} . By the Borel functional calculus, we can define $F := \text{sgn}(\mathcal{D})$. F is called the Hilbert transform. For a function $f \in L^1(\mathbb{R})$ (which is complex valued) we have the (not necessarily everywhere defined) pointwise multiplication operator M_f , the operator

$$\bar{d}f := [F, M_f] \quad (1.12)$$

is an operator on $L^2(\mathbb{R})$ called the *quantised differential* of f .

$\bar{d}f$ is supposed to play the role of df in 17th century analysis. We use the symbol $\bar{d}f$ instead of df since in modern mathematics the symbol df has been taken to mean the exterior differential of f . It must be emphasised that these are very different objects.

Similar to the case of functions $f : \mathbb{R} \rightarrow \mathbb{C}$, we can also study functions on the circle. Let

$$\mathbb{T} := \{z \in \mathbb{C} : |z| = 1\}. \quad (1.13)$$

Then we have a differentiation operator

$$\mathcal{D} := \frac{1}{2\pi i} \frac{d}{d\theta}, \quad (1.14)$$

and $F := \text{sgn}(\mathcal{D})$. Then we can define quantised differentials $\bar{d}f := [F, M_f]$.

Remark 1.7. We have chosen to call the quantity $[F, M_f]$ a quantised differential, and to denote it $\bar{d}f$.

This terminology is not universal. In particular, the book [1] calls $[F, M_f]$ a quantised derivative and denotes it df .

We shall use the notation $\bar{d}f$ to prevent confusion with the exterior derivative df , and we call it a differential to be consistent with the idea of an “infinitesimal increment”.

It is not easy to motivate the definition $\bar{d}f := [F, M_f]$. Instead, we shall show that $\bar{d}f$ satisfies all the properties anticipated for a differential.

The two questions that we shall attempt to answer are as follows:

1. In what sense is it true that if f is smoother than g , then $\bar{d}f$ is smaller than $\bar{d}g$?
2. In what sense is it true that if φ is a function that is smooth on the range of a function f , then $\bar{d}\varphi(f) = \varphi'(f)\bar{d}f$?

In order to answer those questions, it is informative to give an explanation of the origin of the definition of $\bar{d}f$.

1.5 Non-commutative geometry

1.5.1 Introduction to the non-commutative world

Non-commutative geometry is a relatively new topic in mathematics. Non-commutative geometry is best thought of not as a collection of results, but instead as a perspective on mathematics.

It is difficult to give a completely satisfying general definition of non-commutative geometry, but one can say that non-commutative geometry is the study of non-commutative algebras which are somehow similar to algebras of functions on geometric spaces, using the methods and language of geometry.

The key idea which underlies most of non-commutative geometry is the duality between geometric spaces and algebras.

Example 1.1. *Let X be a compact Hausdorff space. Let $C(X)$ be the algebra of continuous complex valued functions on X . $C(X)$ naturally carries the structure of a commutative unital C^* -algebra. In fact, for any commutative unital C^* -algebra \mathcal{A} , there is a compact Hausdorff space K such that \mathcal{A} is isometrically $*$ -isomorphic to $C(K)$.*

Given a continuous function $f : X \rightarrow Y$ between compact Hausdorff spaces X and Y , there is a pull-back function $f_ : C(Y) \rightarrow C(X)$ defined by $f_*(h) = h \circ f$ for $h \in C(Y)$. Since $\text{id}_X = \text{id}_{C(X)}$, and $(f \circ g)_* = g_* \circ f_*$, the mapping $X \mapsto C(X)$ is a functor.*

*Let **CHTop** be the category of compact Hausdorff spaces with morphisms as continuous functions, and let **CUC*Alg** be the category of commutative unital C^* -algebras with morphisms as continuous $*$ -algebra homomorphisms.*

Thus we have a contravariant functor,

$$C : \mathbf{CHTop} \rightarrow \mathbf{CUC^*Alg}. \quad (1.15)$$

This effects an equivalence of categories,

$$\mathbf{CUC^*Alg} \cong \mathbf{CHTop}^{Op}. \quad (1.16)$$

Let $\mathbf{UC}^*\mathbf{Alg}$ be the category of unital C^* -algebras, which are not necessarily commutative. Inspired by the duality between commutative unital C^* -algebras and compact Hausdorff spaces, we define the category of (potentially) non-commutative compact Hausdorff spaces to be $\mathbf{UC}^*\mathbf{Alg}^{Op}$.

1.5.2 A brief introduction to Quantum Mechanics

Non-commutative geometry can be thought of simply as the study of non-commutative algebras using geometric language. However, much research in non-commutative geometry is inspired by quantum mechanics. It is therefore instructive to give a brief description of quantum mechanics.

Definition 1.8. A *quantum mechanical system* is a pair $(\mathcal{A}, \mathcal{H})$ where \mathcal{H} is a complex separable Hilbert space and \mathcal{A} is a $*$ -algebra of (possibly densely defined and unbounded) operators on \mathcal{H} . Denote the inner product on \mathcal{H} by (\cdot, \cdot) and $\|\psi\|^2 := (\psi, \psi)$.

A self-adjoint element of \mathcal{A} is called an *observable*. The elements of \mathcal{H} are called *states*.

Typically, we identify together elements of \mathcal{H} which differ by a nonzero scale factor, and the element $0 \in \mathcal{H}$ is ignored entirely. So technically we work over the *projective Hilbert space* $\mathbb{P}\mathcal{H}$.

To specify *the state of the system* $(\mathcal{A}, \mathcal{H})$ is the same as specifying some $\psi \in \mathcal{H}$.

We think of $(\mathcal{A}, \mathcal{H})$ as encoding a physical system. Typically we think of an observable as a measurable property of a system. The states correspond to potential configurations of the system.

Given an observable A , the potential range of values is the spectrum $\sigma(A)$. Unlike in classical mechanics, quantum mechanics can only make predictions that are probabilistic. The second difference to classical mechanics is that the act of observation changes the state of the system.

The link between $\sigma(A)$ and A is provided by the spectral theorem,

Theorem 1.9 (The Spectral Theorem). *Let $(\mathcal{A}, \mathcal{H})$ be a quantum mechanical system. Let $A \in \mathcal{A}$ be an observable. Then there is a projection valued measure E_A on $\sigma(A)$ such that*

$$A = \int_{\sigma(A)} \lambda dE_A(\lambda).$$

Using the spectral theorem, we may state our first postulate.

Postulate 1.10. Let $(\mathcal{A}, \mathcal{H})$ be a quantum mechanical system, in a state ψ .

Let $A \in \mathcal{A}$ be an observable, with associated spectral measure E_A .

For some Borel set $\Delta \subseteq \sigma(A)$, the probability that the observed value of A lies in Δ is given by

$$P_A(\Delta; \psi) := \frac{(\psi, E_A(\Delta)\psi)}{\|\psi\|^2} = \frac{\|E_A(\Delta)\psi\|^2}{\|\psi\|^2}.$$

(Note that this is the same for any scalar multiple of ψ , and is undefined for $\psi = 0$.)

Suppose that A is now observed, and the value is known to lie in the set $\Delta \subseteq \sigma(A)$. Then the state of the system changes to

$$\frac{E_A(\Delta)\psi}{\|E_A(\Delta)\psi\|}.$$

Remark 1.11. There are two extraordinary features of this postulate.

1. The state of the system changes upon observation.
2. The order of observation is important, since for any observables A and B , and $\Sigma \times \Delta \subseteq \sigma(A) \times \sigma(B)$, in general,

$$\frac{E_A(\Sigma)E_B(\Delta)\psi}{\|E_A(\Sigma)E_B(\Delta)\psi\|} \neq \frac{E_B(\Delta)E_A(\Sigma)\psi}{\|E_B(\Delta)E_A(\Sigma)\psi\|}.$$

Theorem 1.12. Let $(\mathcal{A}, \mathcal{H})$ be a quantum mechanical system which has state ψ . If $A \in \mathcal{A}$ is an observable, then the expected value of the observed value of A is

$$E(A; \psi) = \frac{(\psi, A\psi)}{\|\psi\|^2}$$

Proof. First consider the case when A is positive. Then $\sigma(A) \subseteq [0, \infty)$, and

$$E(A; \psi) := \int_{\sigma(A)} \lambda dP_A(\lambda; \psi).$$

By the monotone convergence theorem, this is

$$E(A; \psi) = \lim_{n \rightarrow \infty} \sum_{k=0}^{2^{2n}} \frac{k}{2^n} P_A \left(\left[\frac{k}{2^n}, \frac{k+1}{2^n} \right); \psi \right).$$

Now we use the definition of P_A in postulate 1.10:

$$\begin{aligned} E(A; \psi) &= \lim_{n \rightarrow \infty} \sum_{k=0}^{2^{2n}} \frac{k}{2^n} \frac{(\psi, E_A\left(\left[\frac{k}{2^n}, \frac{k+1}{2^n}\right]\right) \psi)}{\|\psi\|^2} \\ &= \frac{1}{\|\psi\|^2} \left(\psi, \lim_{n \rightarrow \infty} \left(\sum_{k=0}^{2^{2n}} \frac{k}{2^n} E_A\left(\left[\frac{k}{2^n}, \frac{k+1}{2^n}\right]\right) \right) \psi \right). \end{aligned}$$

Now, using $A = \int_{\sigma(A)} \lambda \, dE_A(\lambda)$, we get

$$E(A; \psi) = \frac{(\psi, A\psi)}{\|\psi\|^2}.$$

This completes the proof for $A \geq 0$. For general A , express A as a difference of two positive operators. \square

1.5.3 Non-commutative spaces

Inspired by quantum mechanics, we define the category of non-commutative spaces to be the opposite category of the category of complex $*$ -algebras.

Chapter 2

Quantised Differentials on \mathbb{R} and \mathbb{T}

2.1 The Quantised Differential

Definition 2.1. For $\varphi \in L^1(\mathbb{T})$, the *quantised differential* of φ is the (potentially unbounded, densely defined) linear operator

$$\mathring{d}\varphi := [F, M_\varphi] = [\mathbf{P}_+, M_\varphi].$$

In this chapter, we will discuss alternative descriptions of $\mathring{d}\varphi$.

We require an alternative description of $d\varphi$. For this, we introduce the Hardy spaces.

Definition 2.2. For $p \geq 2$, define

$$H^p(\mathbb{T}) := \mathbf{P}_+ L^p(\mathbb{T})$$

$$H_-^p(\mathbb{T}) := \mathbf{P}_- L^p(\mathbb{T}).$$

So that $L^2(\mathbb{T}) = H^2(\mathbb{T}) \oplus H_-^2(\mathbb{T})$.

Hence, we may consider the quantised derivative $\mathring{d}\varphi$ as a function from $H^2(\mathbb{T}) \oplus H_-^2(\mathbb{T})$.

Lemma 2.3. Let $\varphi \in L^2(\mathbb{T})$ and define $\varphi_+ := \mathbf{P}_+ \varphi$ and $\varphi_- := \mathbf{P}_- \varphi$. Then $\mathring{d}\varphi : H^2(\mathbb{T}) \oplus H_-^2(\mathbb{T}) \rightarrow H^2(\mathbb{T}) \oplus H_-^2(\mathbb{T})$ may be written as

$$\mathring{d}\varphi(f \oplus g) = (\mathbf{P}_+ M_{\varphi_+})g \oplus -(\mathbf{P}_- M_{\varphi_-})f$$

for $f \in H^2(\mathbb{T})$ and $g \in H_-^2(\mathbb{T})$.

Proof. This is a simple computation. Let $f \in H^2(\mathbb{T})$ and $g \in H_-^2(\mathbb{T})$. Then,

$$\bar{d}\varphi(f+g) = [\mathbf{P}_+, M_{\varphi_+} + M_{\varphi_-}](f+g).$$

Hence,

$$\begin{aligned} \bar{d}\varphi(f+g) &= [\mathbf{P}_+, M_{\varphi_+}]f + [\mathbf{P}_+, M_{\varphi_-}]f + [\mathbf{P}_+, M_{\varphi_+}]g + [\mathbf{P}_+, M_{\varphi_-}]g \\ &= (\mathbf{P}_+ M_{\varphi_-})f + (\mathbf{P}_+ M_{\varphi_+})g - M_{\varphi_-}f \end{aligned}$$

since $\mathbf{P}_+f = f$ and $\mathbf{P}_+g = 0$.

By the identity $\mathbf{P}_+ = I - \mathbf{P}_-$, we find

$$\bar{d}\varphi(f+g) = (\mathbf{P}_+ M_{\varphi_+})g - (\mathbf{P}_- M_{\varphi_-})f.$$

□

The problem of determining the boundedness of $\bar{d}\varphi$ is then reduced to the problem of determining the boundedness of operators of the form $\mathbf{P}_+ M_\psi : H_-^2(\mathbb{T}) \rightarrow H^2(\mathbb{T})$ and $\mathbf{P}_- M_\psi : H^2(\mathbb{T}) \rightarrow H_-^2(\mathbb{T})$ for $\psi \in L^2(\mathbb{T})$. We may simplify this further with the following lemma:

Lemma 2.4. *Let $\psi \in L^2(\mathbb{T})$. Then*

$$(\mathbf{P}_+ M_\psi)^* = \mathbf{P}_- M_{\bar{\psi}}.$$

and therefore $\mathbf{P}_+ M_\psi$ is bounded if and only if $\mathbf{P}_- M_{\bar{\psi}}$ is.

Proof. Let $e_k(z) = z^k$.

This is again a simple computation. Let $m, n \in \mathbb{Z}$ with $m \geq 0$ and $n < 0$. Then,

$$\begin{aligned} \langle (\mathbf{P}_+ M_\psi)e_n, e_m \rangle &= \int_{\mathbb{T}} \sum_{k > -n} \hat{\psi}(k) \zeta^{k+n-m} d\mathbf{m}(\zeta) \\ &= \hat{\psi}(m-n). \end{aligned}$$

Similarly,

$$\begin{aligned} \langle e_n, (\mathbf{P}_- M_{\bar{\psi}})e_m \rangle &= \int_{\mathbb{T}} \sum_{k > m} \hat{\varphi}(k) \zeta^{n-m+k} d\mathbf{m}(\zeta) \\ &= \hat{\psi}(m-n). \end{aligned}$$

Hence, $(\mathbf{P}_+ M_\psi)^* = \mathbf{P}_- M_{\bar{\psi}}$.

□

Therefore, we only need to study operators of the form $\mathbf{P}_- M_\psi$. So for $\psi \in L^2(\mathbb{T})$, define

$$H_\psi := \mathbf{P}_- M_\psi : H^2 \rightarrow H_-^2.$$

We study these operators using the Fourier transform. Use the standard basis $\{z^n\}_{n \geq 0}$ on $H^2(\mathbb{T})$ and the standard basis with negative indices $\{z^{-n}\}_{n \geq 0}$ on $H_-^2(\mathbb{T})$.

Let $\psi \in L^2(\mathbb{T})$. Then in the bases above, H_ψ has matrix representation with (n, k) th entry $\hat{\psi}(-n - k)$.

This means that H_ψ is represented by a *Hankel matrix*. So we require results on the boundedness of Hankel matrices

2.2 Integral representation of the quantised differential on \mathbb{T}

The following theorem allows us to describe the quantised differential as an integral operator.

Lemma 2.5.

$$\text{p.v.} \int_{\mathbb{T}} \frac{1}{\tau - 1} d\mathbf{m}(\tau) = -\frac{1}{2}$$

where the principal value is defined to be

$$\lim_{\varepsilon \rightarrow 0} \int_{|\tau - 1| > \varepsilon} \frac{1}{\tau - 1} d\mathbf{m}(\tau).$$

Proof. Note that

$$\text{p.v.} \int_{\mathbb{T}} \frac{1}{\tau - 1} d\mathbf{m}(\tau) = \text{p.v.} \int_{\text{Im}(\tau) > 0} \frac{1}{\bar{\tau} - 1} + \frac{1}{\tau - 1} d\mathbf{m}(\tau).$$

Hence,

$$\text{p.v.} \int_{\mathbb{T}} \frac{1}{\tau - 1} d\mathbf{m}(\tau) = \text{p.v.} \int_{\text{Im}(\tau) > 0} 2 \text{Re} \left(\frac{1}{\tau - 1} \right) d\mathbf{m}(\tau).$$

However, if $\tau = \exp(i\theta) \neq 1$, then

$$\begin{aligned} \operatorname{Re} \left(\frac{1}{\tau - 1} \right) &= \operatorname{Re} \left(\frac{e^{-i\theta/2}}{2i \sin(\theta/2)} \right) \\ &= -\frac{1}{2}. \end{aligned}$$

Hence,

$$\text{p.v.} \int_{\mathbb{T}} \frac{1}{\tau - 1} d\mathbf{m}(\tau) = 2 \text{p.v.} \int_{\operatorname{Im}(\tau) > 0} -\frac{1}{2} d\mathbf{m}(\tau) = -\frac{1}{2}.$$

□

Theorem 2.6. *Let $\varphi \in L^2(\mathbb{T})$. Then*

$$\mathbf{P}_+ \varphi(z) = \text{p.v.} \int_{\mathbb{T}} \frac{\varphi(\tau)}{1 - \bar{\tau}z} d\mathbf{m}(\tau) + \frac{1}{2} \varphi(z)$$

and hence,

$$H\varphi(z) = \text{p.v.} \int_{\mathbb{T}} \frac{\varphi(\tau)}{1 - \bar{\tau}z} d\mathbf{m}(\tau).$$

where in both equations, the principal value means that the integral is to be taken along the set $\{\zeta : |\zeta - z| > \varepsilon\}$ and then consider the limit $\varepsilon \rightarrow 0$.

Proof. It is sufficient to check this on the basis elements $e_n(z) = z^n$ for $n \in \mathbb{Z}$.

First let $n \geq 0$. Then

$$\text{p.v.} \int_{\mathbb{T}} \frac{\tau^n}{1 - \bar{\tau}z} d\mathbf{m}(\tau) = \text{p.v.} \int_{\mathbb{T}} \frac{z^n \tau^n}{1 - \bar{\tau}} d\mathbf{m}(\tau)$$

by translation invariance. Hence,

$$\begin{aligned} \text{p.v.} \int_{\mathbb{T}} \frac{\tau^n}{1 - \bar{\tau}z} d\mathbf{m}(\tau) &= z^n \text{p.v.} \int_{\mathbb{T}} \frac{\tau^{n+1}}{\tau - 1} d\mathbf{m}(\tau) \\ &= z^n \text{p.v.} \int_{\mathbb{T}} \frac{\tau^{n+1} - 1}{\tau - 1} + \frac{1}{\tau - 1} d\mathbf{m}(\tau) \\ &= z^n \text{p.v.} \int_{\mathbb{T}} 1 + \tau + \tau^2 + \cdots + \tau^n d\mathbf{m}(\tau) + z^n \text{p.v.} \int_{\mathbb{T}} \frac{1}{\tau - 1} d\mathbf{m}(\tau) \\ &= z^n + z^n \text{p.v.} \int_{\mathbb{T}} \frac{1}{\tau - 1} d\mathbf{m}(\tau) \\ &= \frac{1}{2} z^n \end{aligned}$$

where the last step follows from lemma 2.5.

Suppose $n > 0$, then

$$\text{p.v.} \int_{\mathbb{T}} \frac{\tau^{-n}}{1 - \bar{\tau}z} d\mathbf{m}(\tau) = z^{-n} \text{p.v.} \int_{\mathbb{T}} \frac{\tau^{1-n}}{\tau - 1} d\mathbf{m}(\tau)$$

by translation invariance. Hence,

$$\begin{aligned} \text{p.v.} \int_{\mathbb{T}} \frac{\tau^{-n}}{1 - \bar{\tau}z} d\mathbf{m}(\tau) &= z^{-n} \text{p.v.} \int_{\mathbb{T}} \frac{1}{\tau^n - \tau^{n-1}} d\mathbf{m}(\tau) \\ &= z^{-n} \text{p.v.} \int_{\mathbb{T}} \frac{\tau^n}{1 - \tau} \\ &= -\frac{1}{2} z^{-n}. \end{aligned}$$

Hence,

$$\text{p.v.} \int_{\mathbb{T}} \frac{\tau^n}{1 - \bar{\tau}z} d\mathbf{m}(\tau) = \begin{cases} \frac{1}{2} z^n & \text{if } n \geq 0 \\ -\frac{1}{2} z^n & \text{if } n < 0. \end{cases}$$

So the result follows. □

So we have the following integral form of the quantised derivative. Let $\varphi, f \in L^2(\mathbb{T})$. Then

$$\bar{\partial}\varphi(f)(z) = ([H, M_\varphi]f)(z) = \text{p.v.} \int_{\mathbb{T}} \frac{\varphi(z) - \varphi(\tau)}{1 - \bar{\tau}z} f(\tau) d\mathbf{m}(\tau).$$

2.3 The Cayley Transform

2.3.1 Notation

$\mathbb{H} = \{z \in \mathbb{C} : \Im(z) > 0\}$ denotes the upper half plane, and $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ denotes the open unit ball. $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$.

We use normalised Haar measure on \mathbb{T} , denoted \mathbf{m} . Lebesgue measure on \mathbb{R} is denoted λ , and two dimensional Lebesgue measure on \mathbb{C} is denoted \mathbf{m}_2 .

Throughout these notes, ω denotes the *Cayley transform*. $\omega : \mathbb{C} \setminus \{1\} \rightarrow \mathbb{C}$, and

$$\omega(\zeta) = i \frac{1 + \zeta}{1 - \zeta}, \quad \zeta \in \mathbb{D}.$$

For a Banach space E , and a measure space (X, Σ, μ) , we define

$$\|f\|_{L^p(X;E)} = \left(\int_X \|f\|_E^p d\mu \right)^{1/p}$$

for $p \in (0, \infty)$, and

$$\|f\|_{L^\infty(X;E)} = \sup_{x \in X} \|f(x)\|_E$$

for a weakly measurable $f : X \rightarrow E$. We define $L^p(X;E)$ as the set of measurable $f : X \rightarrow E$ with $\|f\|_{L^p(X;E)} < \infty$. As usual, we identify together functions on a measure space (X, Σ, μ) which agree μ -almost everywhere.

$L^0(X;E)$ denotes the set of all (μ -almost everywhere equivalence classes of) weakly measurable functions from X to E .

When X is a set with counting measure, we denote $L^p(X;E)$ as $\ell^p(X;E)$.

Suppose $\zeta \in \mathbb{T}$. Provided that $\zeta \neq 1$, we see that $\omega(\zeta)$ is defined, and ω maps $\mathbb{T} \setminus \{1\}$ smoothly to \mathbb{R} . Thus for $f \in L^0(\mathbb{R};E)$, we can define $\tilde{f} \in L^0(\mathbb{T};E)$ by

$$\tilde{f} := f \circ \omega^{-1}.$$

Thus we can define the important operator $U : L^0(\mathbb{T};E) \rightarrow L^0(\mathbb{R};E)$,

$$(Uf)(x) = \frac{1}{\sqrt{\pi}} \frac{(f \circ \omega^{-1})(x)}{x + i}.$$

It is the purpose of these notes to collate various results concerning the images of certain subspaces of $L^0(\mathbb{T};E)$ under U .

Chapter 3

Basic Properties of Hankel Operators

3.1 Definition of a Hankel matrix

A Hankel matrix is an infinite matrix $\{M_{j,k}\}_{j,k \geq 0}$ whose (j,k) th entry depends only on $j+k$. If $a = \{a_j\}_{j \geq 0}$, Let $M_a = \{a_{j+k}\}_{j,k \geq 0}$ be the Hankel matrix with (j,k) th entry a_{j+k} . That is,

$$M_a = \begin{pmatrix} a_0 & a_1 & a_2 & a_3 \cdots \\ a_1 & a_2 & a_3 & a_4 \cdots \\ a_2 & a_3 & a_4 & a_5 \cdots \\ a_3 & a_4 & a_5 & a_6 \cdots \\ \vdots & \vdots & \vdots & \vdots \end{pmatrix}.$$

An infinite matrix does not necessarily define an operator on $\ell^2(\mathbb{N})$, however any infinite matrix can be identified with a linear operator on the dense subset $c_{00}(\mathbb{N}) \subset \ell^2(\mathbb{N})$ of sequences of finite support.

For a sequence $a \in c_{00}(\mathbb{N})$, and an infinite matrix $M = (M_{j,k})_{j,k \geq 0}$, we define $Ma \in c_{00}(\mathbb{N})$ as $Ma = \{\sum_{k=0}^{\infty} M_{j,k} a_k\}_{j=0}^{\infty}$.

Hence we shall interchangeably talk about infinite matrices and linear operators $c_{00}(\mathbb{N}) \rightarrow c_{00}(\mathbb{N})$.

Since $c_{00}(\mathbb{N})$ is dense in $\ell^2(\mathbb{N})$, if an infinite matrix considered as an operator on $c_{00}(\mathbb{N})$ is bounded on $c_{00}(\mathbb{N})$ in the ℓ^2 -norm, then the matrix extends uniquely to an operator on $\ell^2(\mathbb{N})$. Conversely, any operator on $c_{00}(\mathbb{N})$ which extends to a bounded operator on $\ell^2(\mathbb{N})$ is bounded on $c_{00}(\mathbb{N})$ in the ℓ^2 -norm, and the extension to $\ell^2(\mathbb{N})$ is unique.

Denote the inner product on $\ell^2(\mathbb{N})$ as $(a, b) := \sum_{n=0}^{\infty} \overline{a_n} b_n$, which we note is linear in the second argument.

We let $\mathbb{T} = \{\zeta \in \mathbb{C} : |\zeta| = 1\}$. \mathbb{T} is a compact topological group, and we let \mathbf{m} be the normalised Haar measure (or arc length measure). Let $z : \mathbb{T} \rightarrow \mathbb{T}$ be the identity function. From now on, denote $L^p(\mathbb{T}) := L^p(\mathbb{T}, \mathbf{m})$.

Given $\varphi \in L^1(\mathbb{T}, \mathbf{m})$. For $n \in \mathbb{Z}$, the n th Fourier coefficient is defined as

$$\hat{\varphi}(n) := \int_{\mathbb{T}} z^{-n} \varphi \, d\mathbf{m}.$$

See [?] for the elementary properties of the Fourier transform.

It is proved in [?], that for any $f \in L^1(\mathbb{T}, \mathbf{m})$, we have

$$f = \lim_{r \rightarrow 1^-} \sum_{n \in \mathbb{Z}} \hat{f}(n) r^n z^n.$$

where the limit is in the L^1 sense. Hence we define the Riesz projection \mathbf{P}_+ , for $f \in L^1(\mathbb{T}, \mathbf{m})$,

$$\mathbf{P}_+ f = \lim_{r \rightarrow 1^-} \sum_{n \geq 0} \hat{f}(n) r^n z^n.$$

This limit exists in L^1 by the dominated convergence theorem.

For $p \in [1, \infty]$, we define the p th Hardy space $H^p(\mathbb{T})$ as the image of $L^p(\mathbb{T})$ under \mathbf{P}_+ .

A *polynomial* on \mathbb{T} is a finite linear combination of the *monomials* $\{z^n\}_{n \in \mathbb{Z}}$. We call the space of polynomials $P(\mathbb{T})$. An *analytic polynomial* is a polynomial consisting only of non-negative powers of z , we denote $P_+(\mathbb{T})$ for the space of analytic polynomials.

The Fourier transform gives a vector space isomorphism between $P(\mathbb{T})$ and $c_{00}(\mathbb{Z})$, and $P_A(\mathbb{T})$ and $c_{00}(\mathbb{N})$.

3.2 Bounded Hankel operators

It is of interest to determine when a Hankel operator defines a bounded linear operator on $\ell^2(\mathbb{N})$. This is answered completely by the *Nehari theorem*, which we cover now.

Theorem 3.1. *Let $a = \{a_j\}_{j=0}^{\infty}$ be a sequence. Then the associated Hankel matrix M_a defines a bounded linear operator on $\ell^2(\mathbb{N})$ if and only if there exists $\psi \in L^{\infty}(\mathbb{T}, \mathbf{m})$ such that*

$$\hat{\psi}(m) = a_m$$

for $m \geq 0$.

Proof. Suppose first that there is $\psi \in L^\infty(\mathbb{T})$ such that $\hat{\psi}(m) = a_m$ for $m \geq 0$.

Choose $f, h \in P_A(\mathbb{T})$, so that $\hat{f}, \hat{h} \in c_{00}(\mathbb{N})$.

Let $g \in P_A(\mathbb{T})$ be given by $g = \sum_{n=0}^{\infty} \overline{\hat{g}(n)} z^n$. Let $q = fg$.

Then we compute,

$$\begin{aligned} (\hat{h}, M_a \hat{f}) &= \sum_{j,k \geq 0} \overline{\hat{h}(j)} a_{j+k} \hat{f}(k) \\ &= \sum_{j,k \geq 0} \overline{\hat{h}(j)} \hat{\psi}(j+k) \hat{f}(k) \\ &= \sum_{j \geq 0} \hat{\psi}(j) \sum_{k=0}^j \overline{\hat{g}(j-k)} \hat{f}(k) \\ &= \sum_{j \geq 0} \hat{\psi}(j) \hat{q}(j) \\ &= \int_{\mathbb{T}} \psi(\zeta) q(\bar{\zeta}) d\mathbf{m}(\zeta). \end{aligned}$$

Hence,

$$\begin{aligned} |(\hat{h}, M_a \hat{f})| &\leq \|\psi\|_\infty \|q\|_1 \\ &\leq \|\psi\|_\infty \|f\|_2 \|h\|_2 \\ &= \|\psi\|_\infty \|\hat{f}\|_2 \|\hat{h}\|_2. \end{aligned}$$

And thus M_a is bounded on $\ell^2(\mathbb{N})$.

Conversely, suppose that M_a is bounded on $\ell^2(\mathbb{N})$.

Let \mathcal{L} be the linear functional on $P(\mathbb{T})$ defined by

$$\mathcal{L}(q) := \sum_{n \geq 0} a_n \hat{q}(n).$$

If $a \in \ell^1(\mathbb{N})$, then \mathcal{L} is bounded on $H^1(\mathbb{T})$, since the inverse fourier transform of a is in $L^\infty(\mathbb{T})$. Now let us prove in this case that $\|\mathcal{L}\| \leq \|M_a\|$.

Let $q \in H^1(\mathbb{T})$, with $\|q\|_1 \leq 1$. Then $q = fg$ for some $f, g \in H^2(\mathbb{T})$ with $\|f\|_2, \|g\|_2 \leq 1$.

Then we can compute,

$$\begin{aligned} |\mathcal{L}(q)| &= \left| \sum_{n=0}^{\infty} a_n \sum_{m=0}^n \hat{f}(m) \hat{g}(n-m) \right| \\ &= \sum_{n,m \geq 0} a_{n+m} \hat{f}(n) \hat{g}(m) \\ &= (M_a \hat{f}, \hat{g}). \end{aligned}$$

And hence,

$$|\mathcal{L}(q)| \leq \|M_a\| \|f\|_2 \|g\|_2 \leq \|M_a\|.$$

So $\mathcal{L}(q)$ is bounded on $H^1(\mathbb{T})$ whenever $a \in \ell^1(\mathbb{N})$.

Now we consider a to be an arbitrary sequence such that M_a is bounded. Let $r \in (0, 1)$ and

$$a^{(r)} = \{r^j a_j\}_{j \geq 0}.$$

Then $a^{(r)} \in \ell^1(\mathbb{N})$,

Now we can see that $M_{a^{(r)}} = D_r M_a D_r$, where D_r is multiplication by the sequence $\{r^j\}_{j \geq 0}$. Since $\|D_r\| \leq 1$, we must have $\|M_{a^{(r)}}\| \leq \|M_a\|$. Since $a^{(r)} \in \ell^1(\mathbb{N})$, we have that the linear functional

$$\mathcal{L}_r(q) := \sum_{n=0}^{\infty} a_n r^n \hat{q}(n)$$

is bounded on $H^1(\mathbb{T})$, and the functionals $\{\mathcal{L}_r\}_{r \in (0,1)}$ converge strongly to \mathcal{L} , and are uniformly bounded. Hence \mathcal{L} is continuous on $H^1(\mathbb{T})$.

Now by the Hahn-Banach theorem, since \mathcal{L} is a linear functional on the subspace $H^1(\mathbb{T})$ which has norm bounded by $\|M_a\|$, it is the restriction of a linear functional on $L^1(\mathbb{T})$, with norm bounded by $\|M_a\|$. Hence $\mathcal{L}(q) = (\psi, q)$ for some $\psi \in L^\infty(\mathbb{T})$ with $\|\psi\|_\infty \leq \|M_a\|$. This proves the result. \square

The key idea of this theorem is that it relates the sequence defining a Hankel operator with a function on \mathbb{T} . We can state this in a slightly more elegant way using the following result of Fefferman [?],

Proposition 3.2. *The space $\text{BMO}(\mathbb{T})$ is defined as the set of measurable functions f on \mathbb{T} (modulo almost-everywhere equivalence) such that*

$$\sup_I \int_I \left| f - \frac{1}{m(I)} \int_I f \, d\mathbf{m} \right| d\mathbf{m} < \infty$$

where I is taken over all arcs in \mathbb{T} .

Then

$$\text{BMO}(\mathbb{T}) = L^\infty(\mathbb{T}) + \mathbf{P}_+ L^\infty(\mathbb{T}) = \{f + \mathbf{P}_+ g : f, g \in L^\infty(\mathbb{T})\}.$$

Using this description of $\text{BMO}(\mathbb{T})$, we can prove the following,

Corollary 3.3. *Let $a = \{a_j\}_{j=0}^\infty$ be a sequence. Then M_a defines a bounded operator on $\ell^2(\mathbb{N})$ if and only if*

$$\varphi := \sum_{n=0}^{\infty} a_n z^n \in \text{BMO}(\mathbb{T}) \cap H^1(\mathbb{T}).$$

Proof. By theorem 3.1, M_a is bounded if and only if $\varphi = \mathbf{P}_+ \psi$ for some $\psi \in L^\infty(\mathbb{T})$. Hence if M_a is bounded, then $\varphi \in \text{BMO}(\mathbb{T}) \cap H^1(\mathbb{T})$.

Conversely, if $\varphi \in \text{BMO}(\mathbb{T}) \cap H^1(\mathbb{T})$, then $\varphi = f + \mathbf{P}_+ g$ for $f, g \in L^\infty$. Thus $\varphi = \mathbf{P}_+ \varphi = \mathbf{P}_+(f + g)$, so M_a is bounded. \square

From now on, we are no longer interested in Hankel matrices M_a defined by an arbitrary sequence a , we are only interested in those matrices M_a such that a arises from the fourier transform of a function.

Definition 3.4. Let $\varphi \in H^1(\mathbb{T})$. Let Γ_φ be the Hankel matrix with (i, j) th entry $\hat{\varphi}(i + j)$.

3.3 Finite Rank Hankel operators

The strongest condition that we can put on a Hankel matrix Γ_φ is that it is a finite rank operator on $\ell^2(\mathbb{N})$. The problem of determining φ such that Γ_φ is finite rank was solved by Kronecker, as follows.

Theorem 3.5. *Let $\varphi \in H^1(\mathbb{T})$, and Γ_φ be the associated Hankel matrix. Then Γ_φ defines a bounded operator on $\ell^2(\mathbb{N})$ if and only if φ is a rational function.*

Proof. Suppose that $\text{rank}(\Gamma_\varphi) = n$. Then the first $n + 1$ columns of the matrix of Γ_φ are linearly dependent. Let B denote the backward shift operator, $B(a_0, a_1, \dots) := (a_1, a_2, \dots)$ and let F be the forward shift operator, $F(a_0, a_1, \dots) := (0, a_0, a_1, \dots)$. Let $a = \hat{\varphi}$.

Hence there exist complex scalars $\{c_0, c_1, \dots, c_n\}$ not all equal to zero such that

$$c_0 a + c_1 B a + \dots + c_n B^n a = 0.$$

Now let $n, k \geq 0$. It is elementary that

$$F^n B^k a = F^{n-k} a - F^{n-k}(a_0, a_1, \dots, a_{k-1}, 0, 0, \dots).$$

So hence we have,

$$\begin{aligned} 0 &= F^n \sum_{k=0}^n c_k B^k a \\ &= \sum_{k=0}^n c_k F^n B^k a \\ &= \sum_{k=0}^n c_k F^{n-k} a - p \end{aligned}$$

where p is a finitely supported sequence.

Let $q = (c_n, c_{n-1}, \dots, c_0, 0, 0, \dots)$. Then we have,

$$0 = q * a - p.$$

Where the $*$ is convolution. Therefore, if we take the inverse fourier transform,

$$\varphi \check{a} = \check{p}.$$

And hence φ is a quotient of two polynomials.

Conversely, suppose that φ is a rational function. Suppose that $\varphi = p/q$, where $p, q \in P(\mathbb{T})$. Let $n = \max\{\deg p, \deg q\}$. If

$$q = \sum_{k=0}^n c_{n-k} z^k$$

then since $\varphi q = p$, we have

$$\sum_{k=0}^n c_k F^{n-k} a = p.$$

Now multiply by B^n ,

$$\begin{aligned} B^n \sum_{k=0}^n c_k F^{n-k} a &= \sum_{k=0}^n c_k B^k a \\ &= 0. \end{aligned}$$

Let $m \leq n$ be the largest number for which $c_m \neq 0$. Then $B^m a$ is a linear combination of the $B^k a$ with $k \leq m-1$,

$$B^m a = \sum_{k=0}^{m-1} d_k B^k a$$

for some coefficients d_k .

We now proceed by induction to show that any row is a linear combination of the first n rows.

Let $k > m$. Then we have,

$$\begin{aligned} B^k a &= B^{k-m} B^m a \\ &= \sum_{j=0}^{m-1} d_j B^{k-m+1} a. \end{aligned}$$

Since $k-m+j < k$, we have that the terms on the right hand side are linear combinations of the first m rows by the inductive hypothesis. Hence $\text{rank}(\Gamma_\varphi) \leq m$. \square

3.4 Compactness of Hankel Operators

If $\varphi \in L^1(\mathbb{T})$, we are interested in conditions on φ such that Γ_φ is compact.

Our first result shows that Hankel matrices are continuous in their symbol.

Proposition 3.6. *Let $\varphi \in L^\infty(\mathbb{T})$, then*

$$\|\Gamma_\varphi\| \leq \|\varphi\|_\infty.$$

and therefore if $\varphi \in C(\mathbb{T})$, then Γ_φ is compact.

Proof. It was shown in the proof of theorem 3.1 that if g, f are sequences of finite support, then

$$|(g, \Gamma_\varphi f)| \leq \|\varphi\|_\infty \|g\|_2 \|f\|_2.$$

Hence $\|\Gamma_\varphi\| \leq \|\varphi\|_\infty$. \square

Proposition 3.7. *The class $\text{VMO}(\mathbb{T})$ is the set of measurable functions $f : \mathbb{T} \rightarrow \mathbb{C}$ such that*

$$\lim_{m(I) \rightarrow 0} \int_I \left| f - \frac{1}{m(I)} \int_I f \, d\mathbf{m} \right| d\mathbf{m} = 0$$

where the limit is over all arcs $I \subseteq \mathbb{T}$.

It is a result of Fefferman [?] that,

$$\text{VMO}(\mathbb{T}) = C(\mathbb{T}) + \mathbf{P}_+ C(\mathbb{T}).$$

Corollary 3.8. *If $\varphi \in C(\mathbb{T})$, then Γ_φ is compact.*

Proof. Since Γ_φ is finite rank for φ a polynomial, and $\|\Gamma_\varphi\| \leq \|\varphi\|_\infty$, the result follows. \square

3.5 Hankel Operators of Trace Class

We recall the definition of the \mathcal{L}^1 norm on $\mathcal{B}(\mathcal{H})$,

Definition 3.9. Let \mathcal{H} be a separable Hilbert space, and let $T \in \mathcal{B}(\mathcal{H})$. For a non-negative integer n , we define the n th singular value,

$$s_n(T) := \inf\{\|T - F\| : F \in \mathcal{B}(\mathcal{H}), \text{rank}(F) \leq n\}.$$

For $p \in (0, \infty)$, we define the \mathcal{L}^p norm of T as

$$\|T\|_p = \left(\sum_{n=0}^{\infty} |s_n(T)|^p \right)^{1/p}$$

with the convention that $\|T\|_p = \infty$ if the sum does not converge. The space \mathcal{L}^p is the set of $T \in \mathcal{B}(\mathcal{H})$ such that $\|T\|_p < \infty$.

In particular we are interested in the case $p = 1$. We are interested in finding conditions on a function φ holomorphic in the unit disc such that Γ_φ is in \mathcal{L}^1 .

Lemma 3.10. *Let \mathcal{H} be a separable Hilbert space. Suppose that $x, y \in \mathcal{H}$, and let T be the rank one operator defined by $T(\zeta) = \langle x, \zeta \rangle y$. Then $\|T\|_1 = \|x\| \|y\|$.*

The answer is provided by the *Besov classes*. Many different definitions of Besov spaces can be found, but the one of most relevance to us is given below.

Definition 3.11. We define a sequence of polynomials $\{W_n\}_{n \in \mathbb{Z}} \subset P(\mathbb{T})$ as follows. First,

$$W_0 = z^{-1} + 1 + z.$$

And now for $n > 0$, we define W_n by asserting that $\widehat{W}(2^n) = 1$, $\widehat{W}(2^{n-1}) = 0$, $\widehat{W}(2^{n+1}) = 0$, and \widehat{W} is a linear increasing function between 2^{n-1} and 2^n , and a linear decreasing

function between 2^n and 2^{n+1} . We assert that $\widehat{W}(n)$ is symmetric in n , and is zero for all values not already defined.

Now, for $p, q > 0$, and $s \geq 0$, we define the *Besov class* $B_{pq}^s(\mathbb{T})$ to be the space of distributions f on \mathbb{T} such that

$$\sum_{n \geq 0} 2^{n sq} \|W_n * f\|_p^q < \infty.$$

We shall denote B_{pp}^s as B_p^s .

In particular, we are going to prove that if $\varphi \in B_1^1(\mathbb{T})$, then $\Gamma_\varphi \in \mathcal{L}^1$. First, we need a lemma.

Lemma 3.12. *Let $f \in P_A(\mathbb{T})$ be an analytic polynomial of degree at most m . Then,*

$$\|\Gamma_f\|_1 \leq (m+1)\|f\|_1.$$

Proof. Let $\zeta \in \mathbb{T}$, now define the following elements of $\ell^2(\mathbb{N})$,

$$x_\zeta(j) = \begin{cases} \zeta^j, & 0 \leq j \leq m, \\ 0, & j > m \end{cases}$$

$$y_\zeta(j) = \begin{cases} f(\zeta)\zeta^{-k}, & 0 \leq k \leq m, \\ 0, & k > m. \end{cases}$$

That is, $x_\zeta = (1, \zeta, \zeta^2, \dots, \zeta^m, 0, 0, \dots)$ and $y_\zeta = f(\zeta)\overline{x_\zeta}$.

Let A_ζ be the rank one operator, $A_\zeta(x) = (x_\zeta, x)y_\zeta$, so that $\|A_\zeta\|_1 = \|x_\zeta\|_2\|y_\zeta\|_2 = (m+1)|f(\zeta)|$

Then we have a componentwise equality of infinite matrices,

$$\Gamma_f = \int_{\mathbb{T}} A_\zeta d\mathbf{m}(\zeta).$$

Hence, $\|\Gamma_f\|_1 \leq (m+1)\|f\|_1$ by the triangle inequality. □

Theorem 3.13. *Let $\varphi \in B_1^1$. Then $\Gamma_\varphi \in \mathcal{L}^1$.*

Proof. We have the following L^∞ -convergent sequence,

$$\varphi = \sum_{n \geq 0} W_n * \varphi.$$

Hence,

$$\Gamma_\varphi = \sum_{n \geq 0} \Gamma_{W_n * \varphi}.$$

So since the degree of W_n is 2^{n+1} , we have

$$\|\Gamma_\varphi\|_1 \leq \sum_{n \geq 0} 2^{n+1} \|W_n * \varphi\|_1.$$

□

This proves the sufficiency of the condition $\varphi \in B_1^1$ so that Γ_φ is trace class. The proof of the necessity of this condition is more difficult,

Theorem 3.14. *Let φ be a function holomorphic in the unit disc. Then if $\Gamma_\varphi \in \mathcal{L}^1$, then $\varphi \in B_1^1$.*

Proof. Define a pair of sequences of polynomials $\{Q_n\}_{n=0}^\infty$ as follows,

$$\widehat{Q}_n(k) = \begin{cases} 0, & k \leq 2^{n-1} \\ 1 - \frac{|k-2^n|}{2^{n-1}}, & 2^{n-1} \leq k \leq 2^n + 2^{n-1}, \\ 0 & k \geq 2^n + 2^{n-1}. \end{cases}$$

and a sequence $\{R_n\}_{n=0}^\infty$,

$$\widehat{R}_n(k) = \begin{cases} 0, & k \leq 2^n \\ 1 - \frac{|k-2^n-2^{n-1}|}{2^{n-1}}, & 2^n \leq k \leq 2^{n+1} \\ 0, & k \geq 2^{n+1}. \end{cases}$$

This is a decomposition of the sequence $\{W_n\}_{n=0}^\infty$, given by $W_n = Q_n + \frac{1}{2}R_n$.

First we prove that

$$\sum_{n \geq 0} 2^{2n+1} \|Q_{2n+1} * \varphi\|_1 < \infty.$$

To this end, we wish to construct an operator B such that

$$\langle \Gamma_\varphi, B \rangle = \sum_{n \geq 0} 2^{2n} \|Q_{2n+1} * \varphi\|_1.$$

Now define the sequence of squares, for $n \geq 1$,

$$S_n = [2^{2n-1}, 2^{2n-1} + 2^{2n} - 1] \times [2^{2n-1} + 1, 2^{2n-1} + 2^{2n}].$$

Note that this sequence is pairwise disjoint.

Let $\{\psi_n\}_{n=0}^\infty$ be a sequence in $L^\infty(\mathbb{T})$, yet to be defined with $\|\psi_n\|_\infty \leq 1$. Now we define the matrix $B = \{B_{j,k}\}_{j,k \geq 0}$ by

$$B_{j,k} = \begin{cases} \widehat{\psi_n}(j+k), (j,k) \in S_n, n \geq 1, \\ 0, (j,k) \notin \bigcup_{n \geq 1} S_n. \end{cases}$$

We wish to prove that B is bounded, and in fact $\|B\| \leq 1$. Let $\{e_n\}_{n \geq 0}$ be the standard basis for $\ell^2(\mathbb{N})$, with $e_n(m) = \delta_{n,m}$. Define the subspaces

$$\begin{aligned} \mathcal{H}_n &= \{e_j : 2^{2n-1} \leq j \leq 2^{2n-1} + 2^{2n} - 1\}, \\ \mathcal{H}'_n &= \{e_j : 2^{2n-1} + 1 \leq j \leq 2^{2n-1} + 2^{2n}\}. \end{aligned}$$

Let P_n and P'_n be the orthogonal projection onto \mathcal{H}_n and \mathcal{H}'_n respectively. So that

$$B = \sum_{n \geq 1} P'_n \Gamma_{\psi_n} P_n,$$

where P_n and P'_n are the orthogonal projections onto \mathcal{H}_n and \mathcal{H}'_n respectively.

Now since the spaces $\{\mathcal{H}_n\}_{n \geq 1}$ are pairwise orthogonal, as are the spaces $\{\mathcal{H}'_n\}_{n \geq 1}$, we have

$$\begin{aligned} \|B\| &\leq \sup_{n \geq 1} \|P'_n \Gamma_{\psi_n} P_n\| \\ &\leq \sup_n \|\Gamma_{\psi_n}\| \\ &\leq \sup_n \|\psi_n\|_\infty \\ &\leq 1. \end{aligned}$$

Now, we compute

$$\begin{aligned} \langle \Gamma_\varphi, B \rangle &= \sum_{n \geq 1} \langle \Gamma_\varphi, P'_n \Gamma_{\psi_n} P_n \rangle \\ &= \sum_{n \geq 1} \sum_{j=2^{2n}}^{2^{2n}+2^{2n+1}} (2^{2n} - |j - 2^{2n+1}| \widehat{\bar{\varphi}}(j) \widehat{\psi_n}(j)) \\ &= \sum_{n \geq 1} 2^{2n} (Q_{2n+1} * \varphi, \psi_n). \end{aligned}$$

Now, using the sharpness of Hölder's inequality, we can choose a sequence $\{\psi_n\}_{n=0}^\infty$ so that $\langle Q_{2n+1} * \varphi, \psi_n \rangle$ is arbitrarily close to $\|Q_{2n+1} * \varphi\|_1$. Hence,

$$\sum_{n \geq 1} 2^{2n+1} \|Q_{2n+1} * \varphi\|_1 = 2 \langle \Gamma_\varphi, B \rangle \leq 2 \|\Gamma_\varphi\|_1.$$

In exactly the same way, we may prove that

$$\sum_{n \geq 1} 2^{2n} \|Q_{2n} * \varphi\|_1 < \infty$$

that

$$\sum_{n \geq 0} 2^{2n+1} \|R_{2n+1} * \varphi\|_1 < \infty$$

and

$$\sum_{n \geq 1} 2^{2n} \|R_{2n} * \varphi\|_1 < \infty$$

and therefore that $\varphi \in B_1^1$. □

Remark 3.15. We thus conclude that

$$\frac{1}{6} \sum_{n \geq 1} 2^n \|W_n * \varphi\|_1 \leq \|\Gamma_\varphi\|_1 \leq 2 \sum_{n \geq 0} 2^n \|W_n * \varphi\|_1.$$

3.6 Interpolation

Lemma 3.16. *Let K be the real interpolation functor, described in [?]. Then*

$$K(\mathcal{L}^1, \mathcal{K})_{\theta, q} = \mathcal{L}_{p, q}.$$

for $p = (1 - \theta)^{-1}$.

Corollary 3.17. *Hence, we have that $\Gamma_\varphi \in \mathcal{L}_{p, q}$ if and only if $\varphi \in (B_1^1, \text{VMO})_{\theta, q}$, where $p = (1 - \theta)^{-1}$.*

Chapter 4

Abstract Differential Algebra and Spectral Triples

4.1 Introduction

For several decades now, mathematicians have been attempting to find analogues of theorems of differential and algebraic geometry in noncommutative algebra. The biggest obstacle to learning this topic is that many of the definitions were arrived at through many years of hard work, and may seem unmotivated at first. The purpose of these notes is to explain the algebra of Connes Differentials, which were invented by Alain Connes as a noncommutative generalisation of the exterior algebra bundle on a manifold.

4.2 Classical Differential Algebra

Let M be an n dimensional manifold. The cotangent bundle $\Omega^1(M)$ is a rank n vector bundle on M . We build higher bundles by wedge products,

$$\Omega^p(M) := \bigwedge_p \Omega^1(M).$$

and define $\Omega^0(M) := C^\infty(M)$. See that $\dim_{\mathbb{R}}(\Omega^p(M)) = \binom{n}{p}$.

The exterior algebra bundle is the direct sum of all the $\Omega^p(M)$,

$$\Omega(M) := \bigoplus_{p=0}^{\infty} \Omega^p(M).$$

$\Omega(M)$ is a *graded algebra*.

In general, if A is an algebra over a ring R , we say that A is \mathbb{N} -graded if there exists a decomposition into submodules $A^{(p)}$,

$$A = \bigoplus_{p=0}^{\infty} A^{(p)}$$

such that $A^{(n)}A^{(m)} \subseteq A^{(n+m)}$.

In the case $A = \Omega(M)$, we have $R = \mathbb{R}$, and $A^{(p)} = \Omega^p(M)$.

The exterior derivative, $d : \Omega(M) \rightarrow \Omega(M)$ acts on the grading by,

$$d : \Omega^p(M) \rightarrow \Omega^{p+1}(M).$$

and $d^2 = 0$. Hence we have a sequence,

$$0 \rightarrow \Omega^0(M) \xrightarrow{d} \Omega^1(M) \xrightarrow{d} \cdots \xrightarrow{d} \Omega^n(M) \xrightarrow{d} 0.$$

Denote d_p as the restriction of d to $\Omega^p(M)$. Then we have the *de Rham cohomology* spaces,

$$H_{dR}^p(M) = \frac{\ker(d_p)}{\operatorname{im}(d_{p-1})}$$

This is a sequence of real vector spaces, and their dimensions are topological invariants of M .

The maps d_p satisfy a graded version of Leibniz's rule, for $a \in \Omega^n(M)$ and $b \in \Omega^m(M)$, we have:

$$d_{n+m}(ab) = d_n(a)b + (-1)^n a d_m(b)$$

4.3 Abstract Differential Algebra

4.3.1 Graded Differential Algebras

We now take the ideas of the previous section and move them to a more abstract setting. Let R be a commutative ring, and let A be an \mathbb{N} -graded algebra over R , with decomposition

$$A = \bigoplus_{p=0}^{\infty} A^{(p)}$$

There is also an R -linear map $d : A \rightarrow A$ such that,

$$d : A^{(p)} \rightarrow A^{(p+1)}.$$

and $d^2 = 0$. If we denote the restriction of d to $A^{(p)}$ as d_p , we require that the maps d_p satisfy a graded Leibniz rule, for $a \in A^{(n)}$ and $b \in A^{(m)}$,

$$d_{n+m}(ab) = d_n(a)b + (-1)^n ad_m(b).$$

A pair (A, d) satisfying these conditions is called a *differential graded algebra*.

Thus we have a sequence,

$$0 \rightarrow A^{(0)} \xrightarrow{d} A^{(1)} \xrightarrow{d} \dots \xrightarrow{d} A^{(n)} \xrightarrow{d} \dots$$

The quotient R -modules,

$$H_{dR}^p(M) := \frac{\ker(d_p)}{\operatorname{im}(d_{p-1})}.$$

are the de Rham cohomology modules for the graded differential algebra (A, d) .

4.3.2 Kähler Differentials

Given an R -algebra A , we would like to be able to build an algebra of differential forms over A , in a manner analogous to how $\Omega^1(M)$ is constructed from $C^\infty(M)$. It turns out that there is a good way of doing this, called the algebra of *Kähler differentials*. This is simplest in the commutative case, which we briefly outline here.

Let R be a commutative ring, and let A be a unital commutative R -algebra. The module $\Omega_{\text{com}}^1(A)$ of Kähler differentials is defined as

$$\Omega_{\text{com}}^1(A) := \frac{A \otimes_R A}{\langle c \otimes (ab) - (ca) \otimes b - (bc) \otimes a \rangle}$$

The idea here is that $\Omega_{\text{com}}^1(A)$ is the left A -module spanned by all symbols of the form adb , where $d(ab) = adb + bda$. We think of $a \otimes b$ as adb .

More precisely, we let $d : A \rightarrow \Omega_{\text{com}}^1(A)$ be given by

$$da := 1_A \otimes a$$

Where 1_A is the unit in A .

The utility of $\Omega_{\text{com}}^1(A)$ is that it allows us to study all derivations on A .

In full abstraction, a derivation on A is a map $\theta : A \rightarrow M$, where M is some left A -module, such that θ satisfies the leibniz rule,

$$\theta(ab) = a\theta(b) + b\theta(a).$$

We see that d is a derivation on A to the A -module $\Omega_{\text{com}}^1(A)$. It is in fact universal with this property,

Theorem 4.1. *Let A be a unital commutative R -algebra, and let $\theta : A \rightarrow M$ be a derivation to some left A -module M . There exists a unique R -linear map $\Omega(\theta)$ such that the following diagram commutes,*

$$\begin{array}{ccc} A & \xrightarrow{d} & \Omega_{\text{com}}^1(A) \\ & \searrow \theta & \downarrow \Omega(\theta) \\ & & M \end{array}$$

In other words, there is an isomorphism of R -modules,

$$D(A, M) \cong \text{Hom}_R(\Omega_{\text{com}}^1(A), M).$$

Where $D(A, M)$ is the set of derivations from A to M . Note that this universal property defines $\Omega_{\text{com}}^1(A)$ up to unique isomorphism.

Proof. Basically, $\Omega(\theta)$ maps adb to $a\theta(b)$. Checking the universal property is routine. \square

We would now like to create a similar algebra of differentials for a non-commutative associative algebra A over R . In the noncommutative case, we must restrict attention to derivations that take values in A -bimodules, rather than left A modules.

Definition 4.2. Let A be an associative unital algebra over a commutative ring R . Let $m : A \otimes A \rightarrow A$ be the multiplication map. We define

$$\Omega^1(A) = \ker(m)$$

This is an A bimodule.

This the motivation behind this definition is not at all clear. However, this does agree with the commutative case and this provides the appropriate definition for noncommutative Kähler differentials. To see this, we define the map $d : A \rightarrow \Omega^1(A)$, by

$$d(a) = 1_A \otimes a - a \otimes 1_A.$$

We see that d is a derivation. In fact, $\Omega^1(A)$ should be thought of as the space of all linear combinations of terms of the form $ad(b)$.

$\Omega^1(A)$ satisfies the same universal property as Ω_{com}^1 . Namely, if M is an A -bimodule, and $\theta : A \rightarrow M$ is a derivation, then there exists a unique R -linear map $\Omega(\theta)$ such that the following diagram commutes,

$$\begin{array}{ccc} A & \xrightarrow{d} & \Omega^1(A) \\ & \searrow \theta & \downarrow \Omega(\theta) \\ & & M \end{array}$$

4.3.3 Universal Differential Algebra

Given an associative unital algebra R over a commutative ring R , we define

$$\Omega^p(A) := \bigotimes_{A,p} \Omega^1(A).$$

And the algebra,

$$\Omega A = \bigoplus_p \Omega^p(A).$$

We extend the function $d : A \rightarrow \Omega^1(A)$ to ΩA by

$$d(ada_1da_2 \cdots da_n) = dada_1da_2 \cdots da_n.$$

Theorem 4.3. ΩA is the “largest” graded differential algebra generated by A .

If (Γ, Δ) is a graded differential algebra, with grading $\Gamma = \bigoplus_n \Gamma^{(n)}$, and $\rho : A \rightarrow \Gamma^{(0)}$ is an algebra homomorphism, then ρ extends uniquely to a morphism $\Omega A \rightarrow \Gamma$ such that the following diagram commutes,

$$\begin{array}{ccc} \Omega^p(A) & \xrightarrow{\rho} & \Gamma^{(p)} \\ \downarrow d & & \downarrow \Delta \\ \Omega^{p+1}(A) & \xrightarrow{\rho} & \Gamma^{(p+1)} \end{array}$$

The Insufficiency of Kähler Differentials

We now have a good definition of $\Omega(A)$ for any potentially non-commutative algebra A , and it would seem that this is the final word on generalising the algebra of differential forms to the noncommutative setting.

However, this is incorrect, as the following proposition shows:

Proposition 4.4. *Let $A = C^\infty(\mathbb{R})$. Then if f and g do not satisfy some polynomial relation, then in the algebra of Kähler differentials df and dg are algebraically independent over A . In particular, $d \exp(x) \neq \exp(x)dx$.*

Hence, the usual algebra of differential forms is not isomorphic to the algebra of Kähler differentials, but is instead a quotient of it.

So in order to produce a theory of differential algebra in the noncommutative setting, we need to introduce more structure on the algebra A . Thus we are led to the concept of a spectral triple.

Non-commutative Geometry

4.3.4 Spectral Triples

Non-commutative geometry is an attempt to generalise theorems of differential geometry to non-commutative algebras. The basic object of study in non-commutative differential geometry is the *spectral triple*. A spectral triple plays the role of a “manifold”. The definition is as follows.

Definition 4.5. A spectral triple is a triple, $(\mathcal{A}, \mathcal{H}, \mathcal{D})$. Where, \mathcal{A} is a $*$ -algebra of bounded operators on a Hilbert space \mathcal{H} , and \mathcal{D} is a densely defined unbounded self adjoint operator on \mathcal{H} , such that $[\mathcal{D}, a] \in \mathcal{B}(\mathcal{H})$ for all $a \in \mathcal{A}$ and $(\mathcal{D} - \lambda)^{-1}$ is compact for all $\lambda \in \mathbb{C} \setminus \mathbb{R}$.

This is supposed to be an abstract version of a spin manifold. The commutative case is when M is a Riemannian manifold with a spin bundle S , then we have $\mathcal{A} = C^\infty(M)$, acting by pointwise multiplication on the bundle $\mathcal{H} = L^2(S)$, and \mathcal{D} is the dirac operator.

It can be shown, under fairly general conditions, that when \mathcal{A} is commutative, then such a spin manifold can be constructed. Hence a spectral triple deserves the name of “non-commutative manifold”.

4.3.5 Connes Differentials

Given a spectral triple, $(\mathcal{A}, \mathcal{H}, \mathcal{D})$, we would like to construct an “exterior algebra” on \mathcal{A} . Connes does this by identifying the 1-form da with $[\mathcal{D}, a]$.

Since $[D, a]$ is a derivation on \mathcal{A} , by the universal property we have a map, $\pi : \Omega\mathcal{A} \rightarrow \mathcal{B}(\mathcal{H})$ given by $\pi(ada_1da_2 \cdots da_n) = a[D, a_1][D, a_2] \cdots [D, a_n]$.

One may then naïvely define the algebra of differential forms as $\pi(\Omega\mathcal{A})$, but this does not work since there exists $a \in \Omega\mathcal{A}$ such that $\pi(a) = 0$ but $\pi(da) \neq 0$. These are called “junk forms” and we must factor them out to get a good differential algebra. Hence, define

Theorem 4.6. *Let J_0 be the graded ideal of $\Omega\mathcal{A}$ defined by*

$$J_0^{(p)} = \{a \in \Omega^p(\mathcal{A}) : \pi(a) = 0\}$$

And define $J^{(p)} = J_0^{(p)} + dJ_0^{(p)}$. Then $J = \bigoplus_p J^{(p)}$.

Now we can define the algebra of Connes’ forms,

$$\Omega_{\mathcal{D}}\mathcal{A} = \frac{\Omega\mathcal{A}}{J} \cong \frac{\pi(\Omega\mathcal{A})}{\pi(dJ_0)}$$

$\Omega_{\mathcal{D}}\mathcal{A}$ is naturally graded by the gradings on $\Omega\mathcal{A}$ and J , with the space of p -forms being $\Omega_{\mathcal{D}}^p\mathcal{A} = \Omega^p(\mathcal{A})/J^{(p)}$.

Since J is a differential ideal, the operator d on $\Omega\mathcal{A}$ extends to $\Omega_{\mathcal{D}}\mathcal{A}$.

Chapter 5

Weak \mathcal{L}^p membership of quantised differentials

5.1 Introduction

We have already given a thorough description of the quantised differentials $\bar{d}f$ when f is a function on the circle. It so happens that we can find analogous, although weaker, results in far greater generality.

5.2 The setting

We shall consider spectral triples $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ such that \mathcal{A} is contained in a semi-finite Von Neumann algebra \mathcal{N} which is contained in $\mathcal{B}(\mathcal{H})$.

From now on, fix a semifinite Von Neumann algebra \mathcal{N} with a faithful normal trace τ .

Formally,

Definition 5.1 (Spectral Triple). A (semifinite) spectral triple is a triple $(\mathcal{A}, \mathcal{H}, \mathcal{D})$, where \mathcal{H} is a Hilbert space with $\mathcal{N} \subseteq \mathcal{B}(\mathcal{H})$, and $\mathcal{A} \subseteq \mathcal{N}$ is a $*$ -algebra.

\mathcal{D} is a densely defined unbounded operator on \mathcal{H} satisfying the following two properties:

1. $[\mathcal{D}, a]$ is densely defined and extends to a bounded operator in \mathcal{N} for all $a \in \mathcal{A}$.
2. $(\lambda - \mathcal{D})^{-1}$ is τ -compact for all $\lambda \in \mathbb{C} \setminus \mathbb{R}$.

A spectral triple $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ can be either *even* or *odd*:

- We say that $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ is even if there exists a $\mathbb{Z}/2\mathbb{Z}$ grading on the linear operators on \mathcal{H} such that \mathcal{A} is even and \mathcal{D} is odd. Equivalently, there is an operator Γ on \mathcal{H} with $\Gamma^2 = 1$ and $\Gamma^* = \Gamma$ such that $a\Gamma = \Gamma a$ for all $a \in \mathcal{A}$ and $\mathcal{D}\Gamma = -\Gamma\mathcal{D}$.
- If no such operator Γ exists, then we say that $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ is odd.

To define the ideal $\mathcal{L}^{p,\infty}$ we need to first define the generalised singular values,

Definition 5.2 (Generalised Singular Values). Let $T \in \mathcal{N}$, and $t \in [0, \infty)$. We define

$$\mu(t, T) := \inf\{\|TE\| : E \text{ is a projection, with } \tau(1 - E) \leq t\}. \quad (5.1)$$

Definition 5.3 ($\mathcal{L}^{p,\infty}$ ideal). Let $\Psi_p : [0, \infty) \rightarrow [0, \infty)$ for $p > 0$ be defined as

$$\Psi(t) := \begin{cases} t, & \text{for } t \in [0, 1] \\ t^{1-\frac{1}{p}} & \text{for } t \geq 1. \end{cases} \quad (5.2)$$

Now define the subset $\mathcal{L}^{p,\infty} \subseteq \mathcal{N}$ as

$$\mathcal{L}^{p,\infty} = \left\{ T \in \mathcal{N} : \|T\|_{p,\infty} := \sup_{t>0} \frac{1}{\Psi_p(t)} \int_0^t \mu(t, T) dt < \infty \right\}. \quad (5.3)$$

Definition 5.4 (Quantum Differentiability). A spectral triple $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ is called QC^k for $k \geq 0$ if \mathcal{A} is contained in the domain of the operator δ^k , where $\delta(a) = [|\mathcal{D}|, a]$.

Definition 5.5 (Summability). A spectral triple $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ is called (p, ∞) summable for $p > 0$ if $(1 + \mathcal{D}^2)^{-1/2} \in \mathcal{L}^{p,\infty}$.

5.3 The Theorem

Theorem 5.6. Let $k \geq 0$ and $p \geq 1$, and let $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ be a (p, ∞) summable QC^k spectral triple such that \mathcal{D} is invertible. Let $F = \text{sgn}(\mathcal{D})$ such that $\mathcal{D} = F|\mathcal{D}|$. Then for all $a \in \mathcal{A}$,

$$[F, a] \in \mathcal{L}^{p,\infty}. \quad (5.4)$$

Proof. Since $\mathcal{D} = |\mathcal{D}|F$, we can compute

$$[\mathcal{D}, a] = |\mathcal{D}|[F, a] + [|\mathcal{D}|, a]F. \quad (5.5)$$

Or in other words,

$$da = |\mathcal{D}|da + \delta(a)F. \quad (5.6)$$

Since \mathcal{D} is invertible, $|\mathcal{D}|$ is invertible. Thus,

$$\bar{d}a = |\mathcal{D}|^{-1}da + |\mathcal{D}|^{-1}\delta(a)F.$$

By assumption, $da, \delta(a) \in \mathcal{B}(\mathcal{H})$. Thus, since $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ is (p, ∞) -summable, $|\mathcal{D}|^{-1} \in \mathcal{L}^{p, \infty}$. Hence, $\bar{d}a \in \mathcal{L}^{p, \infty}$.

□

Chapter 6

The Chain Rule in Quantised Calculus

6.1 Introduction

In classical analysis, the chain rule states that if M, N and P are manifolds, and the maps $f : M \rightarrow N$ and $g : N \rightarrow P$ are smooth, then

$$d(f \circ g) = df \circ dg.$$

Where both sides of the equation are maps $TM \rightarrow TP$.

Remark 6.1. The chain rule is simply a consequence of the functoriality of the tangent bundle construction, $M \mapsto TM$.

It is desirable to find a noncommutative generalisation of this identity.

6.2 The setting

We let $\mathcal{L}_0^{p,\infty}$ be the closure of the finite rank operators in the $\mathcal{L}^{p,\infty}$ -metric topology.

Let $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ be a spectral triple, satisfying the following properties:

1. For all $a \in \mathcal{A}$, $da \in \mathcal{L}^{p,\infty}$.
2. \mathcal{A} is closed under the holomorphic functional calculus.
3. Let $\mathcal{A}_0 \subseteq \mathcal{B}(\mathcal{H})$ be the collection of all T such that $[\text{sgn}(\mathcal{D}), T]$ is finite rank. Then \mathcal{A} is contained within the norm-closure of \mathcal{A}_0 .

6.3 The Commutator Lemma

Lemma 6.2. *Let $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ be as above, and $a \in \mathcal{A}$. Then*

$$[\bar{d}a, a] \in \mathcal{L}_0^{p, \infty}.$$

Proof. By our assumptions on $(\mathcal{A}, \mathcal{H}, \mathcal{D})$ there exists a sequence $\{a_n\}_{n=0}^\infty$ with $\|a_n - a\| \rightarrow 0$, and $\bar{d}a_n$ is finite rank. Hence $[\bar{d}a_n, a]$ is finite rank, and

$$[\bar{d}a, a_n] = \bar{d}([a_n, a]) + [\bar{d}a, a_n].$$

□

Chapter 7

Higher Dimensions

7.1 Introduction

Let $d > 0$ be a positive integer. Let \mathcal{A}_λ be the d -dimensional non-commutative torus, that is the universal C^* -algebra generated by unitaries $\{U_j\}_{j=1}^d$ subject to the commutation relation $U_j U_k = \lambda_{jk} U_k U_j$ for a unimodular scalar constant λ_{jk} , where we assume that $\lambda_{jj} = 1$ for all j , and $\lambda_{jk} = \lambda_{kj}^{-1}$.

\mathcal{A}_λ has a unique normalised trace τ . The Hilbert space $L^2(\mathcal{A}_\lambda, \tau)$ has orthonormal basis $\{u(n)\}_{n \in \mathbb{Z}^d}$ where $u(n) = U_1^{n_1} U_2^{n_2} \cdots U_d^{n_d}$.

Define $\omega_{j,k}$ for $j, k \in \mathbb{Z}^d$ to be such that $u(j)u(k) = \omega_{j,k}u(k+j)$.

Given $f \in L^2(\mathcal{A}_\lambda, \tau)$, we define

$$\hat{f}(n) := \tau(fu(n)^*). \quad (7.1)$$

Then we have

$$f = \sum_{k \in \mathbb{Z}^d} \hat{f}(k)u(k), \quad (7.2)$$

which converges in the L^2 -sense. Note that $\tau(f) = \hat{f}(0)$.

For a function $F : \mathbb{Z}^d \rightarrow \mathbb{R}$, define $F(D)u(k) = F(k)u(k)$ as a densely defined linear operator on $L^2(\mathcal{A}_\lambda, \tau)$.

It is easy to see that $F(D)$ is bounded on $L^2(\mathcal{A}_\lambda, \tau)$ if and only if $F \in \ell^\infty(\mathbb{Z}^d)$, with operator norm $\|F\|_\infty$.

We shall identify $u(n) \in \mathcal{A}_\lambda$ with the multiplication operator $M_{u(n)}f = u(n)f$ for $f \in L^2(\mathcal{A}_\lambda, \tau)$

7.2 Commutators

Let $n \in \mathbb{Z}^d$. We are interested in $[F(D), u(n)]$.

Let $f \in L^2(\mathcal{A}_\lambda, \tau)$.

Then,

$$u(n)f = \sum_{k \in \mathbb{Z}^d} \hat{f}(k)u(n)u(k) \quad (7.3)$$

$$= \sum_{k \in \mathbb{Z}^d} \hat{f}(k)\omega_{n,k}u(n+k). \quad (7.4)$$

Hence,

$$F(D)u(n)f = \sum_{k \in \mathbb{Z}^d} \hat{f}(k)\omega_{n,k}F(n+k)u(n+k). \quad (7.5)$$

Now we compute,

$$u(n)F(D)f = \sum_{k \in \mathbb{Z}^d} \hat{f}(k)F(k)\omega_{n,k}u(n+k). \quad (7.6)$$

Thus,

$$[F(D), u(n)]f = \sum_{k \in \mathbb{Z}^d} \hat{f}(k)(F(n+k) - F(k))\omega_{n,k}u(n+k). \quad (7.7)$$

We can simplify this further,

$$[F(D), u(n)]f = \sum_{k \in \mathbb{Z}^d} \hat{f}(k)(F(n+k) - F(k))u(n)u(k) \quad (7.8)$$

$$= u(n) \sum_{k \in \mathbb{Z}^d} \hat{f}(k)F(n+k)u(k) - u(n) \sum_{k \in \mathbb{Z}^d} \hat{f}(k)F(k)u(k) \quad (7.9)$$

$$= u(n)F(n+D)f - u(n)F(D)f \quad (7.10)$$

$$= u(n)F(n+D)f - u(n)F(D)f \quad (7.11)$$

So $[F(D), u(n)] = u(n)(F(n+D) - F(D))$.

7.3 Hilbert-Schmidt Commutators

It is of interest to ask when $[F(D), u(n)] \in \mathcal{L}^2$. This is equivalent to,

$$\|[F(D), u(n)]\|_2^2 := \sum_{j,k \in \mathbb{Z}^d} |\tau(u(j)^*[F(D), u(n)]u(k))|^2 < \infty. \quad (7.12)$$

So we compute,

$$[F(D), u(n)]u(k) = u(n)F(n+D)u(k) - u(n)F(D)u(k) \quad (7.13)$$

$$= u(n)F(n+k)u(k) - u(n)F(k)u(k) \quad (7.14)$$

$$= (F(n+k) - F(k))u(n)u(k). \quad (7.15)$$

Hence,

$$|\tau(u(j)^*[F(D), u(n)]u(k))| = |F(n+k) - F(k)| |\tau(u(j)^*u(n)u(k))| \quad (7.16)$$

$$= |F(n+k) - F(k)| \delta_{j,n+k} \quad (7.17)$$

$$(7.18)$$

So therefore

$$\|[F(D), u(n)]\|_2^2 = \sum_{j,k \in \mathbb{Z}^d} |F(n+k) - F(k)|^2 \delta_{j,n+k} \quad (7.19)$$

$$= \sum_{k \in \mathbb{Z}^d} |F(n+k) - F(k)|^2. \quad (7.20)$$

So we have necessary and sufficient conditions for $[F(D), u(n)] \in \mathcal{L}^2$.

7.4 Membership of \mathcal{L}^p

We can see that since $F(D)$ and $F(n+D)$ are self adjoint,

$$[F(D), u(n)]^* = (F(n+D) - F(D))u(n)^* \quad (7.21)$$

Hence,

$$[F(D), u(n)]^*[F(D), u(n)] = (F(n+D) - F(D))^2. \quad (7.22)$$

Hence,

$$|[F(D), u(n)]| = |F(n+D) - F(D)|. \quad (7.23)$$

Now,

$$\mathrm{Tr}(|[F(D), u(n)]|^p) = \sum_{k \in \mathbb{Z}^d} \tau(u(k)^* |F(D+n) - F(D)|^p u(k)) \quad (7.24)$$

$$= \sum_{k \in \mathbb{Z}^d} \tau(u(k)^* |F(k+n) - F(k)|^p u(k)) \quad (7.25)$$

$$= \sum_{k \in \mathbb{Z}^d} |F(k+n) - F(k)|^p. \quad (7.26)$$

Hence, for any $p > 0$, we have $[F(D), u(n)] \in \mathcal{L}^p$ if and only if $\{F(k+n) - F(k)\} \in \ell^p(\mathbb{Z}^d)$ with an equality of (quasi-)norms.

7.5 Double commutators

We are also interested in the properties of the double commutators,.

$$[[F(D), u(n)], u(n)]$$

for $n \in \mathbb{Z}^d$. By equation 7.11, this is

$$[[F(D), u(n)], u(n)] = [F(D), u(n)]u(n) - u(n)[F(D), u(n)] \quad (7.27)$$

$$= u(n)(F(n+D) - F(D))u(n) - u(n)u(n)(F(D+n) - F(D)) \quad (7.28)$$

$$= (F(D) - F(D-n))u(n)^2 - (F(D-n) - F(D-2n))u(n)^2 \quad (7.29)$$

$$= (F(D-2n) - 2F(D-n) + F(D))u(n)^2 \quad (7.30)$$

$$= u(n)(F(D+n) - 2F(D) + F(D-n))u(n). \quad (7.31)$$

Hence,

$$|[[F(D), u(n)], u(n)]| = |F(D+2n) - 2F(D+n) + F(D)|.$$

So we have $[[F(D), u(n)], u(n)] \in \mathcal{L}^p$ if and only if $\{F(k+n) - 2F(k) + F(k-n)\}_{k \in \mathbb{Z}^d} \in \ell^p(\mathbb{Z}^d)$.

Appendix A

Classical Harmonic Analysis

A.1 Introduction

Given a function $f : \mathbb{T} \rightarrow \mathbb{C}$, we have an associated fourier series,

$$f \sim \sum_{n \in \mathbb{Z}} \hat{f}(n) z^n \tag{A.1}$$

where $z : \mathbb{T} \rightarrow \mathbb{T}$ is the identity function, and

$$\hat{f}(n) = \int_{\mathbb{T}} z^{-n} f \, d\mathbf{m} \tag{A.2}$$

where \mathbf{m} is the normalised Haar (or arc length) measure on \mathbb{T} . Implicitly f is sufficiently regular so that these fourier coefficients exist.

I have used the symbol “ \sim ” rather than an equals sign, since in general we do not have equality. In general, for $f \in L^1(\mathbb{T}, \mathbf{m})$, the fourier series might diverge almost everywhere if it is interpreted as a sum.

However, if one turns to alternative methods of summation, it is possible to interpret \sim as an equality.

We have already explored in a previous set of notes the concept of Cesàro summation. Here we consider Abel summation.

A.2 Abel summation

Abel summation is inspired by Abel’s theorem, which we prove now.

Proposition A.1. *Suppose that*

$$f(z) = \sum_{n=0}^{\infty} a_n z^n$$

is a power series converging in $\{z \in \mathbb{C} : |z| < 1\}$ such that the coefficients come from a Banach space X . Suppose further that the sum

$$\sum_{n=0}^{\infty} a_n$$

converges. Then,

$$\lim_{z \rightarrow 1^-} f(z) = \sum_{n=0}^{\infty} a_n.$$

where by $z \rightarrow 1^-$, we mean that z is restricted to the subset of the unit disc where $|1 - z| \leq M(1 - |z|)$ for some constant M .

Proof. Assume without loss of generality that

$$\sum_{n=0}^{\infty} a_n = 0.$$

Now define

$$s_k = \sum_{n=0}^k a_n$$

and $s_{-1} = 0$. Then we have

$$f(z) = \sum_{n=0}^{\infty} (s_n - s_{n-1}) z^n.$$

so

$$f(z) = (1 - z) \sum_{k=0}^{\infty} s_k z^k.$$

Let $\varepsilon > 0$, and choose n large enough such that $\|s_k\| < \varepsilon$ for $k > n$. Then we have

$$\left\| (1 - z) \sum_{k=n}^{\infty} s_k z^k \right\| \leq \varepsilon |1 - z| \sum_{k=n}^{\infty} |z|^k = \varepsilon |1 - z| \frac{|z|^n}{1 - |z|} \leq M\varepsilon.$$

When z is sufficiently close to 1, we have

$$\left\| (1 - z) \sum_{k=0}^{n-1} s_k z^k \right\| < \varepsilon.$$

Hence, for z sufficiently close to 1, we have

$$\|f(z)\| < (M + 1)\varepsilon.$$

□

With this in mind, we define the Abel summation method.

Definition A.2. Let $\{a_k\}_{k=0}^{\infty} \subset X$ be a sequence in a Banach space X . Suppose that for all $r \in (1 - \varepsilon, 1)$, for some $\varepsilon > 0$ we have

$$\sum_{k=0}^{\infty} a_k r^k$$

exists. Then we define the Abel sum, denoted by,

$$A - \sum_{k=0}^{\infty} a_k$$

as

$$A - \sum_{k=0}^{\infty} a_k := \lim_{r \rightarrow 1^-} \sum_{k=0}^{\infty} a_k r^k.$$

if this limit exists.

Abel's theorem automatically implies that the Abel sum of a series agrees with the usual sum, however there are series which are summable in the Abel sense but not the classical sense. It is easy to see that,

$$A - \sum_{n=0}^{\infty} (-1)^n = \lim_{r \rightarrow 1^-} \sum_{n=0}^{\infty} (-r)^n = \lim_{r \rightarrow 1^-} \frac{1}{1+r} = \frac{1}{2}.$$

A.3 The Poisson Kernel

To sum a Fourier series in the classical sense, we use the Dirichlet kernel, and we use the Fejér kernel for Cesàro summation. For Abel summation, we use the Poisson kernel.

Given $f \in L^1(\mathbb{T}, \mathbf{m})$, define

$$A_r f := \sum_{n \in \mathbb{Z}} r^{|n|} \hat{f}(n) z^n.$$

with $r \in (0, 1)$. $A_r f$ exists since the fourier coefficients of f are bounded. By definition,

$$A - \sum_{n \in \mathbb{Z}} \hat{f}(n) z^n = \lim_{r \rightarrow 1^-} A_r f.$$

where the limit is taken in an appropriate Banach space.

Like with classical and Cesàro sums, Abel sums can be constructed with a convolution.

Proposition A.3. *We can write,*

$$A_r f = P_r * f.$$

where

$$P_r = 1 + \frac{rz}{1-rz} + \frac{r}{z-r}$$

Proof. The Fourier coefficients of $A_r f$ are the Fourier coefficients of f multiplied by the coefficients of

$$\sum_{n \in \mathbb{Z}} r^{|n|} z^n.$$

Hence define

$$P_r = \sum_{n \in \mathbb{Z}} r^{|n|} z^n.$$

The result follows from summing this geometric series. \square

The reason for the superiority of Abel summation over classical summation is that the Poisson kernels form an approximate identity.

Proposition A.4. *The Poisson kernels $\{P_r\}_{r \in (0,1)}$ form an approximate identity.*

Proof. By the formula,

$$P_r = \sum_{n \in \mathbb{Z}} r^{|n|} z^n,$$

we have

$$\int_{\mathbb{T}} P_r d\mathbf{m} = 1.$$

To complete the proof, we convert to coordinates. Let $z = \exp(2\pi i\theta)$, for $\theta \in (-1/2, 1/2)$, and we can regard P_r as a function of θ .

Then we have

$$\begin{aligned} P_r(\theta) &= 1 + \frac{re^{2\pi i\theta}}{1-re^{2\pi i\theta}} + \frac{re^{-2\pi i\theta}}{1-re^{-2\pi i\theta}} \\ &= 1 + \frac{re^{2\pi i\theta}(1-re^{-2\pi i\theta}) + re^{-2\pi i\theta}(1-re^{2\pi i\theta})}{(1-re^{2\pi i\theta})(1-re^{-2\pi i\theta})} \\ &= 1 + \frac{2r \cos(2\pi\theta) - 2r^2}{1+r^2-2r \cos(2\pi\theta)} \\ &= \frac{1-r^2}{1-2r \cos(2\pi\theta) + r^2} \end{aligned}$$

Let $\delta > 0$. Now we estimate,

$$\int_{\delta}^{1/2} P_r(\theta) d\theta \leq \int_{\delta}^{1/2} \frac{1 - r^2}{1 - 2r \cos(2\pi\delta) + r^2} d\theta \leq \frac{1 - r^2}{1 - 2r \cos(2\pi\delta) + r^2}$$

Hence this integral vanishes as $r \rightarrow \infty$.

Moreover, we have

$$1 - 2r \cos(2\pi\theta) + r^2 \geq (1 - r)^2.$$

Hence $P_r \geq 0$. Therefore, $\|P_r\|_1 = 1$.

Thus the Poisson kernels form an approximate identity. □

As a consequence of this, we have

1. If $f \in C(\mathbb{T})$, then $A_r f \rightarrow f$ uniformly.
2. If $f \in L^p(\mathbb{T}, \mathbf{m})$, for $1 \leq p < \infty$, we have $A_r f \rightarrow f$ in the L^p sense.

Hence, if $f \in L^1(\mathbb{T}, \mathbf{m})$, we have

$$f(\zeta) = A - \sum_{n \in \mathbb{Z}} \hat{f}(n) \zeta^n$$

for almost all $\zeta \in \mathbb{T}$.

A.4 Fourier Series of Measures

Suppose μ is a complex Borel regular measure on \mathbb{T} . Associated to μ is a fourier series,

$$\mu \sim \sum_{n \in \mathbb{Z}} \hat{\mu}(n) z^n$$

where

$$\hat{\mu}(n) = \int_{\mathbb{T}} z^{-n} d\mu.$$

It was proved in “Cesàro convergence of Fourier series” that this series can be interpreted as a sequence of measures converging in the Cesàro sense and the weak* topology to μ . Here we wish to analyse the Abel sums of the Fourier series of μ . Define for $r \in (0, 1)$,

$$A_r \mu = \sum_{n \in \mathbb{Z}} r^{|n|} \hat{\mu}(n) z^n.$$

This series converges because μ is bounded, so the Fourier coefficients are bounded.

Recall that if $f \in C(\mathbb{T})$, we define $f * \mu$ as the function

$$(f * \mu)(\zeta) := \int_{\mathbb{T}} f(\zeta \tau^{-1}) d\mu(\tau).$$

This allows us to compute $A_r \mu$ for a measure μ :

Proposition A.5. *If μ is a complex Borel regular measure on \mathbb{T} , then*

$$A_r \mu = P_r * \mu.$$

Proof. We compute,

$$\begin{aligned} (P_r * \mu)(\zeta) &= \int_{\mathbb{T}} \sum_{n \in \mathbb{Z}} r^{|n|} \zeta^n \tau^{-n} d\mu(\tau) \\ &= \sum_{n \in \mathbb{Z}} \zeta^n r^{|n|} \int_{\mathbb{T}} \tau^{-n} d\mu(\tau). \end{aligned}$$

The interchange of the integral and summation is justified by the uniform convergence of the sum defining P_r . Hence we have $A_r \mu = P_r * \mu$. \square

Recall that a sequence of measures $\{\mu_n\}_{n=0}^{\infty}$ converges in the weak* sense if for any $f \in C(\mathbb{T})$, we have $\int_{\mathbb{T}} f d\mu_n \rightarrow \int_{\mathbb{T}} f d\mu$. Any $f \in L^1(\mathbb{T})$ can be interpreted as a measure μ_f , by $d\mu_f = f d\mathbf{m}$.

Proposition A.6. *Let μ be a complex Borel regular measure on \mathbb{T} . The functions $\{A_r \mu\}_{r \in (0,1)}$, interpreted as measures, converge in the weak* sense to μ as $r \rightarrow 1^-$.*

Proof. Let $f \in C(\mathbb{T})$. We compute,

$$\begin{aligned} \int_{\mathbb{T}} f d\mu_{A_r \mu} &= \int_{\mathbb{T}} f P_r * \mu d\mathbf{m} \\ &= \int_{\mathbb{T}} \int_{\mathbb{T}} f(\zeta) P_r(\zeta \tau^{-1}) d\mu(\tau) d\mathbf{m}(\zeta). \end{aligned}$$

However note that $P_r(\tau) = P_r(\tau^{-1})$, so by Fubini's theorem,

$$\begin{aligned} \int_{\mathbb{T}} f d\mu_{A_r \mu} &= \int_{\mathbb{T}} \int_{\mathbb{T}} P_r(\tau \zeta^{-1}) f(\zeta) d\mathbf{m}(\zeta) d\mu(\tau) \\ &= \int_{\mathbb{T}} P_r * f d\mu. \end{aligned}$$

However, since f is continuous, we have that $P_r * f \rightarrow f$ uniformly as $r \rightarrow 1^-$. Hence

$$\lim_{r \rightarrow 1^-} \int_{\mathbb{T}} f d\mu_{A_r \mu} = \int_{\mathbb{T}} f d\mu$$

So $A_r \mu$ converges in the weak* sense to μ . \square

A.5 Harmonic functions on \mathbb{D}

Abel summation is much more heavily studied than Cesàro summation. The reason for this is that there is a close connection between the Abel sums $A_r f$ for a function f on \mathbb{T} , and functions on $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$ which are *harmonic*.

Given $f \in L^1(\mathbb{T})$, define $\tilde{f} : \mathbb{D} \rightarrow \mathbb{C}$ by

$$\tilde{f}(r\zeta) = A_r f(\zeta) = (P_r * f)(\zeta)$$

for $r \in [0, 1]$ and $\zeta \in \mathbb{T}$.

Note that, by Young's convolution inequality,

$$\left(\int_{\mathbb{T}} |\tilde{f}(r\zeta)|^p d\mathbf{m}(\zeta) \right)^{1/p} \leq \|P_r\|_1 \|f\|_p.$$

Hence,

$$\sup_{r \in [0, 1)} \left(\int_{\mathbb{T}} |\tilde{f}(r\zeta)|^p d\mathbf{m}(\zeta) \right)^{1/p} \leq \|f\|_p.$$

Appendix B

Lorentz Spaces

B.1 Introduction

Lorentz spaces form a broad generalisation of L^p spaces and weak L^p spaces. These notes cover the basic definitions.

B.2 Weak L^p -spaces

Let (X, \mathcal{A}, μ) be a σ -finite measure space. Given a measurable function $f : X \rightarrow \mathbb{C}$, we may define the distribution function

$$d_f(\alpha) = \mu\{x \in X : |f(x)| \geq \alpha\}.$$

for $\alpha \geq 0$.

By Markov's inequality, for $p > 0$,

$$d_f(\alpha) \leq \frac{\|f\|_p^p}{\alpha^p}.$$

The linear span of the class of functions f for which

$$d_f(\alpha) \leq \frac{C^p}{\alpha^p}$$

for some constant C is called the weak L^p space or $L^{p,w}(X)$. Given $f \in L^{p,w}(X)$, define

$$\|f\|_{p,w} := \sup_{\alpha > 0} \alpha d_f(\alpha)^{1/p}.$$

B.3 Non-increasing Rearrangements

Again let (X, \mathcal{A}, μ) be a σ -finite measure space, and $f : X \rightarrow \mathbb{C}$ is measurable. Then define, for $t \geq 0$,

$$f^*(t) = \inf\{s \geq 0 : d_f(s) \leq t\}.$$

f^* is called the non-increasing rearrangement of f .

Lemma B.1.

$$\left(\int_X |f|^p d\mu \right)^{1/p} = \left(\int_0^\infty (f^*)^p d\mu \right)^{1/p}$$

Lemma B.2.

$$\sup_{t>0} t^q f^*(t) = \sup_{\alpha>0} \alpha d_f(\alpha)^{1/p}.$$

Inspired by the above two results is the following definition,

Definition B.3 (Lorentz spaces). Let $p, q > 0$. For a σ -finite measure space (X, \mathcal{A}, μ) , and a measurable function $f : X \rightarrow \mathbb{C}$, define

$$\|f\|_{p,q} = \left(\int_0^\infty (t^{1/p} f^*(t))^q \frac{dt}{t} \right)^{1/q}$$

and

$$\|f\|_{p,\infty} = \sup_{t \geq 0} t^{1/p} f^*(t).$$

B.4 Discrete spaces

When X is a set with counting measure, the $L^{p,q}$ spaces are denoted $\ell^{p,q}(X)$.

Appendix C

Interpolation

C.1 Introduction

The method of interpolation is a very powerful one in analysis, and it allows many results to be obtained from “edge” cases.

Historically, interpolation is motivated by the Riesz-Thorin theorem, which we state here.

Theorem C.1. *Let $p_0, q_0 \in [1, \infty]$ and $p_1, q_1 \in [1, \infty]$ with $p_0 \neq p_1$ and $q_0 \neq q_1$. Suppose that (U, μ) and (V, ν) are measure spaces. Let T be an operator such that*

$$T : L^{p_0}(U) \rightarrow L^{q_0}(V)$$

with norm M_0 and

$$T : L^{p_1}(U) \rightarrow L^{q_1}(V)$$

with norm M_1 .

Let $\theta \in (0, 1)$, and $p_\theta^{-1} = \theta p_0^{-1} + (1 - \theta)p_1^{-1}$, and $q_\theta^{-1} = \theta q_0^{-1} + (1 - \theta)q_1^{-1}$.

Then,

$$T : L^{p_\theta}(U) \rightarrow L^{q_\theta}(V)$$

with norm $M \leq M_0^\theta M_1^{1-\theta}$.

This theorem can be proved directly, however it is more insightful to prove it using abstract interpolation theory. The purpose of these notes is to introduce this theory.

C.2 Abstract Interpolation Theory

Let \mathbf{NLS} be the category of normed linear spaces with morphisms given by bounded linear maps.

Definition C.2. A pair $X, Y \in \mathbf{NLS}$ is called a compatible pair if X and Y are both subspaces of a topological vector space U .

Definition C.3. The category \mathbf{NLS}_1 is the category of compatible pairs (X, Y) of normed spaces, where a morphism $T : (X_1, Y_1) \rightarrow (X_2, Y_2)$ is a linear map $T : X_1 + Y_1 \rightarrow X_2 + Y_2$ such that $T : X_1 \rightarrow X_2$ and $T : Y_1 \rightarrow Y_2$ is bounded.

Proposition C.4. Let $\Delta : \mathbf{NLS}_1 \rightarrow \mathbf{NLS}$ be the function that maps (X, Y) to $X \cap Y$, where $X \cap Y$ is given the norm,

$$\|x\|_{X \cap Y} = \max\{\|x\|_X, \|x\|_Y\}.$$

Let $\Sigma : \mathbf{NLS}_1 \rightarrow \mathbf{NLS}$ be given by $\Sigma((X, Y)) = X + Y$, where $X + Y$ is given the norm,

$$\|x\|_{X+Y} = \inf\{\|x_1\|_X + \|x_2\|_Y : x = x_1 + x_2, x_1 \in X, x_2 \in Y\}.$$

Then Δ and Σ are functors.

Definition C.5. An interpolation functor is a functor $\mathcal{F} : \mathbf{NLS}_1 \rightarrow \mathbf{NLS}$ such that for $(X, Y) \in \mathbf{NLS}_1$, we have

$$\Delta((X, Y)) \subseteq \mathcal{F}((X, Y)) \subseteq \Sigma((X, Y))$$

We say that an interpolation functor \mathcal{F} is *uniform* if for any morphism $T : (X_1, Y_1) \rightarrow (X_2, Y_2)$ in \mathbf{NLS}_1 , we have

$$\|\mathcal{F}(T)\| \leq C \max\{\|T\|_{X_1 \rightarrow X_2}, \|T\|_{Y_1 \rightarrow Y_2}\}.$$

for some constant $C > 0$. If $C = 1$, we say that \mathcal{F} is *exact*.

We say that an interpolation functor \mathcal{F} is of exponent $\theta \in (0, 1)$ if

$$\|\mathcal{F}(T)\| \leq C \|T\|_{X_1 \rightarrow X_2}^\theta \|T\|_{Y_1 \rightarrow Y_2}^{1-\theta}.$$

for some constant $C > 0$. If $C = 1$, we say that \mathcal{F} is *exact of exponent θ* .

C.3 Real Interpolation: The K Method

Definition C.6. Let $(X_0, X_1) \in \mathbf{NLS}_1$. For $x \in X_0 + X_1$ and $t > 0$, define

$$K(x, t; X_0, X_1) = \inf \{ \|x_0\| + t\|x_1\| : x_0 \in X_0, x_1 \in X_1, x = x_0 + x_1 \}.$$

Then for $\theta \in (0, 1)$ and $q \in [1, \infty)$, define

$$\|x\|_{\theta, q; K} := \left(\int_0^\infty (t^{-\theta} K(x, t; X_0, X_1))^q \frac{dt}{t} \right)^{1/q}$$

and for $\theta \in [0, 1]$,

$$\|x\|_{\theta, \infty; K} := \sup_{t>0} t^{-\theta} K(x, t; X_0, X_1).$$

The space $K_{\theta, q}(X_0, X_1)$ is the set of $x \in X_0 + X_1$ such that $\|x\|_{\theta, q; K} < \infty$.

Proposition C.7. *The function $(X, Y) \mapsto K_{\theta, q}(X, Y)$ is an interpolation functor.*

C.4 Complex Interpolation

Definition C.8. Let (X_0, X_1) be a compatible pair of Banach spaces. Let $\mathcal{S} := \{z \in \mathbb{C} : \Re(z) \in (0, 1)\}$.

Define the set $\mathcal{F}(X_0, X_1)$ to be the space of functions $f : \overline{\mathcal{S}} \rightarrow X_0 + X_1$ which are complex differentiable in \mathcal{S} , continuous on $\overline{\mathcal{S}}$ and bounded on $\partial\mathcal{S}$.

It is true that $\mathcal{F}(X_0, X_1)$ is a Banach space under the norm,

$$\|f\|_{\mathcal{F}(X_0, X_1)} = \max \left\{ \sup_{t \in \mathbb{R}} \|f(it)\|, \sup_{t \in \mathbb{R}} \|f(1+it)\| \right\}.$$

Let $\theta \in (0, 1)$. Then, define

$$(X_0, X_1)_\theta = \{f(\theta) : f \in \mathcal{F}(X_0, X_1)\}.$$

We define the norm,

$$\|x\|_{(X_0, X_1)_\theta} = \inf \{ \|f\|_{\mathcal{F}(X_0, X_1)} : x = f(\theta) \}.$$

Proposition C.9. *The mapping $(X_0, X_1) \rightarrow (X_0, X_1)_\theta$ is an interpolation functor, which is exact of exponent θ .*

C.5 Using interpolation

Once we know how to describe certain interpolation spaces, results such as the Riesz-Thorin theorem become immediate.

Proposition C.10. *Let X be a measure space. Let $p_0, p_1 \in [1, \infty]$, and $p_\theta^{-1} = \theta p_0^{-1} + (1 - \theta)p_1^{-1}$ for $\theta \in (0, 1)$. Then*

$$(L^{p_0}(X), L^{p_1}(X))_\theta = L^{p_\theta}(X).$$

Another immediate corollary:

Proposition C.11. *Let G be a locally compact abelian group equipped with bi-variant Haar measure μ , and $p \in [1, \infty]$. For $\varphi \in L^1(G, \mu)$, the map $Tf = \varphi * f$ is bounded on $L^p(G, \mu)$, with norm less than or equal to $\|\varphi\|_1$.*

Proof. To prove the case $p = 1$, we let $f \in L^p(G)$, and compute,

$$\begin{aligned} \|\varphi * f\|_1 &= \int_G \left| \int_G \varphi(x - y) f(y) d\mu(y) \right| d\mu(x) \\ &\leq \int_G \int_G |\varphi(x - y)| |f(y)| d\mu(y) d\mu(x) \\ &= \|\varphi\|_1 \|f\|_1 \end{aligned}$$

where the interchange of integrals is justified by Tonelli's theorem.

Now for $p = \infty$, we compute,

$$\|\varphi * f\|_\infty = \text{ess-sup}_{x \in G} \left| \int_G \varphi(x - y) f(y) d\mu(y) \right|$$

By Hölder's inequality, this can be bounded by $\|\varphi\|_1 \|f\|_\infty$.

The rest of the cases follow from complex interpolation of the pair $(L^1(G, \mu), L^\infty(G, \mu))$. □

Appendix D

Group Actions on Von Neumann Algebras

D.1 Introduction

There is a beautiful and extraordinary generalisation of the Fourier transform in the theory of Von Neumann algebras relating to the action of compact groups. The purpose of these notes is to give a basic exposition of this idea.

D.2 Compact Group actions

Let G be a compact abelian group equipped with normalised Haar measure μ , and let \mathcal{M} be a Von Neumann algebra.

Recall that a group action on a Von Neumann algebra is a group action $\alpha : G \times \mathcal{M} \rightarrow \mathcal{M}$ such that for all $g \in G$ the function $a \mapsto \alpha(g, a)$ is an algebra homomorphism.

Suppose that G acts on \mathcal{M} in a way that is

1. *ergodic*: the only projections in \mathcal{M} fixed by G are 0 and 1 and
2. *free*: there is no nontrivial projection $p \in \mathcal{M}$ such that some $g \in G$ not equal to the identity fixes all of $p\mathcal{M}p$.

We then have the following result:

Proposition D.1. *Suppose that G is a compact abelian group that acts freely and ergodically on the Von Neumann algebra \mathcal{M} , by the action $\alpha : G \times \mathcal{M} \rightarrow \mathcal{M}$. Then there is a set*

$$\{u(p) : p \in \widehat{G}\}$$

of unitary eigenoperators for the action indexed by the dual group \widehat{G} , and the map $p \mapsto u(p)$ is a representation of \widehat{G} .

Proof. Let $x \in \mathcal{M}$ and $p \in \widehat{G}$. Define

$$\hat{x}(p) := \int_G \alpha(s, x) p(s)^{-1} d\mu(s).$$

Let $g \in G$, then we can compute,

$$\begin{aligned} \alpha(g, \hat{x}(p)) &= \int_G \alpha(sg, x) p(s)^{-1} d\mu(s) \\ &= p(g) \hat{x}(p). \end{aligned}$$

Hence we have that $\hat{x}(p)$ is an eigenoperator for the action of G . Hence, for $x, y \in \mathcal{M}$, we have that $\hat{x}(p) \hat{y}(p)^*$ is a fixed point of G , hence a scalar multiple of 1.

Then define

$$u(p) = \frac{\hat{x}(p)}{\|\hat{x}(p)\|}.$$

Hence u is unitary, and for each $y \in \mathcal{M}$, we have $\hat{y}(p)$ is a scalar multiple of $u(p)$. \square

Example D.1. *The prototypical example of this decomposition is for $\mathcal{M} = L^\infty(\mathbb{T})$, and $G = \mathbb{T}$ with Haar measure acting on \mathcal{M} by translation. Then $\widehat{G} = \mathbb{Z}$, and for $f \in L^\infty(\mathbb{T})$ and $n \in \mathbb{Z}$, we have*

$$\hat{f}(n)(\zeta) = \int_{\mathbb{T}} f(\tau \zeta) \tau^{-n} d\mathbf{m}(\tau)$$

This is simply the n th Fourier coefficient times ζ^n . Hence we have

$$u(n)(\zeta) = \zeta^n.$$

So the system of unitaries is the set of monomials on \mathbb{T} .

The important feature of this system of unitaries is that it spans \mathcal{M} . First we recall the definition of the σ -weak topology.

Definition D.2. Let \mathcal{H} be a Hilbert space, and let $\{\xi_j\}_{j=0}^\infty$ and $\{\eta_j\}_{j=0}^\infty$ be sequences of elements of \mathcal{H} such that

$$\sum_{j=0}^{\infty} \|\xi_j\|^2 < \infty$$

$$\sum_{j=0}^{\infty} \|\eta_j\|^2 < \infty.$$

Then for $a \in \mathcal{B}(\mathcal{H})$, define the semi-norm,

$$\left| \sum_{j=0}^{\infty} \langle \xi_j, a\eta_j \rangle \right|.$$

This system of semi-norms defines the σ -weak topology on $\mathcal{B}(\mathcal{H})$.

Proposition D.3. Let G be a compact abelian group acting freely and ergodically on a Von Neumann algebra \mathcal{M} . Let

$$\mathcal{P} = \{u(p) : p \in \widehat{G}\}$$

be the corresponding unitary eigenoperators of G . Then the span of \mathcal{P} is dense in the σ -weak topology on \mathcal{M} .

Proof. (sketch: See [?] 8.1.6 for details) This follows from the claim that the Arveson spectrum $\text{Sp}(\alpha)$ of α is \widehat{G} . We claim that since

$$\text{Sp}^\perp(\alpha) = \{s \in G : \alpha(s, x) = x \text{ for all } x \in \mathcal{M}\}$$

we have that $\text{Sp}^\perp(\alpha) = \{0\}$ because α is free, from which it follows that $\text{Sp}(\alpha) = \widehat{G}$.

Consequently, the system $\{u(p) : p \in \widehat{G}\}$ is σ -weakly dense in \mathcal{M} . \square

D.3 The space $\mathcal{L}^2(\mathcal{M}, \tau)$

Suppose that τ is a faithful trace on \mathcal{M} such that $\tau(1) = 1$, and τ is invariant under the action of G , that is for all $g \in G$ and $x \in \mathcal{M}$ we have $\tau(\alpha(g, x)) = \tau(x)$. Then the map $(x, y) \mapsto \tau(x^*y)$ is an inner product on \mathcal{M} , and the completion of \mathcal{M} in this inner product is denoted $\mathcal{L}^2(\mathcal{M}, \tau)$.

We require the following lemma from Pedersen [?], Theorem 3.6.5,

Lemma D.4. *A state φ on a Von Neumann algebra is normal if and only if it is σ -weakly continuous.*

Since \mathcal{M} carries a group action, such a trace exists.

Lemma D.5. *Let \mathcal{M} be a Von Neumann algebra, and let G act on \mathcal{M} freely and ergodically. Then there is a G -invariant state on \mathcal{M} , and it is a faithful normal trace.*

Proof. Define for $x \in \mathcal{M}$,

$$\tau(x) = \int_G \alpha(s, x) d\mu(s) = \hat{x}(0).$$

See that $\alpha(g, \tau(x)) = \tau(x)$, so $\tau(x)$ is G -invariant so by ergodicity must be in $\mathbb{C}1$. If we identify $\mathbb{C}1$ with \mathbb{C} , we can think of $\tau(x)$ as a scalar, so $\tau(x)$ is a state. By the continuity of the group action, we have that τ is normal, and if $\tau(x^*x) = 0$, we must have $\alpha(s, x)^* \alpha(s, x) = 0$ for all $s \in G$, so $x = 0$. Hence τ is faithful.

To prove that τ is a trace it is sufficient to prove that

$$\tau(u(p)u(q)) = \tau(u(q)u(p))$$

for all $p, q \in \widehat{G}$.

Let $g \in G$, then we have

$$\begin{aligned} \tau(u(p)u(q)) &= \tau(\alpha(g, u(p)u(q))) \\ &= p(g)q(g)\tau(u(p)u(q)). \end{aligned}$$

If $p \neq q^{-1}$, we can find g such that $p(g)q(g) \neq 1$. Thus, $\tau(u(p)u(q)) = 0$.

Otherwise, if $p = q^{-1}$ there is a scalar λ_p with $|\lambda_p| = 1$ such that

$$\tau(u(p)u(q)) = \lambda_p \tau(u(p)u(p)^*) = \lambda_p.$$

Hence $\tau(u(p)u(q)) = \tau(u(q)u(p))$.

Hence, since τ is normal, it is σ -weakly continuous. Since the set $\{u(p) : p \in \widehat{G}\}$ is σ -weakly dense in \mathcal{M} , we have that $\tau(xy) = \tau(yx)$ for all $x, y \in \mathcal{M}$.

□

Proposition D.6. *Let $\mathcal{P} := \{u(p) : p \in \widehat{G}\}$ be the set of unitary eigenoperators corresponding to the action of G . Then $\{u(p) : p \in \widehat{G}\}$ is an orthonormal basis for $\mathcal{L}^2(\mathcal{M}, \tau)$.*

Proof. By standard Hilbert space theory, it is sufficient to prove that \mathcal{P} is orthonormal and has dense span.

First we prove ortho-normality. Let $p, q \in \widehat{G}$. Then for all $g \in G$,

$$\begin{aligned}\tau(u(p)^*u(q)) &= \tau(p(g)^{-1}q(g)u(p)^*u(q)) \\ &= p(g)^{-1}q(g)\tau(u(p)^*u(q)).\end{aligned}$$

If $p \neq q$, there is some $g \in G$ such that $p(g)^{-1}q(g) \neq 1$, so we conclude that $\tau(u(p)^*u(q)) = 0$. If $p = q$, then by unitarity we have

$$\tau(u(p)^*u(q)) = \tau(1) = 1.$$

Hence \mathcal{P} is orthonormal.

To prove that the span of \mathcal{P} is dense in $\mathcal{L}^2(\mathcal{M}, \tau)$, it is sufficient to prove that it is dense in \mathcal{M} in the norm $\|x\|_2 := \sqrt{\tau(x^*x)}$. This follows from the σ -weak continuity of τ . \square

Remark D.7. In fact, for any $p \geq 1$, we have that the norm $\|x\|_p = (\tau(|x|^p))^{1/p}$ is σ -weakly continuous, so $\{u(p) : p \in \widehat{G}\}$ is dense in $\mathcal{L}^p(M, \tau)$, defined as the completion of \mathcal{M} in the norm $\|\cdot\|_p$.

D.4 Non-commutative Harmonic Analysis

(The following is entirely original and therefore suspect)

We now fix $G = \mathbb{T}^d$, so that $\widehat{G} = \mathbb{Z}^d$. It is of interest to prove certain results that are analogous to classical results of harmonic analysis. In particular, one known classical result is that if $f \in L^1(\mathbb{T}^d, \mathbf{m})$, then the Fourier coefficients $\hat{f}(n)$ vanish as $\|n\| \rightarrow \infty$. We now prove an analogy of this result.

Let \mathbf{m} denote the normalised Haar measure on \mathbb{T}^d .

From now on, denote the action of $t \in \mathbb{T}^d$ on $x \in \mathcal{M}$ as $\alpha_t(x)$.

Fix \mathcal{M} a Von Neumann Algebra, and let \mathbb{T}^d act on \mathcal{M} freely and ergodically. Let τ be the unique normalised faithful \mathbb{T}^d -invariant trace on \mathcal{M} . Let

$$\mathcal{P} := \{u(n) : n \in \mathbb{Z}^d\}$$

be the spanning system of unitary eigenoperators of the group action.

For $a \in \mathcal{L}^1(\mathcal{M}, \tau)$ and $n \in \mathbb{Z}^d$, define

$$\hat{a}(n) := \tau(au(n)^*)$$

For any $a \in \mathcal{L}^2(\mathcal{M}, \tau)$, we have that

$$a = \sum_{n \in \mathbb{Z}^d} \hat{a}(n)u(n)$$

where the convergence is in the \mathcal{L}^2 norm. This is an isomorphism with $\ell^2(\mathbb{Z}^d)$. In general, for $N = (N_1, \dots, N_d) \in \mathbb{Z}^d$, define

$$S_N a := \sum_{n=-N}^N \hat{a}(n)u(n)$$

where the summation runs over all multi-indices $n = (n_1, \dots, n_d)$ such that for each $1 \leq j \leq d$, we have $-N_j \leq n_j \leq N_j$. and

$$\sigma_N a := \frac{1}{N_1 N_2 \dots N_d} \sum_{n=(1, \dots, 1)}^N S_n a.$$

and the summation runs over multi-indices $n = (n_1, \dots, n_d)$ such that for each j , $1 \leq n_j \leq N_j$.

Now we define the subspaces of “continuous” and “uniformly continuous” functions in \mathcal{M} .

Definition D.8. Define $\mathcal{C}(\mathcal{M})$ to be the closure of the span of \mathcal{P} in the norm topology of \mathcal{M} .

Define $\mathcal{U}(\mathcal{M})$ to be the set of elements of \mathcal{M} such that for any $\varepsilon > 0$ there exists $\delta > 0$ such that

$$\sup_{|t| < \delta} \|\alpha_t(x) - x\| < \varepsilon.$$

Where, for $t \in \mathbb{T}^d$, If $t = (e^{2\pi i \varphi_1}, \dots, e^{2\pi i \varphi_d})$, we denote

$$|t| = \max\{|\varphi_1|, \dots, |\varphi_d|\}.$$

Definition D.9. Let $n = (n_1, \dots, n_d) \in \mathbb{Z}^d$, and $t = (t_1, \dots, t_d) \in \mathbb{T}^d$. Then we introduce the notation

$$t^d := t_1^{n_1} \dots t_d^{n_d}.$$

Lemma D.10. *We have the inclusion,*

$$\mathcal{C}(\mathcal{M}) \subseteq \mathcal{U}(\mathcal{M}).$$

Proof. First let $u(n) \in \mathcal{P}$. Then for $t \in \mathbb{T}^d$, we have

$$\alpha_t(u(n)) = t^n u(n)$$

Hence

$$\|\alpha_t(u(n)) - u(n)\| = |t^n - 1| \|u(n)\|.$$

So since $\lim_{t \rightarrow 1} |t^n - 1| = 0$, we have that for any $\varepsilon > 0$ there exists $\delta > 0$ such that

$$\sup_{|t| < \delta} \|\alpha_t(u(n)) - u(n)\| < \varepsilon.$$

Now let a be in the linear span of \mathcal{P} . Since a is only a finite linear combination of terms of the form $u(n)$, $n \in \mathbb{Z}$, we have that for any $\varepsilon > 0$ there exists a $\delta > 0$ such that

$$\sup_{|t| < \delta} \|\alpha_t(a) - a\| < \varepsilon.$$

Hence the linear span of \mathcal{P} is contained in $\mathcal{U}(\mathcal{M})$. We will be finished if we can show that $\mathcal{U}(\mathcal{M})$ is closed in the norm topology.

Suppose that $\{x_n\}_{n=0}^\infty$ is a sequence in $\mathcal{U}(\mathcal{M})$ converging in the norm topology to $x \in \mathcal{M}$. Then let $t \in \mathbb{T}^d$, by the triangle inequality,

$$\|\alpha_t(x) - x\| < \|\alpha_t(x) - \alpha_t(x_n)\| + \|\alpha_t(x_n) - x_n\| + \|x_n - x\|$$

By assumption, the group action is continuous in the norm topology, so there is a constant C such that $\|\alpha_t(x) - \alpha_t(x_n)\| < C\|x - x_n\|$. Hence the result follows. \square

Definition D.11. Suppose that $\varphi \in L^1(\mathbb{T}^d, \mathbf{m})$, and $x \in \mathcal{M}$. Define the *convolution*

$$\varphi * x = \int_{\mathbb{T}^d} \varphi(t) \alpha_t(x) d\mathbf{m}(t).$$

It can be proved that

$$S_N x = D_N * x$$

where

$$D_N(t) = \prod_{j=1}^d \frac{t_j^{N_j+1/2} - t_j^{-N_j-1/2}}{t_j^{1/2} - t_j^{-1/2}}$$

and

$$\sigma_N x = F_N * x$$

where

$$F_N(t) = \prod_{j=1}^d \frac{t_j^{N_j} - 2 + t_j^{-N_j}}{N(t_j^{1/2} - t_j^{-1/2})^2}$$

where the fractional powers in the formulae for D_N and F_N are defined in a principal value sense. See [?] for proofs.

Now we define approximate identities,

Definition D.12. A net $\{\Phi_\lambda\}_{\lambda \in \Lambda} \subset L^1(\mathbb{T}^d, \mathbf{m})$ is called an approximation to the identity if,

1. For all λ , we have $\int_{\mathbb{T}^d} \Phi_\lambda d\mathbf{m} = 1$.
2. We have $\sup_\lambda \int_{\mathbb{T}^d} |\Phi_\lambda| d\mathbf{m} < \infty$
3. For any $\delta > 0$, we have $\lim_{\lambda \in \Lambda} \int_{|t| > \delta} |\Phi_\lambda| d\mathbf{m} = 0$.

It can be proved that the sequence $\{F_n\}_{n \in \mathbb{N}^d}$ is an approximate identity (see [?]).

Approximate identities are so named because of the following two results:

Proposition D.13. Let $x \in \mathcal{U}(\mathcal{M})$, and let $\{\Phi_\lambda\}_{\lambda \in \Lambda}$ be an approximate identity. Then we have

$$\lim_{\lambda \in \Lambda} \|\Phi_\lambda * x - x\| = 0.$$

Proof. Using the fact that $\int_{\mathbb{T}} \Phi_\lambda d\mu = 1$, we compute,

$$\Phi_n * x - x = \int_{\mathbb{T}} \Phi_n(t)(\alpha_t(x) - x) d\mu(t).$$

Let $\varepsilon > 0$. Choose δ small enough such that

$$\sup_{|t| < \delta} \|\alpha_t(x) - x\| < \varepsilon.$$

Now estimate,

$$\|\Phi_\lambda * x - x\| \leq \int_{|t| < \delta} |\Phi_\lambda(t)| \|\alpha_t(x) - x\| d\mu(t) + (C + 1) \int_{|t| \geq \delta} |\Phi_\lambda(t)| \|x\| d\mu(t)$$

where C is a constant such that $\|\alpha_t(x)\| < C\|x\|$. So taking the limit over λ , we have

$$\lim_{\lambda \in \Lambda} \|\Phi_\lambda * x - x\| < \varepsilon \sup_{\lambda \in \Lambda} \int_{\mathbb{T}} |\Phi_n| d\mathbf{m}.$$

But ε is arbitrary, so the result follows. \square

Proposition D.14. *We have that*

$$\mathcal{C}(\mathcal{M}) = \mathcal{U}(\mathcal{M}).$$

Proof. Let $a \in \mathcal{U}(\mathcal{M})$. Since $\{F_n\}_{n \in \mathbb{N}^d}$ is an approximate identity, we have that $F_n * a \rightarrow a$ in the norm topology. But $F_n * a \in \text{span}(\mathcal{P})$. Hence $a \in \mathcal{C}(\mathcal{M})$. \square

As usual, define $\|x\|_p = \tau(|x|^p)^{1/p}$. We require the following lemma:

Lemma D.15. *Let $p \geq 1$. Suppose $\varphi \in L^1(\mathbb{T}^d, \mathbf{m})$, and $x \in \mathcal{L}^p(\mathcal{M}, \tau)$. Then*

$$\|\varphi * x\|_p \leq \|\varphi\|_1 \|x\|_p.$$

Proof. First we establish the case $p = 1$. This is a computation,

$$\|\varphi * x\|_1 \leq \int_{\mathbb{T}^d} |\varphi(t)| \|\alpha_t(a)\|_1 d\mathbf{m}(t).$$

Similarly, we define $\|x\|_\infty = \|x\|$. Thus the $p = \infty$ case,

$$\|\varphi * x\|_\infty \leq \|\varphi\|_1 \|x\|.$$

Hence by interpolation, the result follows. \square

Proposition D.16. *If $\{\Phi_\lambda\}_{\lambda \in \Lambda}$ is an approximate identity, and $x \in \mathcal{L}^p(\mathcal{M}, \tau)$ and $p \geq 1$ then*

$$\lim_{\lambda \in \Lambda} \|\Phi_\lambda * x - x\|_p = 0.$$

Proof. Let $\varepsilon > 0$. Since $\mathcal{U}(\mathcal{M})$ contains the linear span of \mathcal{P} , and the linear span of \mathcal{P} is dense in $\mathcal{L}^p(\mathcal{M}, \tau)$ in the $\|\cdot\|_p$ norm, we can find $y \in \mathcal{U}(\mathcal{M})$ such that $\|x - y\|_p < \varepsilon$. Hence,

$$\begin{aligned} \|\Phi_\lambda * x - x\|_p &\leq \|\Phi_\lambda * x - \Phi_\lambda * y\|_p + \|\Phi_\lambda * y - y\|_p + \|y - x\|_p \\ &\leq \sup_{\mu \in \Lambda} \|\Phi_\mu\|_1 \varepsilon + \|\Phi_\lambda * y - y\|_p + \varepsilon. \end{aligned}$$

Now take the limit over λ , and thus we obtain the result. \square

At last we can prove the following result:

Proposition D.17. *Let $x \in \mathcal{L}^1(\mathcal{M}, \tau)$. Then we have $\hat{x}(n) \rightarrow 0$ as $\|n\| \rightarrow \infty$.*

Proof. For $n = (n_1, \dots, n_d) \in \mathbb{Z}^d$, denote $|n| := (n_1, \dots, n_d)$. We have that $\tau(\sigma_{|N|-1}xu(N)^*) = 0$. Hence,

$$\begin{aligned} |\hat{x}(n)| &= |\tau(xu(n)^*)| \\ &\leq |\tau((x - \sigma_{|n|-1}x)u(n)^*)| \\ &\leq \|u(n)^*\| \|x - \sigma_{|n|-1}x\|_1 \\ &= \|x - F_{|n|-1} * x\|_1. \end{aligned}$$

But the right hand side vanishes as $\|n\| \rightarrow \infty$. \square

D.5 The Operators δ_j

Let \mathcal{M} be a Von Neumann Algebra, and let \mathbb{T}^d act freely and ergodically on \mathcal{M} .

The generalised differentiation operator δ_j is defined as the *infinitesimal generator* of the action of \mathbb{T}^d , as follows,

Definition D.18. Let $j \in \{1, \dots, d\}$.

For $t \in \mathbb{T}$, we have an action α_t^j on \mathcal{M} which is the action of the j th coordinate of \mathbb{T}^d on \mathcal{M}

For $x \in \mathcal{M}$, we define

$$\delta_j(x) = \lim_{t \rightarrow 1} \frac{\alpha_t^j(x) - x}{|t|}$$

where $|t|$ is the minimal normalised arc length between t and 1.

The limit is in the sense of the norm topology on \mathcal{M} .

This limit may not exist for all x . Let $\text{Dom}(\delta_j)$ be the set of all x such that $\delta_j(x)$ exists.

Note that $\text{Dom}(\delta_j)$ is automatically a vector space.

Lemma D.19. We have $\mathcal{P} \subset \text{Dom}(\delta_j)$.

Proof. For $t \in \mathbb{T}$, and $u(n) \in \mathcal{P}$ with $n = (n_1, \dots, n_d)$, we have $\alpha_t^j u(n) = t_j^{n_j} u(n)$.

Hence,

$$\frac{\alpha_t^j(u(n)) - u(n)}{|t|} = \frac{t_j^{n_j} - 1}{|t|} u(n).$$

We parametrise \mathbb{T} by $t \mapsto \exp(i\theta)$, for $\theta \in [0, 2\pi)$. Then we have

$$\frac{t^{n_j} - 1}{|t|} = \frac{\exp(in_j\theta) - 1}{\theta}.$$

Hence,

$$\begin{aligned} \delta_j(u(n)) &= \lim_{t \rightarrow 1} \frac{t^{n_j} - 1}{|t|} u(n) \\ &= \lim_{\theta \rightarrow 0} \frac{\exp(in_j\theta) - 1}{\theta} u(n) \\ &= in_j u(n). \end{aligned}$$

Hence, $u(n) \in \text{Dom}(\delta_j)$. □

Proposition D.20. *Suppose that $x \in \text{Dom}(\delta_j)$. Then*

$$\widehat{\delta_j(x)}(n) = in_j \hat{x}(n).$$

Proof. By definition, $\text{Dom}(\delta_j) \subseteq \mathcal{M} \subseteq \mathcal{L}^1(\mathcal{M}, \tau)$.

Let $x \in \text{Dom}(\delta_j)$.

Hence, we have $F_n * x \rightarrow x$ in the \mathcal{L}^1 sense, and since $\delta_j(x) \in \mathcal{L}^1(\mathbb{T})$, we have $F_n * \mathcal{D}(x) \rightarrow \mathcal{D}(x)$ in the \mathcal{L}^1 sense.

Note that since $F_n * x$ is in the linear span of \mathcal{P} , we have $F_n * x \in \text{Dom}(\delta_j)$. See that

$$\delta_j(F_n * x) = \frac{1}{n_1 n_2 \cdots n_d} \sum_{k=0}^n \delta_j(D_k * x).$$

where the sum is over multi-indices $k = (k_1, \dots, k_d)$ with each $1 \leq k_m \leq n_m$. We also have,

$$\begin{aligned} \delta_j(D_k * x) &= \sum_{j=-k}^k \hat{x}(j) \mathcal{D}(u(j)) \\ &= \sum_{j=-k}^k ij \hat{x}(j) u(j). \end{aligned}$$

Now we have,

$$\begin{aligned} \widehat{\delta_j(x)}(n) &= \lim_{k \rightarrow \infty} \tau(F_k * xu(-n)) \\ &= \lim_{k \rightarrow \infty} in_j \hat{x}(n). \end{aligned}$$

□

Now define

$$\mathcal{D}_j := \frac{1}{i} \delta_j.$$

We define $\text{Dom}(\mathcal{D}_j) = \text{Dom}(\delta_j)$.

Hence we have the formula,

$$\widehat{\mathcal{D}_j x}(n) = n_j \hat{x}(n).$$

Appendix E

Dirac Operators

E.1 Introduction

This chapter is intended to give an introduction to the relationship between the Dirac operator and the exterior algebra. The general philosophy is that any expressions involving coordinates are to be avoided.

Throughout these notes, (M, g) is a Riemannian manifold.

E.2 Music, Clifford bundles and Modules

A metric g on a manifold M gives us a canonical isomorphism between T^*M and TM , called \sharp , pronounced “sharp”. For $x \in M$, given a linear functional $\omega \in T_x^*M$ we define $\sharp\omega$ to be the unique vector such that $\omega(v) = g(\sharp\omega, v)$ for all $v \in T_xM$. This is called the “musical isomorphism”.

The Clifford bundle of (M, g) is a vector bundle on M defined as follows.

Definition E.1. Let $x \in M$. The clifford algebra at x , $\text{Cliff}_x(M, g)$ is defined as the free associative unital algebra generated by T_xM modulo the relation

$$uv + vu = -2g(u, v)1$$

where $u, v \in \text{Cliff}_x(M, g)$, and 1 is the identity in $\text{Cliff}_x(M, g)$.

The clifford bundle $\text{Cliff}(M, g)$ is the vector bundle on M whose fibres are $\text{Cliff}_x(M, g)$.

Let's not care about the topology on $\text{Cliff}(M, g)$ at the moment.

There is clearly an embedding $TM \hookrightarrow \text{Cliff}(M, g)$.

Now for a vector bundle V on M , we say that V is a clifford module if there is a right multiplication map $\gamma : \text{Cliff}(M, g) \otimes V \rightarrow V$.

A *connection* on V is a linear map

$$\nabla : V \rightarrow T^*M \otimes V.$$

satisfying the Leibniz rule, for $f \in C^\infty(M)$ and $v \in V$,

$$\nabla(fv) = df \otimes v + f\nabla(v).$$

Now we may define a *Dirac operator*. Suppose V is a clifford bundle with connection ∇ . Then the composition of linear maps,

$$V \xrightarrow{\nabla} T^*M \otimes V \xrightarrow{\sharp \otimes I} TM \otimes V \xrightarrow{\gamma} V$$

is called the Dirac operator associated with V and ∇ .

E.3 Relationship with differentials

Suppose we have a clifford bundle V with connection ∇ .

Via the musical isomorphism, we may regard any differential form $\omega \in \Gamma(T^*M)$ as an operator on $\Gamma(V)$, since $\sharp(\omega)$ is an element of $\Gamma(\text{Cliff}(M, g))$, it may act on V .

Similarly, by pointwise multiplication, any $f \in C^\infty(M)$ is an operator on $\Gamma(V)$.

Theorem E.2. *We have an equality of operators on $\Gamma(V)$,*

$$[D, f] = df.$$

for any $f \in C^\infty(M)$.

Proof. Let $f \in C^\infty(M)$ and $v \in \Gamma(V)$. Let us compute $D(fv)$.

By definition,

$$D(fv) = (\gamma \circ (\sharp \otimes I) \circ \nabla)(fv).$$

By the Leibniz rule,

$$(\sharp \otimes I)\nabla(fv) = \sharp(df) \otimes v + (I \otimes f)(\sharp \otimes 1)\nabla(v).$$

Hence,

$$D(fv) = \gamma(\sharp(df))v + fD(v).$$

Therefore, $[D, f]v = df(v)$.

□

Bibliography

- [1] Alain Connes. *Non-commutative Geometry*. InterEditions Paris, Paris, France, 1994.