

Geodesic Loss Aversion in Predictive Interaction Field Theory: The Kahneman–Tversky Value Function as an Interaction Source Tensor

Chewin Pinmook

Independent Theoretical Development

chewinp@gmail.com

May 2026

Abstract

Predictive Interaction Field Theory (PIFT) models behavioral dynamics as geodesic flow on a Riemannian manifold $(\mathcal{M}, g_{\mu\nu})$ whose curvature is sourced by the Expected Free Energy $\mathcal{G} \approx \mathcal{A} + \mathcal{R}$ (Ambiguity plus Relative Risk). This paper establishes a principled, first-principles derivation showing that the Kahneman–Tversky (KT) value function $V_{\text{KT}}(\Delta)$ is exactly recoverable as the geodesic arc-length on the PIFT behavioral manifold. Rather than fitting the metric post hoc, we begin from an information-geometric motivaton and define the one-dimensional *Interaction Source Tensor* component as $T_{11}(\Delta) = [\text{d}V_{\text{KT}}/\text{d}\Delta]^2$, which represents the squared marginal psychological sensitivity—the natural Fisher-information density of the value function. Under the isometric coupling condition $g_{11} = T_{11}$ (realised in the behaviorally determined vacuum solution of the PIFT field equations), the arc-length functional $L(x) = \int_0^x \sqrt{g_{11}(\Delta)} \text{d}\Delta$ satisfies $L(x) \equiv V_{\text{KT}}(x)$ for all $x \in \mathbb{R} \setminus \{0\}$. We demonstrate that the singularity of g_{11} at the gain/loss boundary $\Delta = 0$ is integrable for all $\alpha \in (0, 1)$, provide an intrinsic geometric explanation of the Framing Effect, prove that the metric-level loss-aversion ratio equals $\lambda^2 = 5.0625$ —the square of the KT behavioral ratio $\lambda = 2.25$ —and confirm numerical agreement at machine-precision RMSE. The Bianchi identity is shown to hold in the distributional sense. These results establish that Prospect Theory is not merely analogous to Riemannian geometry but is isometrically embeddable into it, with loss aversion emerging as a curvature asymmetry whose magnitude is uniquely determined by the source tensor.

Keywords: Predictive Interaction Field Theory, Prospect Theory, Kahneman–Tversky value function, loss aversion, Riemannian geodesic, interaction source tensor, Fisher information geometry, Expected Free Energy, integrable singularity.

Contents

1	Introduction	4
2	Background: PIFT Metric Structure	5
2.1	Interaction manifold	5
2.2	PIFT field equations	5
2.3	Geodesic arc-length as value functional	5
3	The Behavioral Source Tensor	6
3.1	Information-geometric motivation	6
3.2	Pullback metric interpretation	6
3.3	Definition of the source tensor component	6
4	Isometric Coupling and the Induced Metric	7
4.1	Behavioral vacuum solution	7
4.2	Explicit metric component	7
5	Geodesic Arc-Length and Exact Recovery of V_{KT}	8
6	Singularity Analysis at $\Delta = 0$	8
6.1	Geometric interpretation of the singularity	8
7	Geometric Loss-Aversion Ratio	9
7.1	Two levels of loss aversion	9
7.2	Christoffel analysis	10
8	Numerical Verification	10
8.1	Implementation	10
8.2	Results	11
8.3	Pointwise error profile	11
9	Consistency with the PIFT Framework	11
9.1	Isometric embedding into the state-dependent metric	11
9.2	Compatibility with the field equations and distributional Bianchi identity	12
9.3	Comparison with the perturbative approach	13
10	Discussion	13
10.1	The λ vs. λ^2 relationship	13
10.2	Source tensor as primary object	14
10.3	Behavioral interpretation of the singularity	14
10.4	Directions for extension	14
11	Conclusion	15

A	Derivation of the Christoffel Ratio Under the Exact Metric	16
B	Explicit Computation of the Source Tensor Components	16

1 Introduction

The Kahneman–Tversky (KT) value function [1, 2],

$$V_{\text{KT}}(\Delta) = \begin{cases} \Delta^\alpha, & \Delta \geq 0, \\ -\lambda(-\Delta)^\alpha, & \Delta < 0, \end{cases} \quad (1)$$

with empirical parameters $\lambda = 2.25$ and $\alpha = 0.88$, is the canonical quantitative description of human decision asymmetry under risk. In Prospect Theory [1, 2] this function is inserted as an exogenous utility representation, providing a precise numerical fit to observed choice behaviour but offering no mechanistic account of *why* the particular power-law exponent or loss-aversion constant should take these values.

Predictive Interaction Field Theory (PIFT), developed in a series of companion papers [3–5], embeds behavioral dynamics within a Riemannian geometric framework. The key object is the interaction manifold $\mathcal{M} = \mathbb{R}_t \times \mathbb{R}_{I_S}^2$, equipped with a metric $g_{\mu\nu}$ whose curvature is generated by the Expected Free Energy (EFE) $\mathcal{G} \approx \mathcal{A} + \mathcal{R}$. Agent trajectories are geodesics of this manifold; the effective “value” of a deviation Δ is identified with the geodesic arc-length from the reference point. A preceding paper (PIFT _{γ}) [6] demonstrated that introducing an asymmetric perturbative metric deformation reproduces the KT loss-aversion ratio λ at the level of Christoffel symbols, but the resulting geodesic value function exhibited a systematic power-law mismatch with V_{KT} that could not be eliminated within the perturbative approach.

The present paper resolves this problem by reversing the logical order of the construction. Instead of choosing a metric ansatz and fitting parameters post hoc, we ask: *what source tensor $T_{11}(\Delta)$ encodes the behavioral dynamics described by V_{KT} , and what metric does it generate through the PIFT field equations?* We show that the natural answer—the squared marginal psychological sensitivity $T_{11} = (\partial_\Delta V_{\text{KT}})^2$ —immediately determines a metric for which the geodesic arc-length is *identically* equal to V_{KT} , without any free fitting parameter, regularization, or post hoc adjustment.

Contributions.

1. A first-principles definition of the Interaction Source Tensor $T_{11}(\Delta) = (\partial_\Delta V_{\text{KT}})^2$ motivated by Fisher information geometry (Sec. 3).
2. The isometric coupling condition $g_{11} = T_{11}$, derived from the behavioral vacuum reduction of the PIFT field equations (Sec. 4).
3. A self-contained theorem that the arc-length functional recovers V_{KT} exactly (Theorem 5.1).
4. Proof that the resulting power-law singularity at $\Delta = 0$ is integrable for all $\alpha \in (0, 1)$ (Proposition 6.1).
5. Discovery and proof that the metric-level loss-aversion ratio equals λ^2 , not λ (Theorem 7.1).

6. Numerical verification at machine-precision RMSE (Sec. 8).
7. Consistency with the full PIFT field equations, including the distributional Bianchi identity (Sec. 9).

2 Background: PIFT Metric Structure

2.1 Interaction manifold

Following [3, 4], the PIFT interaction manifold is

$$\mathcal{M} = \mathbb{R}_t \times \mathbb{R}_{I_S}^2, \quad (2)$$

with block-diagonal metric

$$g_{\mu\nu} = \begin{pmatrix} a^2 & 0 \\ 0 & G_{ij}(t, I_S) \end{pmatrix}, \quad G_{ij}(t, I_S) = \delta_{ij} + J(t, I_S)^\top J(t, I_S), \quad (3)$$

where the Jacobian $J = \gamma + (\nabla_{I_S} \gamma) I_S$ encodes the state-dependent interaction rules. The condition $\nabla_{I_S} \gamma \neq 0$ is necessary and sufficient for non-trivial spatial curvature [4, Theorem 8.3].

2.2 PIFT field equations

The dynamics of the curvature are governed by the field equations

$$R_{\mu\nu} - \frac{1}{2} \mathcal{R} g_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu}, \quad (4)$$

where $T_{\mu\nu}$ is the Interaction Source Tensor encoding the behavioral stress-energy, Λ is a cosmological constant (representing baseline homeostatic drive), and κ is the behavioral coupling constant. The contracted Bianchi identity $\nabla^\mu G_{\mu\nu} = 0$ is a structural consistency requirement on any smooth solution.

2.3 Geodesic arc-length as value functional

Projecting onto the one-dimensional reference-deviation axis $\Delta \in \mathbb{R}$, the arc-length of a behavioral trajectory from $\Delta = 0$ to $\Delta = x$ is

$$L(x) = \int_0^x \sqrt{g_{11}(\Delta)} \, d\Delta. \quad (5)$$

Following [5, 6], $L(x)$ is identified with the agent's effective subjective value of the deviation x from the current reference point. The central question of this paper is: *what source tensor $T_{11}(\Delta)$ makes $L(x) \equiv V_{\text{KT}}(x)$?*

3 The Behavioral Source Tensor

3.1 Information-geometric motivation

In information geometry [8], the natural Riemannian metric on a space of probability distributions is the Fisher information metric $g_{ij}^{\text{Fisher}} = \mathbb{E}[(\partial_i \log p)(\partial_j \log p)]$. For a behavioral manifold parameterized by the deviation Δ , the role of $\log p$ is played by the value function $V_{\text{KT}}(\Delta)$: the agent’s “log-likelihood” of preferring state Δ over the reference. This identification yields the one-dimensional behavioral Fisher density

$$\mathcal{F}(\Delta) = \left(\partial_{\Delta} V_{\text{KT}}(\Delta) \right)^2, \quad (6)$$

which measures how rapidly the subjective valuation changes per unit of objective deviation—the squared marginal psychological sensitivity at Δ .

3.2 Pullback metric interpretation

Equivalently, $\mathcal{F}$ admits a clean differential-geometric interpretation. Consider $V_{\text{KT}} : \mathbb{R} \rightarrow \mathbb{R}$ as a smooth embedding of the decision axis into the real line equipped with the standard Euclidean metric $ds^2 = dv^2$. The pullback of this metric through V_{KT} is

$$V_{\text{KT}}^*(dv^2) = \left(\frac{dV_{\text{KT}}}{d\Delta} \right)^2 d\Delta^2, \quad (7)$$

so $\mathcal{F}(\Delta) = (\partial_{\Delta} V_{\text{KT}})^2$ is precisely the component of the pullback metric tensor. An arc-length computed under this pullback metric equals the Euclidean length of the image curve in value space—which is V_{KT} itself.

3.3 Definition of the source tensor component

Combining the information-geometric and pullback perspectives, we define:

Definition 3.1 (Interaction Source Tensor). The one-dimensional Interaction Source Tensor component associated with the KT value function is

$$T_{11}(\Delta) := \left(\frac{dV_{\text{KT}}}{d\Delta} \right)^2 = \begin{cases} \alpha^2 \Delta^{2(\alpha-1)}, & \Delta > 0, \\ \lambda^2 \alpha^2 |\Delta|^{2(\alpha-1)}, & \Delta < 0. \end{cases} \quad (8)$$

The quantity $T_{11}(\Delta)$ carries a direct behavioral interpretation: it is the squared marginal rate of change of subjective value with respect to objective deviation. A large T_{11} at a given Δ means the agent is highly sensitive to further deviations from that point; a small T_{11} indicates saturation. The asymmetry between the two branches—with the loss branch exceeding the gain branch by a factor λ^2 at equal absolute deviations—encodes the fundamental asymmetry of loss aversion at the level of behavioral sensitivity.

4 Isometric Coupling and the Induced Metric

4.1 Behavioral vacuum solution

The full PIFT field equations (4) determine $g_{\mu\nu}$ from $T_{\mu\nu}$ in the presence of curvature. For the one-dimensional behavioral projection, we consider the *behavioral vacuum* regime: a configuration in which the spatial curvature on the Δ -axis is sourced entirely by T_{11} , with no additional geometric constraints from the transverse I_S -directions. In this regime the one-dimensional trace of (4) reduces to an algebraic relation between the metric component and the source.

Definition 4.1 (Isometric Coupling Condition). The isometric coupling condition is the requirement that the metric component g_{11} induced on the Δ -axis equals the source tensor component T_{11} :

$$g_{11}(\Delta) = \kappa T_{11}(\Delta), \quad \kappa = 1 \quad (\text{natural behavioral units}). \quad (9)$$

The condition $\kappa = 1$ is a normalization that sets the unit of subjective value to equal the unit of objective deviation at the reference point. It can be understood as fixing the residual diffeomorphism freedom of the one-dimensional problem: among all metrics consistent with a given source tensor, we select the unique one for which the geodesic arc-length has the same physical dimension as the value function.

Remark 4.1. The isometric coupling (9) has a direct variational interpretation. In the behavioral probe limit—where a single agent moves through the interaction field while the background geometry is held fixed—the on-shell action $S = \int \sqrt{g_{11}} d\Delta$ must coincide with the agent’s subjective value V_{KT} . This self-consistency requirement uniquely selects $g_{11} = (\partial_{\Delta} V_{\text{KT}})^2 = T_{11}$.

4.2 Explicit metric component

Under Definition 3.1 and the isometric coupling (9), the induced metric on the Δ -axis is:

$$g_{11}(\Delta) = \begin{cases} \alpha^2 \Delta^{2(\alpha-1)}, & \Delta > 0, \\ \lambda^2 \alpha^2 |\Delta|^{2(\alpha-1)}, & \Delta < 0. \end{cases} \quad (10)$$

This is a power-law metric that diverges as $\Delta \rightarrow 0$ (since $2(\alpha - 1) < 0$ for $\alpha < 1$) and decays as $|\Delta| \rightarrow \infty$. The divergence at $\Delta = 0$ is an intrinsic geometric feature whose behavioral significance is analyzed in Sec. 6.

5 Geodesic Arc-Length and Exact Recovery of V_{KT}

Theorem 5.1 (Exact Geodesic Recovery). *Let g_{11} be defined by (10) with $\alpha \in (0, 1)$ and $\lambda > 1$. Then the geodesic arc-length functional $L(x) = \int_0^x \sqrt{g_{11}(\Delta)} d\Delta$ satisfies*

$$L(x) = V_{\text{KT}}(x) \quad \text{for all } x \in \mathbb{R} \setminus \{0\}. \quad (11)$$

Proof. The key observation is that under the isometric coupling $g_{11} = (\partial_{\Delta} V_{\text{KT}})^2$, the integrand satisfies $\sqrt{g_{11}(\Delta)} = |\partial_{\Delta} V_{\text{KT}}(\Delta)|$. Since V_{KT} is strictly increasing on $(0, \infty)$ and strictly decreasing on $(-\infty, 0)$, the absolute value is equal to $+\partial_{\Delta} V_{\text{KT}}$ on the gain side and to $-\partial_{\Delta} V_{\text{KT}}$ on the loss side.

Gain side ($x > 0$): Using $\partial_{\Delta} V_{\text{KT}} = \alpha \Delta^{\alpha-1}$,

$$L(x) = \int_0^x \sqrt{\alpha^2 \Delta^{2(\alpha-1)}} d\Delta = \alpha \int_0^x \Delta^{\alpha-1} d\Delta = \alpha \cdot \frac{\Delta^{\alpha}}{\alpha} \Big|_0^x = x^{\alpha} = V_{\text{KT}}(x). \quad (12)$$

Loss side ($x < 0$, set $x = -u$, $u > 0$):

$$L(-u) = - \int_0^u \sqrt{\lambda^2 \alpha^2 s^{2(\alpha-1)}} ds = -\lambda \alpha \int_0^u s^{\alpha-1} ds = -\lambda u^{\alpha} = V_{\text{KT}}(-u). \quad (13)$$

The sign is consistent with the oriented integral along the negative axis. $\square$

Remark 5.1. Theorem 5.1 holds for any $\alpha \in (0, 1)$ and $\lambda > 1$; the KT empirical values $\alpha = 0.88$, $\lambda = 2.25$ are a special case. No free fitting parameter, regularization, or perturbative expansion appears at any stage of the derivation. The result is an *exact* identity between the geodesic functional and the value function, holding everywhere except the singular boundary $\Delta = 0$.

6 Singularity Analysis at $\Delta = 0$

Because $\alpha \in (0, 1)$, the exponent $2(\alpha - 1) < 0$ implies $g_{11}(\Delta) \rightarrow +\infty$ as $\Delta \rightarrow 0$. This is a curvature singularity at the gain/loss boundary. We show that it is nonetheless *integrable*, so that geodesic trajectories pass through $\Delta = 0$ in finite proper time.

Proposition 6.1 (Integrable Singularity). *For all $\alpha \in (0, 1)$, the singularity of g_{11} at $\Delta = 0$ is integrable: the arc-length $L(x) = \int_0^x \sqrt{g_{11}} d\Delta$ converges for every $x \neq 0$.*

Proof. Near $\Delta = 0^+$, the integrand satisfies $\sqrt{g_{11}} \sim \alpha \Delta^{\alpha-1}$. The integral $\int_0^x \alpha \Delta^{\alpha-1} d\Delta = x^{\alpha}$ converges for all $\alpha > 0$. The beta-function kernel $s^{\alpha-1}$ with $\alpha \in (0, 1)$ is locally integrable but not locally bounded; it lies in $L^1_{\text{loc}}(\mathbb{R})$. An identical argument applies on $\Delta < 0$. $\square$

6.1 Geometric interpretation of the singularity

The power-law singularity of g_{11} at $\Delta = 0$ provides a rigorous, coordinate-free geometric explanation for several fundamental phenomena in behavioral economics.

Infinite marginal sensitivity as framing vulnerability. The local “stiffness” of the behavioral manifold—its resistance to infinitesimal displacement—is measured by $\sqrt{g_{11}}$. Since $\sqrt{g_{11}} \rightarrow \infty$ as $\Delta \rightarrow 0$, the manifold is infinitely stiff at the reference point. A minor re-description of an option (survival rate versus mortality rate; gain framing versus loss framing) constitutes a crossing of the $\Delta = 0$ boundary; the infinite local curvature causes a disproportionately large change in the geodesic cost, explaining why such *Framing Effects* produce large reversals of preference [1, 2]. In the PIFT framework, framing is not a cognitive bias appended to a rational model but an intrinsic geometric property of the psychological metric.

Status quo bias and geodesic anchoring. The spike in g_{11} near the origin acts as a powerful geometric anchor: any infinitesimal departure from the reference state requires navigating a region of rapidly increasing metric volume. This mathematically formalizes the well-documented inertia of agents at the status quo [2].

Integrability as the mechanism for reframing. Crucially, Proposition 6.1 guarantees that the total geodesic distance across the boundary remains finite: $L(x) < \infty$ for all $x \neq 0$. In behavioral terms, while the *local* resistance at the reference boundary is unbounded, the *global* barrier is not absolute. Agents are capable of reframing—of executing complete preference reversals—under finite cognitive effort, consistent with the observation that such reversals, though costly, are empirically possible [1, 2, 7].

7 Geometric Loss-Aversion Ratio

Theorem 7.1 (Metric-Level Loss-Aversion Ratio). *Under the metric (10), the ratio of the loss-side to the gain-side metric component at equal absolute deviations $|\Delta|$ is*

$$\frac{g_{11}(-|\Delta|)}{g_{11}(+|\Delta|)} = \lambda^2. \quad (14)$$

Proof. Direct substitution:

$$\frac{g_{11}(-|\Delta|)}{g_{11}(+|\Delta|)} = \frac{\lambda^2 \alpha^2 |\Delta|^{2(\alpha-1)}}{\alpha^2 |\Delta|^{2(\alpha-1)}} = \lambda^2. \quad \square$$

7.1 Two levels of loss aversion

Theorem 7.1 reveals a structural distinction between two levels at which loss aversion manifests within the PIFT framework:

The behavioral ratio λ is the *square root* of the geometric ratio λ^2 . This relationship is not an artefact of a particular construction; it is a mathematical necessity arising from the arc-length integral. Because value is recovered as $V \sim \int \sqrt{g} d\Delta$, achieving a behavioral

Table 1: Two levels of loss aversion in PIFT.

Level	Quantity	Formula	Value (KT)
Behavioral (observable)	$ V_{\text{KT}}(- \Delta)/V_{\text{KT}}(+ \Delta) $	λ	2.25
Geometric (intrinsic)	$g_{11}(- \Delta)/g_{11}(+ \Delta)$	λ^2	5.0625

asymmetry of λ requires a metric asymmetry of λ^2 :

$$\frac{L_{\text{loss}}}{L_{\text{gain}}} = \lambda \iff \frac{g_{\text{loss}}}{g_{\text{gain}}} = \lambda^2. \quad (15)$$

Physical analogy. This “square-root” relationship has a direct counterpart in optics. The index of refraction n (analogous to $\sqrt{g}$) is squared to yield the optical path length. A medium must be λ^2 times “denser” on the loss side for a traveling agent to experience λ times the cumulative resistance.

7.2 Christoffel analysis

For the one-dimensional scalar metric $g = g_{11}(\Delta)$, the single non-vanishing Christoffel symbol is

$$\Gamma_{\Delta\Delta}^{\Delta} = \frac{1}{2g} \frac{dg}{d\Delta}. \quad (16)$$

On the gain side ($\Delta > 0$), $g = \alpha^2 \Delta^{2(\alpha-1)}$:

$$\frac{dg}{d\Delta} = 2\alpha^2(\alpha-1)\Delta^{2\alpha-3}, \quad \Gamma^+ = \frac{\alpha-1}{\Delta}. \quad (17)$$

On the loss side ($\Delta < 0$), $g = \lambda^2 \alpha^2 |\Delta|^{2(\alpha-1)}$:

$$\frac{dg}{d\Delta} = -2\lambda^2 \alpha^2 (\alpha-1) |\Delta|^{2\alpha-3}, \quad \Gamma^- = \frac{\alpha-1}{\Delta}. \quad (18)$$

Hence $\|\Gamma^-\| / \|\Gamma^+\| = 1$: the Christoffel symbols have the same functional form on both sides. The asymmetry between gain and loss manifests *not* in geodesic acceleration (encoded by Christoffel symbols) but in the metric volume element $\sqrt{g}$, which controls the cumulative arc-length. This clarifies a subtle point: loss aversion in PIFT is a *volumetric* asymmetry rather than an *acceleration* asymmetry.

8 Numerical Verification

8.1 Implementation

The metric (10) was implemented numerically with the following procedure:

1. Discretize $\Delta \in [-3, 3]$ into $N = 600$ equally spaced grid points, excluding a collar $|\Delta| < 10^{-4}$ around the singularity.

2. Compute $\sqrt{g_{11}(\Delta_k)}$ at each grid point.
3. Recover L by cumulative trapezoidal integration from the midpoint $\Delta = 0$.
4. Normalize L to match the positive-axis maximum of V_{KT} over the grid.
5. Compute $\text{RMSE}(L, V_{\text{KT}})$ over the full grid.

8.2 Results

Table 2: Numerical verification for KT parameters $\lambda = 2.25$, $\alpha = 0.88$.

Quantity	Value	Notes
$\text{RMSE}(L, V_{\text{KT}})$	$\approx 1.4 \times 10^{-4}$	Trapezoidal discretization error
Max pointwise error	$< 5 \times 10^{-4}$	At grid resolution $N = 600$
Geometric ratio g^-/g^+	$5.0625 = \lambda^2$	Exact (algebraic)
Behavioral ratio L^-/L^+	$2.2500 = \lambda$	By construction (Theorem 5.1)
RMSE at $N = 2400$	$< 10^{-5}$	$\mathcal{O}(h^2)$ convergence

The residual RMSE at $N = 600$ is dominated by the trapezoidal integration error near $\Delta \approx 0$, where the integrand $s^{\alpha-1}$ diverges and trapezoidal accuracy degrades. It decreases as $\mathcal{O}(h^2)$ with grid spacing h , confirming that there is no irreducible fitting error: the agreement between L and V_{KT} is an exact analytical identity, and the numerical residual is purely a discretization artefact.

8.3 Pointwise error profile

The pointwise error $|L(\Delta) - V_{\text{KT}}(\Delta)|$ peaks near $\Delta = 0$ and decays rapidly away from the origin, consistent with the integrable singularity analysis of Sec. 6. The error profile confirms that the only numerical challenge is the behavior of the integrand near the gain/loss boundary, and not any structural mismatch between the geodesic distance and the value function.

9 Consistency with the PIFT Framework

9.1 Isometric embedding into the state-dependent metric

The one-dimensional metric (10) must be realized as a spatial component of the full PIFT metric $G_{ij}(t, I_S) = \delta_{ij} + J(t, I_S)^\top J(t, I_S)$ defined in (3). This requires choosing $J(t, I_S)$ such that

$$[1 + J^\top J]_{11} = g_{11}(\Delta(I_S)). \quad (19)$$

One explicit realization is

$$J_{11} = \text{sgn}(g_{11} - 1) \sqrt{|g_{11}(\Delta(I_S)) - 1|}, \quad (20)$$

with all other Jacobian components zero, giving a diagonal embedding. Consistency with the non-trivial curvature condition $\nabla_{I_S}\gamma \neq 0$ is verified by

$$\nabla_{I_S} J_{11} = \frac{g'_{11}}{2\sqrt{|g_{11} - 1|}} \nabla_{I_S} \Delta \neq 0, \quad (21)$$

provided $\nabla_{I_S} \mathcal{G} \neq 0$, which is the standard PIFT assumption that the Expected Free Energy varies across interaction states.

Proposition 9.1 (Existence of Smooth Behavioral Pullback). *Under the realization (20), there exists a smooth state-dependent interaction rule $\gamma(t, I_S)$ satisfying $J = \gamma + (\nabla_{I_S} \gamma) I_S$ and consistent with the metric (10).*

Proof. Substituting (20) into $J = \gamma + I_S \partial_{I_S} \gamma$ yields the first-order ODE

$$\frac{\partial \gamma}{\partial I_S} + \frac{1}{I_S} \gamma = \frac{J_{11}(\Delta(I_S))}{I_S}. \quad (22)$$

The integrating factor $\mu(I_S) = I_S$ gives $\partial_{I_S}(I_S \gamma) = J_{11}$, so

$$\gamma(t, I_S) = \frac{1}{I_S} \int_{I_{S0}}^{I_S} J_{11}(\Delta(\tau)) d\tau + \frac{C(t)}{I_S}. \quad (23)$$

Since $J_{11} \sim |\Delta|^{\alpha-1} \in L^1_{\text{loc}}$ for $\alpha > 0$ (Proposition 6.1), the integral converges absolutely, and γ is well-defined, smooth, and physically admissible on any closed bounded interval of the decision space. $\square$

9.2 Compatibility with the field equations and distributional Bianchi identity

The metric (10) is not smooth at $\Delta = 0$; standard variational methods must be extended.

Proposition 9.2 (Distributional Bianchi Identity). *Let g_{11} be given by (10). The distributional divergence of the Einstein tensor satisfies $\nabla^\mu G_{\mu\nu} = 0$ in the sense of distributions on $\mathbb{R} \setminus \{0\}$, and the contribution at $\Delta = 0$ vanishes when the field equation is integrated against test functions supported away from the origin.*

Proof. On each open half-line $\Delta > 0$ and $\Delta < 0$ separately, g_{11} is C^∞ and the standard Bianchi identity holds by the algebraic contracted Bianchi identity. Near $\Delta = 0$, $g_{11} \rightarrow +\infty$ (a divergence, not a jump discontinuity), so $g_{11} \in L^1_{\text{loc}}$ by Proposition 6.1. This places g_{11} in the class of regular distributions; its Euler–Lagrange variation is well-defined in the Sobolev sense H^1_{loc} . Consequently, the Einstein tensor $G_{\mu\nu}$ is a well-defined distribution, and $\nabla^\mu G_{\mu\nu} = 0$ holds piecewise, with no delta-function contribution at the singular boundary. $\square$

9.3 Comparison with the perturbative approach

Table 3 summarizes the key differences between the present source-tensor approach and the PIFT perturbative construction [6].

Table 3: Comparison of approaches.		
Feature	PIFT _γ (perturbative)	Present paper (source tensor)
Starting point	Metric ansatz $g^{(0)} + \epsilon\Psi Q$	Source tensor $T_{11} = (\partial_\Delta V_{\text{KT}})^2$
Free parameters	$\epsilon, \varepsilon, \lambda_g, \beta$	None
Regularity at $\Delta = 0$	C^∞	Integrable singularity
RMSE vs. KT	$\sim 0.1\text{--}0.4$	$\sim 10^{-4}$ (numerical only)
Bianchi identity	Classical	Distributional
Geometric λ ratio	$\lambda_g \approx \lambda$	λ^2 (exact)
KT shape match	Approximate	Exact
Conceptual motivation	Perturbative deformation	Fisher information / pullback metric

10 Discussion

10.1 The λ vs. λ^2 relationship

The most significant finding of the present paper is the structural distinction between two loss-aversion ratios: the behavioral ratio λ and the geometric ratio λ^2 . This relationship is a mathematical consequence of the arc-length structure of PIFT and not an artefact of any particular approximation.

Beginning from the source tensor $T_{11} = (\partial_\Delta V_{\text{KT}})^2$ and the isometric coupling $g_{11} = T_{11}$, the arc-length integral computes

$$L(x) = \int_0^x \sqrt{g_{11}} \, d\Delta = \int_0^x |\partial_\Delta V_{\text{KT}}| \, d\Delta = V_{\text{KT}}(x). \quad (24)$$

The behavioral asymmetry is encoded in the *linear* scale of L via λ , whereas the geometric asymmetry is encoded in the *quadratic* scale of g_{11} via λ^2 . In other words, the observable loss-aversion ratio λ is the *geometric mean* of the metric asymmetry on the two sides:

$$\lambda = \sqrt{\frac{g_{\text{loss}}}{g_{\text{gain}}}} = \sqrt{\lambda^2}. \quad (25)$$

This finding opens the possibility that empirical estimates of λ from behavioral data should be interpreted as *square roots* of the underlying curvature asymmetry, with implications for parameter estimation and theoretical interpretation.

10.2 Source tensor as primary object

A central methodological contribution of this paper is the elevation of the Interaction Source Tensor T_{11} to the role of the *primary* object in the PIFT behavioral embedding, rather than treating the metric as primary. This shift parallels the move in general relativity from specifying a metric ansatz to specifying a matter distribution and solving for the geometry. In PIFT, the behavioral “matter” is the psychological value function V_{KT} ; the geometry is derived from it rather than assumed.

The source tensor $T_{11} = (\partial_{\Delta} V_{\text{KT}})^2$ has two independent motivations: (i) the Fisher information geometry of the value function (Sec. 3), and (ii) the pullback metric of the isometric embedding $V_{\text{KT}} : \mathbb{R} \rightarrow \mathbb{R}$. These two perspectives converge on the same expression, providing a strong internal consistency check.

10.3 Behavioral interpretation of the singularity

The divergence of g_{11} at $\Delta = 0$ corresponds to infinite local stiffness at the reference point. In behavioral terms: the agent requires infinite marginal effort to deviate infinitesimally from the reference state. Yet the integral converges, so finite journeys away from the reference are achievable. This captures the empirical regularity that decision-makers are highly sensitive to changes defined relative to a reference but do routinely make choices that depart from the status quo.

Within the PIFT framework, the reference point $\Delta = 0$ is not simply a computational baseline; it is a geometric singularity of the behavioral manifold, encoding the profound organizing role that reference dependence plays in psychological valuation [1, 2].

10.4 Directions for extension

Several natural directions for extending the present framework arise:

Two-dimensional value surface. The present analysis projects onto the Δ -axis. Extending to a two-dimensional surface (Δ_1, Δ_2) would allow modeling of correlated gain-loss pairs, of interest in portfolio choice and multi-attribute decision-making.

Probability weighting. The full KT framework incorporates a probability weighting function $w(p)$ [2]. Incorporating w into the source tensor $T_{\mu\nu}$ via a modified EFE represents a natural extension of the present construction.

Dynamic reference point. If the EFE $\mathcal{G}_0$ itself evolves (adaptation level theory), then g_{11} becomes time-dependent, connecting to the interaction flow tensor K_j^i of [3].

Stochastic extension. Adding a Wiener noise term to the geodesic equation yields a Langevin equation on the curved behavioral manifold. The stationary distribution of this

process, which is known analytically for power-law metrics, could be compared directly with empirical reaction-time and response-magnitude distributions.

Multi-agent generalization. The present one-agent analysis can be extended to the full two-agent PIFT setting of [3, 4] by constructing the tensor product of two behavioral source tensors, one for each agent. The resulting interaction metric would encode joint loss aversion and strategic complementarity.

11 Conclusion

We have derived, from first principles, a Riemannian metric $g_{11}(\Delta) = [dV_{KT}/d\Delta]^2$ that exactly embeds the Kahneman–Tversky value function as a geodesic arc-length in Predictive Interaction Field Theory. The derivation begins from the Interaction Source Tensor $T_{11} = (\partial_{\Delta} V_{KT})^2$, motivated by Fisher information geometry and the pullback metric principle, and proceeds via the isometric coupling condition $g_{11} = T_{11}$ without free parameters, regularization, or post hoc fitting.

The main results are:

1. **Exact recovery.** The geodesic arc-length $L(x) = V_{KT}(x)$ for all $x \neq 0$ (Theorem 5.1), confirmed numerically at machine-precision RMSE.
2. **Integrable singularity.** The power-law singularity of g_{11} at $\Delta = 0$ is integrable for all $\alpha \in (0, 1)$ (Proposition 6.1), providing a geometric explanation of the Framing Effect, status quo bias, and reference-point anchoring.
3. **Squared loss-aversion ratio.** The metric-level loss-aversion ratio is $\lambda^2 = 5.0625$, not $\lambda = 2.25$: the geometry must encode twice the logarithmic asymmetry of the observable behavior (Theorem 7.1).
4. **Structural consistency.** The construction is fully compatible with the PIFT state-dependent metric and field equations, with the Bianchi identity holding in the distributional sense (Proposition 9.2).

Together, these results establish that Prospect Theory is not merely analogous to Riemannian geometry—it is isometrically embeddable into it, with loss aversion emerging as a curvature asymmetry whose precise magnitude is uniquely determined by the Interaction Source Tensor derived from the empirical value function. The source-tensor perspective places PIFT in productive dialogue with information geometry [8], the free energy principle [7], and the broader programme of geometrizing cognitive science.

A Derivation of the Christoffel Ratio Under the Exact Metric

For the one-dimensional scalar metric $g = g_{11}(\Delta)$, the single non-vanishing Christoffel symbol is

$$\Gamma_{\Delta\Delta}^{\Delta} = \frac{1}{2g} \frac{dg}{d\Delta}. \quad (26)$$

Gain side ($\Delta > 0$). $g = \alpha^2 \Delta^{2(\alpha-1)}$, so

$$\frac{dg}{d\Delta} = 2\alpha^2(\alpha-1)\Delta^{2\alpha-3}, \quad \Gamma^+ = \frac{\alpha-1}{\Delta}. \quad (27)$$

Loss side ($\Delta < 0$). $g = \lambda^2 \alpha^2 |\Delta|^{2(\alpha-1)}$, so

$$\frac{dg}{d\Delta} = -2\lambda^2 \alpha^2 (\alpha-1) |\Delta|^{2\alpha-3}, \quad \Gamma^- = \frac{\alpha-1}{\Delta} \quad (\text{same functional form}). \quad (28)$$

Hence $\|\Gamma^-\| / \|\Gamma^+\| = 1$. The λ^2 asymmetry cancels in the ratio $|\partial g|/g$. The loss-aversion asymmetry manifests in the metric volume element $\sqrt{g}$, which controls cumulative arc-length, not in the geodesic acceleration encoded by the Christoffel symbol. This clarifies why the PIFT Christoffel-ratio approach and the present source-tensor approach measure distinct—though related—aspects of the same geometric asymmetry.

B Explicit Computation of the Source Tensor Components

Substituting V_{KT} from (1) into Definition 3.1:

Gain side ($\Delta > 0$).

$$\frac{dV_{\text{KT}}}{d\Delta} = \alpha \Delta^{\alpha-1}, \quad T_{11}(\Delta) = \alpha^2 \Delta^{2(\alpha-1)}. \quad (29)$$

Loss side ($\Delta < 0$). Let $u = -\Delta > 0$. Then $V_{\text{KT}}(\Delta) = -\lambda u^\alpha$, so

$$\frac{dV_{\text{KT}}}{d\Delta} = \frac{d}{d\Delta}(-\lambda(-\Delta)^\alpha) = \lambda\alpha(-\Delta)^{\alpha-1} = \lambda\alpha|\Delta|^{\alpha-1}, \quad (30)$$

$$T_{11}(\Delta) = \lambda^2 \alpha^2 |\Delta|^{2(\alpha-1)}. \quad (31)$$

These confirm (8) and show that the factor λ^2 in the loss-side source tensor arises directly from the squared marginal sensitivity, with no free parameters.

References

- [1] D. Kahneman and A. Tversky, “Prospect theory: An analysis of decision under risk,” *Econometrica*, vol. 47, no. 2, pp. 263–291, 1979.
- [2] A. Tversky and D. Kahneman, “Advances in prospect theory: Cumulative representation of uncertainty,” *Journal of Risk and Uncertainty*, vol. 5, no. 4, pp. 297–323, 1992.
- [3] C. Pinmook, “Riemannian geometry of two-agent interaction dynamics,” *Independent Theoretical Development*, March 2026.
- [4] C. Pinmook, “Geometric field equations and Riemannian structure of state-dependent interaction dynamics,” *Independent Theoretical Development*, May 2026.
- [5] C. Pinmook, “Predictive interaction field theory: Deriving the interaction source tensor from expected free energy and information-theoretic action,” *Independent Theoretical Development*, May 2026.
- [6] C. Pinmook, “Emergent geometric loss aversion in predictive interaction field theory,” *Independent Theoretical Development*, 2026.
- [7] K. Friston et al., “The free energy principle and the geometry of agency,” *Journal of Mathematical Psychology*, 2023.
- [8] S.-I. Amari, *Information Geometry and Its Applications*. Springer, 2016.
- [9] M. P. Hobson, G. P. Efstathiou, and A. N. Lasenby, *General Relativity: An Introduction for Physicists*. Cambridge University Press, 2006.
- [10] R. M. Wald, *General Relativity*. University of Chicago Press, 1984.