



ENGINEERING MATHEMATICS - I

Taylor's and Maclaurin's Series

Dr. Anitha
Science and Humanities

ENGINEERING MATHEMATICS - I

UNIT 2 : Partial Differentiation

Session : 8

Sub Topic : Taylor's and Maclaurin's Series of a function of Two Variables

Dr. Anitha

Department of Science and Humanities

ENGINEERING MATHEMATICS - I

Intuition behind Taylor's and Maclaurin's Series

- Suppose you want to calculate the value of $e^{\tan(\sqrt[3]{3})}$, the first thing you look for is the calculator.
- A calculator expresses any complicated function in terms of its maclaurin's series and find its value.
- Suppose we need to evaluate an integral of the form $\int e^{x^2} dx$ or $\int \cos(x^2) dx$ all the methods we know so far fails.
- If you want to evaluate the following limit

$$\lim_{x \rightarrow 0} \frac{\sin(x) - x}{x^4}$$



Taylor's and Maclaurin's Series for a function of single variable

If a function $f(x)$ has continuous derivatives of order $(n+1)$, then this function can be expressed as a polynomial at a point $x=a$ in the form

$$f(x)$$

$$= f(a) + (x - a)f'(a) + \frac{(x - a)^2}{2!} f''(a) + \dots + \frac{(x - a)^n}{n!} f^n(a) + \dots$$

If the function is expanded about the origin we get Maclaurin's Series

$$f(x) = f(0) + (x)f'(0) + \frac{(x)^2}{2!} f''(0) + \dots + \frac{(x)^n}{n!} f^n(0) + \dots$$

ENGINEERING MATHEMATICS - I

Standard Maclaurin's Series for a function of single variable



1. $(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3 + \dots, |x| < 1$
2. $e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots, x \in R$
3. $\log_e(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots, |x| < 1$
4. $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$
5. $\cos x = 1 - \frac{x^2}{2} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$
6. $\sinh x = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots$
7. $\cosh x = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots$
8. $\tan^{-1} x = x - \frac{x^3}{3!} + \frac{x^5}{5!} + \dots$
9. $\sin^{-1} x = x + \frac{1}{2} \frac{x^3}{3} + \frac{1 \cdot 3}{2 \cdot 4} \frac{x^5}{5} + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \frac{x^7}{7} + \dots, -1 \leq x \leq 1$
10. $\cos^{-1} x = \frac{\pi}{2} - x - \frac{1}{2} \frac{x^3}{3} - \frac{1 \cdot 3}{2 \cdot 4} \frac{x^5}{5} - \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \frac{x^7}{7} - \dots, -1 \leq x \leq 1$

ENGINEERING MATHEMATICS - I

Applications of Taylor's and Maclaurin's Series for a function of single variable



1. Expressing any complicated function in terms of a polynomial
2. Evaluating definite integrals which cannot be evaluated by any other methods.
3. Understanding asymptotic behavior of the function
4. Understanding the growth of function
5. Solving differential equations

ENGINEERING MATHEMATICS - I

Applications of Taylor's and Maclaurin's Series for a function of single variable




6. In physics, Maclaurin's series is used to approximate the Lorentz factor in special relativity.
7. The most famous equation $E = mc^2$ is an approximation for low velocities.
8. In the motion of a pendulum an approximation of $\sin(\theta) \cong \theta$ which comes from Taylor's series.
9. Used in power flow analysis of electrical power system.
10. Can be used to find the Generating functions.

Source: <https://math.stackexchange.com/questions/218421/what-are-the-practical-applications-of-the-taylor-series>



ENGINEERING MATHEMATICS - I

Applications of Taylor's and Maclaurin's Series for a function of single variable

$n=4$



$n=6$



Taylor's expansion of a function of two variables

Statement:

Taylor's series of $f(x, y)$ about (a, b) is given by

$$\begin{aligned} f(x, y) &= f(a, b) + \frac{1}{1!} \{ (x - a)f_x(a, b) + (y - b)f_y(a, b) \} \\ &+ \frac{1}{2!} \{ (x - a)^2 f_{xx}(a, b) + 2(x - a)(y - b)f_{xy}(a, b) \end{aligned}$$

Statement:

Maclaurin's series for a function of two variables is given by

$$f(x, y) = f(0,0) + \frac{1}{1!} \{x f_x(0,0) + y f_y(0,0)\} + \frac{1}{2!} \{x^2 f_{xx}(0,0) + 2xy f_{xy}(0,0) + y^2 f_{yy}(0,0)\} + \dots$$

Problems on Taylor's and Maclaurin's expansion




Figure 3: Graph of $f(x, y) = xe^y + 1$ and its 1st-degree Taylor polynomial, $L(x, y) = 1 + x + y$




Figure 4: Graph of $f(x, y) = xe^y + 1$ and its 2nd-degree Taylor polynomial, $L(x, y) = 1 + x + xy + \frac{y^2}{2}$

ENGINEERING MATHEMATICS - I

Applications of Euler's Theorem of a multivariate function

1. Taylor's series is used in different optimization techniques by expressing it as a linear or quadratic expression.
2. Distances can be measured accurately using sine law and Taylor's series.
3. To calculate variance of complicated configurations
4. To visualize complicated functions.





Dr. Anitha

Department of Science and Humanities

nanitha@pes.edu

Extn 730