



ENGINEERING MATHEMATICS - I

APARNA B. S

Department of Science and Humanities

Class Content

- **Series of Positive terms**
- **Conditions for convergence of Series of Positive terms**
- **Comparison test**
- **Limit form of comparison test**
- **P Series (statement)**
- **Examples**

Series of Positive terms

If all the terms of the series $\sum a_n$ are positive i.e., $a_n > 0$, $\forall n$, then the series $\sum a_n$ is called a series of positive terms.

Results on Series of Positive terms:

1. If c is a constant and $\sum a_n$ converges to l , then $\sum (ca_n)$ converges to cl .
2. If $\sum_{n=1}^{n=\infty} a_n$ and $\sum_{n=1}^{n=\infty} b_n$ converges to l and m respectively then
 - a. $\sum (a_n + b_n)$ converges to $(l + m)$.
 - b. $\sum (a_n - b_n)$ converges to $(l - m)$.

Series of Positive terms

If all the terms of the series $\sum a_n$ are positive i.e., $a_n > 0$, $\forall n$, then the series $\sum a_n$ is called a series of positive terms.

Results on Series of Positive terms:

1. If c is a constant and $\sum a_n$ converges to l , then $\sum (ca_n)$ converges to cl .
2. If $\sum_{n=1}^{n=\infty} a_n$ and $\sum_{n=1}^{n=\infty} b_n$ converges to l and m respectively then
 - a. $\sum (a_n + b_n)$ converges to $(l + m)$.
 - b. $\sum (a_n - b_n)$ converges to $(l - m)$.

Series of Positive terms

If all the terms of the series $\sum a_n$ are positive i.e., $a_n > 0$, $\forall n$, then the series $\sum a_n$ is called a series of positive terms.

Results on Series of Positive terms:

1. If c is a constant and $\sum a_n$ converges to l , then $\sum (ca_n)$ converges to cl .
2. If $\sum_{n=1}^{n=\infty} a_n$ and $\sum_{n=1}^{n=\infty} b_n$ converges to l and m respectively then
 - a. $\sum (a_n + b_n)$ converges to $(l + m)$.
 - b. $\sum (a_n - b_n)$ converges to $(l - m)$.

Series of Positive terms

If all the terms of the series $\sum a_n$ are positive i.e., $a_n > 0, \forall n$, then the series $\sum a_n$ is called a series of positive terms.

Results on Series of Positive terms:

1. If c is a constant and $\sum a_n$ converges to l , then $\sum (ca_n)$ converges to cl .
2. If $\sum_{n=1}^{n=\infty} a_n$ and $\sum_{n=1}^{n=\infty} b_n$ converges to l and m respectively then
 - a. $\sum (a_n + b_n)$ converges to $(l + m)$.
 - b. $\sum (a_n - b_n)$ converges to $(l - m)$.

Conditions for Convergence of a Series of positive terms

Necessary condition for the convergence of a series of positive terms :

If a series of positive terms $\sum a_n$ is convergent then $\lim_{n \rightarrow \infty} a_n = 0$

REMARK

- 1. Converse of the above statement is not true while the contrapositive of the statement is true. i.e if $\lim_{n \rightarrow \infty} a_n \neq 0$, then the series of positive terms is not convergent.*
- 2. The nature of a series is not altered by adding a finite number of terms to the series or by removing a finite terms from the beginning of the series .*

Conditions for Convergence of a Series of positive terms

Necessary condition for the convergence of a series of positive terms :

If a series of positive terms $\sum a_n$ is convergent then $\lim_{n \rightarrow \infty} a_n = 0$

REMARK

- 1. Converse of the above statement is not true while the contrapositive of the statement is true. i.e if $\lim_{n \rightarrow \infty} a_n \neq 0$, then the series of positive terms is not convergent.*
- 2. The nature of a series is not altered by adding a finite number of terms to the series or by removing a finite terms from the beginning of the series .*

Conditions for Convergence of a Series of positive terms

Necessary condition for the convergence of a series of positive terms :

If a series of positive terms $\sum a_n$ is convergent then $\lim_{n \rightarrow \infty} a_n = 0$

REMARK

- 1. Converse of the above statement is not true while the contrapositive of the statement is true. i.e if $\lim_{n \rightarrow \infty} a_n \neq 0$, then the series of positive terms is not convergent.*
- 2. The nature of a series is not altered by adding a finite number of terms to the series or by removing a finite terms from the beginning of the series .*

Conditions for Convergence of a Series of positive terms

Necessary condition for the convergence of a series of positive terms :

If a series of positive terms $\sum a_n$ is convergent then $\lim_{n \rightarrow \infty} a_n = 0$

REMARK

- 1. Converse of the above statement is not true while the contrapositive of the statement is true. i.e if $\lim_{n \rightarrow \infty} a_n \neq 0$, then the series of positive terms is not convergent.*
- 2. The nature of a series is not altered by adding a finite number of terms to the series or by removing a finite terms from the beginning of the series .*

Comparison test

TEST (Comparison Test)

Suppose that $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are two series of positive terms i.e $a_n \geq 0$ and $b_n \geq 0$. Then,

1) If $\sum_{n=1}^{\infty} b_n$ is convergent and $a_n \leq k b_n$ for all n ,

where k is a positive constant,

$\Rightarrow \sum_{n=1}^{\infty} a_n$ is convergent.

2) If $\sum_{n=1}^{\infty} b_n$ is divergent and $a_n \geq k b_n$ for all n ,

where k is a positive constant

$\Rightarrow \sum_{n=1}^{\infty} a_n$ is divergent.

Comparison test

TEST (Comparison Test)

Suppose that $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are two series of positive terms i.e $a_n \geq 0$ and $b_n \geq 0$. Then,

1) If $\sum_{n=1}^{\infty} b_n$ is convergent and $a_n \leq k b_n$ for all n ,

where k is a positive constant,

$\Rightarrow \sum_{n=1}^{\infty} a_n$ is convergent.

2) If $\sum_{n=1}^{\infty} b_n$ is divergent and $a_n \geq k b_n$ for all n ,

where k is a positive constant

$\Rightarrow \sum_{n=1}^{\infty} a_n$ is divergent.

Comparison test

TEST (Comparison Test)

Suppose that $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are two series of positive terms i.e $a_n \geq 0$ and $b_n \geq 0$. Then,

1) If $\sum_{n=1}^{\infty} b_n$ is convergent and $a_n \leq k b_n$ for all n ,

where k is a positive constant,

$\Rightarrow \sum_{n=1}^{\infty} a_n$ is convergent.

2) If $\sum_{n=1}^{\infty} b_n$ is divergent and $a_n \geq k b_n$ for all n ,

where k is a positive constant

$\Rightarrow \sum_{n=1}^{\infty} a_n$ is divergent.

Comparison test

TEST (Comparison Test)

Suppose that $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are two series of positive terms i.e $a_n \geq 0$ and $b_n \geq 0$. Then,

1) If $\sum_{n=1}^{\infty} b_n$ is convergent and $a_n \leq k b_n$ for all n ,

where k is a positive constant,

$\Rightarrow \sum_{n=1}^{\infty} a_n$ is convergent.

2) If $\sum_{n=1}^{\infty} b_n$ is divergent and $a_n \geq k b_n$ for all n ,

where k is a positive constant

$\Rightarrow \sum_{n=1}^{\infty} a_n$ is divergent.

Comparison test

REMARK

This test is a useful to justify whether a given series, is convergent or divergent, whenever the nature of another series is known.

Comparison test

REMARK

This test is a useful to justify whether a given series, is convergent or divergent, whenever the nature of another series is known.

Comparison test

REMARK

This test is a useful to justify whether a given series, is convergent or divergent, whenever the nature of another series is known.

Comparison test

REMARK

This test is a useful to justify whether a given series, is convergent or divergent, whenever the nature of another series is known.

Limit Comparison Test

TEST (Limit Comparison Test)

Suppose that $\sum_{i=1}^{\infty} a_i$ and $\sum_{i=1}^{\infty} b_i$ are series with all terms positive. 1)

$\lim_{i \rightarrow \infty} \frac{a_i}{b_i} = c > 0 \implies \sum_{i=1}^{\infty} a_i$ and $\sum_{i=1}^{\infty} b_i$ either both converge, or both diverge.

2) $\lim_{i \rightarrow \infty} \frac{a_i}{b_i} = 0$ and $\sum_{i=1}^{\infty} a_i$ converges
 \implies the series $\sum_{i=1}^{\infty} b_i$ converges

3) $\lim_{i \rightarrow \infty} \frac{a_i}{b_i} = \infty$ and $\sum_{i=1}^{\infty} b_i$ diverges
 \implies the series $\sum_{i=1}^{\infty} a_i$ diverges.

Limit Comparison Test

TEST (Limit Comparison Test)

Suppose that $\sum_{i=1}^{\infty} a_i$ and $\sum_{i=1}^{\infty} b_i$ are series with all terms positive. 1)

$\lim_{i \rightarrow \infty} \frac{a_i}{b_i} = c > 0 \implies \sum_{i=1}^{\infty} a_i$ and $\sum_{i=1}^{\infty} b_i$ either both converge, or both diverge.

2) $\lim_{i \rightarrow \infty} \frac{a_i}{b_i} = 0$ and $\sum_{i=1}^{\infty} a_i$ converges
 \implies the series $\sum_{i=1}^{\infty} b_i$ converges

3) $\lim_{i \rightarrow \infty} \frac{a_i}{b_i} = \infty$ and $\sum_{i=1}^{\infty} b_i$ diverges
 \implies the series $\sum_{i=1}^{\infty} a_i$ diverges.

Limit Comparison Test

TEST (Limit Comparison Test)

Suppose that $\sum_{i=1}^{\infty} a_i$ and $\sum_{i=1}^{\infty} b_i$ are series with all terms positive. 1)

$\lim_{i \rightarrow \infty} \frac{a_i}{b_i} = c > 0 \implies \sum_{i=1}^{\infty} a_i$ and $\sum_{i=1}^{\infty} b_i$ either both converge, or both diverge.

2) $\lim_{i \rightarrow \infty} \frac{a_i}{b_i} = 0$ and $\sum_{i=1}^{\infty} a_i$ converges
 \implies the series $\sum_{i=1}^{\infty} b_i$ converges

3) $\lim_{i \rightarrow \infty} \frac{a_i}{b_i} = \infty$ and $\sum_{i=1}^{\infty} b_i$ diverges
 \implies the series $\sum_{i=1}^{\infty} a_i$ diverges.

Limit Comparison Test

TEST (Limit Comparison Test)

Suppose that $\sum_{i=1}^{\infty} a_i$ and $\sum_{i=1}^{\infty} b_i$ are series with all terms positive. 1)

$\lim_{i \rightarrow \infty} \frac{a_i}{b_i} = c > 0 \implies \sum_{i=1}^{\infty} a_i$ and $\sum_{i=1}^{\infty} b_i$ either both converge, or both diverge.

2) $\lim_{i \rightarrow \infty} \frac{a_i}{b_i} = 0$ and $\sum_{i=1}^{\infty} a_i$ converges
 \implies the series $\sum_{i=1}^{\infty} b_i$ converges

3) $\lim_{i \rightarrow \infty} \frac{a_i}{b_i} = \infty$ and $\sum_{i=1}^{\infty} b_i$ diverges
 \implies the series $\sum_{i=1}^{\infty} a_i$ diverges.

Remarks

- > To know whether $\sum a_n$ is convergent, we have to compare $\sum a_n$ with another series of the same order and this series can be determined as

$$b_n = \frac{\text{Highest degree term in } n \text{ in the Numerator of } a_n}{\text{Highest degree term in } n \text{ in the Denominator of } a_n}$$

REMARK

This is one of the most powerful tests, because it squeezes the two series “in the limit”. Just be sure to use it right!

REMARK

We will mainly be using only the limit form of the comparison test for solving problems.

Remarks

- > To know whether $\sum a_n$ is convergent, we have to compare $\sum a_n$ with another series of the same order and this series can be determined as

$$b_n = \frac{\text{Highest degree term in } n \text{ in the Numerator of } a_n}{\text{Highest degree term in } n \text{ in the Denominator of } a_n}$$

REMARK

This is one of the most powerful tests, because it squeezes the two series “in the limit”. Just be sure to use it right!

REMARK

We will mainly be using only the limit form of the comparison test for solving problems.

Remarks

- > To know whether $\sum a_n$ is convergent, we have to compare $\sum a_n$ with another series of the same order and this series can be determined as

$$b_n = \frac{\text{Highest degree term in } n \text{ in the Numerator of } a_n}{\text{Highest degree term in } n \text{ in the Denominator of } a_n}$$

REMARK

This is one of the most powerful tests, because it squeezes the two series “in the limit”. Just be sure to use it right!

REMARK

We will mainly be using only the limit form of the comparison test for solving problems.

Remarks

- > To know whether $\sum a_n$ is convergent, we have to compare $\sum a_n$ with another series of the same order and this series can be determined as

$$b_n = \frac{\text{Highest degree term in } n \text{ in the Numerator of } a_n}{\text{Highest degree term in } n \text{ in the Denominator of } a_n}$$

REMARK

This is one of the most powerful tests, because it squeezes the two series “in the limit”. Just be sure to use it right!

REMARK

We will mainly be using only the limit form of the comparison test for solving problems.

A useful test

$\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges **if** $p > 1$, **and** diverges **if** $p \leq 1$

A useful test

$\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges **if** $p > 1$, **and** diverges **if** $p \leq 1$

A useful test

$\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges **if** $p > 1$, **and** diverges **if** $p \leq 1$

A useful test

$\sum_{n=1}^{\infty} \frac{1}{n^p}$ converges **if** $p > 1$, **and** diverges **if** $p \leq 1$

Examples on Comparision test

Problem 1: Discuss the convergence of the series $\frac{1}{1.2} + \frac{1}{2.3} + \frac{1}{3.4} + \dots$

Examples on Comparision test

Problem 1: Discuss the convergence of the series $\frac{1}{1.2} + \frac{1}{2.3} + \frac{1}{3.4} + \dots$

Examples on Comparision test

Problem 1: Discuss the convergence of the series $\frac{1}{1.2} + \frac{1}{2.3} + \frac{1}{3.4} + \dots$

Examples on Comparision test

Problem 1: Discuss the convergence of the series $\frac{1}{1.2} + \frac{1}{2.3} + \frac{1}{3.4} + \dots$

Example for Comparison Test

Soln:

$$\text{Here } a_n = \frac{1}{n(n+1)} = \frac{1}{n^2 \left(1 + \frac{1}{n}\right)}.$$

Let us compare $\sum a_n$ with $\sum b_n$ where $b_n = \frac{1}{n^2}$. $\frac{a_n}{b_n} = \frac{1}{1 + \frac{1}{n}} \therefore \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{1}{1 + \frac{1}{n}} = 1$

which is finite and $\neq 0$, $\therefore \sum a_n$ and $\sum b_n$ behave alike.

Since $\sum b_n = \sum \frac{1}{n^2}$ is of the form $\sum \frac{1}{n^p}$ with $p = 2 > 1$, $\sum b_n$ is convergent

$\Rightarrow \sum a_n$ is convergent.

Example for Comparison Test

Soln:

$$\text{Here } a_n = \frac{1}{n(n+1)} = \frac{1}{n^2 \left(1 + \frac{1}{n}\right)}.$$

Let us compare $\sum a_n$ with $\sum b_n$ where $b_n = \frac{1}{n^2}$. $\frac{a_n}{b_n} = \frac{1}{1 + \frac{1}{n}} \therefore \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{1}{1 + \frac{1}{n}} = 1$

which is finite and $\neq 0$, $\therefore \sum a_n$ and $\sum b_n$ behave alike.

Since $\sum b_n = \sum \frac{1}{n^2}$ is of the form $\sum \frac{1}{n^p}$ with $p = 2 > 1$, $\sum b_n$ is convergent

$\Rightarrow \sum a_n$ is convergent.

Example for Comparison Test

Soln:

$$\text{Here } a_n = \frac{1}{n(n+1)} = \frac{1}{n^2 \left(1 + \frac{1}{n}\right)}.$$

Let us compare $\sum a_n$ with $\sum b_n$ where $b_n = \frac{1}{n^2}$. $\frac{a_n}{b_n} = \frac{1}{1 + \frac{1}{n}} \therefore \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{1}{1 + \frac{1}{n}} = 1$

which is finite and $\neq 0$, $\therefore \sum a_n$ and $\sum b_n$ behave alike.

Since $\sum b_n = \sum \frac{1}{n^2}$ is of the form $\sum \frac{1}{n^p}$ with $p = 2 > 1$, $\sum b_n$ is convergent
 $\Rightarrow \sum a_n$ is convergent.

Example for Comparison Test

Soln:

$$\text{Here } a_n = \frac{1}{n(n+1)} = \frac{1}{n^2 \left(1 + \frac{1}{n}\right)}.$$

Let us compare $\sum a_n$ with $\sum b_n$ where $b_n = \frac{1}{n^2}$. $\frac{a_n}{b_n} = \frac{1}{1 + \frac{1}{n}} \therefore \lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{1}{1 + \frac{1}{n}} = 1$

which is finite and $\neq 0$, $\therefore \sum a_n$ and $\sum b_n$ behave alike.

Since $\sum b_n = \sum \frac{1}{n^2}$ is of the form $\sum \frac{1}{n^p}$ with $p = 2 > 1$, $\sum b_n$ is convergent

$\Rightarrow \sum a_n$ is convergent.

Example for Comparison Test

Problem 2: Discuss the convergence of the series $\frac{1}{1.2.3} + \frac{3}{2.3.4} + \frac{5}{3.4.5} + \dots$

Example for Comparison Test

Problem 2: Discuss the convergence of the series $\frac{1}{1.2.3} + \frac{3}{2.3.4} + \frac{5}{3.4.5} + \dots$

Example for Comparison Test

Problem 2: Discuss the convergence of the series $\frac{1}{1.2.3} + \frac{3}{2.3.4} + \frac{5}{3.4.5} + \dots$

Example for Comparison Test

Problem 2: Discuss the convergence of the series $\frac{1}{1.2.3} + \frac{3}{2.3.4} + \frac{5}{3.4.5} + \dots$

Example for Comparison Test

Soln:

$$\begin{aligned}\text{Here } a_n &= \frac{T_n \text{ of } 1, 3, 5, \dots}{n(n+1)(n+2)} = \frac{2n-1}{n(n+1)(n+2)} \\ &= \frac{n(2 - \frac{1}{n})}{n^3(1 + \frac{1}{n})(1 + \frac{2}{n})} \\ &= \frac{(2 - \frac{1}{n})}{n^2(1 + \frac{1}{n})(1 + \frac{2}{n})}.\end{aligned}$$

Example for Comparison Test

Soln:

$$\begin{aligned}\text{Here } a_n &= \frac{T_n \text{ of } 1, 3, 5, \dots}{n(n+1)(n+2)} = \frac{2n-1}{n(n+1)(n+2)} \\ &= \frac{n(2 - \frac{1}{n})}{n^3(1 + \frac{1}{n})(1 + \frac{2}{n})} \\ &= \frac{(2 - \frac{1}{n})}{n^2(1 + \frac{1}{n})(1 + \frac{2}{n})}.\end{aligned}$$

Example for Comparison Test

Soln:

$$\begin{aligned}\text{Here } a_n &= \frac{T_n \text{ of } 1, 3, 5, \dots}{n(n+1)(n+2)} = \frac{2n-1}{n(n+1)(n+2)} \\ &= \frac{n(2 - \frac{1}{n})}{n^3(1 + \frac{1}{n})(1 + \frac{2}{n})} \\ &= \frac{(2 - \frac{1}{n})}{n^2(1 + \frac{1}{n})(1 + \frac{2}{n})}.\end{aligned}$$

Example for Comparison Test

Soln:

$$\begin{aligned}\text{Here } a_n &= \frac{T_n \text{ of } 1, 3, 5, \dots}{n(n+1)(n+2)} = \frac{2n-1}{n(n+1)(n+2)} \\ &= \frac{n(2 - \frac{1}{n})}{n^3(1 + \frac{1}{n})(1 + \frac{2}{n})} \\ &= \frac{(2 - \frac{1}{n})}{n^2(1 + \frac{1}{n})(1 + \frac{2}{n})}.\end{aligned}$$

Example for Comparison Test

Let us compare $\sum a_n$ with $\sum b_n$ where $b_n = \frac{1}{n^2}$

$$\frac{a_n}{b_n} = \frac{\left(2 - \frac{1}{n}\right)}{\left(1 + \frac{1}{n}\right)\left(1 + \frac{2}{n}\right)}.$$

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\left(2 - \frac{1}{n}\right)}{\left(1 + \frac{1}{n}\right)\left(1 + \frac{2}{n}\right)} = \frac{2}{(1)(1)} = 2 \text{ which is finite and } \neq 0.$$

Example for Comparison Test

Let us compare $\sum a_n$ with $\sum b_n$ where $b_n = \frac{1}{n^2}$

$$\frac{a_n}{b_n} = \frac{\left(2 - \frac{1}{n}\right)}{\left(1 + \frac{1}{n}\right)\left(1 + \frac{2}{n}\right)}.$$

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\left(2 - \frac{1}{n}\right)}{\left(1 + \frac{1}{n}\right)\left(1 + \frac{2}{n}\right)} = \frac{2}{(1)(1)} = 2 \text{ which is finite and } \neq 0.$$

Example for Comparison Test

Let us compare $\sum a_n$ with $\sum b_n$ where $b_n = \frac{1}{n^2}$

$$\frac{a_n}{b_n} = \frac{\left(2 - \frac{1}{n}\right)}{\left(1 + \frac{1}{n}\right)\left(1 + \frac{2}{n}\right)}.$$

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\left(2 - \frac{1}{n}\right)}{\left(1 + \frac{1}{n}\right)\left(1 + \frac{2}{n}\right)} = \frac{2}{(1)(1)} = 2 \text{ which is finite and } \neq 0.$$

Example for Comparison Test

Let us compare $\sum a_n$ with $\sum b_n$ where $b_n = \frac{1}{n^2}$

$$\frac{a_n}{b_n} = \frac{\left(2 - \frac{1}{n}\right)}{\left(1 + \frac{1}{n}\right)\left(1 + \frac{2}{n}\right)}.$$

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\left(2 - \frac{1}{n}\right)}{\left(1 + \frac{1}{n}\right)\left(1 + \frac{2}{n}\right)} = \frac{2}{(1)(1)} = 2 \text{ which is finite and } \neq 0.$$

Example for Comparison Test

$\therefore \sum a_n$ and $\sum b_n$ behave alike.

Since $\sum b_n = \sum \frac{1}{n^2}$ is of the form $\sum \frac{1}{n^p}$ with $p = 2 > 1$,

$\sum b_n$ is convergent

$\Rightarrow \sum a_n$ is convergent.

Example for Comparison Test

$\therefore \sum a_n$ and $\sum b_n$ behave alike.

Since $\sum b_n = \sum \frac{1}{n^2}$ is of the form $\sum \frac{1}{n^p}$ with $p = 2 > 1$,

$\sum b_n$ is convergent

$\Rightarrow \sum a_n$ is convergent.

Example for Comparison Test

$\therefore \sum a_n$ and $\sum b_n$ behave alike.

Since $\sum b_n = \sum \frac{1}{n^2}$ is of the form $\sum \frac{1}{n^p}$ with $p = 2 > 1$,

$\sum b_n$ is convergent

$\Rightarrow \sum a_n$ is convergent.

Example for Comparison Test

$\therefore \sum a_n$ and $\sum b_n$ behave alike.

Since $\sum b_n = \sum \frac{1}{n^2}$ is of the form $\sum \frac{1}{n^p}$ with $p = 2 > 1$,

$\sum b_n$ is convergent

$\Rightarrow \sum a_n$ is convergent.

ENGINEERING MATHEMATICS-I



Problems :

1. Discuss the convergence or divergence of the following :

$$1. \frac{1}{a \cdot 1^2 + b} + \frac{1}{a \cdot 2^2 + b} + \frac{1}{a \cdot 3^2 + b} + \dots \dots \dots \infty$$

$$n^{\text{th}} \text{ term : } a_n = \frac{1}{a \cdot n^2 + b} = \frac{1}{n^2 \left(a + \frac{b}{n^2} \right)}$$

$\sum b_n = \sum \frac{1}{n^2}$ (p series with $p=2 > 1$ hence convergent)

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{1}{a + \frac{b}{n^2}} = \frac{1}{a} = \text{finite and } \neq 0$$

$\therefore \sum b_n$ is convergent $\therefore \sum a_n$ and $\sum b_n$ behave alike
 $\therefore \sum a_n$ is convergent.

ENGINEERING MATHEMATICS-I

$$2. \frac{1}{1^p} + \frac{1}{3^p} + \frac{1}{5^p} + \frac{1}{7^p} + \dots \dots \infty$$

$$\Rightarrow n^{\text{th}} \text{ term : } a_n = \frac{1}{(2n-1)^p} = \frac{1}{n^p \left(2 - \frac{1}{n}\right)^p}$$

$$\text{choosing } b_n = \frac{1}{n^p}$$

$$\sum b_n = \sum \frac{1}{n^p} \text{ converges } p > 1 \text{ and diverges } p \leq 1 \quad \rightarrow p \text{ series}$$

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{1}{\left(2 - \frac{1}{n}\right)^p} = \frac{1}{2^p} \text{ which is finite and non zero}$$

$$\because \sum b_n \text{ converges } \therefore \sum a_n \text{ converges}$$

ENGINEERING MATHEMATICS-I

$\sum b_n$ diverges $\Rightarrow \sum a_n$ also diverges for $p \leq 1$ and
converges $p > 1$

3. $\sum \frac{\sqrt{n+1} - \sqrt{n}}{n^p}$

$$\Rightarrow \frac{\sqrt{n+1} - \sqrt{n}}{n^p} \cdot \frac{\sqrt{n+1} + \sqrt{n}}{\sqrt{n+1} + \sqrt{n}} = a_n$$

On simplifying

$$\frac{1}{n^{p+1/2} \left(\sqrt{1 + \frac{1}{n}} + 1 \right)}$$

Choose $V_n = \frac{1}{n^{p+1/2}}$

Convergent

$p + 1/2 > 1$ divergent $p + 1/2 \leq 1$

ENGINEERING MATHEMATICS-I



$$\lim_{n \rightarrow \infty} \frac{a_n}{\sqrt[n]{n}} = \frac{1}{2}$$

$\therefore \sum a_n$ and $\sum \sqrt[n]{n}$ behave alike

$\sum a_n$ converges for $p + \frac{1}{2} > 1$

diverges for $p + \frac{1}{2} \leq 1$

$\therefore \sum a_n$ is convergent if $p > \frac{1}{2}$

divergent if $p \leq \frac{1}{2}$



THANK YOU

Aparna B. S

Department of Science and Humanities

aparnabs@pes.edu