



ENGINEERING MATHEMATICS - I

Partial Differentiation

Dr. Anitha
Science and Humanities

ENGINEERING MATHEMATICS - I

UNIT 2 : Partial Differentiation

Session : 5

Sub Topic : Implicit Functions, Problems

Dr. Anitha

Department of Science and Humanities

Consider an implicit function $x^3y + 3xy^2 = 3$

Then,

$$x^3 \frac{dy}{dx} + 3yx^2 + 3y^2 + 6xy \frac{dy}{dx} = 0,$$

$$\frac{dy}{dx} = -\frac{3y(x+y)}{x(x^2+6y)}$$

Similarly, if $z = f(x_1, x_2, x_3, \dots, x_n)$

total differential of z is defined by

$$dz = f_{x_1} dx_1 + f_{x_2} dx_2 + f_{x_3} dx_3 + \dots + f_{x_n} dx_n$$

If $z = f(x, y)$ is a function of two independent variables, then


$$dz = f_x(x, y)dx + f_y(x, y)dy$$

If z represents a level surface, i.e. if $z = f(x, y) = c$, then

$$dz = 0, \Rightarrow \frac{dy}{dx} = -\frac{f_y(x, y)}{f_x(x, y)}$$


The total derivative is also called the differential.

The differential is used in integration by substitution and in approximation. It is also useful to calculate the Errors in approximation.



Branch diagram for differentiating $z = f(x, y)$ w.r.t x .

Setting $\frac{dz}{dx} = 0$ leads to a simple computation formula for implicit differentiation



ENGINEERING MATHEMATICS - I

Implicit function of three variables

Let $f(x, y, z) = 0$ be the equation of an implicit function of three variables x, y, z . Suppose y and z are functions of x , then f is a function of one independent variable x and y, z are intermediate variables.

Keeping z constant, differentiating w.r.t. x , we get


$$\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial x} = 0$$

Solving, $\frac{\partial y}{\partial x} = -\frac{f_x}{f_y}$ provided $f_y \neq 0$.

Similarly, differentiating w.r.t. x , holding y constant

$$\frac{\partial f}{\partial x} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial x} = 0,$$

Solving, $\frac{\partial z}{\partial x} = -\frac{f_x}{f_z}$, if $f_z \neq 0$.



<https://www.intmath.com/vectors/3d-grapher.php>

ENGINEERING MATHEMATICS - I

Implicit Functions -Problems

1. For the curve $xe^y + ye^x = 0$, find the equation of the tangent line at the origin.

Solution:

$$xe^y + ye^x = 0 \dots\dots\dots(1)$$

Differentiating w.r.t x,


$$\frac{dy}{dx} = -\frac{ye^x + e^y}{xe^y + e^x}$$

Slope of the tangent at origin

$$\frac{dy}{dx} \text{ at } (0,0) = -1$$

Equation of the tangent line at(0,0),

$$y + x = 0$$



<https://www.intmath.com/vectors/3d-grapher.php>

2. Find $\frac{dy}{dx}$ if $(\cos x)^y = (\sin y)^x$

Solution:

$$\text{Let } f(x, y) = (\cos x)^y - (\sin y)^x = 0$$

$$\frac{dy}{dx} = -\frac{f_x}{f_y}$$

$$= \left[\frac{y(\cos x)^{y-1}(-\sin x) - (\sin y)^x \log(\sin y)}{(\cos x)^y \log(\cos x) - x(\sin y)^{x-1}(\cos y)} \right]$$

$$= \frac{\sin y (y \sin x + \cos x \log(\sin y))}{\cos x (\sin y \log(\cos x) - x \cos y)}$$

3. Find $\frac{\partial y}{\partial x}$ and $\frac{\partial z}{\partial x}$ at (0,1,2) at $z^3 + xy - y^2z = 6$.

Solution:

$$f = z^3 + xy - y^2z - 6 = 0$$

Keeping z as constant

$$\frac{\partial y}{\partial x} = \frac{-\frac{\partial f}{\partial x}}{\frac{\partial f}{\partial y}} = \frac{-y}{x-2yz}$$

$$\frac{\partial y}{\partial x} \text{ at } (0,1,2) = \frac{1}{4}$$

Contd.....

Keeping y as constant

$$\begin{aligned}\frac{\partial z}{\partial x} &= -\frac{\frac{\partial f}{\partial x}}{\frac{\partial f}{\partial z}} \\ &= \frac{-y}{3z^2 - y^2}\end{aligned}$$

$$\frac{\partial z}{\partial x} \text{ at } (0,1,2) = \frac{-1}{11}$$



Dr. Anitha

Department of Science and Humanities

nanitha@pes.edu

Extn 730