

Task 6: Investigating Fossil Dating — Complete Study Pack

Course: ATAR Human Biology Units 3 & 4 (ATHBY), Kent Street SHS, 2026
Sources used: KSSHS_Course_Outline_(1)_ATHBY_2026_v2.docx (Weeks 26–27 row) and hp12wa8e_11_0therevidenceofevolution.pdf (Human Perspectives ATAR Units 3 & 4, **Chapter 11 "Other Evidence of Evolution", pp. 295–321**)

1. What the outline actually says about Task 6

Field	Detail
Task	Task 6: Investigating Fossil dating
Timeline	Weeks 26–27 (Semester 2, Unit 4)
Topic	Evidence for evolution
Textbook	CH 11 (pp. 295–321) , Review Questions pp. 320–321 , plus AYKQ

Syllabus content examined by this task (verbatim from the outline):

- Evolutionary relationships between groups can be represented using phylogenetic trees (*construction of phylogenetic trees is NOT required*).
- The fossil record is incomplete and cannot represent the entire biodiversity of a time or location, due to many factors affecting fossil formation, the persistence of fossils, and accessibility of fossilised remains.
- Sequencing a fossil record requires a **combination** of relative and absolute dating techniques to place fossils onto the geological timeline.
- Both relative techniques (**stratigraphy, index fossils**) and absolute techniques (**radiocarbon, potassium-argon**) have **limitations of application**.

Learning intentions listed for these weeks:

- Fossils and fossil **formation**
- **Problems and limitations** of the fossil record
- **Application and limitations** of relative dating: stratigraphy and index fossils
- **Application and limitations** of absolute dating: radiocarbon and potassium-argon
- **How these techniques are represented graphically** and used to accurately date a fossil

Note: the outline names the task but does **not** state its weighting, duration, or format. Because it is called an "Investigating..." task and the syllabus dot point about **graphical representation** is explicitly listed, expect a **Science Inquiry / data-analysis style task**: decay curves, stratigraphic columns, half-life calculations, and evaluation of reliability, validity and limitations. It maps directly onto textbook **Activity 11.1 (Investigating radioisotope methods of dating)** and **Activity 11.2 (Investigating stratigraphy)** — do both of these before the task.

Also expect the Week 1 **Science Inquiry Skills** to be assessed alongside the content: hypotheses and predictions, analysing data, identifying trends/patterns, discussing measurement error, instrument accuracy, procedure and sample size as sources of limitation, and justifying conclusions with evidence.

2. Fossils (Ch 11.1)

Fossil: any preserved trace left by an organism that lived long ago. Forms: bones, teeth, shells, footprints, burrows, faeces (coprolites), impressions of part or all of a plant/animal. For human ancestors: usually bones, teeth, occasionally footprints.

Artefact: an object deliberately made or modified by humans — stone tools, beads, carvings, charcoal from cooking fires, cave paintings. (An artefact is *made*; a fossil is *organic remains/traces*.)

Four conditions required for fossilisation

1. **Rapid burial** of the material (drifting sand, river mud, volcanic ash, deliberate burial by other members of the species).
2. Presence of **hard body parts** (bone, teeth, shell).
3. **Absence of decay organisms** (decomposers) / low oxygen.
4. A **long period of stability** — remains left undisturbed.

Effect of soil type

- **Wet, acidic soil** → minerals in bone dissolve → **no bone fossilisation**.
- **Acidic + oxygen-free (e.g. peat bogs)** → **soft tissue** preserved (skin, hair, gut contents), but bones demineralised.
- **Alkaline soil** → **best bone fossils**: minerals are not dissolved; new minerals (lime, iron oxide) are deposited in bone pores, replacing the organic matter (~35% of bone by weight) → bone becomes **petrified** but structural detail is preserved.

Where human-ancestor fossils are found (and why)

- Edges of **ancient lakes and rivers** — sediment builds up quickly during floods/slowed flow → rapid burial.
- **Caves** — often limestone (calcium carbonate) deposited around remains; roof/wall collapse buries bodies.
- **Volcanically active areas** — usually heat destroys remains, **but** East African **ash falls** have preserved many hominin fossils.

Excavation ("a dig")

Site surveyed and marked into sections → **small hand tools** used so material isn't damaged → removed soil **sieved** so tiny fragments aren't lost → photographs at every stage to record positions → each item labelled and catalogued → in the lab: bones scraped clean, fragments pieced together, measurements taken, plaster casts or latex moulds made.

Reconstruction problem

Whole skeletons are rare (e.g. *Homo ergaster* skull from Kenya, ~1.8 My, reconstructed from fragments with missing areas estimated). Reconstructions are **approximations based on expert judgement**, so different scientists interpret differently → controversy. The only resolution is finding more fossils.

3. Dating fossils (Ch 11.2)

- **Absolute date** — the **actual age in years** of the specimen.
- **Relative date** — whether a specimen is **older or younger** than another specimen (or than the rock/soil it's in). Establishes a **sequence** of events.
- Ages are given as **years BP** = "before present" (e.g. 45 000 years BP = 45 000 years old).

3.1 Potassium-argon dating (absolute)

- Potassium (**K**) is one of the most abundant elements in Earth's crust; occurs as three isotopes — K-39 (20 n), K-40 (21 n), K-41 (22 n). **Isotopes** = atoms of the same element (same protons) with different neutron numbers.
- **K-40 is radioactive** and decays to **calcium-40** and **argon-40** at an extremely slow but **constant** rate.
- As the rock ages: **K-40 decreases, Ar-40 increases** → measuring the **K-40 : Ar-40 ratio** gives the rock's age (Figure 11.6 = parent-decay / daughter-buildup curves crossing at 1 half-life).
- **Half-life of K-40 = 1.25×10^9 years (1.25 billion = 1250 million years)**. ⚠ The chapter text contains a **typo**: "1250 billion years (1.25×10^9 years)". The bracketed figure is correct (1.25 billion); Activity 11.1B quotes "around 1300 million years". Use **1.25 billion years** and expect either figure to be accepted.
- **Argon is a gas** — it escapes from molten lava, so the "clock" starts when the lava **solidifies**; trapped Ar-40 in solid lava must therefore have come from K-40 decay.

Limitations of potassium-argon dating

- Only works on **suitable rock types** — volcanic/lava deposits (not all rocks are suitable).
- Only useful for material **older than ~200 000 years** (theoretical limit 100 000–200 000). After 100 000 years only **0.0053%** of K-40 has decayed — below reliable detection limits.
- It dates the **rock, not the fossil** — you need volcanic rock **of the same age** adjacent to (e.g. burying) the fossil, and the fossil's age is then **inferred**.
- Cannot be used directly on bone or any organic material.
- Contamination or argon leakage/atmospheric argon inclusion reduces accuracy.

3.2 Radiocarbon (carbon-14) dating (absolute)

- **C-14 is produced in the upper atmosphere** by cosmic radiation acting on nitrogen, at about the same rate as it decays; it **decays back to nitrogen-14**.
- Atmospheric ratio $\approx$ **1 C-14 atom : 10^{12} C-12 atoms**.
- Plants fix atmospheric CO₂ in **photosynthesis** → C-14 enters plant tissue → animals eat plants → C-14 enters animal tissue. **At death, intake stops** but decay continues at a fixed rate.
- Measuring radiation released (or counting atoms) gives the **C-14 : C-12 ratio** → age.
- **Half-life = 5730 ± 40 years**. Ratio falls 1 → 0.5 → 0.25 → 0.125 (per 10^{12}) each 5730 years.
- **Conventional method** needs $\geq$ **3 g** of organic material.
- **AMS (accelerator mass spectrometry) radiocarbon dating** breaks the sample into constituent atoms and counts isotope atoms — works on as little as **100 µg (0.0001 g)**. This allows accurate dating of **cave-painting pigments**, which often contain charcoal or organic binders (honey, milk, blood, oil seed).

Limitations of radiocarbon dating

- Material must contain **organic (carbon) compounds** — i.e. it must once have been living.
- Maximum useful age $\approx$ **60 000 years** (by ~70 000 years only **0.021%** of C-14 remains — too little to measure accurately).
- The atmospheric **C-14 : C-12 ratio is not constant** (it was once assumed to be) → dates must be treated with caution. **Corrections** are possible for roughly the **last 9000 years** using independent evidence such as **tree-ring dating (dendrochronology)**.
- Contamination with older or modern carbon skews results; small samples increase error.
- Because bones are often mineralised/petrified (organic matter replaced), fossil bone is frequently undatable by C-14 — the associated **charcoal from hearths** is dated instead and the artefact age deduced.

3.3 Table 11.1 — Useful range of common absolute dating methods

Method	Material used	Useful range (years BP)
Tree growth rings (dendrochronology)	Wood	Up to 9 000
Carbon-14	Carbon compounds	Up to 60 000
Protactinium	Sea sediments	Up to 250 000
Uranium-thorium	Sea sediments, coral	Up to 600 000
Potassium-argon	Volcanic deposits	200 000 and earlier
Electron spin resonance	Calcium carbonate (shells, coral, teeth), quartz, flint	Up to 100 000, possibly 300 000
Fission tracks	Minerals and glass	100 years ago to 4550 million
Thermoluminescence	Sediments, lava, ceramics	300 years ago to 100 000

(Only C-14 and K-Ar are examinable in detail; the rest show that methods overlap and are chosen to suit the sample and age range.)

3.4 Relative dating — stratigraphy

Stratigraphy = the study of layers (strata). Two uses:

(a) Principle of superposition — in undisturbed sedimentary rock, **upper layers are younger than those beneath**, so fossils/artefacts in upper layers are younger. *Limitations / when it fails:*

- **Distortions of Earth's crust** (folding, faulting, tilting, overturning) can invert or offset a sequence.
- Fossils/artefacts can be **buried later** by animals (burrowing) or by early humans (digging, graves) → the specimen may be **younger** than layers above it.
- Erosion can remove layers; material can wash or fall down cracks.

(b) Correlation of rock strata — matching layers between different areas (sometimes thousands of km apart) by examining the rock itself and the fossils it contains. **Rocks containing the same fossils are assumed to be the same age.**

Index fossils — fossils that are especially useful for correlation because they:

1. were present on Earth for only a **short, limited period of time**;
2. are **widely (geographically) distributed**;
3. are **abundant** in the strata; and
4. are **easily recognised/identified**. Examples: **trilobites** (lived ~500–300 Mya then went extinct), ammonites, and some **fossil pollen grains**.

Fossil pollen (palynology) — some pollen grains work as index fossils; even when they don't, preserved pollen lets a botanist reconstruct the **type and amount of vegetation**, and hence the **climate**, at the time the deposit formed. This can **confirm or refute** relative dates obtained by other methods.

3.5 Why relative and absolute dating are used together

Absolute methods each need a **particular set of circumstances** (right material, right age range). When those aren't available, relative methods still establish the **sequence**. Combining them lets a fossil be placed on the **geological timeline**: e.g. a layer bracketed between two K-Ar-dated lava flows must be between those ages; a layer correlated by index fossils to a C-14-dated layer elsewhere inherits that age.

4. Problems and limitations of the fossil record (high-frequency exam question)

1. **Fossilisation is a chance event** — the four required conditions rarely coincide, and occur irregularly in time, so there are **gaps**.
 2. Soft-bodied organisms and **soft tissues** (skin, muscle, organs) are rarely preserved → record biased toward **hard parts** (bone, teeth, shell) and toward organisms living in depositional environments.
 3. Acidic/wet soils **destroy** bone; heat (volcanic) destroys remains.
 4. Only a **very small proportion** of existing fossils have been **discovered**: some are buried too deep, some are in **inaccessible** places.
 5. Some are **not recognised** as fossils, or are **destroyed inadvertently** by agriculture, industry, construction, war.
 6. Fossils are usually **fragmentary** → reconstructions are interpretive and disputed.
 7. **Dating is often problematic** — C-14 needs carbon and works only to ~60 000 years; K-Ar needs suitable volcanic material and only works beyond ~200 000 years; so some fossils fall in awkward gaps or lack datable associated material.
 8. Therefore the fossil record **cannot represent the entire biodiversity** of any time or place.
-

5. Graphical representation and calculations (the "investigating" part)

5.1 The two curves you must be able to draw and read

- Carbon-14 decay curve (Fig 11.8):** x-axis = age of sample (thousand years), y-axis = ratio of C-14 to C-12 (1 in $10^{12} \rightarrow 0.5 \rightarrow 0.25 \rightarrow 0.125 \dots$). Each dot = one half-life = 5730 years. Smooth exponential decay curve.
- Potassium-argon parent/daughter curves (Fig 11.6):** x-axis = number of half-lives, y-axis = % of atoms. **Parent (K-40) decays** from 100% downward; **daughter (Ar-40) builds up** from 0% upward; the curves **cross at 50% after one half-life**.

Half-life rules: after n half-lives, remaining fraction = $(1/2)^n \rightarrow 50\%, 25\%, 12.5\%, 6.25\%, 3.125\% \dots$ Age = $n \times$ half-life, where n = number of half-lives elapsed. Parent : daughter ratio 1:1 $\rightarrow$ 1 half-life; 1:3 $\rightarrow$ 2 half-lives; 1:7 $\rightarrow$ 3 half-lives.

5.2 Activity 11.1A — Radiocarbon decay table (worked)

Starting decay rate for living/recently dead material = **15.6 nuclei/s/g**; half-life 5730 y.

Half-lives	Age (years)	Radioactivity (nuclei/s/g)	Fraction remaining
0	0	15.6	100%
1	5 730	7.8	50%
2	11 460	3.9	25%
3	17 190	1.95	12.5%
4	22 920	0.975	6.25%
5	28 650	0.4875	3.125%
6	34 380	0.244	1.56%
7	40 110	0.122	0.78%
8	45 840	0.061	0.39%
9	51 570	0.030	0.20%
10	57 300	0.015	0.10%

Plot decay rate (y) against age (x) to 60 000 years, join with a **smooth curve** (not a ruled line), label axes with units, and read values by interpolation.

Worked answers to Activity 11.1A Q3 (read off the curve; calculated values shown):

- (a)** Charcoal at 8.9 nuclei/s/g $\rightarrow 8.9/15.6 = 0.571$ remaining $\rightarrow \approx$ **0.81 half-lives $\approx$ 4600 years old** (graph read $\sim$ 4500–5000 y).
- (b)** Wood of 9000 years $\rightarrow 9000/5730 = 1.57$ half-lives $\rightarrow 0.337$ remaining $\rightarrow \approx$ **5.3 nuclei/s/g** (graph read $\sim$ 5–5.5).
- (c)** If the wood is much **older** than the 9000-year tools: the wood came from an already-old tree/timber that was **re-used** or was already dead when deposited; older material was **redeposited/washed in**, or the deposits were **disturbed** (burrowing animals, human digging, roof collapse, trampling) mixing layers; sample **contaminated** with old carbon; the tools may have been misdated.
- (d)** If the wood is much **younger**: it was introduced **later** — fell/washed down a crack, carried down by **burrowing animals or plant roots**, buried by later human activity that dug into older deposits; or the sample was **contaminated with modern carbon** (handling, humic acids, recent rootlets).
- (e)** Bone at 1.5 nuclei/s/g $\rightarrow 15.6/1.5 = 10.4 \rightarrow \log_2 10.4 = 3.38$ half-lives $\rightarrow \approx$ **19 400 years old** (graph read $\sim$ 19 000–20 000). (*Only possible if collagen/organic carbon survives.*)
- (f)** Wood tree-ring dated at 4000 y $\rightarrow 4000/5730 = 0.70$ half-lives $\rightarrow 0.616$ remaining $\rightarrow \approx$ **9.6 nuclei/s/g**.

5.3 Activity 11.1B — Potassium-argon (method for the volcano diagram)

Layers **A, C, E, G** = **lava**; layers **B, D, F** = **sediment** deposited between eruptions.

- **Q1 — Why no Ar-40 in layer A?** A is the **youngest/most recent** lava. Argon-40 is a gas that escapes while lava is molten, so the clock starts on solidification; insufficient time has passed for a **detectable** amount of K-40 (half-life 1.25 billion years) to decay to Ar-40.
- **Q2 — Ratio in layer E:** count symbols: ratio = K-40 atoms : Ar-40 atoms. Convert to half-lives — $1:1 = 1$ half-life $\Rightarrow \sim 1.25$ billion y; $1:3 = 2$ half-lives $\Rightarrow \sim 2.5$ billion y. State the ratio in simplest form, then the age.
- **Q3 — What is the material in B, D, F?** **Sedimentary** material (sand/silt/mud/ash weathered and deposited by wind and water) laid down **between eruptions** during the long quiet intervals, then buried by the next lava flow.
- **Q4 — Why no fossils in A, C, E, G?** The **extreme heat of molten lava incinerates/destroys** organisms and their remains, so nothing is preserved in igneous layers.
- **Q5 — Conditions assisting fossilisation:** rapid burial; hard body parts; absence of decomposers/oxygen; long undisturbed period; alkaline sediment.
- **Q6 — Dating layer B at 40 000–70 000 y (too young for K-Ar):** use **radiocarbon dating** on any organic material/charcoal in the layer (valid up to $\sim 60\,000$ y; based on C-14 $\rightarrow$ N-14 decay with 5730-y half-life, measuring the C-14:C-12 ratio); **thermoluminescence** on sediments/ceramics (measures light released from trapped electrons that accumulate since burial/heating; 300–100 000 y); **electron spin resonance** on tooth enamel/shell/quartz (measures trapped unpaired electrons; up to $\sim 100\,000$ – $300\,000$ y); plus **relative dating** — it must be younger than lava layer C and older than lava layer A (superposition), and **index fossils** can correlate it with dated strata elsewhere.

5.4 Activity 11.2 — Stratigraphy (how to answer)

- **How were the sediments formed?** Successive **deposition** of sand/silt/mud by water (rivers, floods, lakes, sea) or wind, compacted over long periods into layers.
- **Why don't the layers in series A and B align?** **Earth movements** — faulting (vertical displacement along a fault), folding/tilting, uplift or subsidence, plus **erosion** removing part of a sequence and later deposition on top (unconformity).
- **Oldest stratum:** the **lowest** layer in the deepest series, by the **principle of superposition**; confirm with **correlation** — where the same index fossils appear in two series, those layers are the same age, so you can order layers across sites.
- **Youngest stratum:** the **topmost** layer, again by superposition, cross-checked by correlation.
- **Are the shared fossils index fossils?** Only if they meet all criteria: existed for a **short period**, **geographically widespread**, **abundant**, and **easily identified**. Sharing a fossil between two sites is consistent with index-fossil use because it lets the strata be **correlated as the same age**.
- **If A4 is C-14 dated at 45 000 y**, layers correlated to it are also $\sim 45\,000$ y; layers **above** are younger and layers **below** are older — that's an inference about **relative** age, not an absolute date.
- **Could layer A6 be K-Ar dated?** Only if it contains **volcanic material (lava/ash)** and is **older than $\sim 200\,000$ years**; sedimentary layers cannot be K-Ar dated.
- **Could dendrochronology date layer C2?** Only if it contains **preserved wood with visible growth rings**, and the age is **under ~ 9000 years**; otherwise no.

6. Phylogenetic trees — what you DO need

The outline says **construction is not required**, but you still need to be able to **interpret** them.

- **Phylogenetic tree (dendrogram)**: a diagram showing the **evolutionary relationships** between organisms descended from a common ancestor. Ancestor at the **base**; descendants at branch tips.
 - **Closer branching = more recent common ancestor = more closely related**. Species alive today are drawn **level**; an **extinct** species' branch **ends lower down**. An axis shows **time**.
 - Trees are built from presence/absence of traits, DNA differences, or amino-acid sequence differences — greatest similarity = branches closest together.
 - Trees show **inferred** relationships only: different researchers can produce different trees from the same data, so a tree is effectively a **hypothesis** that can be revised as new data arrives.
-

7. Comparative anatomy (Ch 11.3) — in the chapter, useful supporting evidence

- **Embryology:** vertebrate embryos are strikingly similar early in development — human and chicken embryos both have **slits and arches in the neck** homologous to fish gill slits (which never become gills), a **tail**, a **two-chambered heart**, and similar brain development → evidence of **common ancestry** with later divergence.
- **Homologous structures:** structures with the **same basic structure** but possibly **different functions** — the classic case is the **vertebrate forelimb** (frog forelimb, whale flipper, horse leg, lion limb, human arm, bat wing, bird wing): the same bones arranged the same way → common ancestor.
- **Vestigial structures:** reduced structures that **no longer perform their original function** (humans may have up to ~90). Examples: **nictitating membrane** (third eyelid remnant at the inner corner of the eye — retains a minor tear-drainage role), **ear muscles**, **wisdom teeth (third molars)**, **pyramidalis muscles**, **coccyx** (fused tail vertebrae), **male nipples**, **muscles at the base of hairs** (goosebumps), and the **appendix**. Natural selection reduces useless organs because maintaining them wastes energy/resources, but harmless remnants aren't fully eliminated.
- **Appendix caution:** once the standard vestigial example, but research indicates it has an **immune role and stores beneficial gut bacteria** → now debated. Lesson: be careful labelling an organ vestigial (Weidersheim's 1893 list of 86 included tonsils, thymus, pituitary and vein valves — all now known to be functional).

8. Glossary (Ch 11) — learn these definitions verbatim-ish

Term	Definition
Absolute date	The actual age (in years) of a fossil or artefact
AMS radiocarbon dating	Technique giving radiocarbon dates for very small samples (from 100 µg)
Artefact	An object made or modified by humans
Carbon-14	The radioactive isotope of carbon
Correlation of rock strata	Matching rock strata from different places
Dating	Determining the age of excavated artefacts or fossils
Dendrogram	See phylogenetic tree
Embryology	Study of early development (in humans, fertilisation to end of week 8)
Fossil	Evidence of, or remains of, an organism that lived long ago
Half-life	Time required for half of a quantity of radioactive material to decay to a stable form
Homologous structure	Structures with similar structure but not necessarily similar function
Index fossil	Fossil of an organism present on Earth for only a short period, useful for relative dating of strata
Isotope	Atoms of the same element with the same protons but different numbers of neutrons
Nictitating membrane	Transparent third eyelid; vestigial in humans
Organic compound	A compound of large carbon-containing molecules
Phylogenetic tree	Diagram showing evolutionary relationships between related organisms
Potassium-argon dating	Dating using the known decay rate of radioactive potassium
Principle of superposition	In a vertical sequence, sedimentary layers on top are younger than those lower down
Radiocarbon dating	Dating using the known decay rate of radioactive carbon
Relative date	Age of a fossil/artefact relative to another (older or younger)
Stratigraphy	Study of the sequence of rock layers as a means of relative dating
Vestigial structure	A reduced structure that appears to have no function (e.g. human ear muscles)

9. Comparison table you should be able to reproduce (Review Q5)

Method	Type	Material	Range	Advantages	Limitations
Carbon-14	Absolute	Once-living organic material (bone collagen, charcoal, wood, pigment binders)	$\leq \sim 60\,000$ y	Actual age in years; dates the specimen or associated hearth charcoal directly; AMS works on tiny samples; calibratable against tree rings for last ~ 9000 y	Needs organic carbon; useless $> \sim 60\,000$ y; atmospheric C-14:C-12 has varied; contamination; conventional method needs ≥ 3 g
Potassium-argon	Absolute	Volcanic rock/lava/ash	$> \sim 200\,000$ y	Dates very ancient material; extremely long half-life covers hominin timescales; valuable in East African volcanic sites	Needs suitable volcanic rock of the same age as the fossil; unreliable $< 100\,000$ – $200\,000$ y (only 0.0053% decay at $100\,000$ y); dates the rock, so fossil age is inferred; argon loss/contamination
Dendrochronology	Absolute	Wood with growth rings	≤ 9000 y	Very precise, annual resolution; used to calibrate radiocarbon	Only preserved ringed wood; short range; needs a reference ring sequence
Thermoluminescence	Absolute	Sediments, lava, ceramics	300 – $100\,000$ y	Fills the gap between C-14 and K-Ar; no organic material needed	Narrower accuracy; sensitive to sample's heat/light history
Electron spin resonance	Absolute	Calcium carbonate (shell, coral, teeth), quartz, flint	$\leq 100\,000$, possibly $300\,000$ y	Works on teeth/enamel, common in hominin sites	Limited materials; assumptions about radiation dose
Stratigraphy (superposition)	Relative	Any layered sedimentary sequence	Any	Simple, always applicable to layered deposits; establishes sequence when absolute dating is impossible	No actual age; invalid if strata are inverted/faulted/eroded or if remains were buried later by animals or humans
Index fossils / correlation	Relative	Widespread, short-lived, abundant, recognisable fossils	Any	Correlates strata across huge distances; more precise relative dating; can transfer an absolute date from one site to another	No actual age; requires the right fossils; assumes rocks with the same fossils are the same age

10. Chapter 11 Review Questions (pp. 320-321) — the set to work through

Recall: 1 (define fossil; five forms) · 2 (fossil vs artefact; index fossil; "index artefacts"?) · 3 (best soils; why soft tissue rare) · 4 (relative vs absolute; why relative is still used) · 5 (**three-column table of methods + advantages + limitations** — see §9) · 6 (principle of superposition; when it fails) · 7 (phylogenetic trees and their purpose) · 8 (homologous structures + example) · 9 (vestigial organs; four human examples; significance to evolution).

Explain: 10 (principle behind radioisotope dating) · 11 (**why K-Ar cannot date fossil bone**) · 12 (how C-14 gets into bodies; meaning of a 5730-y half-life; why some artefacts can't be radiocarbon dated; what AMS is) · 13 (index fossils for correlating strata; uses of fossil pollen) · 14 (embryology as evidence, with examples).

Apply: 15 — cave floor: charcoal hearth at 50 cm with C-14:C-12 = **0.25 in 10^{12}** → that's **two half-lives** → **~11 460 years**, so the adjacent stone tool is ~11 500 y; at 95 cm a human jaw and an animal thigh bone. (b) Same depth/stratum in an undisturbed deposit = same age by superposition/association. (c) The thigh bone could be younger if it was introduced later — a burrowing animal, a predator/scavenger dragging it in, roots or water moving it down, or human digging disturbing the layer. 16 — alkaline coastal dunes + rapid burial by drifting sand + undisturbed = **yes, bones should fossilise** (minerals not dissolved; new minerals petrify the bone). 17 — peat bogs: **acidic, waterlogged, oxygen-free, cold** = decomposers inhibited = soft tissue preserved; **but** the acid dissolves bone minerals, so skeletons are **poorly preserved/demineralised**. 18 — Riversleigh: the fauna list (crocodiles, turtles, lungfish, frogs, snails, rainforest mammals) indicates a wet, warm, well-watered **rainforest with permanent water bodies**; limestone caves and pools gave **rapid burial in calcium carbonate-rich (alkaline) sediment** with low oxygen and long stability. 19 — forelimb modified into a **flipper** (shortened, flattened, thickened bones, elongated digits, fused/streamlined into a paddle), a **wing** (elongated, lightened/hollow bones, greatly extended digits supporting membrane or fused for feathers), an **arm** (mobile shoulder, free digits, opposable thumb for manipulation). 20 — other homologies: vertebrate **skull and jaw bones, vertebral column, hindlimb/pelvic girdle**, mammalian **middle ear bones** (from reptilian jaw bones), **teeth, hearts/circulatory patterns**. 21 — 22 — 23: see §7 (vestigial caution), Darwin's prediction (Africa: great apes most anatomically/behaviourally similar to humans live there, so common ancestors likely lived there), and research on uranium-thorium, ESR and thermoluminescence (§3.3/§9).

11. Exam/task technique for this topic

- **"State/identify"** = one-word or one-line answer. **"Describe"** = features/steps, no reasoning required. **"Explain"** = give the mechanism and the *why*. **"Compare"** = both similarities and differences, with linking words (*whereas, both*). **"Evaluate/Discuss limitations"** = make a judgement supported by evidence and state the constraints.
 - Whenever you name a dating technique, get in the marking points: **what decays into what, the half-life, what is measured (ratio), the age range, the material needed, and one or two limitations**.
 - For any graph: **label both axes with units**, use a **sensible even scale**, plot points accurately, draw a **smooth curve** for decay data, and show **dashed interpolation lines** when reading a value off it.
 - For calculations: show **fraction remaining → number of half-lives → × half-life = age**, and give the answer with **units** and a sensible number of significant figures.
 - For "sequence the fossil record" questions: use **superposition first**, then **index fossils to correlate sites**, then **absolute dates to anchor the sequence**, and finish by stating that a **combination** of methods is needed because each has restricted application.
 - For validity/reliability wording: sources of error include small/contaminated samples, disturbed stratigraphy, assuming a constant atmospheric C-14 ratio, assuming a constant decay rate, instrument detection limits, and dating rock rather than the fossil itself. Improvements: larger samples, repeated/replicate measurements, cross-checking with a second independent method, calibration curves.
-

12. Study checklist

- ☐ Define fossil, artefact, index fossil, half-life, isotope, absolute vs relative date, BP
- ☐ List the four conditions for fossilisation and explain soil-type effects (acidic vs alkaline vs peat)
- ☐ Explain where and why hominin fossils are found; describe excavation methods
- ☐ Explain $K-40 \rightarrow Ca-40 + Ar-40$, half-life 1.25 billion y, what's measured, range $> 200\,000$ y, all limitations
- ☐ Explain C-14 formation and decay to N-14, half-life 5730 y, ratio $1:10^{12}$, range $\leq 60\,000$ y, AMS, all limitations
- ☐ Reproduce Table 11.1 (at least C-14, K-Ar, dendrochronology, TL, ESR ranges)
- ☐ Draw and read a C-14 decay curve; do half-life calculations both directions (activity $\rightarrow$ age, age $\rightarrow$ activity)
- ☐ Draw/interpret the K-40 decay + Ar-40 buildup graph; convert parent:daughter ratios to half-lives
- ☐ Apply the principle of superposition and state three reasons it can fail
- ☐ Explain correlation of strata and the four criteria of an index fossil
- ☐ Explain the uses of fossil pollen
- ☐ List at least six reasons the fossil record is incomplete
- ☐ Interpret a phylogenetic tree (recency of common ancestor, extinct branches, tree = hypothesis)
- ☐ Complete textbook **Activities 11.1, 11.2** (and 11.3 for tree interpretation) and **Review Questions pp. 320-321**

Note on the other files in this repo

Experiment 31.pdf and Exp 32 - Esters.pdf are **Chemistry** practical documents (ester preparation) and the html file is an unrelated web page — none of them relate to Human Biology Task 6.