

# A new proof of non-Cohen-Macaulayness of Bertin's example

Takuma Seno

Bertin's example is famous as the first known Noetherian UFD that is not Cohen-Macaulay. In the example, she employed a ring of invariants and proved that the ring is not Cohen-Macaulay by calculating a homogeneous system of parameter and generators of it. In this talk, we give a new proof by arguments on ring theoretic properties and describe a generalization of our new method of proof.

## REFERENCES

- [1] A. Braun, On the Gorenstein property for modular invariants, *J.Algebra*, **345** (2011), 81–99.
- [2] G. Ellingsrud and T. Skjelbred, Profondeur d'anneaux d'invariants en caractéristique  $p$ , *Compositio Math*, **41** (1980), 233–244.
- [3] G. Kemper, On the Cohen-Macaulay Property of Modular Invariant Rings, *J.Algebra*, **215** (1999), 330–351.
- [4] H. E. A. Eddy Campbell and David L. Wehlau, *Modular Invariant Theory*, Springer, 2011.
- [5] P. Fleischmann and C. Woodcock, Relative invariants, ideal classes and quasicanonical modules of modular rings of invariants, *J. Algebra*, **348** (2011), 110–134.
- [6] R. P. Stanley, Hilbert functions of Graded Algebra, *Adv. in Math*, **28** (1978) 57–83.
- [7] W. Bruns and J. Herzog, *Cohen-Macaulay Rings*, Cambridge University Press, 1993.