

NCCRS OF TORIC SINGULARITIES WITH DIVISOR CLASS RANK ONE

RYU TOMONAGA

The notion of non-commutative crepant resolutions (NCCRs) is introduced in [17] as a virtual space of crepant resolutions of a given Gorenstein normal singularity. After that, NCCRs turn out to have deep connections with Cohen-Macaulay representation theory [9, 23], higher Auslander-Reiten theory [5, 10, 14], Calabi-Yau algebras [11] and additive categorification of cluster algebras [1, 3, 6]. The existence of NCCRs is proven for quotient singularities [10, 16], compound Du Val singularities having crepant resolutions [18], Du Val del Pezzo cones [15, 17], some toric singularities and so on [4, 7, 19]. In these cases, NCCRs give rich information to their geometry, derived categories and Cohen-Macaulay representations.

For toric singularities, it is convenient to restrict ourselves to toric NCCRs which are NCCRs given by a direct sum of divisorial modules. For a Gorenstein toric singularity R , the existence of toric NCCRs is proven if $\dim R \leq 3$ [2], $\text{Cl}(R)$ is torsion, $\text{Cl}(R) \cong \mathbb{Z}$ [17] or some other cases [8, 13, 20, 21]. We remark here that there exists a Gorenstein toric singularity R with $\dim R = 4$ and $\text{Cl}(R) \cong \mathbb{Z}^2$ such that R has an NCCR, but has *no* toric NCCRs [19, 20]. Based on these facts, it is conjectured that all Gorenstein toric singularities have NCCRs.

In this talk, we prove the existence of toric NCCRs of Gorenstein toric singularities R with $\text{rk Cl } R = 1$. Remark that we include the case when $\text{Cl}(R)$ has a torsion. Moreover, we give a classification of toric NCCRs which is new even when $\text{Cl}(R) \cong \mathbb{Z}$. To state our main result, we prepare some notations. Let $\vec{x}_1, \dots, \vec{x}_{d+2} \in \text{Cl}(R)$ be weights. If we put $\pi: \text{Cl}(R) \rightarrow \text{Cl}(R)/\text{Cl}(R)_{\text{tors}}$, then we

may assume that $\pi(\vec{x}_i) \begin{cases} > 0 & (1 \leq i \leq l) \\ < 0 & (l+1 \leq i \leq l+l') \\ = 0 & (l+l'+1 \leq i \leq d+2) \end{cases}$ holds. Here, we can prove $l, l' \geq 2$. We put

$H := G/(\sum_{i=l'+1}^{d+2} \mathbb{Z}\vec{x}_i)$ and let $q: G \rightarrow H$ be the natural surjection. Then we can define a partial order on H as

$$h_1 \geq h_2 \Leftrightarrow h_1 - h_2 \in \sum_{i=1}^l \mathbb{Z}_{\geq 0} q(\vec{x}_i) + \sum_{j=1}^{l'} \mathbb{Z}_{\geq 0} q(-x_{l+j}) \subseteq H \text{ for } h_1, h_2 \in H.$$

Put $p := \sum_{i=1}^l q(\vec{x}_i) = \sum_{j=1}^{l'} q(-x_{l+j}) \in H$. For a finite subset $J \subseteq \text{Cl}(R)$, let $M_J \in \text{ref } R$ denote the direct sum of divisorial modules corresponding to elements in J .

Theorem 0.1. Let R be a Gorenstein toric singularity with $\text{rk Cl}(R) = 1$. In the above notations, we have a bijection between the following sets.

- (1) The set of non-trivial upper sets in H .
- (2) $\{J \subseteq \text{Cl}(R) \mid M_J \text{ gives an NCCR.}\}$

A bijection from (1) to (2) is given by $I \mapsto q^{-1}(J(I))$ where $J(I) := I \cap (I^c + p) \subseteq H$.

According to our classification, we can find R with $\text{Cl}(R) \cong \mathbb{Z}$ having a toric NCCR other than one constructed in [17].

Finally, we investigate Iyama-Wemyss mutations of our toric NCCRs. The operation of mutations of NCCRs is introduced in [12] and they proved that two NCCRs connected by mutations are derived equivalent. Moreover, for compound Du Val singularities, the mutations of NCCRs correspond to the flips of crepant resolutions [22]. For these reasons, it is a fundamental task to compute mutations of a given NCCR. We show that in our setting, Iyama-Wemyss mutations are compatible with mutations of upper sets.

Theorem 0.2. In the notation of Theorem 0.1, let $I \subseteq H$ be a non-trivial upper set. Take a minimal element $m \in I$ and write $\mu_m^-(I) := I \setminus \{m\}$. Put $M := M_{q^{-1}(J(I) \setminus \{m\})}$. Then we have

$$(\mu_M^+)^{l'-1}(M_{q^{-1}(J(I))}) = M_{q^{-1}(J(\mu_m^-(I)))} = (\mu_M^-)^{l-1}(M_{q^{-1}(J(I))}).$$

Combining those our results with combinatorics of upper sets, we obtain the following corollary.

Corollary 0.3. Let R be a Gorenstein toric singularity with $\text{rk Cl}(R) = 1$. Then all toric NCCRs of R are connected by iterated Iyama-Wemyss mutations. In particular, they are all derived equivalent to each other.

REFERENCES

1. Claire Amiot, Osamu Iyama, and Idun Reiten, *Stable categories of Cohen-Macaulay modules and cluster categories: Dedicated to Ragnar-Olaf Buchweitz on the occasion of his sixtieth birthday*, American Journal of Mathematics **137** (2015), no. 3, 813–857.
2. Nathan Broomhead, *Dimer models and Calabi-Yau algebras*, American Mathematical Society **215** (2012), no. 1011.
3. Aslak Bakke Buan, Bethany Marsh, Markus Reineke, Idun Reiten, and Gordana Todorov, *Tilting theory and cluster combinatorics*, Advances in mathematics **204** (2006), no. 2, 572–618.
4. Norihiro Hanihara, *Non-commutative resolutions for Segre products and Cohen-Macaulay rings of hereditary representation type*, arXiv:2303.14625, 2023.
5. ———, *Higher hereditary algebras arising from some toric singularities*, arXiv:2412.19040, 2024.
6. Norihiro Hanihara and Osamu Iyama, *Enhanced Auslander-Reiten duality and tilting theory for singularity categories*, arXiv:2209.14090, 2022.
7. Wahei Hara, *Non-commutative crepant resolution of minimal nilpotent orbit closures of type A and Mukai flops*, Advances in mathematics **318** (2017), 355–410.
8. Akihiro Higashitani and Nakajima Yusuke, *Conic divisorial ideals of hibi rings and their applications to non-commutative crepant resolutions*, Selecta Mathematica **25** (2019), no. 5, 78.
9. Osamu Iyama, *Auslander correspondence*, Advances in mathematics **210** (2007), no. 1, 51–82.
10. ———, *Higher-dimensional Auslander-Reiten theory on maximal orthogonal subcategories*, Advances in mathematics **210** (2007), no. 1, 22–50.
11. Osamu Iyama and Idun Reiten, *Fomin-Zelevinsky mutation and tilting modules over Calabi-Yau algebras*, American Journal of Mathematics **130** (2008), no. 4, 1087–1149.
12. Osamu Iyama and Michael Wemyss, *Maximal modifications and Auslander-Reiten duality for non-isolated singularities*, Inventiones mathematicae **197** (2014), no. 3, 521–586.
13. Koji Matsushita, *Conic divisorial ideals of toric rings and applications to Hibi rings and stable set rings*, (2022).
14. Ryu Tomonaga, *Higher hereditary algebras and toric Fano stacks with Picard number at most two*, in preparation.
15. ———, *Weak del Pezzo surfaces yield 2-hereditary algebras and 3-Calabi-Yau algebras*, arXiv:2510.26199.
16. ———, *Cohen-Macaulay representations of invariant subrings*, arXiv:2403.19282, 2024.
17. Michel Van den Bergh, *Non-commutative crepant resolutions*, The Legacy of Niels Henrik Abel: The Abel Bicentennial, Oslo, 2002. Berlin, Heidelberg: Springer Berlin Heidelberg (2004), 749–770.
18. ———, *Three-dimensional flops and noncommutative rings*, Duke Math. J. **122** (2004), no. 3, 423–455.
19. Špela Špenko and Michel Van den Bergh, *Non-commutative resolutions of quotient singularities for reductive groups*, Inventiones mathematicae **210** (2017), 3–67.
20. ———, *Non-commutative crepant resolutions for some toric singularities i*, International Mathematics Research Notices **21** (2020), 8120–8138.
21. ———, *Non-commutative crepant resolutions for some toric singularities ii*, Journal of Noncommutative Geometry **14** (2020), no. 1, 73–103.
22. Michael Wemyss, *Flops and clusters in the homological minimal model programme*, Inventiones mathematicae **211** (2018), no. 2, 435–521.
23. Yuji Yoshino, *Cohen-Macaulay modules over Cohen-Macaulay rings*, vol. 146, Cambridge University Press, 1990.