

KE-CLOSED SUBCATEGORIES OVER A COMMUTATIVE NOETHERIAN RING

TOSHINORI KOBAYASHI

This talk is based on joint work with S. Saito [KS24, KS25].

Understanding subcategories has been one of the central themes in ring theory. Gabriel [Gab62] achieved a landmark result by classifying Serre subcategories of the category of finitely generated modules over a commutative noetherian ring (or more generally coherent sheaves on a noetherian scheme). Beyond Serre subcategories, much subsequent work has examined other classes of subcategories. One important direction concerns *torsion-free classes* (subcategories closed under subobjects and extensions). For finitely generated modules over a commutative noetherian ring, Takahashi [Tak08] completely classified torsion-free classes in terms of the prime spectrum of the ring.

Motivated by these classification results, we focus on subcategories that are closed under basic categorical operations. Let R be a commutative noetherian ring. Let $\mathbf{mod} R$ denote the category of finitely generated R -modules. A subcategory \mathcal{X} of $\mathbf{mod} R$ is called *KE-closed* respectively, *CE-closed* or *IE-closed*) if \mathcal{X} is closed under kernels and extensions (respectively, under cokernels and extensions, or under images and extensions). It was recently shown by Stanley–Wang [SW11] and Enomoto [Eno] that CE-closed and IE-closed subcategories coincide with Serre subcategories and torsion-free classes, respectively. In contrast, there exists a KE-closed subcategory which is not a torsion-free class.

The aim of this talk is to introduce our classification of KE-closed subcategories of $\mathbf{mod} R$. The outline consists of two parts. First, we associate a function on $\mathrm{Spec} R$ with each KE-closed subcategory of $\mathbf{mod} R$, and show that this function completely determines the original subcategory. Second, we investigate when a function arises from KE-closed subcategories. To this end, we introduce the notion of an n -Bass function for each $n \geq 0$, and we establish a bijection between the set of KE-closed subcategories and the set of 2-Bass functions provided that R is (S_2) -excellent in the sense of Česnavičius [Čes21].

REFERENCES

- [Čes21] K. Česnavičius, *Macaulayfication of Noetherian schemes*, Duke Math. J. **170** (2021), no. 7, 1419–1455.
- [Eno] H. Enomoto, *IE-closed subcategories of commutative rings are torsion-free classes*, preprint (2023), arXiv:2304.03260v2.
- [Gab62] P. Gabriel, *Des catégories abéliennes*, Bull. Soc. Math. France **90** (1962), 323–448.
- [KS24] T. Kobayashi, S. Saito, *When are KE-closed subcategories torsion-free classes?*, Math. Z. **307** (2024), no. 4, Paper No. 65.
- [KS25] T. Kobayashi, S. Saito, *Classifying KE-closed subcategories over a commutative noetherian ring*, preprint, arXiv:2509.05767.
- [SW11] D. Stanley, B. Wang, *Classifying subcategories of finitely generated modules over a Noetherian ring*, J. Pure Appl. Algebra **215** (2011), no. 11, 2684–2693.
- [Tak08] R. Takahashi, *Classifying subcategories of modules over a commutative Noetherian ring*, J. Lond. Math. Soc. (2) **78** (2008), no. 3, 767–782.

SCHOOL OF SCIENCE AND TECHNOLOGY, MEIJI UNIVERSITY, 1-1-1 HIGASHIMITA, TAMAKU, KAWASAKI 214-8571, JAPAN

Email address: tkobayashi@meiji.ac.jp