

Complexity of Reasoning in Kleene and Action Algebras

Stepan Kuznetsov

ESSLI 2022 · Galway, Ireland · Gallimh, Éire

Lecture 3

- Recall that we are developing a finitary and cut-free proof system for deciding inclusion of regular languages: $A \rightarrow B$ should be derivable iff $v(A) \subseteq v(B)$.

- Recall that we are developing a finitary and cut-free proof system for deciding inclusion of regular languages: $A \rightarrow B$ should be derivable iff $v(A) \subseteq v(B)$.
- Such a calculus will be the leftmost fragment of \mathbf{HKA}_∞ (Das & Pous 2017).

- Recall that we are developing a finitary and cut-free proof system for deciding inclusion of regular languages: $A \rightarrow B$ should be derivable iff $v(A) \subseteq v(B)$.
- Such a calculus will be the leftmost fragment of \mathbf{HKA}_∞ (Das & Pous 2017).
- \mathbf{HKA}_∞ is a hypersequent calculus, whose derivable objects are of the form $\Pi \rightarrow \mathcal{X}$, where Π is a sequence of formulae (reg. exp.) and $\mathcal{X} = \Delta_1 \mid \dots \mid \Delta_k$ is a multiset of sequences of formulae.

Leftmost \mathbf{HKA}_∞

The rules of \mathbf{HKA}_∞ , which allow only leftmost proofs, are as follows:

$$\begin{array}{c}
 \frac{\Gamma \rightarrow \mathcal{X} \quad A, A^*, \Gamma \rightarrow \mathcal{X}}{A^*, \Gamma \rightarrow \mathcal{X}} *L \qquad \frac{A, B, \Gamma \rightarrow \mathcal{X}}{A \cdot B, \Gamma \rightarrow \mathcal{X}} \cdot L \\
 \\
 \frac{A, \Gamma \rightarrow \mathcal{X} \quad B, \Gamma \rightarrow \mathcal{X}}{A + B, \Gamma \rightarrow \mathcal{X}} +L \qquad \frac{\Gamma \rightarrow \mathcal{X}}{1, \Gamma \rightarrow \mathcal{X}} 1L \qquad \frac{}{0, \Gamma \rightarrow \mathcal{X}} 0L \\
 \\
 \frac{\Pi \rightarrow \Delta \mid A, A^*, \Delta \mid \mathcal{X}}{\Pi \rightarrow A^*, \Delta \mid \mathcal{X}} *R \qquad \frac{\Pi \rightarrow A, B, \Delta \mid \mathcal{X}}{\Pi \rightarrow A \cdot B, \Delta \mid \mathcal{X}} \cdot R \\
 \\
 \frac{\Pi \rightarrow A, \Delta \mid B, \Delta \mid \mathcal{X}}{\Pi \rightarrow A + B, \Delta \mid \mathcal{X}} +R \qquad \frac{\Pi \rightarrow \Delta \mid \mathcal{X}}{\Pi \rightarrow 1, \Delta \mid \mathcal{X}} 1R \\
 \\
 \frac{}{\varepsilon \rightarrow \varepsilon} Ax \qquad \frac{\Gamma \rightarrow \Delta_1 \mid \dots \mid \Delta_m}{a, \Gamma \rightarrow a, \Delta_1 \mid \dots \mid a, \Delta_m} K \\
 \\
 \frac{\Pi \rightarrow \mathcal{X}}{\Pi \rightarrow \Delta \mid \mathcal{X}} WR \qquad \frac{\Pi \rightarrow \Delta \mid \Delta \mid \mathcal{X}}{\Pi \rightarrow \Delta \mid \mathcal{X}} CR
 \end{array}$$

- HKA_∞ allows non-well-founded proofs, i.e., infinite paths starting from the goal sequent.

- HKA_∞ allows non-well-founded proofs, i.e., infinite paths starting from the goal sequent.
- Such proofs should obey the **fairness condition**: any infinite path should traverse $*L$, turning right, infinitely many times.

- HKA_∞ allows non-well-founded proofs, i.e., infinite paths starting from the goal sequent.
- Such proofs should obey the **fairness condition**: any infinite path should traverse $*L$, turning right, infinitely many times.
- It is illegal to produce an infinite path using only $*R$.

- HKA_∞ allows non-well-founded proofs, i.e., infinite paths starting from the goal sequent.
- Such proofs should obey the **fairness condition**: any infinite path should traverse $*L$, turning right, infinitely many times.
- It is illegal to produce an infinite path using only $*R$.
- In a **circular** proof, each infinite branch has a repeated copy of the sequent already seen before, with an $*L$ between these occurrences.

- HKA_∞ allows non-well-founded proofs, i.e., infinite paths starting from the goal sequent.
- Such proofs should obey the **fairness condition**: any infinite path should traverse $*L$, turning right, infinitely many times.
- It is illegal to produce an infinite path using only $*R$.
- In a **circular** proof, each infinite branch has a repeated copy of the sequent already seen before, with an $*L$ between these occurrences.
- In such cases, further development of the path is replaced by a **backlink**, and the whole structure becomes finite.

Theorem (Das & Pous 2017)

A sequent is derivable in \mathbf{HKA}_∞ iff it is valid in the standard interpretation. Moreover, any valid sequent has a circular leftmost proof.

- The “only if” direction (soundness) was proved yesterday.

Productivity Lemma

- For completeness, we use the following lemma.

Productivity Lemma

- For completeness, we use the following lemma.

Lemma

If there is a subderivation

$$\frac{\frac{\Gamma \rightarrow A^*, \Delta \mid \mathcal{X}}{\vdots \pi}}{\frac{\Gamma \rightarrow \Delta \mid A, A^*, \Delta \mid \mathcal{Y}}{\Gamma \rightarrow A^*, \Delta \mid \mathcal{Y}} *R}$$

where π consists only of right logical rules, and the sequent $\Gamma \rightarrow A^, \Delta \mid \mathcal{Y}$ is valid in the standard interpretation, then so is $\Gamma \rightarrow \mathcal{X}$ (i.e., A^*, Δ there is redundant).*

Productivity Lemma

- For completeness, we use the following lemma.

Lemma

If there is a subderivation

$$\frac{\frac{\Gamma \rightarrow A^*, \Delta \mid \mathcal{X}}{\vdots \pi}}{\frac{\Gamma \rightarrow \Delta \mid A, A^*, \Delta \mid \mathcal{Y}}{\Gamma \rightarrow A^*, \Delta \mid \mathcal{Y}} *R}$$

where π consists only of right logical rules, and the sequent $\Gamma \rightarrow A^, \Delta \mid \mathcal{Y}$ is valid in the standard interpretation, then so is $\Gamma \rightarrow \mathcal{X}$ (i.e., A^*, Δ there is redundant).*

- This was also established yesterday.

Now we are going to prove the following two statements, which give the necessary result:

1. If $v(A) \subseteq v(B)$, then $A \rightarrow B$ has a leftmost proof in \mathbf{HKA}_∞ .
2. Each leftmost proof can be transformed into a circular one.

Constructing Leftmost Proofs

- Let a hypersequent $\Pi \rightarrow \mathcal{X}$ be valid in the standard model (with \cdot interpreted as \cdot and $|$ interpreted as $+$). We shall construct a leftmost proof for it.

Constructing Leftmost Proofs

- Let a hypersequent $\Pi \rightarrow \mathcal{X}$ be valid in the standard model (with \cdot interpreted as \cdot and $|$ interpreted as $+$). We shall construct a leftmost proof for it.
- First (**Stage 1**), we eagerly apply leftmost left logical rules.

Constructing Leftmost Proofs

- Let a hypersequent $\Pi \rightarrow \mathcal{X}$ be valid in the standard model (with \cdot interpreted as \cdot and $|$ interpreted as $+$). We shall construct a leftmost proof for it.
- First (**Stage 1**), we eagerly apply leftmost left logical rules.
- The rules are invertible, so all leaves are valid in the standard interpretation.

Constructing Leftmost Proofs

- Let a hypersequent $\Pi \rightarrow \mathcal{X}$ be valid in the standard model (with \cdot interpreted as \cdot and $|$ interpreted as $+$). We shall construct a leftmost proof for it.
- First (**Stage 1**), we eagerly apply leftmost left logical rules.
- The rules are invertible, so all leaves are valid in the standard interpretation.
- Each leaf is either $\rightarrow \mathcal{X}$ or $a, \Gamma \rightarrow \mathcal{X}$.

Constructing Leftmost Proofs

- Let a hypersequent $\Pi \rightarrow \mathcal{X}$ be valid in the standard model (with \cdot interpreted as \cdot and $|$ interpreted as $+$). We shall construct a leftmost proof for it.
- First (**Stage 1**), we eagerly apply leftmost left logical rules.
- The rules are invertible, so all leaves are valid in the standard interpretation.
- Each leaf is either $\rightarrow \mathcal{X}$ or $a, \Gamma \rightarrow \mathcal{X}$.
- An important thing to mention here is that Stage 1 could generate infinite paths, e.g.:

$$\begin{array}{c}
 \vdots \\
 \hline
 1^*, \Gamma \rightarrow \mathcal{X} \\
 \hline
 \Gamma \rightarrow \mathcal{X} \quad 1, 1^*, \Gamma \rightarrow \mathcal{X} \quad 1L \\
 \hline
 1^*, \Gamma \rightarrow \mathcal{X} \quad *L
 \end{array}$$

Constructing Leftmost Proofs

- Such infinite paths are fine, since they obey the fairness condition (i.e., traverse $*L$ infinitely often).

Constructing Leftmost Proofs

- Such infinite paths are fine, since they obey the fairness condition (i.e., traverse $*L$ infinitely often).
- We shall show that they are actually regular, but by now we do not care about it.

Constructing Leftmost Proofs

- Such infinite paths are fine, since they obey the fairness condition (i.e., traverse $*L$ infinitely often).
- We shall show that they are actually regular, but by now we do not care about it.
- On **Stage 2** we apply right logical rules.

Constructing Leftmost Proofs

- Such infinite paths are fine, since they obey the fairness condition (i.e., traverse $*L$ infinitely often).
- We shall show that they are actually regular, but by now we do not care about it.
- On **Stage 2** we apply right logical rules.
- These rules are also invertible.

Constructing Leftmost Proofs

- Such infinite paths are fine, since they obey the fairness condition (i.e., traverse $*L$ infinitely often).
- We shall show that they are actually regular, but by now we do not care about it.
- On **Stage 2** we apply right logical rules.
- These rules are also invertible.
- However, here we cannot go eagerly, since this would result in proofs violating fairness:

$$\frac{\frac{\vdots}{\Pi \rightarrow \Delta \mid 1^*, \Delta \mid \mathcal{X}}}{\Pi \rightarrow \Delta \mid 1, 1^*, \Delta \mid \mathcal{X}} 1R \quad *R$$

Constructing Leftmost Proofs

- Due to this reason, we impose the following restriction: we apply leftmost right logical rules to all occurrences, except for A^* 's which were already obtained by decomposition ($*R$).

Constructing Leftmost Proofs

- Due to this reason, we impose the following restriction: we apply leftmost right logical rules to all occurrences, except for A^* 's which were already obtained by decomposition ($*R$).
- Now in our leaves we have multisets \mathcal{X} , and each sequent of \mathcal{X} is either b, Δ , or empty, or A^*, Δ , where A^* was obtained by decomposition.

Constructing Leftmost Proofs

- Due to this reason, we impose the following restriction: we apply leftmost right logical rules to all occurrences, except for A^* 's which were already obtained by decomposition ($*R$).
- Now in our leaves we have multisets \mathcal{X} , and each sequent of \mathcal{X} is either b, Δ , or empty, or A^*, Δ , where A^* was obtained by decomposition.
- The latter get weakened out by Productivity Lemma (weakenings form **Stage 3**).

Constructing Leftmost Proofs

- Due to this reason, we impose the following restriction: we apply leftmost right logical rules to all occurrences, except for A^* 's which were already obtained by decomposition ($*R$).
- Now in our leaves we have multisets \mathcal{X} , and each sequent of \mathcal{X} is either b, Δ , or empty, or A^*, Δ , where A^* was obtained by decomposition.
- The latter get weakened out by Productivity Lemma (weakenings form **Stage 3**).
- Recall that our leaves are $\rightarrow \mathcal{X}$ or $a, \Gamma \rightarrow \mathcal{X}$.

Constructing Leftmost Proofs

- Due to this reason, we impose the following restriction: we apply leftmost right logical rules to all occurrences, except for A^* 's which were already obtained by decomposition ($*R$).
- Now in our leaves we have multisets \mathcal{X} , and each sequent of \mathcal{X} is either b, Δ , or empty, or A^*, Δ , where A^* was obtained by decomposition.
- The latter get weakened out by Productivity Lemma (weakenings form **Stage 3**).
- Recall that our leaves are $\rightarrow \mathcal{X}$ or $a, \Gamma \rightarrow \mathcal{X}$.
- In the first case, by validity, we have $\varepsilon \in v(\mathcal{X})$. Thus, \mathcal{X} contains the empty sequence. Everything else gets weakened out.

- For the second case, we weaken all sequences in \mathcal{X} , which are the empty sequence or begin with $b \neq a$.

Constructing Leftmost Proofs

- For the second case, we weaken all sequences in \mathcal{X} , which are the empty sequence or begin with $b \neq a$.
- Next, we apply K (**Stage 4**).

- For the second case, we weaken all sequences in \mathcal{X} , which are the empty sequence or begin with $b \neq a$.
- Next, we apply K (**Stage 4**).
- Above K , we have new valid sequents, on which we again start with Stage 1.

- For the second case, we weaken all sequences in \mathcal{X} , which are the empty sequence or begin with $b \neq a$.
- Next, we apply K (**Stage 4**).
- Above K , we have new valid sequents, on which we again start with Stage 1.
- Notice that there is not (transfinite) induction here, we do not care for reducing any sort of parameter.

Constructing Leftmost Proofs

- For the second case, we weaken all sequences in \mathcal{X} , which are the empty sequence or begin with $b \neq a$.
- Next, we apply K (**Stage 4**).
- Above K , we have new valid sequents, on which we again start with Stage 1.
- Notice that there is not (transfinite) induction here, we do not care for reducing any sort of parameter.
- The only thing we should care about is that the proof never gets stuck, and fairness is maintained.

- To prove fairness, for each infinite path we consider two possibilities.

Constructing Leftmost Proofs

- To prove fairness, for each infinite path we consider two possibilities.
- Case 1: the branch eventually ends in one stage. Then it should be Stage 1, and fairness is maintained (see above).

- To prove fairness, for each infinite path we consider two possibilities.
- Case 1: the branch eventually ends in one stage. Then it should be Stage 1, and fairness is maintained (see above).
- Case 2: the branch traverses Stages 1–4 infinitely.

- To prove fairness, for each infinite path we consider two possibilities.
- Case 1: the branch eventually ends in one stage. Then it should be Stage 1, and fairness is maintained (see above).
- Case 2: the branch traverses Stages 1–4 infinitely.
- In this case, consider the size of the antecedent. This size gets strictly reduced at each Stage 4, thus, it should be restored, which can be done only by $*L$.

- Now let us regularise our leftmost proof.

Making Proofs Circular

- Now let us regularise our leftmost proof.
- For this, we shall need yet another order, now on occurrences of subformulae in the goal sequent.

Making Proofs Circular

- Now let us regularise our leftmost proof.
- For this, we shall need yet another order, now on occurrences of subformulae in the goal sequent.
- We have $A \preccurlyeq B$, if A is either a subformula of B or is located to the left of B . This is a linear (total) order.

Making Proofs Circular

- Now let us regularise our leftmost proof.
- For this, we shall need yet another order, now on occurrences of subformulae in the goal sequent.
- We have $A \preccurlyeq B$, if A is either a subformula of B or is located to the left of B . This is a linear (total) order.
- Moreover, thanks to subformula property, the order can be propagated to occurrences in sequents throughout the proof.

Making Proofs Circular

- Now let us regularise our leftmost proof.
- For this, we shall need yet another order, now on occurrences of subformulae in the goal sequent.
- We have $A \preccurlyeq B$, if A is either a subformula of B or is located to the left of B . This is a linear (total) order.
- Moreover, thanks to subformula property, the order can be propagated to occurrences in sequents throughout the proof.

Lemma

In a leftmost proof, each sequence is strictly increasing w.r.t. \preccurlyeq .

Making Proofs Circular

- Now let us regularise our leftmost proof.
- For this, we shall need yet another order, now on occurrences of subformulae in the goal sequent.
- We have $A \preccurlyeq B$, if A is either a subformula of B or is located to the left of B . This is a linear (total) order.
- Moreover, thanks to subformula property, the order can be propagated to occurrences in sequents throughout the proof.

Lemma

In a leftmost proof, each sequence is strictly increasing w.r.t. \preccurlyeq .

- Corollary: the length of such a sequence is bounded by the size of the goal sequent.

Making Proofs Circular

- Finally, using WR and CR , we can assure that in each antecedent we have no more than 2 copies of a sequence.

Making Proofs Circular

- Finally, using WR and CR , we can assure that in each antecedent we have no more than 2 copies of a sequence.
- Therefore, the total number of sequents which could appear in our proof is finite.

Making Proofs Circular

- Finally, using WR and CR , we can assure that in each antecedent we have no more than 2 copies of a sequence.
- Therefore, the total number of sequents which could appear in our proof is finite.
- Thus, each infinite path will eventually reach a sequent which already occurred earlier.

Making Proofs Circular

- Finally, using WR and CR , we can assure that in each antecedent we have no more than 2 copies of a sequence.
- Therefore, the total number of sequents which could appear in our proof is finite.
- Thus, each infinite path will eventually reach a sequent which already occurred earlier.
- We identify two such instances with an application of $*L$ in between, and make a cycle.

Making Proofs Circular

- Finally, using WR and CR , we can assure that in each antecedent we have no more than 2 copies of a sequence.
- Therefore, the total number of sequents which could appear in our proof is finite.
- Thus, each infinite path will eventually reach a sequent which already occurred earlier.
- We identify two such instances with an application of $*L$ in between, and make a cycle.
- This makes \mathbf{KA}_ω r.e., and by Post's theorem we obtain decidability.

Making Proofs Circular

- Finally, using WR and CR , we can assure that in each antecedent we have no more than 2 copies of a sequence.
- Therefore, the total number of sequents which could appear in our proof is finite.
- Thus, each infinite path will eventually reach a sequent which already occurred earlier.
- We identify two such instances with an application of $*L$ in between, and make a cycle.
- This makes \mathbf{KA}_ω r.e., and by Post's theorem we obtain decidability.
- A more accurate consideration shows that each path reaches its cycle in polynomial number of steps, which makes \mathbf{KA}_ω PSPACE.

Example

$$\begin{array}{c}
 \frac{\varepsilon \rightarrow \varepsilon}{\varepsilon \rightarrow \varepsilon \mid aa, (aa)^* \mid a(aa)^*} \text{ } WR \\
 \frac{\varepsilon \rightarrow (aa)^* \mid a(aa)^*}{\varepsilon \rightarrow (aa)^* + a(aa)^*} \text{ } *R \\
 \frac{\varepsilon \rightarrow (aa)^* + a(aa)^*}{a^* \rightarrow (aa)^* + a(aa)^*} \text{ } +R
 \end{array}
 \qquad
 \begin{array}{c}
 \frac{a^* \rightarrow a, (aa)^* \mid (aa)^*}{\vdots} \\
 \frac{a^* \rightarrow a, (aa)^* \mid (aa)^*}{a, a^* \rightarrow a, a, (aa)^* \mid a, (aa)^*} \text{ } K \\
 \frac{a, a^* \rightarrow a, a, (aa)^* \mid a, (aa)^*}{a, a^* \rightarrow \varepsilon \mid a, a, (aa)^* \mid a, (aa)^*} \text{ } WR \\
 \frac{a, a^* \rightarrow \varepsilon \mid a, a, (aa)^* \mid a, (aa)^*}{a, a^* \rightarrow \varepsilon \mid aa, (aa)^* \mid a(aa)^*} \text{ } \cdot R \\
 \frac{a, a^* \rightarrow \varepsilon \mid aa, (aa)^* \mid a(aa)^*}{a, a^* \rightarrow (aa)^* \mid a(aa)^*} \text{ } *R \\
 \frac{a, a^* \rightarrow (aa)^* \mid a(aa)^*}{a, a^* \rightarrow (aa)^* + a(aa)^*} \text{ } +R \\
 \frac{a, a^* \rightarrow (aa)^* + a(aa)^*}{a^* \rightarrow (aa)^* + a(aa)^*} \text{ } *L
 \end{array}$$

- The situation changes dramatically when we allow *hypotheses*.

- The situation changes dramatically when we allow *hypotheses*.
- Let \mathcal{H} be a finite set of sequents of the form $A \rightarrow B$, and we ask whether a sequent $C \rightarrow D$ follows from \mathcal{H} .

- The situation changes dramatically when we allow *hypotheses*.
- Let \mathcal{H} be a finite set of sequents of the form $A \rightarrow B$, and we ask whether a sequent $C \rightarrow D$ follows from \mathcal{H} .
- We could think of this both syntactically and semantically.

- The situation changes dramatically when we allow *hypotheses*.
- Let \mathcal{H} be a finite set of sequents of the form $A \rightarrow B$, and we ask whether a sequent $C \rightarrow D$ follows from \mathcal{H} .
- We could think of this both syntactically and semantically.
- Syntax: there exists a (possibly infinite) derivation in \mathbf{KA}_ω , where sequents from \mathcal{H} are allowed to be used as axioms.

- The situation changes dramatically when we allow *hypotheses*.
- Let \mathcal{H} be a finite set of sequents of the form $A \rightarrow B$, and we ask whether a sequent $C \rightarrow D$ follows from \mathcal{H} .
- We could think of this both syntactically and semantically.
- Syntax: there exists a (possibly infinite) derivation in \mathbf{KA}_ω , where sequents from \mathcal{H} are allowed to be used as axioms.
 - Cut is also allowed.

- The situation changes dramatically when we allow *hypotheses*.
- Let \mathcal{H} be a finite set of sequents of the form $A \rightarrow B$, and we ask whether a sequent $C \rightarrow D$ follows from \mathcal{H} .
- We could think of this both syntactically and semantically.
- Syntax: there exists a (possibly infinite) derivation in \mathbf{KA}_ω , where sequents from \mathcal{H} are allowed to be used as axioms.
 - Cut is also allowed.
- Semantics: for any interpretation, on a $*$ -continuous Kleene algebra, where all sequents from \mathcal{H} are valid, $C \rightarrow D$ is also valid.

- The situation changes dramatically when we allow *hypotheses*.
- Let \mathcal{H} be a finite set of sequents of the form $A \rightarrow B$, and we ask whether a sequent $C \rightarrow D$ follows from \mathcal{H} .
- We could think of this both syntactically and semantically.
- Syntax: there exists a (possibly infinite) derivation in \mathbf{KA}_ω , where sequents from \mathcal{H} are allowed to be used as axioms.
 - Cut is also allowed.
- Semantics: for any interpretation, on a $*$ -continuous Kleene algebra, where all sequents from \mathcal{H} are valid, $C \rightarrow D$ is also valid.
 - Notice that this is an abstract $*$ -continuous KA, not the standard interpretation.

- There is a **strong soundness-and-completeness** theorem which states that these two notions are equivalent.

- There is a **strong soundness-and-completeness** theorem which states that these two notions are equivalent.
- The proof is by a standard Lindenbaum–Tarski canonical model argument.

- There is a **strong soundness-and-completeness** theorem which states that these two notions are equivalent.
- The proof is by a standard Lindenbaum–Tarski canonical model argument.
- Recall that an abstract $*$ -continuous KA is an algebraic structure in which axioms and rules of \mathbf{KA}_ω are valid.

- There is a **strong soundness-and-completeness** theorem which states that these two notions are equivalent.
- The proof is by a standard Lindenbaum–Tarski canonical model argument.
- Recall that an abstract \ast -continuous KA is an algebraic structure in which axioms and rules of \mathbf{KA}_ω are valid.
- One can consider only algebras of regular subsets of monoids (but not necessarily the free monoid Σ^\ast).

Complexity Results

The following complexity results for entailment in \ast -continuous KAs are due to Kozen (2002):

\mathcal{H} includes only monoid equations	Π_2^0 -complete
\mathcal{H} includes only length-preserving monoid equations	Π_1^0 -complete
\mathcal{H} is arbitrary	Π_1^1 -complete

Upper Bounds

- For arbitrary \mathcal{H} 's, we have one \forall quantifier over infinite structures (* -continuous KAs).

Upper Bounds

- For arbitrary \mathcal{H} 's, we have one \forall quantifier over infinite structures (\ast -continuous KAs).
- This can also be done syntactically, by considering derivations in \mathbf{KA}_ω .

Upper Bounds

- For arbitrary \mathcal{H} 's, we have one \forall quantifier over infinite structures (* -continuous KAs).
- This can also be done syntactically, by considering derivations in \mathbf{KA}_ω .
- For monoid equations, one can show that entailment in arbitrary KAs from such \mathcal{H} is equivalent to validity in $\text{REG}(\Sigma^*)/\mathcal{H}$ (i.e., the factor-algebra of the algebra of regular languages).

Upper Bounds

- For arbitrary \mathcal{H} 's, we have one \forall quantifier over infinite structures (\ast -continuous KAs).
- This can also be done syntactically, by considering derivations in \mathbf{KA}_ω .
- For monoid equations, one can show that entailment in arbitrary KAs from such \mathcal{H} is equivalent to validity in $\text{REG}(\Sigma^\ast)/\mathcal{H}$ (i.e., the factor-algebra of the algebra of regular languages).
- Now the validity of $C \rightarrow D$ is equivalent to

$$\forall x (x \in v(C) \supset \exists y (y \equiv_{\mathcal{H}} x \ \& \ y \in v(D))).$$

Upper Bounds

- For arbitrary \mathcal{H} 's, we have one \forall quantifier over infinite structures (\ast -continuous KAs).
- This can also be done syntactically, by considering derivations in \mathbf{KA}_ω .
- For monoid equations, one can show that entailment in arbitrary KAs from such \mathcal{H} is equivalent to validity in $\text{REG}(\Sigma^\ast)/\mathcal{H}$ (i.e., the factor-algebra of the algebra of regular languages).
- Now the validity of $C \rightarrow D$ is equivalent to

$$\forall x(x \in v(C) \supset \exists y(y \equiv_{\mathcal{H}} x \ \& \ y \in v(D))).$$

- This is Π_2^0 .

Upper Bounds

- For arbitrary \mathcal{H} 's, we have one \forall quantifier over infinite structures (\ast -continuous KAs).
- This can also be done syntactically, by considering derivations in \mathbf{KA}_ω .
- For monoid equations, one can show that entailment in arbitrary KAs from such \mathcal{H} is equivalent to validity in $\text{REG}(\Sigma^\ast)/\mathcal{H}$ (i.e., the factor-algebra of the algebra of regular languages).
- Now the validity of $C \rightarrow D$ is equivalent to

$$\forall x (x \in v(C) \supset \exists y (y \equiv_{\mathcal{H}} x \ \& \ y \in v(D))).$$

- This is Π_2^0 .
- If the equations are length-preserving, then the $\exists y$ quantifier is decidable, thus Π_1^0 .

- In order to prove lower bounds, we encode **Turing computations**, following Davis (1958).

Encoding Turing Machines

- In order to prove lower bounds, we encode **Turing computations**, following Davis (1958).
- A Turing machine (TM) operates on a **tape**, potentially infinite in both directions, and keeps some finite information in its **state**.

Encoding Turing Machines

- In order to prove lower bounds, we encode **Turing computations**, following Davis (1958).
- A Turing machine (TM) operates on a **tape**, potentially infinite in both directions, and keeps some finite information in its **state**.
- The current **configuration** of a TM will be encoded by $\vdash xqy \dashv$, where x, y are words and q is a state.

Encoding Turing Machines

- In order to prove lower bounds, we encode **Turing computations**, following Davis (1958).
- A Turing machine (TM) operates on a **tape**, potentially infinite in both directions, and keeps some finite information in its **state**.
- The current **configuration** of a TM will be encoded by $\vdash xqy \dashv$, where x, y are words and q is a state.
- Equations encoding the TM:

Encoding Turing Machines

- In order to prove lower bounds, we encode **Turing computations**, following Davis (1958).
- A Turing machine (TM) operates on a **tape**, potentially infinite in both directions, and keeps some finite information in its **state**.
- The current **configuration** of a TM will be encoded by $\vdash xqy \dashv$, where x, y are words and q is a state.
- Equations encoding the TM:
 1. for each transition $(p, a) \rightarrow (b, \text{right}, q)$, add $pa = bq$;

Encoding Turing Machines

- In order to prove lower bounds, we encode **Turing computations**, following Davis (1958).
- A Turing machine (TM) operates on a **tape**, potentially infinite in both directions, and keeps some finite information in its **state**.
- The current **configuration** of a TM will be encoded by $\vdash xqy \dashv$, where x, y are words and q is a state.
- Equations encoding the TM:
 1. for each transition $(p, a) \rightarrow (b, \text{right}, q)$, add $pa = bq$;
 2. for each transition $(p, a) \rightarrow (b, \text{left}, q)$, add $cpa = qcb$ for each letter c ;

Encoding Turing Machines

- In order to prove lower bounds, we encode **Turing computations**, following Davis (1958).
- A Turing machine (TM) operates on a **tape**, potentially infinite in both directions, and keeps some finite information in its **state**.
- The current **configuration** of a TM will be encoded by $\vdash xqy \dashv$, where x, y are words and q is a state.
- Equations encoding the TM:
 1. for each transition $(p, a) \rightarrow (b, \text{right}, q)$, add $pa = bq$;
 2. for each transition $(p, a) \rightarrow (b, \text{left}, q)$, add $cpa = qcb$ for each letter c ;
 3. $\vdash \sqcup = \vdash$ and $\sqcup \dashv = \dashv$.

Theorem

If t is the halting state, then $\vdash xqy \dashv \equiv_{\mathcal{H}} \vdash ztw \dashv$ if and only if $\vdash ztw \dashv$ is obtained from $\vdash xqy \dashv$ by the run of our TM.

- This is not obvious, because equiations (unlike TM execution) are undirected.

Theorem

If t is the halting state, then $\vdash xqy \dashv \equiv_{\mathcal{R}} \vdash ztw \dashv$ if and only if $\vdash ztw \dashv$ is obtained from $\vdash xqy \dashv$ by the run of our TM.

- This is not obvious, because equiations (unlike TM execution) are undirected.
- However, this is true, due to determinism of the TM and the fact that t is the halting state.

Theorem

If t is the halting state, then $\vdash xqy \dashv \equiv_{\mathcal{H}} \vdash ztw \dashv$ if and only if $\vdash ztw \dashv$ is obtained from $\vdash xqy \dashv$ by the run of our TM.

- This is not obvious, because equiations (unlike TM execution) are undirected.
- However, this is true, due to determinism of the TM and the fact that t is the halting state.
- An accurate proof is given, e.g., in Davis' textbook.

Theorem

If t is the halting state, then $\vdash xqy \dashv \equiv_{\mathcal{K}} \vdash ztw \dashv$ if and only if $\vdash ztw \dashv$ is obtained from $\vdash xqy \dashv$ by the run of our TM.

- This is not obvious, because equations (unlike TM execution) are undirected.
- However, this is true, due to determinism of the TM and the fact that t is the halting state.
- An accurate proof is given, e.g., in Davis' textbook.
- Thus, using only monoid equations, we can encode the halting problem, which gives Σ_0^1 -hardness even without Kleene star (in particular, for the finitary **KA** theory).

- Using Kleene star, we can do more and encode **totality** for Turing machines.

- Using Kleene star, we can do more and encode **totality** for Turing machines.
- Totality is a Π_2^0 -complete problem, and it means halting on **any** input.

- Using Kleene star, we can do more and encode **totality** for Turing machines.
- Totality is a Π_2^0 -complete problem, and it means halting on **any** input.
- The input will be a natural number n , encoded by a^n .

- Using Kleene star, we can do more and encode **totality** for Turing machines.
- Totality is a Π_2^0 -complete problem, and it means halting on **any** input.
- The input will be a natural number n , encoded by a^n .
- The encoding is as follows:

$$\mathcal{H} \models \vdash q_0 a^* \dashv \preceq \vdash t \dashv.$$

- Using Kleene star, we can do more and encode **totality** for Turing machines.
- Totality is a Π_2^0 -complete problem, and it means halting on **any** input.
- The input will be a natural number n , encoded by a^n .
- The encoding is as follows:

$$\mathcal{H} \models \vdash q_0 a^* \dashv \preceq \vdash t \dashv.$$

- For length-preserving equalities, we shall have totality for linearly bounded Turing machines, which models, e.g., totality of context-free grammars (which is Π_1^0 -hard).

- The more interesting result is Π_1^1 -hardness of entailment from arbitrary \mathcal{H} 's.

- The more interesting result is Π_1^1 -hardness of entailment from arbitrary \mathcal{H} 's.
- This is also due to Kozen (2002), and the Π_1^1 -hard problem encoded is **well-foundedness of infinite recursive graphs** (WF).

- The more interesting result is Π_1^1 -hardness of entailment from arbitrary \mathcal{H} 's.
- This is also due to Kozen (2002), and the Π_1^1 -hard problem encoded is **well-foundedness of infinite recursive graphs** (WF).
- Vertices of the graph G are natural numbers and the fact whether there is an edge (m, n) is determined by a TM.

- The more interesting result is Π_1^1 -hardness of entailment from arbitrary \mathcal{H} 's.
- This is also due to Kozen (2002), and the Π_1^1 -hard problem encoded is **well-foundedness of infinite recursive graphs** (WF).
- Vertices of the graph G are natural numbers and the fact whether there is an edge (m, n) is determined by a TM.
- The WF problem is to determine whether all paths in G which start from 0 are finite.

- In \mathcal{H} , we encode the TM which is used for defining graph G .

- In \mathcal{H} , we encode the TM which is used for defining graph G .
- s is the starting states, t , r , and u are halting states.

- In \mathcal{H} , we encode the TM which is used for defining graph G .
- s is the starting states, t , r , and u are halting states.
- The machine works as follows:

$$\vdash a^m s a^n \dashv \rightarrow_M^* \begin{cases} \vdash a^n t \dashv & \text{if } (m, n) \text{ is an edge} \\ \vdash r \dashv & \text{if not} \end{cases}$$

(u is used when the input is malformed).

- In \mathcal{H} , we encode the TM which is used for defining graph G .
- s is the starting states, t , r , and u are halting states.
- The machine works as follows:

$$\vdash a^m s a^n \dashv \rightarrow_M^* \begin{cases} \vdash a^n t \dashv & \text{if } (m, n) \text{ is an edge} \\ \vdash r \dashv & \text{if not} \end{cases}$$

(u is used when the input is malformed).

- Additionally, we include the inequation

$$t \preceq s a^*$$

into \mathcal{H} .

- Now we claim that WF holds iff \mathcal{H} entails $\vdash t \dashv \preceq \vdash r \dashv$.
- Suppose WF holds. This means
 $0 \in W = \{k \mid \text{all paths from } k \text{ are finite}\}.$
- We show that if $k \in W$, then \mathcal{H} entails $\vdash a^k t \dashv \preceq \vdash r \dashv$.
- In the simple case k has just no outgoing edges.
- Then we entail $\vdash a^k t \dashv \preceq \vdash a^k s a^* \dashv$, and next $\vdash a^k s a^* \dashv \preceq \vdash r \dashv$ is obtained by the ω -rule (no (k, m) is an edge).

- In the general case, we proceed by transfinite induction on W .
- If $k \in W$, then for each edge (k, m) we have $m \in W$, and, moreover, we may apply induction hypothesis to m .
- Thus, we have $\vdash a^m t \dashv \preceq \vdash r \dashv$.
- Next, the desired $\vdash a^k s a^* \dashv \preceq \vdash r \dashv$ is derived by the ω -rule from $\vdash a^k s a^m \dashv \preceq \vdash r \dashv$.
- If (k, m) is not an edge, it is immediate, otherwise by transitivity with $\vdash a^m t \dashv$.

- The “if” direction, from entailment to WF, is performed via a specially constructed $*$ -continuous KA.
- In this $*$ -continuous KA, \mathcal{H} holds, and $\vdash a^k t \dashv \preceq \vdash r \dashv$ implies $k \in W$.
- This KA is a variant of the algebra of languages.
- In fact, for a language A over our alphabet, we define $\sigma(A)$ as the operator which extends A by words obtained using (in)equations from \mathcal{H} .
- In particular, $t \preceq sa^*$ allows adding utv , if there is already $usa^k v$ for any k .
- If $x \equiv_{\mathcal{H}_0} y$, where \mathcal{H}_0 is the monoid part of \mathcal{H} , then x may be replaced by y in one step of σ .
- The closure of A under infinite iteration of σ is denoted by \overline{A} .

- Now the operations and constants of our KA are defined as follows:

$$A \oplus B = \overline{A \cup B}$$

$$0 = \emptyset$$

$$A \odot B = \overline{A \cdot B}$$

$$1 = \{\varepsilon\}$$

$$A^{\oplus} = \overline{A^*}$$

- Now the operations and constants of our KA are defined as follows:

$$A \oplus B = \overline{A \cup B}$$

$$0 = \emptyset$$

$$A \odot B = \overline{A \cdot B}$$

$$1 = \{\varepsilon\}$$

$$A^{\oplus} = \overline{A^*}$$

- Each letter a is interpreted as $\overline{\{a\}}$.

- Now the operations and constants of our KA are defined as follows:

$$A \oplus B = \overline{A \cup B}$$

$$0 = \emptyset$$

$$A \odot B = \overline{A \cdot B}$$

$$1 = \{\varepsilon\}$$

$$A^{\oplus} = \overline{A^*}$$

- Each letter a is interpreted as $\overline{\{a\}}$.
- This is a * -continuous Kleene algebra, and it satisfies \mathcal{H} .

- Now the operations and constants of our KA are defined as follows:

$$\begin{aligned}A \oplus B &= \overline{A \cup B} & 0 &= \emptyset \\A \odot B &= \overline{A \cdot B} & 1 &= \{\varepsilon\} \\A^{\oplus} &= \overline{A^*}\end{aligned}$$

- Each letter a is interpreted as $\overline{\{a\}}$.
- This is a * -continuous Kleene algebra, and it satisfies \mathcal{H} .
- In this algebra, we have $A \preceq B$ if $A \subseteq \overline{B}$.

- Now suppose that $\vdash a^k t \dashv \leq \vdash r \dashv$ is true in this KA.
- This means that $\vdash a^k t \dashv \in \overline{\{\vdash r \dashv\}}$.
- Again, we proceed by transfinite induction on the closure $\overline{\{\vdash r \dashv\}}$.
- $\vdash a^k t \dashv$ was obtained from $\vdash a^k s a^m \dashv$ for all m .
- The latter was either obtained immediately from $\vdash r \dashv$, which means that (k, m) is not an edge, or from $\vdash a^m t \dashv$, to which we apply the induction hypothesis yielding $m \in W$.
- Thus, $k \in W$.

- Thus, as we see, adding possibility to derive from hypotheses increases complexity dramatically.
- And, in particular, now the finitary system **KA** (or circular fragment of **HKA**_∞) is not a complete axiomatisation.