

# Complexity of Reasoning in Kleene and Action Algebras

---

Stepan Kuznetsov

ESSLI 2022 · Galway, Ireland · Gallimh, Éire

Lecture 1

- This course is devoted to complexity results for reasoning in Kleene algebras and their extension, action algebras.

# Complexity of Logics with Kleene Star

- This course is devoted to complexity results for reasoning in Kleene algebras and their extension, action algebras.
- These structures include Kleene star, which is one of the most interesting algebraic operations used in computer science.

# Complexity of Logics with Kleene Star

- This course is devoted to complexity results for reasoning in Kleene algebras and their extension, action algebras.
- These structures include Kleene star, which is one of the most interesting algebraic operations used in computer science.
- Due to its inductive nature, theories involving Kleene star, even in a purely “propositional” (atomic) language, share some features with much more powerful theories like Peano arithmetic.

# Complexity of Logics with Kleene Star

- This course is devoted to complexity results for reasoning in Kleene algebras and their extension, action algebras.
- These structures include Kleene star, which is one of the most interesting algebraic operations used in computer science.
- Due to its inductive nature, theories involving Kleene star, even in a purely “propositional” (atomic) language, share some features with much more powerful theories like Peano arithmetic.
- The methods used will be mostly proof-theoretic: thus, we shall discuss proof theory for algebraic substructural logics including Kleene star.

# About This Course

- The course belongs to the **LoCo** (Logic & Computation) category.

# About This Course

- The course belongs to the **LoCo** (Logic & Computation) category.
- The **Computation** side is connected to complexity results.

# About This Course

- The course belongs to the **LoCo** (Logic & Computation) category.
- The **Computation** side is connected to complexity results.
- The **Logic** side is connected to substructural proof theory used for logics with Kleene star.



## About This Course

- The course belongs to the **LoCo** (Logic & Computation) category.
- The **Computation** side is connected to complexity results.
- The **Logic** side is connected to substructural proof theory used for logics with Kleene star.
- There is also connection to **Language**, as Kleene star is one of the natural operations on formal languages.

## About This Course

- We suppose that students for this course have some background in Logic and Computation.

## About This Course

- We suppose that students for this course have some background in Logic and Computation.
- However, if you see a notion which you are not familiar with, please feel free to ask for definition. In fact, we mostly do not use really advanced concepts, and many basic things could be explained in a minute, if needed.

## About This Course

- We suppose that students for this course have some background in Logic and Computation.
- However, if you see a notion which you are not familiar with, please feel free to ask for definition. In fact, we mostly do not use really advanced concepts, and many basic things could be explained in a minute, if needed.
- We shall try to give mathematical proofs for most of the results, but due to time limitations some of them will be just sketches of proofs, without deep details.

- We shall start today with the most basic example of Kleene algebra, which is the algebra of **regular expressions**.

- We shall start today with the most basic example of Kleene algebra, which is the algebra of **regular expressions**.
- **Kleene algebras** in general are a generalisation of this algebra; an important subclass of those is the class of **\*-continuous** ones.

# Kleene and Action Algebras

- We shall start today with the most basic example of Kleene algebra, which is the algebra of **regular expressions**.
- **Kleene algebras** in general are a generalisation of this algebra; an important subclass of those is the class of **\*-continuous** ones.
- **Action algebras** are Kleene algebras with **divisions** (residuals), which are substructural versions of implication.

# Kleene and Action Algebras

- We shall start today with the most basic example of Kleene algebra, which is the algebra of **regular expressions**.
- **Kleene algebras** in general are a generalisation of this algebra; an important subclass of those is the class of **\*-continuous** ones.
- **Action algebras** are Kleene algebras with **divisions** (residuals), which are substructural versions of implication.
- We shall discuss complexity of atomic (equational, or inequational) theories and also of Horn theories, i.e., entailment of equations from finite sets of equations.



# Kleene and Action Algebras

- We shall start today with the most basic example of Kleene algebra, which is the algebra of **regular expressions**.
- **Kleene algebras** in general are a generalisation of this algebra; an important subclass of those is the class of **\*-continuous** ones.
- **Action algebras** are Kleene algebras with **divisions** (residuals), which are substructural versions of implication.
- We shall discuss complexity of atomic (equational, or inequational) theories and also of Horn theories, i.e., entailment of equations from finite sets of equations.
- Complexity range will be huge: from PSPACE up to  $\Pi_1^1$ -hard.

# Equivalence of Regular Expressions

- Let us start with a classical algorithmic problem.

# Equivalence of Regular Expressions

- Let us start with a classical algorithmic problem.
- A **regular expression** is constructed from letters of an alphabet  $\Sigma$  and constants 0 and 1 using three operations:
  - and + (binary) and \* (unary).

# Equivalence of Regular Expressions

- Let us start with a classical algorithmic problem.
- A **regular expression** is constructed from letters of an alphabet  $\Sigma$  and constants 0 and 1 using three operations:
  - and + (binary) and \* (unary).
- The **standard interpretation** of a regular expression is a formal language, defined as follows:

$$v(a) = \{a\} \quad v(0) = \emptyset \quad v(1) = \{\varepsilon\}$$

$$v(A \cdot B) = v(A) \cdot v(B) = \{xy \mid x \in v(A), y \in v(B)\}$$

$$v(A + B) = v(A) \cup v(B)$$

$$v(A^*) = v(A)^* = \{x_1 \dots x_n \mid n \geq 0, x_i \in v(A)\}$$

# Equivalence of Regular Expressions

- Let us start with a classical algorithmic problem.
- A **regular expression** is constructed from letters of an alphabet  $\Sigma$  and constants 0 and 1 using three operations:
  - and + (binary) and \* (unary).
- The **standard interpretation** of a regular expression is a formal language, defined as follows:

$$v(a) = \{a\} \quad v(0) = \emptyset \quad v(1) = \{\varepsilon\}$$

$$v(A \cdot B) = v(A) \cdot v(B) = \{xy \mid x \in v(A), y \in v(B)\}$$

$$v(A + B) = v(A) \cup v(B)$$

$$v(A^*) = v(A)^* = \{x_1 \dots x_n \mid n \geq 0, x_i \in v(A)\}$$

- **Equivalence problem:** given two reg. exp.  $A$  and  $B$ , determine whether  $v(A) = v(B)$ .

# Equivalence of Regular Expressions

- The set of words  $v(A)$  is called the language defined by reg. exp.  $A$ .

# Equivalence of Regular Expressions

- The set of words  $v(A)$  is called the language defined by reg. exp.  $A$ .
- For example, reg. exp.  $A = a(a + b)^*$  and  $B = a(a^*b)^*a^*$  are equivalent, since they define the same language, which includes all words starting with  $a$ .

# Equivalence of Regular Expressions

- The set of words  $v(A)$  is called the language defined by reg. exp.  $A$ .
- For example, reg. exp.  $A = a(a + b)^*$  and  $B = a(a^*b)^*a^*$  are equivalent, since they define the same language, which includes all words starting with  $a$ .
- On the other hand,  $A = (ab^*)^*$  and  $B = (a + b)^*$  are not equivalent:  $aba \in v(B)$ , but  $aba \notin v(A)$ .



# Equivalence of Regular Expressions

- The set of words  $v(A)$  is called the language defined by reg. exp.  $A$ .
- For example, reg. exp.  $A = a(a + b)^*$  and  $B = a(a^*b)^*a^*$  are equivalent, since they define the same language, which includes all words starting with  $a$ .
- On the other hand,  $A = (ab^*)^*$  and  $B = (a + b)^*$  are not equivalent:  $aba \in v(B)$ , but  $aba \notin v(A)$ .
- The equivalence problem, of course, has standard solutions (e.g., using finite automata), but we shall focus on the logic behind it.

# Equivalence of Regular Expressions

## Theorem (Hunt et al. 1976)

*The equivalence problem for regular expressions is algorithmically decidable.*

- More precisely, it is PSPACE-complete.

# Equivalence of Regular Expressions

## Theorem (Hunt et al. 1976)

*The equivalence problem for regular expressions is algorithmically decidable.*

- More precisely, it is PSPACE-complete.
- We shall discuss a *proof-theoretic* approach to this problem, following Krob (1991), Kozen (1994), and Das & Pous (2017).

# Equivalence of Regular Expressions

## Theorem (Hunt et al. 1976)

*The equivalence problem for regular expressions is algorithmically decidable.*

- More precisely, it is PSPACE-complete.
- We shall discuss a *proof-theoretic* approach to this problem, following Krob (1991), Kozen (1994), and Das & Pous (2017).
- The idea is to give good axiomatizations for *equational theories* of regular expressions (and more).

# Equivalence of Regular Expressions

## Theorem (Hunt et al. 1976)

*The equivalence problem for regular expressions is algorithmically decidable.*

- More precisely, it is PSPACE-complete.
- We shall discuss a *proof-theoretic* approach to this problem, following Krob (1991), Kozen (1994), and Das & Pous (2017).
- The idea is to give good axiomatizations for *equational theories* of regular expressions (and more).
- For convenience, we shall talk about **inequations** of reg. exp.:  
 $A \preceq B$  means  $v(A) \subseteq v(B)$ .

# Equivalence of Regular Expressions

## Theorem (Hunt et al. 1976)

*The equivalence problem for regular expressions is algorithmically decidable.*

- More precisely, it is PSPACE-complete.
- We shall discuss a *proof-theoretic* approach to this problem, following Krob (1991), Kozen (1994), and Das & Pous (2017).
- The idea is to give good axiomatizations for *equational theories* of regular expressions (and more).
- For convenience, we shall talk about **inequations** of reg. exp.:  
 $A \preceq B$  means  $v(A) \subseteq v(B)$ .
  - $A \preceq B$  is the same as  $A + B \equiv B$ .

# Equivalence of Regular Expressions

## Theorem (Hunt et al. 1976)

*The equivalence problem for regular expressions is algorithmically decidable.*

- More precisely, it is PSPACE-complete.
- We shall discuss a *proof-theoretic* approach to this problem, following Krob (1991), Kozen (1994), and Das & Pous (2017).
- The idea is to give good axiomatizations for *equational theories* of regular expressions (and more).
- For convenience, we shall talk about **inequations** of reg. exp.:  
 $A \preceq B$  means  $v(A) \subseteq v(B)$ .
  - $A \preceq B$  is the same as  $A + B \equiv B$ .
  - Inequation is more like implication, and it will be easier to axiomatise.

- The infinitary nature of the Kleene star is most naturally reflected by an  $\omega$ -**rule**:

$$\frac{(A^n \preceq B)_{n=0}^{\infty}}{A^* \preceq B}$$



- The infinitary nature of the Kleene star is most naturally reflected by an  $\omega$ -**rule**:

$$\frac{(A^n \preceq B)_{n=0}^{\infty}}{A^* \preceq B}$$

- Let us formulate an infinitary axiomatization as a **sequent calculus**  $\mathbf{KA}_\omega$ , with sequents of the form  $\Pi \rightarrow B$ , where  $\Pi$  is a sequence of formulae (reg. exp.),  $B$  is a formula.

- The infinitary nature of the Kleene star is most naturally reflected by an  $\omega$ -**rule**:

$$\frac{(A^n \preceq B)_{n=0}^{\infty}}{A^* \preceq B}$$

- Let us formulate an infinitary axiomatization as a **sequent calculus**  $\mathbf{KA}_\omega$ , with sequents of the form  $\Pi \rightarrow B$ , where  $\Pi$  is a sequence of formulae (reg. exp.),  $B$  is a formula.
- “**KA**” stands for “Kleene algebra,” and “ $\omega$ ” is due to the infinitary rule ( $\omega$ -rule).

# Sequent Calculus

- The idea of sequent calculi goes back to Gentzen (1934); proofs in these calculi are much more convenient for analysis, if compared to, say, Hilbert-style proofs.

# Sequent Calculus

- The idea of sequent calculi goes back to Gentzen (1934); proofs in these calculi are much more convenient for analysis, if compared to, say, Hilbert-style proofs.
- Namely, sequent proofs have the property of **analiticity**: premises of a rule can be somehow obtained from its conclusion, which facilitates proof search.

# Sequent Calculus

- The idea of sequent calculi goes back to Gentzen (1934); proofs in these calculi are much more convenient for analysis, if compared to, say, Hilbert-style proofs.
- Namely, sequent proofs have the property of **analiticity**: premises of a rule can be somehow obtained from its conclusion, which facilitates proof search.
- An example of a non-analytic rule, which could have been used in  $\mathbf{KA}_\omega$ , is transitivity syllogism:

$$\frac{A \preceq B \quad B \preceq C}{A \preceq C} \text{Trans}$$

# Sequent Calculus

- The idea of sequent calculi goes back to Gentzen (1934); proofs in these calculi are much more convenient for analysis, if compared to, say, Hilbert-style proofs.
- Namely, sequent proofs have the property of **analiticity**: premises of a rule can be somehow obtained from its conclusion, which facilitates proof search.
- An example of a non-analytic rule, which could have been used in  $\mathbf{KA}_\omega$ , is transitivity syllogism:

$$\frac{A \preceq B \quad B \preceq C}{A \preceq C} \text{Trans}$$

- If we wish to prove  $A \preceq C$  using *Trans*, we have no idea what  $B$  could be.

# Sequent Calculus

- The idea of sequent calculi goes back to Gentzen (1934); proofs in these calculi are much more convenient for analysis, if compared to, say, Hilbert-style proofs.
- Namely, sequent proofs have the property of **analiticity**: premises of a rule can be somehow obtained from its conclusion, which facilitates proof search.
- An example of a non-analytic rule, which could have been used in  $\mathbf{KA}_\omega$ , is transitivity syllogism:

$$\frac{A \preceq B \quad B \preceq C}{A \preceq C} \text{Trans}$$

- If we wish to prove  $A \preceq C$  using *Trans*, we have no idea what  $B$  could be.
- We shall avoid using such rules.

- As we have already said, our calculus will derive *sequents* of the form  $\Pi \rightarrow B$ .



- As we have already said, our calculus will derive *sequents* of the form  $\Pi \rightarrow B$ .
- The meaning of such a sequent, if  $\Pi = A_1, \dots, A_n$ , is the inequation  $A_1 \cdot \dots \cdot A_n \preceq B$ .

- As we have already said, our calculus will derive *sequents* of the form  $\Pi \rightarrow B$ .
- The meaning of such a sequent, if  $\Pi = A_1, \dots, A_n$ , is the inequation  $A_1 \cdot \dots \cdot A_n \preceq B$ .
- If  $\Pi$  is empty, we have  $1 \preceq B$ .

# Sequent Calculus

- As we have already said, our calculus will derive *sequents* of the form  $\Pi \rightarrow B$ .
- The meaning of such a sequent, if  $\Pi = A_1, \dots, A_n$ , is the inequation  $A_1 \cdot \dots \cdot A_n \preceq B$ .
- If  $\Pi$  is empty, we have  $1 \preceq B$ .
- Thus, the sequential comma in  $\Pi$  is a *meta-syntactic* counterpart of product (multiplication);  $\rightarrow$  corresponds to  $\preceq$ .

# Sequent Calculus

- As we have already said, our calculus will derive *sequents* of the form  $\Pi \rightarrow B$ .
- The meaning of such a sequent, if  $\Pi = A_1, \dots, A_n$ , is the inequation  $A_1 \cdot \dots \cdot A_n \preceq B$ .
- If  $\Pi$  is empty, we have  $1 \preceq B$ .
- Thus, the sequential comma in  $\Pi$  is a *meta-syntactic* counterpart of product (multiplication);  $\rightarrow$  corresponds to  $\preceq$ .
- $\Pi$  and  $B$  are called, respectively, the **antecedent** and the **succedent** of the sequent.

Axioms and rules of inference of  $\mathbf{KA}_\omega$  are as follows:

$$\begin{array}{c}
 \overline{A \rightarrow A} \text{ } Id \qquad \overline{\Gamma, 0, \Delta \rightarrow B} \text{ } 0L \\
 \\
 \frac{\Gamma, \Delta \rightarrow B}{\Gamma, 1, \Delta \rightarrow B} \text{ } 1L \qquad \overline{\rightarrow 1} \text{ } 1R \\
 \\
 \frac{\Gamma, A, B, \Delta \rightarrow C}{\Gamma, A \cdot B, \Delta \rightarrow C} \text{ } \cdot L \qquad \frac{\Gamma \rightarrow A \quad \Delta \rightarrow B}{\Gamma, \Delta \rightarrow A \cdot B} \text{ } \cdot R \\
 \\
 \frac{\Gamma, A, \Delta \rightarrow C \quad \Gamma, B, \Delta \rightarrow C}{\Gamma, A + B, \Delta \rightarrow C} \text{ } +L \qquad \frac{\Pi \rightarrow A}{\Pi \rightarrow A + B} \text{ } +R_1 \quad \frac{\Pi \rightarrow B}{\Pi \rightarrow A + B} \text{ } +R_2 \\
 \\
 \frac{(\Gamma, A^n, \Delta \rightarrow B)_{n=0}^\infty}{\Gamma, A^*, \Delta \rightarrow B} \text{ } *L_\omega \qquad \frac{\Gamma_1 \rightarrow A \quad \dots \quad \Gamma_n \rightarrow A}{\Gamma_1, \dots, \Gamma_n \rightarrow A^*} \text{ } *R_n, \text{ } n \geq 0
 \end{array}$$

- Due to the presence of the  $\omega$ -rule,  $*L_\omega$ , proofs in  $\mathbf{KA}_\omega$  could be infinite.

- Due to the presence of the  $\omega$ -rule,  $*L_\omega$ , proofs in  $\mathbf{KA}_\omega$  could be infinite.
- However, each proof should still be *well-founded*, i.e., while infinitary branching is allowed, each path from the root should be finite and reach an axiom leaf.

- Due to the presence of the  $\omega$ -rule,  $*L_\omega$ , proofs in  $\mathbf{KA}_\omega$  could be infinite.
- However, each proof should still be *well-founded*, i.e., while infinitary branching is allowed, each path from the root should be finite and reach an axiom leaf.
- Equivalently (see Aczel 1977), the set of sequents derivable in  $\mathbf{KA}_\omega$  can be defined as the minimal (w.r.t. set inclusion) set of sequents including all axioms and closed under inference rules.



- Due to the presence of the  $\omega$ -rule,  $*L_\omega$ , proofs in  $\mathbf{KA}_\omega$  could be infinite.
- However, each proof should still be *well-founded*, i.e., while infinitary branching is allowed, each path from the root should be finite and reach an axiom leaf.
- Equivalently (see Aczel 1977), the set of sequents derivable in  $\mathbf{KA}_\omega$  can be defined as the minimal (w.r.t. set inclusion) set of sequents including all axioms and closed under inference rules.
- This set is the least fixpoint of the immediate derivability operator, and it is reached by its transfinite iteration up to the *closure ordinal* for  $\mathbf{KA}_\omega$ .

- $\mathbf{KA}_\omega$  admits the cut rule, which is a generalisation of *Trans*:

$$\frac{\Pi \rightarrow A \quad \Gamma, A, \Delta \rightarrow C}{\Gamma, \Pi, \Delta \rightarrow C} \textit{Cut}$$

- $\mathbf{KA}_\omega$  admits the cut rule, which is a generalisation of *Trans*:

$$\frac{\Pi \rightarrow A \quad \Gamma, A, \Delta \rightarrow C}{\Gamma, \Pi, \Delta \rightarrow C} \textit{Cut}$$

- A syntactic proof of this would be rather standard, but involves **transfinite induction**.

- $\mathbf{KA}_\omega$  admits the cut rule, which is a generalisation of *Trans*:

$$\frac{\Pi \rightarrow A \quad \Gamma, A, \Delta \rightarrow C}{\Gamma, \Pi, \Delta \rightarrow C} \text{ Cut}$$

- A syntactic proof of this would be rather standard, but involves **transfinite induction**.
- We are not going to details of this proof now, since on one of the next lectures we shall present such a proof for a more general system,  $\mathbf{ACT}_\omega$  (infinitary action logic).

- $\mathbf{KA}_\omega$  admits the cut rule, which is a generalisation of *Trans*:

$$\frac{\Pi \rightarrow A \quad \Gamma, A, \Delta \rightarrow C}{\Gamma, \Pi, \Delta \rightarrow C} \text{ Cut}$$

- A syntactic proof of this would be rather standard, but involves **transfinite induction**.
- We are not going to details of this proof now, since on one of the next lectures we shall present such a proof for a more general system,  $\mathbf{ACT}_\omega$  (infinitary action logic).
- However, already today we shall obtain a semantic proof of cut admissibility.

- Let us concentrate on the  $*$ -free fragment of  $\mathbf{KA}_\omega$  and compare it with more well-known syntactic system.

- Let us concentrate on the  $*$ -free fragment of  $\mathbf{KA}_\omega$  and compare it with more well-known syntactic system.
- The intuitionistic propositional calculus (**IPC**) in it sequent form (see, e.g., Takeuti 1975) also uses sequents of the form  $\Pi \rightarrow A$ , but now formulae are built using intuitionistic connective:  $\&$  (conjunction),  $\vee$  (disjunction),  $\supset$  (implication), and constants  $\perp$  (falsity) and  $\top$  (truth).

- Let us concentrate on the  $*$ -free fragment of  $\mathbf{KA}_\omega$  and compare it with more well-known syntactic system.
- The intuitionistic propositional calculus (**IPC**) in it sequent form (see, e.g., Takeuti 1975) also uses sequents of the form  $\Pi \rightarrow A$ , but now formulae are built using intuitionistic connective:  $\&$  (conjunction),  $\vee$  (disjunction),  $\supset$  (implication), and constants  $\perp$  (falsity) and  $\top$  (truth).
- In usual formulations of **IPC**, antecedents ( $\Pi$ ) are multisets or even sets of formulae. That is, order does not matter: **IPC** is commutative.



# Substructurality

Axioms and rules of **IPC** are as follows:

$$\begin{array}{c} \overline{A \rightarrow A} \text{ } Id \qquad \overline{\Gamma, \perp, \Delta \rightarrow B} \perp L \\ \frac{\Gamma, \Delta \rightarrow B}{\Gamma, \top, \Delta \rightarrow B} \top L \qquad \overline{\rightarrow \top} \top R \\ \frac{\Gamma, A, B, \Delta \rightarrow C}{\Gamma, A \& B, \Delta \rightarrow C} \& L \qquad \frac{\Gamma \rightarrow A \quad \Delta \rightarrow B}{\Gamma, \Delta \rightarrow A \& B} \& R \\ \frac{\Gamma, A, \Delta \rightarrow C \quad \Gamma, B, \Delta \rightarrow C}{\Gamma, A \vee B, \Delta \rightarrow C} \vee L \qquad \frac{\Pi \rightarrow A}{\Pi \rightarrow A \vee B} \vee R_1 \quad \frac{\Pi \rightarrow B}{\Pi \rightarrow A \vee B} \vee R_2 \\ \frac{\Pi \rightarrow A \quad \Gamma, B, \Delta \rightarrow C}{\Gamma, \Pi, A \supset B, \Delta \rightarrow C} \supset L \qquad \frac{A, \Pi \rightarrow B}{\Pi \rightarrow A \supset B} \supset R \\ \\ \frac{\Gamma, \Delta \rightarrow C}{\Gamma, A, \Delta \rightarrow C} \text{ } Weak \qquad \frac{\Gamma, A, A, \Delta \rightarrow C}{\Gamma, A, \Delta \rightarrow C} \text{ } Contr \qquad \frac{\Gamma, B, A, \Delta \rightarrow C}{\Gamma, A, B, \Delta \rightarrow C} \text{ } Perm \end{array}$$

- Logical rules of  $\mathbf{KA}_\omega$  and  $\mathbf{IPC}$  are similar, up to renaming of operations:  $\cdot$  to  $\&$ ,  $+$  to  $\vee$ ,  $\top$  to  $1$ , and  $\perp$  to  $0$ .

- Logical rules of  $\mathbf{KA}_\omega$  and  $\mathbf{IPC}$  are similar, up to renaming of operations:  $\cdot$  to  $\&$ ,  $+$  to  $\vee$ ,  $\top$  to  $1$ , and  $\perp$  to  $0$ .
- There are, however, significant differences:

- Logical rules of  $\mathbf{KA}_\omega$  and  $\mathbf{IPC}$  are similar, up to renaming of operations:  $\cdot$  to  $\&$ ,  $+$  to  $\vee$ ,  $\top$  to  $1$ , and  $\perp$  to  $0$ .
- There are, however, significant differences:
  1. There is no counterpart of  $\supset$  in  $\mathbf{KA}_\omega$ . Implication is only on the top level,  $\rightarrow$ . This will be fixed in **action logic**.

- Logical rules of  $\mathbf{KA}_\omega$  and  $\mathbf{IPC}$  are similar, up to renaming of operations:  $\cdot$  to  $\&$ ,  $+$  to  $\vee$ ,  $\top$  to  $1$ , and  $\perp$  to  $0$ .
- There are, however, significant differences:
  1. There is no counterpart of  $\supset$  in  $\mathbf{KA}_\omega$ . Implication is only on the top level,  $\rightarrow$ . This will be fixed in **action logic**.
  2.  $\mathbf{KA}_\omega$  is **substructural**, i.e., it does not include structural rules *Weak*, *Contr*, *Perm*.

- Logical rules of  $\mathbf{KA}_\omega$  and  $\mathbf{IPC}$  are similar, up to renaming of operations:  $\cdot$  to  $\&$ ,  $+$  to  $\vee$ ,  $\top$  to  $1$ , and  $\perp$  to  $0$ .
- There are, however, significant differences:
  1. There is no counterpart of  $\supset$  in  $\mathbf{KA}_\omega$ . Implication is only on the top level,  $\rightarrow$ . This will be fixed in **action logic**.
  2.  $\mathbf{KA}_\omega$  is **substructural**, i.e., it does not include structural rules *Weak*, *Contr*, *Perm*.
  3.  $\mathbf{IPC}$  does not have Kleene star. In fact,  $A^*$  trivialises to  $1$  in the presence of *Weak*. Thus, it is important for logics with Kleene star to be substructural.

- Now we have seen some syntactic definitions, and let us take a look at the course in general.

# Course Outline

- Now we have seen some syntactic definitions, and let us take a look at the course in general.
- The topics of the five lectures will be as follows:



# Course Outline

- Now we have seen some syntactic definitions, and let us take a look at the course in general.
- The topics of the five lectures will be as follows:
  1. We shall show that  $\mathbf{KA}_\omega$  is sound and complete w.r.t. various interpretations, and present an alternative, finitary axiomatisation  $\mathbf{KA}$ .

# Course Outline

- Now we have seen some syntactic definitions, and let us take a look at the course in general.
- The topics of the five lectures will be as follows:
  1. We shall show that  $\mathbf{KA}_\omega$  is sound and complete w.r.t. various interpretations, and present an alternative, finitary axiomatisation  $\mathbf{KA}$ .
  2. We are not going to prove equivalence of  $\mathbf{KA}_\omega$  and  $\mathbf{KA}$ , but rather reduce  $\mathbf{KA}_\omega$  to a circular proof system  $\mathbf{LHKA}_\circ$  (Das & Pous 2018). This yields decidability of  $\mathbf{KA}_\omega$ .

# Course Outline

- Now we have seen some syntactic definitions, and let us take a look at the course in general.
- The topics of the five lectures will be as follows:
  1. We shall show that  $\mathbf{KA}_\omega$  is sound and complete w.r.t. various interpretations, and present an alternative, finitary axiomatisation  $\mathbf{KA}$ .
  2. We are not going to prove equivalence of  $\mathbf{KA}_\omega$  and  $\mathbf{KA}$ , but rather reduce  $\mathbf{KA}_\omega$  to a circular proof system  $\mathbf{LHKA}_\omega$  (Das & Pous 2018). This yields decidability of  $\mathbf{KA}_\omega$ .
  3. This lecture is for complexity of reasoning from hypotheses (Horn theories) of Kleene algebras.

- Now we have seen some syntactic definitions, and let us take a look at the course in general.
- The topics of the five lectures will be as follows:
  1. We shall show that  $\mathbf{KA}_\omega$  is sound and complete w.r.t. various interpretations, and present an alternative, finitary axiomatisation  $\mathbf{KA}$ .
  2. We are not going to prove equivalence of  $\mathbf{KA}_\omega$  and  $\mathbf{KA}$ , but rather reduce  $\mathbf{KA}_\omega$  to a circular proof system  $\mathbf{LHKA}_\omega$  (Das & Pous 2018). This yields decidability of  $\mathbf{KA}_\omega$ .
  3. This lecture is for complexity of reasoning from hypotheses (Horn theories) of Kleene algebras.
  4. We shall add implications (divisions) to Kleene algebras, yielding action algebras. Here we prove complexity results for commutative action algebras.

- Now we have seen some syntactic definitions, and let us take a look at the course in general.
- The topics of the five lectures will be as follows:
  1. We shall show that  $\mathbf{KA}_\omega$  is sound and complete w.r.t. various interpretations, and present an alternative, finitary axiomatisation  $\mathbf{KA}$ .
  2. We are not going to prove equivalence of  $\mathbf{KA}_\omega$  and  $\mathbf{KA}$ , but rather reduce  $\mathbf{KA}_\omega$  to a circular proof system  $\mathbf{LHKA}_\circ$  (Das & Pous 2018). This yields decidability of  $\mathbf{KA}_\omega$ .
  3. This lecture is for complexity of reasoning from hypotheses (Horn theories) of Kleene algebras.
  4. We shall add implications (divisions) to Kleene algebras, yielding action algebras. Here we prove complexity results for commutative action algebras.
  5. Here we sketch complexity proofs for the non-commutative case, and also consider action logic with exponentials.

- It is well-known that **IPC** is decidable in polynomial space (in fact, it is PSPACE-complete).

- It is well-known that **IPC** is decidable in polynomial space (in fact, it is PSPACE-complete).
- The key to the PSPACE decidability procedure is the possibility to make proofs in **IPC** having a polymodal bound on depth (Dyckhoff 1992).

- It is well-known that **IPC** is decidable in polynomial space (in fact, it is PSPACE-complete).
- The key to the PSPACE decidability procedure is the possibility to make proofs in **IPC** having a polymodal bound on depth (Dyckhoff 1992).
- Such a proof could be exponential in size, but there is a non-deterministic depth-first search procedure for finding it.



- It is well-known that **IPC** is decidable in polynomial space (in fact, it is PSPACE-complete).
- The key to the PSPACE decidability procedure is the possibility to make proofs in **IPC** having a polymodal bound on depth (Dyckhoff 1992).
- Such a proof could be exponential in size, but there is a non-deterministic depth-first search procedure for finding it.
- And, by Savitch's theorem, NPSPACE is PSPACE.

- It is well-known that **IPC** is decidable in polynomial space (in fact, it is PSPACE-complete).
- The key to the PSPACE decidability procedure is the possibility to make proofs in **IPC** having a polymodal bound on depth (Dyckhoff 1992).
- Such a proof could be exponential in size, but there is a non-deterministic depth-first search procedure for finding it.
- And, by Savitch's theorem, NPSPACE is PSPACE.
- For  $\mathbf{KA}_\omega$ , all of this will not work, due to the presence of the  $\omega$ -rule,  $*L_\omega$ .

- It is well-known that **IPC** is decidable in polynomial space (in fact, it is PSPACE-complete).
- The key to the PSPACE decidability procedure is the possibility to make proofs in **IPC** having a polymodal bound on depth (Dyckhoff 1992).
- Such a proof could be exponential in size, but there is a non-deterministic depth-first search procedure for finding it.
- And, by Savitch's theorem, NPSPACE is PSPACE.
- For  $\mathbf{KA}_\omega$ , all of this will not work, due to the presence of the  $\omega$ -rule,  $*L_\omega$ .
- Proofs are infinite, no “out-of-the-box” finite proof search is possible.

- In general, infinitary systems like  $\mathbf{KA}_\omega$  could have complexity up to  $\Pi_1^1$ .

- In general, infinitary systems like  $\mathbf{KA}_\omega$  could have complexity up to  $\Pi_1^1$ .
- Recall that a set of natural numbers (or finite objects, e.g., sequents) is in  $\Pi_1^1$  if it can be defined by a second-order formula of the form  $\psi(a) = (\forall X \subseteq \mathbb{N}) \varphi(X, a)$ , where  $\varphi$  is an arithmetical formula.

- In general, infinitary systems like  $\mathbf{KA}_\omega$  could have complexity up to  $\Pi_1^1$ .
- Recall that a set of natural numbers (or finite objects, e.g., sequents) is in  $\Pi_1^1$  if it can be defined by a second-order formula of the form  $\psi(a) = (\forall X \subseteq \mathbb{N}) \varphi(X, a)$ , where  $\varphi$  is an arithmetical formula.
- One could expect  $\exists Y \xi(Y, a)$ , where  $Y$  is the infinite proof (i.e.,  $\Sigma_1^1$ ). However, the key issue here is checking that the proof is well-founded, and this is non-arithmetical.

- In general, infinitary systems like  $\mathbf{KA}_\omega$  could have complexity up to  $\Pi_1^1$ .
- Recall that a set of natural numbers (or finite objects, e.g., sequents) is in  $\Pi_1^1$  if it can be defined by a second-order formula of the form  $\psi(a) = (\forall X \subseteq \mathbb{N}) \varphi(X, a)$ , where  $\varphi$  is an arithmetical formula.
- One could expect  $\exists Y \xi(Y, a)$ , where  $Y$  is the infinite proof (i.e.,  $\Sigma_1^1$ ). However, the key issue here is checking that the proof is well-founded, and this is non-arithmetical.
- There is, however, a much better complexity bound for  $\mathbf{KA}_\omega$ , namely,  $\Pi_1^0$ .

# Complexity

- The  $\Pi_1^0$  complexity class allows one  $\forall$  quantifier over *finite* objects.



# Complexity

- The  $\Pi_1^0$  complexity class allows one  $\forall$  quantifier over *finite* objects.
- This class is dual to the class  $\Sigma_1^0$  or *enumerable* (r.e.) sets.

# Complexity

- The  $\Pi_1^0$  complexity class allows one  $\forall$  quantifier over *finite* objects.
- This class is dual to the class  $\Sigma_1^0$  or *enumerable* (r.e.) sets.
- If a logic is described by a calculus with finite proofs, then it belongs to  $\Sigma_1^0$ : provability means *existence* of a finite object (proof).

# Complexity

- The  $\Pi_1^0$  complexity class allows one  $\forall$  quantifier over *finite* objects.
- This class is dual to the class  $\Sigma_1^0$  or *enumerable* (r.e.) sets.
- If a logic is described by a calculus with finite proofs, then it belongs to  $\Sigma_1^0$ : provability means *existence* of a finite object (proof).
- For  $\Pi_1^0$ , *disproving* of a sequent means presenting a finite witness against it. (The negation of  $\forall$  gives  $\exists$ .)

# Complexity

- The  $\Pi_1^0$  complexity class allows one  $\forall$  quantifier over *finite* objects.
- This class is dual to the class  $\Sigma_1^0$  or *enumerable* (r.e.) sets.
- If a logic is described by a calculus with finite proofs, then it belongs to  $\Sigma_1^0$ : provability means *existence* of a finite object (proof).
- For  $\Pi_1^0$ , *disproving* of a sequent means presenting a finite witness against it. (The negation of  $\forall$  gives  $\exists$ .)
- A syntactic proof of the  $\Pi_1^0$  upper bound will be, again, given later for the larger system  $\mathbf{ACT}_\omega$ .

# Complexity

- The  $\Pi_1^0$  complexity class allows one  $\forall$  quantifier over *finite* objects.
- This class is dual to the class  $\Sigma_1^0$  or *enumerable* (r.e.) sets.
- If a logic is described by a calculus with finite proofs, then it belongs to  $\Sigma_1^0$ : provability means *existence* of a finite object (proof).
- For  $\Pi_1^0$ , *disproving* of a sequent means presenting a finite witness against it. (The negation of  $\forall$  gives  $\exists$ .)
- A syntactic proof of the  $\Pi_1^0$  upper bound will be, again, given later for the larger system  $\mathbf{ACT}_\omega$ .
- However, semantically it is obvious: given soundness and completeness (to appear), we disprove  $A \preceq B$  by presenting a word  $w$  such that  $w \in v(A)$  and  $w \notin v(B)$ .

- If we manage to reduce the infinitary proof system for  $\mathbf{KA}_\omega$  to a finitary one, we get  $\Pi_1^0$  and  $\Sigma_1^0$  at the same time.

- If we manage to reduce the infinitary proof system for  $\mathbf{KA}_\omega$  to a finitary one, we get  $\Pi_1^0$  and  $\Sigma_1^0$  at the same time.
- By Post's theorem, this gives decidability.

- If we manage to reduce the infinitary proof system for  $\mathbf{KA}_\omega$  to a finitary one, we get  $\Pi_1^0$  and  $\Sigma_1^0$  at the same time.
- By Post's theorem, this gives decidability.
- Indeed, the algorithm seeks, in parallel, for a proof of the sequent or a witness against it.



- If we manage to reduce the infinitary proof system for  $\mathbf{KA}_\omega$  to a finitary one, we get  $\Pi_1^0$  and  $\Sigma_1^0$  at the same time.
- By Post's theorem, this gives decidability.
- Indeed, the algorithm seeks, in parallel, for a proof of the sequent or a witness against it.
- Exactly one of the searches succeeds.

- If we manage to reduce the infinitary proof system for  $\mathbf{KA}_\omega$  to a finitary one, we get  $\Pi_1^0$  and  $\Sigma_1^0$  at the same time.
- By Post's theorem, this gives decidability.
- Indeed, the algorithm seeks, in parallel, for a proof of the sequent or a witness against it.
- Exactly one of the searches succeeds.
- If we also manage to impose a polynomial boundary on the proof depth, we get PSPACE.

# Soundness and Completeness

- Before going forward, let us discuss relations between syntax and semantics for the infinitary system  $\mathbf{KA}_\omega$ .

# Soundness and Completeness

- Before going forward, let us discuss relations between syntax and semantics for the infinitary system  $\mathbf{KA}_\omega$ .

## Theorem

$\mathbf{KA}_\omega \vdash A \rightarrow B$  iff  $v(A) \subseteq v(B)$  in the standard interpretation.

- Before going forward, let us discuss relations between syntax and semantics for the infinitary system  $\mathbf{KA}_\omega$ .

## Theorem

$\mathbf{KA}_\omega \vdash A \rightarrow B$  iff  $v(A) \subseteq v(B)$  in the standard interpretation.

- Corollary:  $\mathbf{KA}_\omega$  admits *Cut*.

# Soundness and Completeness

- Before going forward, let us discuss relations between syntax and semantics for the infinitary system  $\mathbf{KA}_\omega$ .

## Theorem

$\mathbf{KA}_\omega \vdash A \rightarrow B$  iff  $v(A) \subseteq v(B)$  in the standard interpretation.

- Corollary:  $\mathbf{KA}_\omega$  admits *Cut*.
- Soundness is easy: all axioms are valid and inference rules preserve validity.

# Soundness and Completeness

- Before going forward, let us discuss relations between syntax and semantics for the infinitary system  $\mathbf{KA}_\omega$ .

## Theorem

$\mathbf{KA}_\omega \vdash A \rightarrow B$  iff  $v(A) \subseteq v(B)$  in the standard interpretation.

- Corollary:  $\mathbf{KA}_\omega$  admits *Cut*.
- Soundness is easy: all axioms are valid and inference rules preserve validity.
- For completeness, we invert the left rules and exhaustively consider all words  $x \in v(A)$ .

# Soundness and Completeness

- Before going forward, let us discuss relations between syntax and semantics for the infinitary system  $\mathbf{KA}_\omega$ .

## Theorem

$\mathbf{KA}_\omega \vdash A \rightarrow B$  iff  $v(A) \subseteq v(B)$  in the standard interpretation.

- Corollary:  $\mathbf{KA}_\omega$  admits *Cut*.
- Soundness is easy: all axioms are valid and inference rules preserve validity.
- For completeness, we invert the left rules and exhaustively consider all words  $x \in v(A)$ .
- This is an infinitary brute-force procedure.



## Soundness and Completeness

- Suppose,  $B = (a + b)^* a(ab)^*$  and  $A = (a + b)^* (a + aab)$ . We have  $v(A) \subseteq v(B)$ .

# Soundness and Completeness

- Suppose,  $B = (a + b)^* a(ab)^*$  and  $A = (a + b)^* (a + aab)$ . We have  $v(A) \subseteq v(B)$ .
- Let us construct a proof of  $A \rightarrow B$  in  $\mathbf{KA}_\omega$ .

# Soundness and Completeness

- Suppose,  $B = (a + b)^* a(ab)^*$  and  $A = (a + b)^* (a + aab)$ . We have  $v(A) \subseteq v(B)$ .
- Let us construct a proof of  $A \rightarrow B$  in  $\mathbf{KA}_\omega$ .
- First, we decompose the product:

$$(a + b)^*, a + aab \rightarrow B.$$

The goal sequent  $A \rightarrow B$  is derivable from this by  $\cdot L$ .

# Soundness and Completeness

- Suppose,  $B = (a + b)^* a(ab)^*$  and  $A = (a + b)^* (a + aab)$ . We have  $v(A) \subseteq v(B)$ .
- Let us construct a proof of  $A \rightarrow B$  in  $\mathbf{KA}_\omega$ .
- First, we decompose the product:

$$(a + b)^*, a + aab \rightarrow B.$$

The goal sequent  $A \rightarrow B$  is derivable from this by  $\cdot L$ .

- This sequent will be derived by  $*L_\omega$  from the sequents

$$(a + b)^n, a + aab \rightarrow B$$

for each natural  $n$ .

# Soundness and Completeness

- Suppose,  $B = (a + b)^* a(ab)^*$  and  $A = (a + b)^* (a + aab)$ . We have  $v(A) \subseteq v(B)$ .
- Let us construct a proof of  $A \rightarrow B$  in  $\mathbf{KA}_\omega$ .
- First, we decompose the product:

$$(a + b)^*, a + aab \rightarrow B.$$

The goal sequent  $A \rightarrow B$  is derivable from this by  $\cdot L$ .

- This sequent will be derived by  $*L_\omega$  from the sequents

$$(a + b)^n, a + aab \rightarrow B$$

for each natural  $n$ .

- Finally, each of these sequents is derived using exhaustive applications of  $+L$ .

- Now we actually have to prove  $\Gamma_w \rightarrow B$  for each  $w \in v(A)$ , e.g.,  
 $b, a, a, b, b, a, a, b \rightarrow B$ .

# Soundness and Completeness

- Now we actually have to prove  $\Gamma_w \rightarrow B$  for each  $w \in v(A)$ , e.g.,  
 $b, a, a, b, b, a, a, b \rightarrow B$ .
- Thus, the first stage of proof construction is essentially brute-force along all (infinitely many of) words of  $v(A)$ .

- Now we actually have to prove  $\Gamma_w \rightarrow B$  for each  $w \in v(A)$ , e.g.,  $b, a, a, b, b, a, a, b \rightarrow B$ .
- Thus, the first stage of proof construction is essentially brute-force along all (infinitely many of) words of  $v(A)$ .
- Now for each word we have to prove that it is in  $v(B)$ .



# Soundness and Completeness

- Now we actually have to prove  $\Gamma_w \rightarrow B$  for each  $w \in v(A)$ , e.g.,  $b, a, a, b, b, a, a, b \rightarrow B$ .
- Thus, the first stage of proof construction is essentially brute-force along all (infinitely many of) words of  $v(A)$ .
- Now for each word we have to prove that it is in  $v(B)$ .
- This is done by applying the corresponding right rules, and at steps for  $+$  and  $*$  we choose the correct way.

# Soundness and Completeness

- Now we actually have to prove  $\Gamma_w \rightarrow B$  for each  $w \in v(A)$ , e.g.,  $b, a, a, b, b, a, a, b \rightarrow B$ .
- Thus, the first stage of proof construction is essentially brute-force along all (infinitely many of) words of  $v(A)$ .
- Now for each word we have to prove that it is in  $v(B)$ .
- This is done by applying the corresponding right rules, and at steps for  $+$  and  $*$  we choose the correct way.
- For example, if we wish to prove  $a, a, a \rightarrow b + a^*$ , we first use  $+R_2$  and then  $*R_3$ .

- The standard interpretation can be generalised to **language models**.

## More Models

- The standard interpretation can be generalised to **language models**.
- In a language model,  $v(a)$  could be an arbitrary language, not necessarily  $\{a\}$ .

- The standard interpretation can be generalised to **language models**.
- In a language model,  $v(a)$  could be an arbitrary language, not necessarily  $\{a\}$ .
- Soundness still holds, and completeness gets inherited from the standard interpretation (which is a specific case of language model).

- The standard interpretation can be generalised to **language models**.
- In a language model,  $v(a)$  could be an arbitrary language, not necessarily  $\{a\}$ .
- Soundness still holds, and completeness gets inherited from the standard interpretation (which is a specific case of language model).
- Further generalisation leads to the abstract class of **\*-continuous Kleene algebras**.

## More Models

- The standard interpretation can be generalised to **language models**.
- In a language model,  $v(a)$  could be an arbitrary language, not necessarily  $\{a\}$ .
- Soundness still holds, and completeness gets inherited from the standard interpretation (which is a specific case of language model).
- Further generalisation leads to the abstract class of **\*-continuous Kleene algebras**.
- Basically, a \*-continuous Kleene algebra is any algebraic structure  $(\mathcal{A}; \preceq, \cdot, +, *, 0, 1)$  in which axioms and rules of  $\mathbf{KA}_\omega$  are valid.

## \*-Continuous Kleene Algebras

- Algebraically,  $(\mathcal{A}; \preceq, \cdot, +, *, 0, 1)$  is a \*-continuous KA, if the following holds:



## \*-Continuous Kleene Algebras

- Algebraically,  $(\mathcal{A}; \preceq, \cdot, +, *, 0, 1)$  is a \*-continuous KA, if the following holds:
  - $(\mathcal{A}; \cdot, +, 0, 1)$  is a semiring (this includes associativity for  $\cdot$  and  $+$ , commutativity for  $+$ , distributivity, and the roles of 0 and 1).

## \*-Continuous Kleene Algebras

- Algebraically,  $(\mathcal{A}; \preceq, \cdot, +, *, 0, 1)$  is a \*-continuous KA, if the following holds:
  - $(\mathcal{A}; \cdot, +, 0, 1)$  is a semiring (this includes associativity for  $\cdot$  and  $+$ , commutativity for  $+$ , distributivity, and the roles of 0 and 1).
  - Idempotency:  $a + a = a$ .

## \*-Continuous Kleene Algebras

- Algebraically,  $(\mathcal{A}; \preceq, \cdot, +, *, 0, 1)$  is a \*-continuous KA, if the following holds:
  - $(\mathcal{A}; \cdot, +, 0, 1)$  is a semiring (this includes associativity for  $\cdot$  and  $+$ , commutativity for  $+$ , distributivity, and the roles of 0 and 1).
  - Idempotency:  $a + a = a$ .
  - $a \preceq b$  iff  $a + b = b$ .

## \*-Continuous Kleene Algebras

- Algebraically,  $(\mathcal{A}; \preceq, \cdot, +, *, 0, 1)$  is a \*-continuous KA, if the following holds:
  - $(\mathcal{A}; \cdot, +, 0, 1)$  is a semiring (this includes associativity for  $\cdot$  and  $+$ , commutativity for  $+$ , distributivity, and the roles of 0 and 1).
  - Idempotency:  $a + a = a$ .
  - $a \preceq b$  iff  $a + b = b$ .
  - \*-continuity: for any  $a, b, c$ , we have  $ba^*c = \sup_{\preceq} \{ba^n c \mid n \geq 0\}$ .

## \*-Continuous Kleene Algebras

- Algebraically,  $(\mathcal{A}; \preceq, \cdot, +, *, 0, 1)$  is a \*-continuous KA, if the following holds:
  - $(\mathcal{A}; \cdot, +, 0, 1)$  is a semiring (this includes associativity for  $\cdot$  and  $+$ , commutativity for  $+$ , distributivity, and the roles of 0 and 1).
  - Idempotency:  $a + a = a$ .
  - $a \preceq b$  iff  $a + b = b$ .
  - \*-continuity: for any  $a, b, c$ , we have  $ba^*c = \sup_{\preceq} \{ba^n c \mid n \geq 0\}$ .
- Another example of \*-continuous KAs is the algebra of binary relations over a set, product being composition,  $+$  being union, and  $*$  being the reflexive-transitive closure.

## \*-Continuous Kleene Algebras

- Algebraically,  $(\mathcal{A}; \preceq, \cdot, +, *, 0, 1)$  is a \*-continuous KA, if the following holds:
  - $(\mathcal{A}; \cdot, +, 0, 1)$  is a semiring (this includes associativity for  $\cdot$  and  $+$ , commutativity for  $+$ , distributivity, and the roles of 0 and 1).
  - Idempotency:  $a + a = a$ .
  - $a \preceq b$  iff  $a + b = b$ .
  - \*-continuity: for any  $a, b, c$ , we have  $ba^*c = \sup_{\preceq} \{ba^n c \mid n \geq 0\}$ .
- Another example of \*-continuous KAs is the algebra of binary relations over a set, product being composition,  $+$  being union, and  $*$  being the reflexive-transitive closure.
- $\mathbf{KA}_\omega$  is sound and complete w.r.t. interpretations on \*-continuous KAs.

- It happens to be possible to reformulate  $\mathbf{KA}_\omega$  as a finitary calculus.

## Finite KA theory

- It happens to be possible to reformulate  $\mathbf{KA}_\omega$  as a finitary calculus.
- This gives a  $\Sigma_1^0$  upper bound, and, by Post's theorem, decidability.



# Finite KA theory

- It happens to be possible to reformulate  $\mathbf{KA}_\omega$  as a finitary calculus.
- This gives a  $\Sigma_1^0$  upper bound, and, by Post's theorem, decidability.
- Kozen and Krob prove that  $\mathbf{KA}_\omega$  is equivalent to  $\mathbf{KA}$ , the logic with the following rules for  $\rightarrow^*$ :

$$\begin{array}{c}
 \frac{\rightarrow B \quad A, B \rightarrow B}{A^* \rightarrow B} \qquad \frac{\Gamma \rightarrow A \quad \Delta \rightarrow A^*}{\Gamma, \Delta \rightarrow A^*} \qquad \frac{}{\rightarrow A^*} \\
 \\
 \frac{\rightarrow B \quad B, A \rightarrow B}{A^* \rightarrow B} \qquad \frac{\Gamma \rightarrow A^* \quad \Delta \rightarrow A}{\Gamma, \Delta \rightarrow A^*} \qquad \frac{\Pi \rightarrow A \quad \Gamma, A, \Delta \rightarrow C}{\Gamma, \Pi, \Delta \rightarrow C} \text{ Cut}
 \end{array}$$

# Finite KA theory

- It happens to be possible to reformulate  $\mathbf{KA}_\omega$  as a finitary calculus.
- This gives a  $\Sigma_1^0$  upper bound, and, by Post's theorem, decidability.
- Kozen and Krob prove that  $\mathbf{KA}_\omega$  is equivalent to  $\mathbf{KA}$ , the logic with the following rules for  $\rightarrow^*$ :

$$\begin{array}{c}
 \frac{\rightarrow B \quad A, B \rightarrow B}{A^* \rightarrow B} \qquad \frac{\Gamma \rightarrow A \quad \Delta \rightarrow A^*}{\Gamma, \Delta \rightarrow A^*} \qquad \frac{}{\rightarrow A^*} \\
 \\
 \frac{\rightarrow B \quad B, A \rightarrow B}{A^* \rightarrow B} \qquad \frac{\Gamma \rightarrow A^* \quad \Delta \rightarrow A}{\Gamma, \Delta \rightarrow A^*} \qquad \frac{\Pi \rightarrow A \quad \Gamma, A, \Delta \rightarrow C}{\Gamma, \Pi, \Delta \rightarrow C} \text{ Cut}
 \end{array}$$

- Axioms and rules for other operations are the same as in  $\mathbf{KA}_\omega$ .

**Theorem (Krob 1991, Kozen 1994)**

*KA and  $\text{KA}_\omega$  have the same set of theorems.*

# Abstract Kleene Algebras

- Algebraically, results of Kozen and Kroh mean that the inequational theory for the standard interpretation coincides with that for interpretations of reg. exp. on **arbitrary Kleene algebras**.

# Abstract Kleene Algebras

- Algebraically, results of Kozen and Krob mean that the inequational theory for the standard interpretation coincides with that for interpretations of reg. exp. on **arbitrary Kleene algebras**.
- The easiest way to define an abstract Kleene algebra: an algebraic structure which obeys **KA**.

# Abstract Kleene Algebras

- Algebraically, results of Kozen and Krohn mean that the inequational theory for the standard interpretation coincides with that for interpretations of reg. exp. on **arbitrary Kleene algebras**.
- The easiest way to define an abstract Kleene algebra: an algebraic structure which obeys **KA**.
- More accurately, Kleene algebras are defined as idempotent semirings with the following conditions for Kleene star:

# Abstract Kleene Algebras

- Algebraically, results of Kozen and Krob mean that the inequational theory for the standard interpretation coincides with that for interpretations of reg. exp. on **arbitrary Kleene algebras**.
- The easiest way to define an abstract Kleene algebra: an algebraic structure which obeys **KA**.
- More accurately, Kleene algebras are defined as idempotent semirings with the following conditions for Kleene star:
  1.  $a^*$  is the least  $x$  such that  $1 + ax \preceq x$ ;

# Abstract Kleene Algebras

- Algebraically, results of Kozen and Krob mean that the inequational theory for the standard interpretation coincides with that for interpretations of reg. exp. on **arbitrary Kleene algebras**.
- The easiest way to define an abstract Kleene algebra: an algebraic structure which obeys **KA**.
- More accurately, Kleene algebras are defined as idempotent semirings with the following conditions for Kleene star:
  1.  $a^*$  is the least  $x$  such that  $1 + ax \preceq x$ ;
  2.  $a^*$  is the least  $y$  such that  $1 + ya \preceq y$ .



# Abstract Kleene Algebras

- Algebraically, results of Kozen and Krob mean that the inequational theory for the standard interpretation coincides with that for interpretations of reg. exp. on **arbitrary Kleene algebras**.
- The easiest way to define an abstract Kleene algebra: an algebraic structure which obeys **KA**.
- More accurately, Kleene algebras are defined as idempotent semirings with the following conditions for Kleene star:
  1.  $a^*$  is the least  $x$  such that  $1 + ax \preceq x$ ;
  2.  $a^*$  is the least  $y$  such that  $1 + ya \preceq y$ .
- Thus  $a^*$  should be the left and the right Kleene star at the same time.

# Abstract Kleene Algebras

- Algebraically, results of Kozen and Krob mean that the inequational theory for the standard interpretation coincides with that for interpretations of reg. exp. on **arbitrary Kleene algebras**.
- The easiest way to define an abstract Kleene algebra: an algebraic structure which obeys **KA**.
- More accurately, Kleene algebras are defined as idempotent semirings with the following conditions for Kleene star:
  1.  $a^*$  is the least  $x$  such that  $1 + ax \preceq x$ ;
  2.  $a^*$  is the least  $y$  such that  $1 + ya \preceq y$ .
- Thus  $a^*$  should be the left and the right Kleene star at the same time.
  - There exist left and right KAs.

- $\mathbf{KA}_\omega$  has the same set of theorems as  $\mathbf{KA}$ , and they both axiomatise the inequational of each of the following classes:
  1. the standard interpretation;
  2. interpretations on the algebra of languages, with arbitrary  $v(a)$ ;
  3. interpretations on  $*$ -continuous Kleene algebras;
  4. interpretations on arbitrary Kleene algebras.

- $\mathbf{KA}_\omega$  has the same set of theorems as  $\mathbf{KA}$ , and they both axiomatise the inequational of each of the following classes:
  1. the standard interpretation;
  2. interpretations on the algebra of languages, with arbitrary  $v(a)$ ;
  3. interpretations on  $*$ -continuous Kleene algebras;
  4. interpretations on arbitrary Kleene algebras.
- Moreover, one can take only “left” versions of Kozen's rules (left-handed completeness).

- However, there exist **non- $^*$ -continuous** Kleene algebras.

- However, there exist **non- $\omega$ -continuous** Kleene algebras.
- Moreover, equivalence between **KA** and **KA $_{\omega}$**  holds only for theoremhood, not for entailment of sequents from finite sets of sequents (hypotheses).

- However, there exist **non- $\omega$ -continuous** Kleene algebras.
- Moreover, equivalence between **KA** and **KA $_{\omega}$**  holds only for theoremhood, not for entailment of sequents from finite sets of sequents (hypotheses).
- The lecturer is not aware of a concrete example at the moment, but we shall see this from complexity results further in the course.

- However, there exist **non- $\omega$ -continuous** Kleene algebras.
- Moreover, equivalence between **KA** and **KA $_{\omega}$**  holds only for theoremhood, not for entailment of sequents from finite sets of sequents (hypotheses).
- The lecturer is not aware of a concrete example at the moment, but we shall see this from complexity results further in the course.
- In this course, we are not going to prove equivalence of **KA $_{\omega}$**  and **KA**. Instead of that, we discuss alternative finitary formulations of the logic of Kleene algebras.



- The formulation of  $\mathbf{KA}$  with fixpoint rules, presented above, does not enjoy cut elimination (thus, not that good from the point of view of structural proof theory).

## Circular and Non-Well-Founded Proofs for $\mathbf{KA}_\omega$

- The formulation of  $\mathbf{KA}$  with fixpoint rules, presented above, does not enjoy cut elimination (thus, not that good from the point of view of structural proof theory).
- Das & Pous (2017) introduced a better formalization, based on **circular proofs**.

## Circular and Non-Well-Founded Proofs for $\mathbf{KA}_\omega$

- The formulation of  $\mathbf{KA}$  with fixpoint rules, presented above, does not enjoy cut elimination (thus, not that good from the point of view of structural proof theory).
- Das & Pous (2017) introduced a better formalization, based on **circular proofs**.
- We start with a naïve approach.

## Circular and Non-Well-Founded Proofs for $\mathbf{KA}_\omega$

- The formulation of  $\mathbf{KA}$  with fixpoint rules, presented above, does not enjoy cut elimination (thus, not that good from the point of view of structural proof theory).
- Das & Pous (2017) introduced a better formalization, based on **circular proofs**.
- We start with a naïve approach.
- Replace the rules for  $*$  with the following ones:

$$\frac{\Gamma, \Delta \rightarrow B \quad \Gamma, A, A^*, \Delta \rightarrow B}{\Gamma, A^*, \Delta \rightarrow B} *L \qquad \frac{}{\rightarrow A^*} *R_0 \qquad \frac{\Gamma \rightarrow A \quad \Delta \rightarrow A^*}{\Gamma, \Delta \rightarrow A^*} *R$$

and allow infinite (non-well-founded) paths in derivations, provided the following **correctness condition** holds.

## Circular and Non-Well-Founded Proofs for $\mathbf{KA}_\omega$

- The formulation of **KA** with fixpoint rules, presented above, does not enjoy cut elimination (thus, not that good from the point of view of structural proof theory).
- Das & Pous (2017) introduced a better formalization, based on **circular proofs**.
- We start with a naïve approach.
- Replace the rules for  $*$  with the following ones:

$$\frac{\Gamma, \Delta \rightarrow B \quad \Gamma, A, A^*, \Delta \rightarrow B}{\Gamma, A^*, \Delta \rightarrow B} *L \qquad \frac{}{\rightarrow A^*} *R_0 \qquad \frac{\Gamma \rightarrow A \quad \Delta \rightarrow A^*}{\Gamma, \Delta \rightarrow A^*} *R$$

and allow infinite (non-well-founded) paths in derivations, provided the following **correctness condition** holds.

- Correctness condition: for every infinite branch, there is a trace of  $A^*$ , which undergoes  $*L$  infinitely many times.

## Non-Well-Founded Proofs for $\mathbf{KA}_{\omega}$

- Let us denote the system with non-well-founded proofs by  $\mathbf{KA}_{\infty}$ .

## Non-Well-Founded Proofs for $\mathbf{KA}_{\omega}$

- Let us denote the system with non-well-founded proofs by  $\mathbf{KA}_{\infty}$ .
- Das & Pous (2018) prove that  $\mathbf{KA}_{\infty}$  enjoys cut elimination.

## Non-Well-Founded Proofs for $\mathbf{KA}_{\omega}$

- Let us denote the system with non-well-founded proofs by  $\mathbf{KA}_{\infty}$ .
- Das & Pous (2018) prove that  $\mathbf{KA}_{\infty}$  enjoys cut elimination.
- We shall not need this result, but rather consider only the cut-free version of  $\mathbf{KA}_{\infty}$ .



## Non-Well-Founded Proofs for $\mathbf{KA}_{\omega}$

- Let us denote the system with non-well-founded proofs by  $\mathbf{KA}_{\infty}$ .
- Das & Pous (2018) prove that  $\mathbf{KA}_{\infty}$  enjoys cut elimination.
- We shall not need this result, but rather consider only the cut-free version of  $\mathbf{KA}_{\infty}$ .
- Moreover, we require  $\Pi$  to be non-empty in  $*R$  (otherwise the rule is meaningless).

## Non-Well-Founded Proofs for $\mathbf{KA}_{\omega}$

- Let us denote the system with non-well-founded proofs by  $\mathbf{KA}_{\infty}$ .
- Das & Pous (2018) prove that  $\mathbf{KA}_{\infty}$  enjoys cut elimination.
- We shall not need this result, but rather consider only the cut-free version of  $\mathbf{KA}_{\infty}$ .
- Moreover, we require  $\Pi$  to be non-empty in  $*R$  (otherwise the rule is meaningless).
- Under these assumptions, no correctness condition is needed!

## Non-Well-Founded Proofs for $\mathbf{KA}_{\omega}$

- Let us denote the system with non-well-founded proofs by  $\mathbf{KA}_{\infty}$ .
- Das & Pous (2018) prove that  $\mathbf{KA}_{\infty}$  enjoys cut elimination.
- We shall not need this result, but rather consider only the cut-free version of  $\mathbf{KA}_{\infty}$ .
- Moreover, we require  $\Pi$  to be non-empty in  $*R$  (otherwise the rule is meaningless).
- Under these assumptions, no correctness condition is needed!
  - Indeed, in all rules, except  $*L$ , the premises are strictly simpler than the conclusion.

## Non-Well-Founded Proofs for $\mathbf{KA}_{\omega}$

- Let us denote the system with non-well-founded proofs by  $\mathbf{KA}_{\infty}$ .
- Das & Pous (2018) prove that  $\mathbf{KA}_{\infty}$  enjoys cut elimination.
- We shall not need this result, but rather consider only the cut-free version of  $\mathbf{KA}_{\infty}$ .
- Moreover, we require  $\Pi$  to be non-empty in  $*R$  (otherwise the rule is meaningless).
- Under these assumptions, no correctness condition is needed!
  - Indeed, in all rules, except  $*L$ , the premises are strictly simpler than the conclusion.
  - Thus, each infinite branch traverses  $*L$  infinitely many times.

## Non-Well-Founded Proofs for $\mathbf{KA}_{\omega}$

- Let us denote the system with non-well-founded proofs by  $\mathbf{KA}_{\infty}$ .
- Das & Pous (2018) prove that  $\mathbf{KA}_{\infty}$  enjoys cut elimination.
- We shall not need this result, but rather consider only the cut-free version of  $\mathbf{KA}_{\infty}$ .
- Moreover, we require  $\Pi$  to be non-empty in  $*R$  (otherwise the rule is meaningless).
- Under these assumptions, no correctness condition is needed!
  - Indeed, in all rules, except  $*L$ , the premises are strictly simpler than the conclusion.
  - Thus, each infinite branch traverses  $*L$  infinitely many times.
  - Moreover, the number of  $A^*$ 's is finite, so by pigeonhole principle there will be a trace.

## Theorem (Das & Pous 2018)

$\mathbf{KA}_\omega$  and  $\mathbf{KA}_\infty$  (its cut-free version) derive the same set of theorems.

## Theorem (Das & Pous 2018)

$\mathbf{KA}_\omega$  and  $\mathbf{KA}_\infty$  (its cut-free version) derive the same set of theorems.

- The direction from  $\mathbf{KA}_\omega$  to  $\mathbf{KA}_\infty$  is easy: we just model the  $\omega$ -rule by an infinite branch:

$$\frac{\Gamma, \Delta \rightarrow B}{\Gamma, A^*, \Delta \rightarrow B} *L \quad \frac{\Gamma, A, \Delta \rightarrow B \quad \frac{\Gamma, A, A, \Delta \rightarrow B \quad \frac{\Gamma, A, A, A^*, \Delta \rightarrow B}{\Gamma, A, A, A^*, \Delta \rightarrow B} \vdots}{\Gamma, A, A, A^*, \Delta \rightarrow B} *L}{\Gamma, A, A^*, \Delta \rightarrow B} *L$$

## Theorem (Das & Pous 2018)

$\mathbf{KA}_\omega$  and  $\mathbf{KA}_\infty$  (its cut-free version) derive the same set of theorems.

- The direction from  $\mathbf{KA}_\omega$  to  $\mathbf{KA}_\infty$  is easy: we just model the  $\omega$ -rule by an infinite branch:

$$\frac{\Gamma, \Delta \rightarrow B}{\Gamma, A^*, \Delta \rightarrow B} *L \quad \frac{\Gamma, A, \Delta \rightarrow B \quad \frac{\Gamma, A, A, A^*, \Delta \rightarrow B}{\Gamma, A, A, A^*, \Delta \rightarrow B} *L \quad \frac{\Gamma, A, A, A^*, \Delta \rightarrow B}{\Gamma, A, A, A^*, \Delta \rightarrow B} *L \quad \vdots}{\Gamma, A, A, A^*, \Delta \rightarrow B} *L$$

- The  $*R_n$  rules is several iterations of  $*R$ .



- The opposite direction is based on the following lemma:

## Lemma

*If  $\mathbf{KA}_\infty \vdash \Gamma, A^*, \Delta \rightarrow B$ , then  $\mathbf{KA}_\infty \vdash \Gamma, A^n, \Delta \rightarrow B$  for each  $n$ .*

- This lemma is proved by induction on  $n$ .
  - We replace  $A^*$  with  $A^n$  and go upwards the proof. At the points of  $*L$  we refer to the induction hypothesis.
- Now we may eagerly apply the  $\omega$ -rule, and translate finitary rules to  $\mathbf{KA}_\omega$ .

- As we have shown, infinite proofs correspond to the  $\omega$ -rule.

- As we have shown, infinite proofs correspond to the  $\omega$ -rule.
- In contrast, **induction** (fixpoint rules) correspond to **circular** proofs.

- As we have shown, infinite proofs correspond to the  $\omega$ -rule.
- In contrast, **induction** (fixpoint rules) correspond to **circular** proofs.
- In a circular derivation, each infinite path at some point reaches **the same** sequent as already seen below.

- As we have shown, infinite proofs correspond to the  $\omega$ -rule.
- In contrast, **induction** (fixpoint rules) correspond to **circular** proofs.
- In a circular derivation, each infinite path at some point reaches **the same** sequent as already seen below.
- At this point, there is no need to develop the derivation further, but one can rather refer backwards via a **backlink**.

- As we have shown, infinite proofs correspond to the  $\omega$ -rule.
- In contrast, **induction** (fixpoint rules) correspond to **circular** proofs.
- In a circular derivation, each infinite path at some point reaches **the same** sequent as already seen below.
- At this point, there is no need to develop the derivation further, but one can rather refer backwards via a **backlink**.
- Alternatively, circular proofs can be defined as **regular** non-well-founded proofs, i.e., proofs with a finite number of non-isomorphic subtrees.

- As we have shown, infinite proofs correspond to the  $\omega$ -rule.
- In contrast, **induction** (fixpoint rules) correspond to **circular** proofs.
- In a circular derivation, each infinite path at some point reaches **the same** sequent as already seen below.
- At this point, there is no need to develop the derivation further, but one can rather refer backwards via a **backlink**.
- Alternatively, circular proofs can be defined as **regular** non-well-founded proofs, i.e., proofs with a finite number of non-isomorphic subtrees.
- We consider circular proofs with cuts, obeying the correctness condition.

## Circular Proofs for $\mathbf{KA}_\omega$

- As we have shown, infinite proofs correspond to the  $\omega$ -rule.
- In contrast, **induction** (fixpoint rules) correspond to **circular** proofs.
- In a circular derivation, each infinite path at some point reaches **the same** sequent as already seen below.
- At this point, there is no need to develop the derivation further, but one can rather refer backwards via a **backlink**.
- Alternatively, circular proofs can be defined as **regular** non-well-founded proofs, i.e., proofs with a finite number of non-isomorphic subtrees.
- We consider circular proofs with cuts, obeying the correctness condition.
- Let us denote this system by  $\mathbf{KA}_{\omega\circ}$ .



# Circular Derivation: Example

$$\begin{array}{c}
 \frac{a \rightarrow a \quad (a+b)^* \rightarrow a^*(ba^*)^*}{a, (a+b)^* \rightarrow aa^*(ba^*)^*} \quad \frac{aa^*(ba^*)^* \rightarrow a^*(ba^*)^*}{a, (a+b)^* \rightarrow a^*(ba^*)^*} \quad \frac{a, (a+b)^* \rightarrow a^*(ba^*)^*}{\rightarrow a^*(ba^*)^*} \\
 \\
 \frac{b \rightarrow b \quad (a+b)^* \rightarrow a^*(ba^*)^*}{b, (a+b)^* \rightarrow ba^*(ba^*)^*} \quad \frac{ba^*(ba^*)^* \rightarrow (ba^*)^*}{b, (a+b)^* \rightarrow (ba^*)^*} \quad \frac{b, (a+b)^* \rightarrow ba^*(ba^*)^* \quad b, (a+b)^* \rightarrow (ba^*)^*}{b, (a+b)^* \rightarrow a^*(ba^*)^*} \\
 \\
 \frac{a+b, (a+b)^* \rightarrow a^*(ba^*)^*}{(a+b)^* \rightarrow a^*(ba^*)^*} \quad *L
 \end{array}$$

- $KA$  and  $KA_{\circ}$  derive the same set of theorems.

- $\mathbf{KA}$  and  $\mathbf{KA}_{\circlearrowleft}$  derive the same set of theorems.
- Unfortunately, cut elimination for  $\mathbf{KA}_{\circlearrowleft}$  again fails: examples include  $(a + b)^* \rightarrow a^*(ba^*)^*$  (see above) and even  $aa^* \rightarrow a^*a$ .

- $\mathbf{KA}$  and  $\mathbf{KA}_{\circlearrowleft}$  derive the same set of theorems.
- Unfortunately, cut elimination for  $\mathbf{KA}_{\circlearrowleft}$  again fails: examples include  $(a + b)^* \rightarrow a^*(ba^*)^*$  (see above) and even  $aa^* \rightarrow a^*a$ .
- A sequent may have a circular proof with cuts, but unravelling it to an infinite proof and eliminating cuts yields a non-circular proof.

- $\mathbf{KA}$  and  $\mathbf{KA}_{\circlearrowleft}$  derive the same set of theorems.
- Unfortunately, cut elimination for  $\mathbf{KA}_{\circlearrowleft}$  again fails: examples include  $(a + b)^* \rightarrow a^*(ba^*)^*$  (see above) and even  $aa^* \rightarrow a^*a$ .
- A sequent may have a circular proof with cuts, but unravelling it to an infinite proof and eliminating cuts yields a non-circular proof.
- Furthermore, it is unclear how to reduce  $\mathbf{KA}_{\infty}$  to  $\mathbf{KA}_{\circlearrowleft}$  (even with cuts).

- $\mathbf{KA}$  and  $\mathbf{KA}_{\circlearrowleft}$  derive the same set of theorems.
- Unfortunately, cut elimination for  $\mathbf{KA}_{\circlearrowleft}$  again fails: examples include  $(a + b)^* \rightarrow a^*(ba^*)^*$  (see above) and even  $aa^* \rightarrow a^*a$ .
- A sequent may have a circular proof with cuts, but unravelling it to an infinite proof and eliminating cuts yields a non-circular proof.
- Furthermore, it is unclear how to reduce  $\mathbf{KA}_{\infty}$  to  $\mathbf{KA}_{\circlearrowleft}$  (even with cuts).
- However, it is possible to construct a cut-free circular calculus for  $\mathbf{KA}$ , but with a more involved sequential syntax.

- $\mathbf{KA}$  and  $\mathbf{KA}_{\circlearrowleft}$  derive the same set of theorems.
- Unfortunately, cut elimination for  $\mathbf{KA}_{\circlearrowleft}$  again fails: examples include  $(a + b)^* \rightarrow a^*(ba^*)^*$  (see above) and even  $aa^* \rightarrow a^*a$ .
- A sequent may have a circular proof with cuts, but unravelling it to an infinite proof and eliminating cuts yields a non-circular proof.
- Furthermore, it is unclear how to reduce  $\mathbf{KA}_{\infty}$  to  $\mathbf{KA}_{\circlearrowleft}$  (even with cuts).
- However, it is possible to construct a cut-free circular calculus for  $\mathbf{KA}$ , but with a more involved sequential syntax.
- We shall discuss this system tomorrow.