

Complexity of Reasoning in Kleene and Action Algebras

Stepan Kuznetsov

ESSLI 2022 · Galway, Ireland · Gallimh, Éire

Lecture 4

Divide and Conquer

- Now let us add **division** operators to Kleene algebras.

Divide and Conquer

- Now let us add **division** operators to Kleene algebras.
- $A \preceq C / B \iff A \cdot B \preceq C \iff B \preceq A \setminus C$

Divide and Conquer

- Now let us add **division** operators to Kleene algebras.
- $A \preceq C / B \iff A \cdot B \preceq C \iff B \preceq A \setminus C$
- Divisions have a natural language interpretation:

$$A \setminus B = \{\beta \mid (\forall \alpha \in A) \alpha\beta \in B\}$$

$$B / A = \{\beta \mid (\forall \alpha \in A) \beta\alpha \in B\}$$

Divide and Conquer

- Now let us add **division** operators to Kleene algebras.
- $A \preceq C / B \iff A \cdot B \preceq C \iff B \preceq A \setminus C$
- Divisions have a natural language interpretation:

$$A \setminus B = \{\beta \mid (\forall \alpha \in A) \alpha\beta \in B\}$$

$$B / A = \{\beta \mid (\forall \alpha \in A) \beta\alpha \in B\}$$

- Kleene algebras with divisions (residuated Kleene algebras) are called **action algebras**.

Divide and Conquer

- Now let us add **division** operators to Kleene algebras.
- $A \preceq C / B \iff A \cdot B \preceq C \iff B \preceq A \setminus C$
- Divisions have a natural language interpretation:

$$A \setminus B = \{\beta \mid (\forall \alpha \in A) \alpha\beta \in B\}$$

$$B / A = \{\beta \mid (\forall \alpha \in A) \beta\alpha \in B\}$$

- Kleene algebras with divisions (residuated Kleene algebras) are called **action algebras**.
- Adding conjunction—intersection yields **action lattices**.

Divide and Conquer

- Now let us add **division** operators to Kleene algebras.
- $A \preceq C / B \iff A \cdot B \preceq C \iff B \preceq A \setminus C$
- Divisions have a natural language interpretation:

$$A \setminus B = \{\beta \mid (\forall \alpha \in A) \alpha\beta \in B\}$$

$$B / A = \{\beta \mid (\forall \alpha \in A) \beta\alpha \in B\}$$

- Kleene algebras with divisions (residuated Kleene algebras) are called **action algebras**.
- Adding conjunction—intersection yields **action lattices**.
- The logic of all action lattices is action logic **ACT**.

Divide and Conquer

- Now let us add **division** operators to Kleene algebras.
- $A \preceq C / B \iff A \cdot B \preceq C \iff B \preceq A \setminus C$
- Divisions have a natural language interpretation:

$$A \setminus B = \{\beta \mid (\forall \alpha \in A) \alpha\beta \in B\}$$

$$B / A = \{\beta \mid (\forall \alpha \in A) \beta\alpha \in B\}$$

- Kleene algebras with divisions (residuated Kleene algebras) are called **action algebras**.
- Adding conjunction—intersection yields **action lattices**.
- The logic of all action lattices is action logic **ACT**.
- The logic of $*$ -continuous action lattices is infinitary action logic **ACT** $_{\omega}$.

- An **action lattice** is an algebraic structure $(\mathcal{A}; \preceq, \backslash, /, \vee, \wedge, 0, 1, *)$, where:

- An **action lattice** is an algebraic structure $(\mathcal{A}; \preceq, \backslash, /, \vee, \wedge, 0, 1, *)$, where:
 1. $(\mathcal{A}; \preceq, \vee, \wedge)$ is a lattice, 0 is its least element;

- An **action lattice** is an algebraic structure $(\mathcal{A}; \preceq, \backslash, /, \vee, \wedge, 0, 1, *)$, where:
 1. $(\mathcal{A}; \preceq, \vee, \wedge)$ is a lattice, 0 is its least element;
 2. $(\mathcal{A}; \cdot, 1)$ is a monoid;

- An **action lattice** is an algebraic structure $(\mathcal{A}; \preceq, \backslash, /, \vee, \wedge, 0, 1, \cdot)$, where:
 1. $(\mathcal{A}; \preceq, \vee, \wedge)$ is a lattice, 0 is its least element;
 2. $(\mathcal{A}; \cdot, 1)$ is a monoid;
 3. \backslash and $/$ are residuals of the product w.r.t. \preceq :
$$A \preceq C / B \iff A \cdot B \preceq C \iff B \preceq A \backslash C;$$

- An **action lattice** is an algebraic structure $(\mathcal{A}; \preceq, \backslash, /, \vee, \wedge, 0, 1, *)$, where:
 1. $(\mathcal{A}; \preceq, \vee, \wedge)$ is a lattice, 0 is its least element;
 2. $(\mathcal{A}; \cdot, 1)$ is a monoid;
 3. \backslash and $/$ are residuals of the product w.r.t. \preceq :
$$A \preceq C / B \iff A \cdot B \preceq C \iff B \preceq A \backslash C;$$
 4. $A^* = \min_{\preceq} \{b \mid 1 \preceq b \text{ and } a \cdot b \preceq b\}.$

- An **action lattice** is an algebraic structure $(\mathcal{A}; \preceq, \backslash, /, \vee, \wedge, 0, 1, *)$, where:
 1. $(\mathcal{A}; \preceq, \vee, \wedge)$ is a lattice, 0 is its least element;
 2. $(\mathcal{A}; \cdot, 1)$ is a monoid;
 3. \backslash and $/$ are residuals of the product w.r.t. \preceq :
$$A \preceq C / B \iff A \cdot B \preceq C \iff B \preceq A \backslash C;$$
 4. $A^* = \min_{\preceq} \{b \mid 1 \preceq b \text{ and } a \cdot b \preceq b\}.$
- An action lattice is $*$ -continuous, if $a^* = \sup_{\preceq} \{a^n \mid n \geq 0\}.$

- In the presence of divisions, many things regularise:

- In the presence of divisions, many things regularise:
 1. Product is monotone w.r.t. \preceq (Lambek 1958).

- In the presence of divisions, many things regularise:
 1. Product is monotone w.r.t. \preceq (Lambek 1958).
 2. The left Kleene star is always the right one.

- In the presence of divisions, many things regularise:
 1. Product is monotone w.r.t. \preceq (Lambek 1958).
 2. The left Kleene star is always the right one.
 3. The zero element is the annihilator for product.

- In the presence of divisions, many things regularise:
 1. Product is monotone w.r.t. \preceq (Lambek 1958).
 2. The left Kleene star is always the right one.
 3. The zero element is the annihilator for product.
- The following **Pratt's normality theorem** holds: if there exists $\sup_{\preceq} \{a^n \mid n \geq 0\}$, then it should coincide with a^* .

- In the presence of divisions, many things regularise:
 1. Product is monotone w.r.t. \preceq (Lambek 1958).
 2. The left Kleene star is always the right one.
 3. The zero element is the annihilator for product.
- The following **Pratt's normality theorem** holds: if there exists $\sup_{\preceq} \{a^n \mid n \geq 0\}$, then it should coincide with a^* .
- This means that if the lattice is complete (allows infinite inf and sup), then it is * -continuous.

- In the presence of divisions, many things regularise:
 1. Product is monotone w.r.t. \leq (Lambek 1958).
 2. The left Kleene star is always the right one.
 3. The zero element is the annihilator for product.
- The following **Pratt's normality theorem** holds: if there exists $\sup_{\leq} \{a^n \mid n \geq 0\}$, then it should coincide with a^* .
- This means that if the lattice is complete (allows infinite inf and sup), then it is * -continuous.
- In particular, this holds for finite action lattices.

History and Motivation

- The first (probably) appearance of divisions (residuals) connected to a partial order is due to Krull (1924), who introduced them in algebra, for ideals in rings.

History and Motivation

- The first (probably) appearance of divisions (residuals) connected to a partial order is due to Krull (1924), who introduced them in algebra, for ideals in rings.
- **Residuated lattices** appear in the work of Ward & Dilworth (1939).

History and Motivation

- The first (probably) appearance of divisions (residuals) connected to a partial order is due to Krull (1924), who introduced them in algebra, for ideals in rings.
- **Residuated lattices** appear in the work of Ward & Dilworth (1939).
- Later on, Lambek (1958) introduced the Lambek calculus for defining natural language syntax.

History and Motivation

- The first (probably) appearance of divisions (residuals) connected to a partial order is due to Krull (1924), who introduced them in algebra, for ideals in rings.
- **Residuated lattices** appear in the work of Ward & Dilworth (1939).
- Later on, Lambek (1958) introduced the Lambek calculus for defining natural language syntax.
- The Lambek calculus is a basic substructural logic; on the connection of substructural logics and residuated lattices see Galatos et al. (2007).

History and Motivation

- Action algebras were introduced by Pratt (1991); Kozen (1994) added \wedge and introduced action lattices.

History and Motivation

- Action algebras were introduced by Pratt (1991); Kozen (1994) added \wedge and introduced action lattices.
- The motivation is in better properties of this class of algebras if compared with Kleene algebras.

History and Motivation

- Action algebras were introduced by Pratt (1991); Kozen (1994) added \wedge and introduced action lattices.
- The motivation is in better properties of this class of algebras if compared with Kleene algebras.
- Namely, Kleene algebras do not form a finitely based variety (Redko 1964, Conway 1971), i.e., they cannot be axiomatised by a finite set of universally valid equations.

History and Motivation

- Action algebras were introduced by Pratt (1991); Kozen (1994) added \wedge and introduced action lattices.
- The motivation is in better properties of this class of algebras if compared with Kleene algebras.
- Namely, Kleene algebras do not form a finitely based variety (Redko 1964, Conway 1971), i.e., they cannot be axiomatised by a finite set of universally valid equations.
- In contrast, action algebras do. Namely, as shown by Pratt, the condition for Kleene star can be replaced by “**pure induction**”

$$(A \setminus A)^* = A \setminus A,$$

and monotonicity: $A^* \preceq (A + B)^*$.

Standard Examples of Action Lattices

Standard Examples of Action Lattices

- $\mathcal{P}(\Sigma^*)$, the algebra of formal languages:
 - the lattice structure is set-theoretic;
 - \cdot is pairwise concatenation, $1 = \{\varepsilon\}$;
 - $x / y = \{u \in \Sigma^* \mid (\forall v \in y) uv \in x\}$,
 $y \setminus x = \{u \in \Sigma^* \mid (\forall v \in y) vu \in x\}$;
 - $x^* = \{u_1 \dots u_n \mid n \geq 0, u_i \in x\}$.

Standard Examples of Action Lattices

- $\mathcal{P}(\Sigma^*)$, the algebra of formal languages:
 - the lattice structure is set-theoretic;
 - \cdot is pairwise concatenation, $1 = \{\varepsilon\}$;
 - $x / y = \{u \in \Sigma^* \mid (\forall v \in y) uv \in x\}$,
 $y \setminus x = \{u \in \Sigma^* \mid (\forall v \in y) vu \in x\}$;
 - $x^* = \{u_1 \dots u_n \mid n \geq 0, u_i \in x\}$.
- $\mathcal{P}(W \times W)$, the algebra of relations:
 - the lattice structure is set-theoretic;
 - \cdot is composition of relations, 1 is the diagonal;
 - $x / y = \{\langle a, b \rangle \mid (\forall \langle b, c \rangle \in y) \langle a, c \rangle \in x\}$,
 $y \setminus x = \{\langle b, c \rangle \mid (\forall \langle a, b \rangle \in y) \langle a, c \rangle \in x\}$;
 - x^* is the reflexive-transitive closure of x .

Standard Examples of Action Lattices

- $\mathcal{P}(\Sigma^*)$, the algebra of formal languages:
 - the lattice structure is set-theoretic;
 - \cdot is pairwise concatenation, $1 = \{\varepsilon\}$;
 - $x / y = \{u \in \Sigma^* \mid (\forall v \in y) uv \in x\}$,
 $y \setminus x = \{u \in \Sigma^* \mid (\forall v \in y) vu \in x\}$;
 - $x^* = \{u_1 \dots u_n \mid n \geq 0, u_i \in x\}$.
- $\mathcal{P}(W \times W)$, the algebra of relations:
 - the lattice structure is set-theoretic;
 - \cdot is composition of relations, 1 is the diagonal;
 - $x / y = \{\langle a, b \rangle \mid (\forall \langle b, c \rangle \in y) \langle a, c \rangle \in x\}$,
 $y \setminus x = \{\langle b, c \rangle \mid (\forall \langle a, b \rangle \in y) \langle a, c \rangle \in x\}$;
 - x^* is the reflexive-transitive closure of x .

These action lattices are *distributive* (as lattices) and * -continuous.

- However, non- \ast -continuous action lattices exist.

Action Lattices

- However, non- \ast -continuous action lattices exist.
- This is an example, modifying Kozen's one (1990) by adding divisions and avoiding suprema:

$$\begin{array}{ccccccc} \bullet & \mapsto & \longleftrightarrow & \longleftrightarrow & \dots & \bullet \\ \perp & \mathbb{N} & \mathbb{Z} & \mathbb{Z} & & \top \end{array}$$

Action Lattices

- However, non- \ast -continuous action lattices exist.
- This is an example, modifying Kozen's one (1990) by adding divisions and avoiding suprema:

$$\begin{array}{ccccccc} \bullet & \longmapsto & \longleftrightarrow & \longleftrightarrow & \dots & \bullet \\ \perp & \mathbb{N} & \mathbb{Z} & \mathbb{Z} & & \top \end{array}$$

- Multiplication: componentwise addition in the middle;
 $\perp \cdot x = x \cdot \perp = \perp$; $\top \cdot y = y \cdot \top = \top$ for $y \neq \perp$.

Action Lattices

- However, non- $*$ -continuous action lattices exist.
- This is an example, modifying Kozen's one (1990) by adding divisions and avoiding suprema:

$$\begin{array}{ccccccc} \bullet & \mapsto & \longleftrightarrow & \longleftrightarrow & \dots & \bullet \\ \perp & \mathbb{N} & \mathbb{Z} & \mathbb{Z} & & \top \end{array}$$

- Multiplication: componentwise addition in the middle;
 $\perp \cdot x = x \cdot \perp = \perp$; $\top \cdot y = y \cdot \top = \top$ for $y \neq \perp$.
- Unit: $1 = (0, 0)$.

Action Lattices

- However, non- \ast -continuous action lattices exist.
- This is an example, modifying Kozen's one (1990) by adding divisions and avoiding suprema:

$$\begin{array}{ccccccc} \bullet & \longmapsto & \longleftrightarrow & \longleftrightarrow & \dots & \bullet \\ \perp & \mathbb{N} & \mathbb{Z} & \mathbb{Z} & & \top \end{array}$$

- Multiplication: componentwise addition in the middle;
 $\perp \cdot x = x \cdot \perp = \perp$; $\top \cdot y = y \cdot \top = \top$ for $y \neq \perp$.
- Unit: $1 = (0, 0)$.
- One can show that this monoid is residuated.

Action Lattices

- However, non- \ast -continuous action lattices exist.
- This is an example, modifying Kozen's one (1990) by adding divisions and avoiding suprema:

$$\begin{array}{ccccccc} \bullet & \longmapsto & \longleftrightarrow & \longleftrightarrow & \dots & \bullet \\ \perp & \mathbb{N} & \mathbb{Z} & \mathbb{Z} & & \top \end{array}$$

- Multiplication: componentwise addition in the middle;
 $\perp \cdot x = x \cdot \perp = \perp$; $\top \cdot y = y \cdot \top = \top$ for $y \neq \perp$.
- Unit: $1 = (0, 0)$.
- One can show that this monoid is residuated.
- $x^\ast = \top$ for $x \succ (0, 0)$; $\perp^\ast = (0, 0)^\ast = (0, 0)$.

Action Lattices

- However, non- \ast -continuous action lattices exist.
- This is an example, modifying Kozen's one (1990) by adding divisions and avoiding suprema:

$$\begin{array}{ccccccc} \bullet & \longmapsto & \longleftrightarrow & \longleftrightarrow & \dots & \bullet \\ \perp & \mathbb{N} & \mathbb{Z} & \mathbb{Z} & & \top \end{array}$$

- Multiplication: componentwise addition in the middle;
 $\perp \cdot x = x \cdot \perp = \perp$; $\top \cdot y = y \cdot \top = \top$ for $y \neq \perp$.
- Unit: $1 = (0, 0)$.
- One can show that this monoid is residuated.
- $x^* = \top$ for $x \succ (0, 0)$; $\perp^* = (0, 0)^* = (0, 0)$.
- $\sup\{(0, 1)^n \mid n \geq 0\} = \sup\{(0, n) \mid n \geq 0\}$ does not exist.

- However, non- \ast -continuous action lattices exist.
- This is an example, modifying Kozen's one (1990) by adding divisions and avoiding suprema:

$$\begin{array}{ccccccc}
 \bullet & \longmapsto & \longleftrightarrow & \longleftrightarrow & \dots & \bullet \\
 \perp & \mathbb{N} & \mathbb{Z} & \mathbb{Z} & & \top
 \end{array}$$

- Multiplication: componentwise addition in the middle;
 $\perp \cdot x = x \cdot \perp = \perp$; $\top \cdot y = y \cdot \top = \top$ for $y \neq \perp$.
- Unit: $1 = (0, 0)$.
- One can show that this monoid is residuated.
- $x^\ast = \top$ for $x \succ (0, 0)$; $\perp^\ast = (0, 0)^\ast = (0, 0)$.
- $\sup\{(0, 1)^n \mid n \geq 0\} = \sup\{(0, n) \mid n \geq 0\}$ does not exist.
- Extra properties: *commutativity* and *linearity* of \preceq .

- Moreover, unlike the situation with Kleene algebras, for action lattices the **inequational theories** differ in the general case and in the * -continuous one.

- Moreover, unlike the situation with Kleene algebras, for action lattices the **inequational theories** differ in the general case and in the $*$ -continuous one.
- This will follow from complexity considerations below, but there is also an explicit example.

- Moreover, unlike the situation with Kleene algebras, for action lattices the **inequational theories** differ in the general case and in the * -continuous one.
- This will follow from complexity considerations below, but there is also an explicit example.
- This example is based on the following **induction-in-the-middle** rule:

$$\frac{\rightarrow B \quad A \rightarrow B \quad A, B, A \rightarrow B}{A^* \rightarrow B} *L_{\text{mid}}$$

- Moreover, unlike the situation with Kleene algebras, for action lattices the **inequational theories** differ in the general case and in the $*$ -continuous one.
- This will follow from complexity considerations below, but there is also an explicit example.
- This example is based on the following **induction-in-the-middle** rule:

$$\frac{\rightarrow B \quad A \rightarrow B \quad A, B, A \rightarrow B}{A^* \rightarrow B} *L_{\text{mid}}$$

- This allows deriving the following sequent:

$$(p \wedge q \wedge (p \setminus q) \wedge (p / q))^+ \rightarrow p,$$

which can be falsified on a non- $*$ -continuous action algebra.

Theorem (Buszkowski & Palka 2007)

ACT_ω is Π_1^0 -complete.

Theorem (K. 2019–20)

ACT is Σ_1^0 -complete.

Theorem (Buszkowski & Palka 2007)

ACT_ω is Π_1^0 -complete.

Theorem (K. 2019–20)

ACT is Σ_1^0 -complete.

- For simplicity, in what follows we shift to the commutative situation, and consider CommACT_ω and CommACT .

Theorem (Buszkowski & Palka 2007)

ACT_ω is Π_1^0 -complete.

Theorem (K. 2019–20)

ACT is Σ_1^0 -complete.

- For simplicity, in what follows we shift to the commutative situation, and consider CommACT_ω and CommACT .
- In the commutative case, we have only one division:
$$A \setminus B \equiv B / A.$$

A commutative action algebra is an action algebra satisfying $ab = ba$. Whereas action logic in general is neutral as to whether ab combines a and b sequentially or concurrently, commutative action logic in effect commits to concurrency.

Pratt (1991)

Multiplicative-additive Lambek Calculus

- The core of **ACT** and **ACT** _{ω} is **MALC**, the inequational theory of residuated lattices.

Multiplicative-additive Lambek Calculus

- The core of **ACT** and **ACT** _{ω} is **MALC**, the inequational theory of residuated lattices.
- We consider its commutative modification **CommMALC**, that is, left-hand sides are multisets.

Multiplicative-additive Lambek Calculus

- The core of **ACT** and **ACT**_ω is **MALC**, the inequational theory of residuated lattices.
- We consider its commutative modification **CommMALC**, that is, left-hand sides are multisets.
- Axioms and rules: $\overline{A \rightarrow A} \text{ } Id \quad \overline{\Gamma, 0 \rightarrow B} \text{ } 0L$

$$\frac{\Gamma \rightarrow B}{\Gamma, 1 \rightarrow B} 1L \quad \frac{}{\rightarrow 1} 1R \quad \frac{\Gamma, A, B \rightarrow C}{\Gamma, A \cdot B \rightarrow C} \cdot L \quad \frac{\Gamma \rightarrow A \quad \Delta \rightarrow B}{\Gamma, \Delta \rightarrow A \cdot B} \cdot R$$

$$\frac{\Gamma, A \rightarrow C \quad \Gamma, B \rightarrow C}{\Gamma, A \vee B \rightarrow C} \vee L \quad \frac{\Pi \rightarrow A}{\Pi \rightarrow A \vee B} \vee R_1 \quad \frac{\Pi \rightarrow B}{\Pi \rightarrow A \vee B} \vee R_2$$

$$\frac{\Gamma, A \rightarrow C}{\Gamma, A \wedge B \rightarrow C} \wedge L_1 \quad \frac{\Gamma, B \rightarrow C}{\Gamma, A \wedge B \rightarrow C} \wedge L_2 \quad \frac{\Pi \rightarrow A \quad \Pi \rightarrow B}{\Pi \rightarrow A \wedge B} \wedge R$$

$$\frac{\Pi \rightarrow A \quad \Gamma, B \rightarrow C}{\Gamma, \Pi, A \setminus B \rightarrow C} \setminus L \quad \frac{A, \Pi \rightarrow B}{\Pi \rightarrow A \setminus B} / L$$

- In **CommACT**_ω:

$$\frac{(\Gamma, A^n \rightarrow C)_{n=0}^{\infty}}{\Gamma, A^* \rightarrow C} *L_{\omega} \qquad \frac{\Gamma_1 \rightarrow A \quad \dots \quad \Gamma_n \rightarrow A}{\Gamma_1, \dots, \Gamma_n \rightarrow A^*} *R_n, \, n \geq 0$$

Rules for Kleene Star

- In **CommACT**_ω:

$$\frac{(\Gamma, A^n \rightarrow C)_{n=0}^{\infty}}{\Gamma, A^* \rightarrow C} *L_{\omega} \qquad \frac{\Gamma_1 \rightarrow A \quad \dots \quad \Gamma_n \rightarrow A}{\Gamma_1, \dots, \Gamma_n \rightarrow A^*} *R_n, n \geq 0$$

- In **CommACT**:

$$\frac{\rightarrow B \quad A, B \rightarrow B}{A^* \rightarrow B} \qquad \frac{}{\rightarrow A^*} \qquad \frac{\Gamma \rightarrow A \quad \Delta \rightarrow A^*}{\Gamma, \Delta \rightarrow A^*}$$

$$\frac{\Pi \rightarrow A \quad \Gamma, A \rightarrow C}{\Gamma, \Pi \rightarrow C} \textit{Cut}$$

Minsky Machines

- A Minsky machine \mathcal{M} operates several *counters* (registers), which hold natural numbers.

Minsky Machines

- A Minsky machine \mathcal{M} operates several *counters* (registers), which hold natural numbers.
- Instructions of \mathcal{M} can be of the following sorts:

$\text{INC}(p, r, q)$

being in state p , increase register r by 1
and move to state q ;

$\text{JZDEC}(p, r, q_0, q_1)$

being in state p , check whether the value of r is 0:
if yes, move to state q_0 ,
if no, decrease r by 1 and move to state q_1 .

Minsky Machines

- A Minsky machine \mathcal{M} operates several *counters* (registers), which hold natural numbers.
- Instructions of \mathcal{M} can be of the following sorts:

$\text{INC}(p, r, q)$

being in state p , increase register r by 1
and move to state q ;

$\text{JZDEC}(p, r, q_0, q_1)$

being in state p , check whether the value of r is 0:
if yes, move to state q_0 ,
if no, decrease r by 1 and move to state q_1 .

- We consider only *deterministic* Minsky machines.

Minsky Machines

- A Minsky machine \mathcal{M} operates several *counters* (registers), which hold natural numbers.
- Instructions of \mathcal{M} can be of the following sorts:

$\text{INC}(p, r, q)$

being in state p , increase register r by 1
and move to state q ;

$\text{JZDEC}(p, r, q_0, q_1)$

being in state p , check whether the value of r is 0:
if yes, move to state q_0 ,
if no, decrease r by 1 and move to state q_1 .

- We consider only *deterministic* Minsky machines.
- Two counters are sufficient for a Σ_1^0 -complete halting problem (Minsky 1961).

Minsky Machines

- A Minsky machine \mathcal{M} operates several *counters* (registers), which hold natural numbers.
- Instructions of \mathcal{M} can be of the following sorts:

$\text{INC}(p, r, q)$

being in state p , increase register r by 1
and move to state q ;

$\text{JZDEC}(p, r, q_0, q_1)$

being in state p , check whether the value of r is 0:
if yes, move to state q_0 ,
if no, decrease r by 1 and move to state q_1 .

- We consider only *deterministic* Minsky machines.
- Two counters are sufficient for a Σ_1^0 -complete halting problem (Minsky 1961).
- ... thus, *non-halting* is Π_1^0 -complete.

Minsky Machines

- A Minsky machine \mathcal{M} operates several *counters* (registers), which hold natural numbers.
- Instructions of \mathcal{M} can be of the following sorts:

$\text{INC}(p, r, q)$

being in state p , increase register r by 1
and move to state q ;

$\text{JZDEC}(p, r, q_0, q_1)$

being in state p , check whether the value of r is 0:
if yes, move to state q_0 ,
if no, decrease r by 1 and move to state q_1 .

- We consider only *deterministic* Minsky machines.
- Two counters are sufficient for a Σ_1^0 -complete halting problem (Minsky 1961).
- ... thus, *non-halting* is Π_1^0 -complete.
- Sometimes it is more convenient to use three counters.

Encoding Minsky Instructions

- Each instruction I of M is encoded by a formula A_I :

$$A_{\text{INC}}(p,r,q) = p \setminus (q \cdot r)$$

$$A_{\text{JZDEC}}(p,r,q_0,q_1) = ((p \cdot r) \setminus q_1) \wedge (p \setminus (q_0 \vee z_r)).$$

Encoding Minsky Instructions

- Each instruction I of M is encoded by a formula A_I :

$$A_{\text{INC}}(p,r,q) = p \setminus (q \cdot r)$$

$$A_{\text{JZDEC}}(p,r,q_0,q_1) = ((p \cdot r) \setminus q_1) \wedge (p \setminus (q_0 \vee z_r)).$$

- Moreover, we add three extra formulae: $N_r = z_r \setminus z_r$ for each counter r (i.e., a, b, or c).

Encoding Minsky Instructions

- Each instruction I of M is encoded by a formula A_I :

$$A_{\text{INC}}(p,r,q) = p \setminus (q \cdot r)$$

$$A_{\text{JZDEC}}(p,r,q_0,q_1) = ((p \cdot r) \setminus q_1) \wedge (p \setminus (q_0 \vee z_r)).$$

- Moreover, we add three extra formulae: $N_r = z_r \setminus z_r$ for each counter r (i.e., a, b, or c).
- The encoding is due to Lincoln et al. 1992.

Encoding Minsky Instructions

- Each instruction I of M is encoded by a formula A_I :

$$A_{\text{INC}}(p,r,q) = p \setminus (q \cdot r)$$

$$A_{\text{JZDEC}}(p,r,q_0,q_1) = ((p \cdot r) \setminus q_1) \wedge (p \setminus (q_0 \vee z_r)).$$

- Moreover, we add three extra formulae: $N_r = z_r \setminus z_r$ for each counter r (i.e., a, b, or c).
- The encoding is due to Lincoln et al. 1992.
- However, we now consider *non-halting* instead of halting, and model it using Kleene star instead of exponential.

Encoding Minsky Instructions

- Each instruction I of M is encoded by a formula A_I :

$$A_{\text{INC}}(p,r,q) = p \setminus (q \cdot r)$$

$$A_{\text{JZDEC}}(p,r,q_0,q_1) = ((p \cdot r) \setminus q_1) \wedge (p \setminus (q_0 \vee z_r)).$$

- Moreover, we add three extra formulae: $N_r = z_r \setminus z_r$ for each counter r (i.e., a, b, or c).
- The encoding is due to Lincoln et al. 1992.
- However, we now consider *non-halting* instead of halting, and model it using Kleene star instead of exponential.
- Also, in succedents of our sequents we now have to represent an *arbitrary* configuration of the Minsky machine being encoded, which is also implemented using Kleene star.

$$E = \bigwedge_I A_I \wedge N_a \wedge N_b \wedge N_c$$

Encoding Infinite Execution

$$E = \bigwedge_I A_I \wedge N_a \wedge N_b \wedge N_c \quad \leftarrow \text{this formula encodes the machine}$$

Encoding Infinite Execution

$$E = \bigwedge_I A_I \wedge N_a \wedge N_b \wedge N_c \quad \leftarrow \text{this formula encodes the machine}$$

$$D = (a^* \cdot b^* \cdot c^* \cdot \bigvee_{q \in Q} q) \vee (b^* \cdot c^* \cdot z_a) \vee (a^* \cdot c^* \cdot z_b) \vee (a^* \cdot b^* \cdot z_c)$$

Encoding Infinite Execution

$$E = \bigwedge_I A_I \wedge N_a \wedge N_b \wedge N_c \quad \leftarrow \text{this formula encodes the machine}$$

$$D = (a^* \cdot b^* \cdot c^* \cdot \bigvee_{q \in Q} q) \vee (b^* \cdot c^* \cdot z_a) \vee (a^* \cdot c^* \cdot z_b) \vee (a^* \cdot b^* \cdot z_c)$$

↑

this formula encodes any valid configuration

Encoding Infinite Execution

$$E = \bigwedge_I A_I \wedge N_a \wedge N_b \wedge N_c \quad \leftarrow \text{this formula encodes the machine}$$

$$D = (a^* \cdot b^* \cdot c^* \cdot \bigvee_{q \in Q} q) \vee (b^* \cdot c^* \cdot z_a) \vee (a^* \cdot c^* \cdot z_b) \vee (a^* \cdot b^* \cdot z_c)$$

↑

this formula encodes any valid configuration

z_a, z_b, z_c are for zero check in JZDEC, run in parallel with the main execution.

Encoding Infinite Execution

$E = \bigwedge_I A_I \wedge N_a \wedge N_b \wedge N_c$ \leftarrow this formula encodes the machine

$D = (a^* \cdot b^* \cdot c^* \cdot \bigvee_{q \in Q} q) \vee (b^* \cdot c^* \cdot z_a) \vee (a^* \cdot c^* \cdot z_b) \vee (a^* \cdot b^* \cdot z_c)$
 \uparrow

this formula encodes any valid configuration

z_a, z_b, z_c are for zero check in JZDEC, run in parallel with the main execution.

Lemma

$E^*, a^a, b^b, c^c, q \rightarrow D$ is derivable in $\mathbf{CommACT}_\omega$ iff the machine runs infinitely starting from (q, a, b, c) .

- $E^*, a^a, b^b, c^c, q \rightarrow D$ is derivable if and only if so is $E^n, a^a, b^b, c^c, q \rightarrow D$ for any $n \geq 0$.

- $E^*, a^a, b^b, c^c, q \rightarrow D$ is derivable if and only if so is $E^n, a^a, b^b, c^c, q \rightarrow D$ for any $n \geq 0$.
- This corresponds to n steps of execution.

Encoding Infinite Computation

- $E^*, a^a, b^b, c^c, q \rightarrow D$ is derivable if and only if so is $E^n, a^a, b^b, c^c, q \rightarrow D$ for any $n \geq 0$.
- This corresponds to n steps of execution.
- Since our machine is deterministic, partial computations form an infinite one. (In the non-deterministic case, use König's lemma.)

Encoding Infinite Computation

- $E^*, a^a, b^b, c^c, q \rightarrow D$ is derivable if and only if so is $E^n, a^a, b^b, c^c, q \rightarrow D$ for any $n \geq 0$.
- This corresponds to n steps of execution.
- Since our machine is deterministic, partial computations form an infinite one. (In the non-deterministic case, use König's lemma.)
- Base case: $n = 0$, and $a^a, b^b, c^c, q \rightarrow D$ is derivable ($((q, a, b, c)$ is a valid configuration).

Encoding $\text{INC}(p, a, q)$

$$\begin{array}{c}
 \frac{p \rightarrow p \quad \frac{E^{k-1}, a^{a+1}, b^b, c^c, q \rightarrow D}{E^{k-1}, a^a, b^b, c^c, q \cdot a \rightarrow D} \cdot L}{E^{k-1}, A_{\text{INC}(p,a,q)}, a^a, b^b, c^c, p \rightarrow D} \setminus L (A_{\text{INC}(p,a,q)} = p \setminus (q \cdot a)) \\
 \hline
 E^k, a^a, b^b, c^c, p \rightarrow D \quad \wedge L \text{ several times}
 \end{array}$$

Encoding JZDEC(p, a, q)

- $a \neq 0$

$$\begin{array}{c}
 \frac{p \rightarrow p \quad a \rightarrow a}{p, a \rightarrow p \cdot a} \cdot R \quad E^{k-1}, a^{a-1}, b^b, c^c, q_1 \rightarrow D \\
 \hline
 E^{k-1}, (p \cdot a) \setminus q_1, a^a, b^b, c^c, p \rightarrow D \\
 \hline
 E^{k-1}, A_{\text{JZDEC}(p, a, q_0, q_1)}, a^a, b^b, c^c, p \rightarrow D \quad \wedge L \\
 \hline
 E^k, a^a, b^b, c^c, p \rightarrow D \quad \wedge L \text{ several times}
 \end{array}$$

Encoding JZDEC(p, a, q)

- $a \neq 0$

$$\begin{array}{c}
 \frac{p \rightarrow p \quad a \rightarrow a}{p, a \rightarrow p \cdot a} \cdot R \quad E^{k-1}, a^{a-1}, b^b, c^c, q_1 \rightarrow D \quad \backslash L \\
 \hline
 E^{k-1}, (p \cdot a) \backslash q_1, a^a, b^b, c^c, p \rightarrow D \quad \wedge L \\
 \hline
 E^{k-1}, A_{\text{JZDEC}(p, a, q_0, q_1)}, a^a, b^b, c^c, p \rightarrow D \quad \wedge L \text{ several times} \\
 \hline
 E^k, a^a, b^b, c^c, p \rightarrow D
 \end{array}$$

- $a = 0$

$$\begin{array}{c}
 \frac{p \rightarrow p \quad \frac{E^{k-1}, b^b, c^c, q_0 \rightarrow D \quad \frac{(z_a \backslash z_a)^{k-1}, b^b, c^c, z_a \rightarrow D}{E^{k-1}, b^b, c^c, z_a \rightarrow D} \wedge L \text{ s.t.}}{E^{k-1}, q_0 \vee z_a, b^b, c^c \rightarrow D} \vee L}{E^{k-1}, p \backslash (q_0 \vee z_a), b^b, c^c, p \rightarrow D} \backslash L \\
 \hline
 E^{k-1}, A_{\text{JZDEC}(p, a, q_0, q_1)}, b^b, c^c, p \rightarrow D \quad \wedge L \\
 \hline
 E^k, b^b, p \rightarrow D \quad \wedge L \text{ several times}
 \end{array}$$

- As usual, proving the backwards direction, from derivation to computation, is technically more involved.

Backwards Implication

- As usual, proving the backwards direction, from derivation to computation, is technically more involved.
- Again, we reduce to finite proofs of sequents $E^n, a^a, b^b, c^c, q \rightarrow D$.

Backwards Implication

- As usual, proving the backwards direction, from derivation to computation, is technically more involved.
- Again, we reduce to finite proofs of sequents $E^n, a^a, b^b, c^c, q \rightarrow D$.
- In fact, it is possible that such a proof does not directly correspond to a counter machine computation: it could include “subprograms.”

Backwards Implication

- For example, let the machine include the following instructions: $\text{INC}(p, a, q)$ and $\text{JZDEC}(q, a, p, p)$, and consider a 4-step execution starting from $(p, 0, 0, 0)$.
- This execution has the following “non-canonical” representation:

$$\frac{\frac{\frac{p \rightarrow p}{E, p \setminus (q \cdot a), p \rightarrow p} \wedge L \quad \frac{\frac{\frac{q, a \rightarrow q \cdot a \quad p \rightarrow p}{(q \cdot a) \setminus p, a, q \rightarrow p} \setminus L \quad E, q, a \rightarrow p}{E, p \setminus (q \cdot a), p \rightarrow p} \wedge L}{E^2, p \rightarrow p} \wedge L \quad \frac{\frac{\frac{q, a \rightarrow q \cdot a \quad p \rightarrow D}{(q \cdot a) \setminus p, q, a \rightarrow D} \setminus L \quad E, q, a \rightarrow D}{E, q \cdot a \rightarrow D} \wedge L}{E, q \cdot a \rightarrow D} \cdot L \quad \frac{E^3, p \setminus (q \cdot a), p \rightarrow D}{E^4, p \rightarrow D} \setminus L \quad \wedge L$$

- Here we perform the INC step, and then start a “subroutine” which performs INC and JZDEC, returning to the same state. Finally, we perform JZDEC.

- Here we perform the INC step, and then start a “subroutine” which performs INC and JZDEC, returning to the same state. Finally, we perform JZDEC.
- The crucial point is that such a “subroutine” could not use the zero branch of JZDEC.

- Here we perform the INC step, and then start a “subroutine” which performs INC and JZDEC, returning to the same state. Finally, we perform JZDEC.
- The crucial point is that such a “subroutine” could not use the zero branch of JZDEC.
- This is due to the fact that D is available only on the main branch.

Backwards Implication

Let \tilde{E}_i denote any formula in the conjunction E or a conjunction of such formulae. The backwards implication is proved by joint induction of 6 statements:

1. Sequents of the form $\tilde{E}_1, \dots, \tilde{E}_k, a^a, b^b, c^c \rightarrow t$, where $t \in Q \cup Z$, are never derivable, neither are sequents of the form $\tilde{E}_1, \dots, \tilde{E}_k, a^a, b^b, c^c \rightarrow t \cdot r$, where $r \in R$.
2. Sequents of the form $\tilde{E}_1, \dots, \tilde{E}_k, z_r, a^a, b^b, c^c \rightarrow t$, where $r \in R$ and $t \in Q \cup Z_{\bar{r}}$, are never derivable, neither are sequents of the form $\tilde{E}_1, \dots, \tilde{E}_k, z_r, a^a, b^b, c^c \rightarrow t \cdot r'$, where $r, r' \in R$ and $t \in Q \cup Z_{\bar{r}}$.
3. If $\tilde{E}_1, \dots, \tilde{E}_k, z_a, a^a, b^b, c^c \rightarrow D$ is derivable, then $a = 0$. Similarly for b and c .

Backwards Implication

4. If $\tilde{E}_1, \dots, \tilde{E}_k, q, a^{a'}, b^{b'}, c^{c'} \rightarrow p$ is derivable, where $p, q \in Q$, then the machine can move from $\langle q, a' + a, b' + b, c' + c \rangle$ to $\langle p, a, b, c \rangle$ in k steps for any a, b, c .
5. If $\tilde{E}_1, \dots, \tilde{E}_k, q, a^{a'}, b^{b'}, c^{c'} \rightarrow p \cdot a$, where $p, q \in Q$, is derivable, then the machine can move from $\langle q, a' + a, b' + b, c' + c \rangle$ to $\langle p, a + 1, b, c \rangle$ in k steps for any a, b, c . Similarly for b and c .
6. If $\tilde{E}_1, \dots, \tilde{E}_k, p, a^a, b^b, c^c \rightarrow D$ is derivable ($p \in Q$), then the machine can perform k steps, starting from $\langle p, a, b, c \rangle$.

- Thus, derivability in **CommACT** is capable of simulating non-halting for counter machines, which yields the commutative version of Buszkowski's theorem:

- Thus, derivability in **CommACT** is capable of simulating non-halting for counter machines, which yields the commutative version of Buszkowski's theorem:

Theorem

CommACT _{ω} is Π_1^0 -hard.

- Thus, derivability in **CommACT** is capable of simulating non-halting for counter machines, which yields the commutative version of Buszkowski's theorem:

Theorem

CommACT _{ω} is Π_1^0 -hard.

- For the Π_1^0 upper bound, there are several methods, one of which we shall discuss tomorrow.

- Thus, derivability in **CommACT** is capable of simulating non-halting for counter machines, which yields the commutative version of Buszkowski's theorem:

Theorem

CommACT _{ω} is Π_1^0 -hard.

- For the Π_1^0 upper bound, there are several methods, one of which we shall discuss tomorrow.
- Next, we aim to prove undecidability of **CommACT**.

- Thus, derivability in **CommACT** is capable of simulating non-halting for counter machines, which yields the commutative version of Buszkowski's theorem:

Theorem

CommACT _{ω} is Π_1^0 -hard.

- For the Π_1^0 upper bound, there are several methods, one of which we shall discuss tomorrow.
- Next, we aim to prove undecidability of **CommACT**.
- This will be done by encoding circular computations by circular proofs.

Circular Proofs for Circular Computations

Lemma

If the machine runs **circularly** starting from (q, a, b, c) , then $E^*, a^a, b^b, c^c, q \rightarrow D$ admits a circular proof, thus, a proof in **CommACT**.

$$\begin{array}{c}
 \frac{E^*, p, a^a, b^b, c^c \rightarrow D}{\vdots} \\
 \frac{p, a^a, b^b, c^c \rightarrow D \quad \frac{E^*, E, p, a^a, b^b, c^c \rightarrow D}{*L}}{E^*, p, a^a, b^b, c^c \rightarrow D} \\
 \vdots \\
 \frac{q \rightarrow D \quad \frac{E^*, E, q \rightarrow D}{*L}}{E^*, q \rightarrow D}
 \end{array}$$

Translating Circular Proofs to CommACT

- Now we have to translate this circular argument into a proof in **CommACT**.

Translating Circular Proofs to CommACT

- Now we have to translate this circular argument into a proof in **CommACT**.
- Recall the rule for * there:

$$\frac{\rightarrow B \quad A, B \rightarrow B}{A^* \rightarrow B}$$

Translating Circular Proofs to CommACT

- Now we have to translate this circular argument into a proof in **CommACT**.
- Recall the rule for $*$ there:

$$\frac{\rightarrow B \quad A, B \rightarrow B}{A^* \rightarrow B}$$

- First we establish $*L$, the decomposition rule:

$$\frac{A^* \rightarrow 1 \vee (A \cdot A^*) \quad \frac{\frac{\Gamma \rightarrow C}{\Gamma, 1 \rightarrow C} \text{ } 1L \quad \frac{\Gamma, A, A^* \rightarrow C}{\Gamma, A \cdot A^* \rightarrow C} \text{ } \cdot L}{\Gamma, 1 \vee (A \cdot A^*) \rightarrow C} \vee L}{\Gamma, A^* \rightarrow C} \text{ } Cut$$

Translating Circular Proofs to CommACT

- Now we have to translate this circular argument into a proof in **CommACT**.
- Recall the rule for $*$ there:

$$\frac{\rightarrow B \quad A, B \rightarrow B}{A^* \rightarrow B}$$

- First we establish $*L$, the decomposition rule:

$$\frac{A^* \rightarrow 1 \vee (A \cdot A^*) \quad \frac{\frac{\Gamma \rightarrow C}{\Gamma, 1 \rightarrow C} \quad 1L \quad \frac{\Gamma, A, A^* \rightarrow C}{\Gamma, A \cdot A^* \rightarrow C} \cdot L}{\Gamma, 1 \vee (A \cdot A^*) \rightarrow C} \vee L}{\Gamma, A^* \rightarrow C} Cut$$

- This will be needed for modelling computation before the cycle.

Translating Circular Proofs to CommACT

- Next, we establish an extended version of induction:

$$\frac{\rightarrow B \quad A \rightarrow B \quad \dots \quad A^{k-1} \rightarrow B \quad A^k, B \rightarrow B}{A^* \rightarrow B}$$

- Next, we establish an extended version of induction:

$$\frac{\rightarrow B \quad A \rightarrow B \quad \dots \quad A^{k-1} \rightarrow B \quad A^k, B \rightarrow B}{A^* \rightarrow B}$$

- This is established by cutting with $A^* \equiv (1 \vee A \vee \dots \vee A^{k-1})(A^k)^*$.

Translating Circular Proofs to CommACT

- Next, we establish an extended version of induction:

$$\frac{\rightarrow B \quad A \rightarrow B \quad \dots \quad A^{k-1} \rightarrow B \quad A^k, B \rightarrow B}{A^* \rightarrow B}$$

- This is established by cutting with $A^* \equiv (1 \vee A \vee \dots \vee A^{k-1})(A^k)^*$.
- Now we have a circular proof of $E^*, p, a^a, b^b, c^c \rightarrow D$ from itself, and let k be the number of $*L$ applications on the cycle.

Translating Circular Proofs to CommACT

- Next, we establish an extended version of induction:

$$\frac{\rightarrow B \quad A \rightarrow B \quad \dots \quad A^{k-1} \rightarrow B \quad A^k, B \rightarrow B}{A^* \rightarrow B}$$

- This is established by cutting with $A^* \equiv (1 \vee A \vee \dots \vee A^{k-1})(A^k)^*$.
- Now we have a circular proof of $E^*, p, a^a, b^b, c^c \rightarrow D$ from itself, and let k be the number of $*L$ applications on the cycle.
- For simplicity, let $F = (p \cdot a^a \cdot b^b \cdot c^c) \setminus D$. Then our sequent is equivalent to $E^* \rightarrow F$.

Translating Circular Proofs to CommACT

- Next, we establish an extended version of induction:

$$\frac{\rightarrow B \quad A \rightarrow B \quad \dots \quad A^{k-1} \rightarrow B \quad A^k, B \rightarrow B}{A^* \rightarrow B}$$

- This is established by cutting with $A^* \equiv (1 \vee A \vee \dots \vee A^{k-1})(A^k)^*$.
- Now we have a circular proof of $E^*, p, a^a, b^b, c^c \rightarrow D$ from itself, and let k be the number of $*L$ applications on the cycle.
- For simplicity, let $F = (p \cdot a^a \cdot b^b \cdot c^c) \setminus D$. Then our sequent is equivalent to $E^* \rightarrow F$.
- From the circular proof, we can easily extract $E^i \rightarrow F$ for $0 \leq i < k$ (by replacing E^* with E^i).

- Next, we replace E with F , making $E^* \rightarrow F$ on top an axiom.

- Next, we replace E with F , making $E^* \rightarrow F$ on top an axiom.
- When we descend to the root, we get $E^k, F \rightarrow F$.

- Next, we replace E with F , making $E^* \rightarrow F$ on top an axiom.
- When we descend to the root, we get $E^k, F \rightarrow F$.
- Applying our extended induction rule, we get $E^* \rightarrow F$, q.e.d.

Translating Circular Proofs to CommACT

- Next, we replace E with F , making $E^* \rightarrow F$ on top an axiom.
- When we descend to the root, we get $E^k, F \rightarrow F$.
- Applying our extended induction rule, we get $E^* \rightarrow F$, q.e.d.
- Due to lack of time, we omitted the zero-checks, which would also involve circular reasoning (or using “pure induction,” $(z_a \setminus z_a)^* = z_a \setminus z_a$).

Translating Circular Proofs to CommACT

- Next, we replace E with F , making $E^* \rightarrow F$ on top an axiom.
- When we descend to the root, we get $E^k, F \rightarrow F$.
- Applying our extended induction rule, we get $E^* \rightarrow F$, q.e.d.
- Due to lack of time, we omitted the zero-checks, which would also involve circular reasoning (or using “pure induction,” $(z_a \setminus z_a)^* = z_a \setminus z_a$).
- In fact, circular proofs can always be rebuilt into inductive ones, but proving this in a general setting is much harder.

Inseparability

- Not any circular derivation, however, corresponds to a circular run of the machine.

Inseparability

- Not any circular derivation, however, corresponds to a circular run of the machine.
- For example, if the machine just increases one counter, $\text{INC}(q, a, q)$, then $E^*, a^a, b^b, c^c, q \rightarrow D$ is also derivable in **CommACT**.

Inseparability

- Not any circular derivation, however, corresponds to a circular run of the machine.
- For example, if the machine just increases one counter, $\text{INC}(q, a, q)$, then $E^*, a^a, b^b, c^c, q \rightarrow D$ is also derivable in **CommACT**.
- In this case $E = q \setminus (q \cdot a)$, and the circular derivation, for $a = b = c = 0$, is as follows:

$$\begin{array}{c}
 \frac{q_S \rightarrow q_S}{\frac{E^*, q_S \rightarrow a^* \cdot q_S}{E^*, E, q_S \rightarrow a^* \cdot q_S} \wedge L} \quad \frac{\frac{E^*, q_S \rightarrow a^* \cdot q_S \quad a \rightarrow a}{E^*, q_S, a \rightarrow a \cdot (a^* \cdot q_S)} \cdot R \quad \frac{a \cdot (a^* \cdot q_S) \rightarrow a^* \cdot q_S}{\frac{E^*, q_S, a \rightarrow a^* \cdot q_S}{E^*, q_S \setminus (q_S \cdot a), q_S \rightarrow a^* \cdot q_S} \cdot L, \setminus L} \text{Cut} \\
 \frac{q_S \rightarrow a^* \cdot q_S}{\frac{E^*, q_S \rightarrow a^* \cdot q_S}{E^*, q_S \rightarrow a^* \cdot q_S} *L} \quad \frac{a^* \cdot q_S \rightarrow D}{\frac{E^*, q_S \rightarrow D}{E^*, q_S \rightarrow D} \text{Cut}}
 \end{array}$$


- Therefore, we use an indirect technique for proving complexity, based on **effective inseparability**.

- Therefore, we use an indirect technique for proving complexity, based on **effective inseparability**.
- Let \mathcal{C} be the set of machines and input data such that the machine works circularly, and let \mathcal{H} be the one where the machine halts.

- Therefore, we use an indirect technique for proving complexity, based on **effective inseparability**.
- Let \mathcal{C} be the set of machines and input data such that the machine works circularly, and let \mathcal{H} be the one where the machine halts.
- There is no decidable set \mathcal{K} which separates \mathcal{C} from \mathcal{H} (i.e., $\mathcal{C} \subseteq \mathcal{K}$ and $\mathcal{H} \cap \mathcal{K} = \emptyset$), therefore **CommACT** is undecidable; but we do not know its complexity yet.

- Therefore, we use an indirect technique for proving complexity, based on **effective inseparability**.
- Let \mathcal{C} be the set of machines and input data such that the machine works circularly, and let \mathcal{H} be the one where the machine halts.
- There is no decidable set \mathcal{K} which separates \mathcal{C} from \mathcal{H} (i.e., $\mathcal{C} \subseteq \mathcal{K}$ and $\mathcal{H} \cap \mathcal{K} = \emptyset$), therefore **CommACT** is undecidable; but we do not know its complexity yet.
- **Folklore:** \mathcal{C} and \mathcal{H} are *effectively* inseparable.

Effective Inseparability



(Here W_u is the u -th r.e. set; f is computable.)

Myhill's Theorem

- (Corollary of) **Myhill's theorem**: if A separates \mathcal{C} and \mathcal{H} and A is r.e., then A is Σ_1^0 -complete.

Myhill's Theorem

- (Corollary of) **Myhill's theorem**: if A separates \mathcal{C} and \mathcal{H} and A is r.e., then A is Σ_1^0 -complete.
- Therefore, we obtain the necessary result:

Myhill's Theorem

- (Corollary of) **Myhill's theorem**: if A separates \mathcal{C} and \mathcal{H} and A is r.e., then A is Σ_1^0 -complete.
- Therefore, we obtain the necessary result:

Theorem

CommACT is Σ_0^1 -complete.

Myhill's Theorem

- (Corollary of) **Myhill's theorem**: if A separates \mathcal{C} and \mathcal{H} and A is r.e., then A is Σ_1^0 -complete.
- Therefore, we obtain the necessary result:

Theorem

CommACT is Σ_0^1 -complete.

- The reasoning in the non-commutative case is similar, however, the encoding of computations is more involved.