## Contents

**Polyfold Theory, verified: the machine-checked edition** **1**

# Polyfold Theory, verified: the machine-checked edition

*A companion to the expanded exposition of Part I of H. Hofer, K. Wysocki and E. Zehnder's* Polyfold and Fredholm Theory *(Springer 2021; the 2014 preprint arXiv 1407.3185 is the text in hand, and its numbering and pages are used, marked as such; the numbers and pages of the book's arXiv version, 1707.08941, are in* `Exposition/BookConcordance.md`*, and the Springer edition is not checked). That exposition is written to be read; this edition is written to be checked. It records, chapter by chapter, exactly which statements of the text have been verified by the Lean 4 proof assistant against the Mathlib library, what each verified statement says when read back in ordinary language, which hypotheses the text left silent, where the formal statement or proof departs from the text's, and what remains unverified and why. Nothing here needs to be read in Lean; every checked statement is restated in prose, and the Lean names are given only so that the claims can be audited.*

**Where to start.** The project has three parts:

- the expanded exposition (`Exposition/Chapter1.md`, …), the book for reading;
- *Polyfold Theory, Checked*, the Lean formalization (`Polyfold/`) and the reports on it;
- the sign-off page, the list of statements a reader is asked to confirm.

This edition is the way into the second part. Its reports each answer one question:

- **What is proved?** This edition: every checked result in plain language, with all its hypotheses, and the list of statements to sign off.
- **How is it proved?** *Polyfold Theory, proof by proof* (`Exposition/ProofByProof.md`): the formal proofs retold as mathematics, with a trace tag on each step naming the Lean declaration that carries it.
- **How does it differ from the text?** *Polyfold Theory beside Lean*, the comparison (`Exposition/polyfold-beside`) each result of the book beside the checked statement, with a verdict (the same, restated, hypotheses made explicit, stronger, narrower) and the differences.

A result has the same number, V1.5 say, in this edition, the comparison and the sign-off page.

---

### 0. Summary

| Chapter | HWZ | Content | Status | Lean lines |
|---|---|---|---|---|
| Chapter 1 | Ch. 1, §§1.1–1.4 | sc-Banach spaces, sc-operators and sc-Fredholm operators, sc-differentiability and its relation to classical differentiability, the chain rule, the degeneracy index | **verified** | 6,895 |
| Chapter 2 | Ch. 2 | sc-smooth retractions and retracts, M-polyfolds, the degeneracy index, tame M-polyfolds, strong bundles | **verified except Proposition 2.2 and Lemma 2.29 as stated; Remark 2.28's negative claim is false** | 15,187 |
| Chapter 3 | Ch. 3 | sc-Fredholm sections and the implicit function theorem | **partly verified** | 54,808 |
| Chapter 4 | Ch. 4, §§4.1–4.4 | manifolds with boundary with corners: $sc^+$-retracts and the characterization, smooth finite-dimensional submanifolds, families and Sard's theorem (vendored) | **verified except Lemma 4.10 and §4.4** | 12,083 |

| Chapter | HWZ | Content | Status | Lean lines |
|---|---|---|---|---|
| Chapter 5 | Ch. 5, §§5.1–5.4 | the Fredholm package for M-polyfolds: auxiliary norms, compactness, sc$^+$-sections and bump functions, transversality, general and good position, homotopies, partitions of unity | **verified except Theorems 5.22, 5.27 and the cited Theorems 5.32, 5.40, 5.43** | 12,100 |

Total: 101,073 lines of Lean in 166 files under `Polyfold/` (`wc -l`, excluding unfinished `WIP_` files). Toolchain: Lean v4.33.1, Mathlib v4.33.1. To reproduce: `lake build`, then `lake env lean -DmaxSynthPendingDepth=3 Polyfold/Test.lean`, which prints the axioms of every main theorem. Edition of 5 November 2026.

---

## 1. How to read a verified statement

**The statements are the contract.** A machine-checked proof guarantees that the *stated* theorem follows from Lean's axioms. It guarantees nothing about whether the statement is the one the authors meant. The reader's job, and the only mathematical judgement this document asks for, is to read each prose restatement and confirm that it says what the text means. Where the restatement is weaker or stronger than the text, or uses a different but equivalent definition, this is flagged as a **Deviation**.

**The setting.** An sc-Banach space is formalized as an indexed family of real Banach spaces $E_m$, $m \geq 0$, with injective compact continuous linear inclusions $E_{m+1} \to E_m$ and the smooth points dense in every level; the text's subspace picture $E_0 \supset E_1 \supset \cdots$ is recovered by identifying $E_m$ with its image in $E_0$, and "$x \in E_m$" for $x \in E_0$ means that $x$ is in that image. A map $f : U \to F$ on a subset $U \subseteq E_0$ is a total function $E_0 \to F_0$ whose values off $U$ play no role; its level maps $f_m : U_m \to F_m$ are determined by the injectivity of the inclusions. "Relatively open in a partial quadrant" enters most theorems only through its consequences on each level, that $U_m$ is the intersection of an open and a convex set and has unique differentiability at each of its points (an *sc-domain*); whenever the text's hypothesis is replaced by this weaker one the item says so.

**Classical differentiability.** $C^k$ on a relatively open subset of a partial quadrant means Mathlib's $C^k$ *within the set*, with derivatives within the set; on these sets derivatives are unique, so the notion agrees with the exposition's Definition 3.C.

**Axioms.** Every verified theorem depends only on the three standard axioms of Lean's Mathlib (`propext`, `Classical.choice`, `Quot.sound`), as `Polyfold/Test.lean` prints; there is no `sorry`.

---

## 2. Chapter 1 (HWZ §§1.1–1.4): verified

Chapter 1 of Part I (preprint pp. 6–17) sets up the sc-calculus: sc-Banach spaces, sc-operators, sc-subspaces and quotients (Definitions 1.1–1.5, Propositions 1.6–1.7), sc-Fredholm operators and their stability under $\mathrm{sc}^+$-perturbations (Definition 1.8, Propositions 1.9–1.10), partial quadrants, $\mathrm{sc}^0$, $\mathrm{sc}^1$, $\mathrm{sc}^k$ and sc-smooth maps (Definitions 1.11–1.15), the relation between sc-differentiability and classical differentiability on the levels (Propositions 1.16–1.19, Corollary 1.20), the chain rule (Theorem 1.21), the degeneracy index of a partial quadrant (Definition 1.23, Lemma 1.24) and boundary recognition (Theorem 1.25). Every numbered statement is verified, Proposition 1.19 and Corollary 1.20 under the text's own hypothesis ($C^{l+1}$ from $U_{m+l}$ to $F_m$ for $l \le k$), by the 2010 paper's induction $(\mathrm{S}_k)$ on elementary maps (V1.37–V1.39). One is verified in a stronger form: Theorem 1.25, which the text quotes from another paper, is proved here for $\mathrm{sc}^1$ germs. Example 1.2, the weighted Sobolev spaces, is not formalized (Mathlib has no Sobolev spaces). The remarks of the text that the exposition turns into lemmas are verified as well (V1.28–V1.36): the invariance of the index under $\mathrm{sc}^+$-perturbations and on every level, the uniqueness of the sc-structure in finite dimensions, sc-splittings as sc-projections, the smooth points of a quotient, the compact-open continuity of $Df$, $\mathrm{sc}^1 = C^1$ in finite dimensions on quadrants with boundary, and the level shift and tangent of a tuple. The setting throughout: an sc-Banach space is an indexed family of real Banach spaces $E_m$ with injective compact continuous linear inclusions $E_{m+1} \to E_m$ and dense smooth points; $\iota_m : E_m \to E_0$ is the composite inclusion, and "$x \in E_m$" for $x \in E_0$ means $x \in \iota_m(E_m)$. A map $f : U \to F$, $U \subseteq E_0$, is a total function $E_0 \to F_0$ whose values off $U$ do not matter, with level maps $f_m : U_m \to F_m$ determined by the injectivity of $\iota_m$. The text's hypothesis "$U$ relatively open in a partial quadrant $C$ of $E$" is, in most theorems, weakened to the *sc-domain* condition it implies: each $U_m$ is the intersection of an open and a convex subset of $E_m$ and has unique differentiability at each of its points. Where a theorem is stated for sc-domains the text's hypothesis is a special case. The infrastructure that Mathlib lacked and that is built here, the Riesz–Schauder theorem and the stability of Fredholm operators under compact perturbations (used without citation in §1.4.1), the compact-open lemma behind §1.2, and sc-domains, has its own items (V1.24–V1.27); so has the machinery of the proof of Proposition 1.19 under the text's hypothesis, the derivative tower, the elementary maps with the induction $(\mathrm{S}_k)$, and the sum rule for $\mathrm{sc}^k$ maps (V1.37–V1.39).

**V1.1 (sc-Banach spaces, sums and shifts; `Polyfold.ScScale`, `Polyfold.ScScale.toBase`, `Polyfold.ScScale.smooth`, `Polyfold.ScScale.smoothLvl`, `Polyfold.ScBanach`, `Polyfold.ScBanach.denseRan` `Polyfold.ScBanach.const`, `Polyfold.ScBanach.prod`, `Polyfold.ScBanach.prod_smoothLvl`, `Polyfold.ScBanach.shift`, `Polyfold.ScBanach.shift_smoothLvl`, `Polyfold.ScBanach.shiftK`, `Polyfold.ScBanach.lvlSet`).** A *scale* consists of real Banach spaces $E_m$, $m \in \mathbb{N}$, and injective compact continuous linear maps $E_{m+1} \to E_m$; $\iota_m : E_m \to E_0$ is the composite, $E_\infty \subseteq E_0$ the subspace of points that lie in $\iota_m(E_m)$ for every $m$, and $E_\infty \cap E_m := \iota_m^{-1}(E_\infty)$. An *sc-Banach space* is a scale in which $E_\infty \cap E_m$ is dense in $E_m$ for every $m$. Consequences: the image of $E_{m+k}$ in $E_m$ is dense. Constructions: for a finite-dimensional normed space $V$ the constant family $E_m = V$, with identity inclusions, is an sc-Banach space all of whose points are smooth; for sc-Banach spaces $E$, $F$, the family $E_m \times F_m$ with the product inclusions is an sc-Banach space $E \oplus F$ whose smooth points at level $m$ are the pairs of smooth points; the family $E_{m+1}$ with the inclusions of $E$ is an sc-Banach space $E^1$ whose smooth points at level $m$ are those of $E$ at level $m+1$; $E^k$ is the $k$-fold iterate. For $U \subseteq E_0$, the level set $U_m := \iota_m^{-1}(U) \subseteq E_m$. *Book:* Definition 1.1, p. 6: a decreasing sequence of Banach spaces $E = E_0 \supset E_1 \supset \cdots$ with compact inclusions and $E_\infty = \bigcap E_m$ dense in every $E_m$; the constant structure on a finite-dimensional space; $(E \oplus F)_m = E_m \oplus F_m$; the filtration $A_m = A \cap E_m$ of a subset and the shift $A^k$. *Hidden*

*hypotheses:* the inclusions are injective (implicit in "$\subset$") and continuous (part of compactness). *Deviation (restated):* the $E_m$ are separate Banach spaces, not subsets of one space; the two pictures are equivalent and nothing in Chapter 1 uses more than the family. The text's remarks that a finite-dimensional space has *only* the constant structure and that the constant family on an infinite-dimensional space is not an sc-structure are V1.31.

**V1.2 (sc-operators and sc-isomorphisms; `Polyfold.ScOperator`, `Polyfold.ScOperator.ext_base`, `Polyfold.IsScOperator`, `Polyfold.ScOperator.isScOperator`, `Polyfold.IsScOperator.toScOperator`, `Polyfold.ScOperator.comp`, `Polyfold.ScOperator.mapsTo_smoothLvl`, `Polyfold.ScIso`, `Polyfold.ScIso.symm`, `Polyfold.ScIso.trans`, `Polyfold.isScOperator_of_mapsTo`, `Polyfold.ScIso.ofBijec`**
An *sc-operator* $T : E \to F$ between sc-Banach spaces is a family of continuous linear maps $T_m : E_m \to F_m$ with $\iota_m^F T_{m+1} = T_m \iota_m^E$; the family is determined by $T_0$, and a continuous linear $T_0 : E_0 \to F_0$ *is an sc-operator* exactly when for each $m$ some continuous linear $T_m : E_m \to F_m$ satisfies $\iota_m T_m = T_0 \iota_m$ (the two formulations correspond). Sc-operators compose, add, and map smooth points to smooth points. An *sc-isomorphism* is a family of continuous linear equivalences $E_m \simeq F_m$ commuting with the inclusions; inverses and composites are sc-isomorphisms. A continuous linear $T_0 : E_0 \to F_0$ with $T_0(E_m) \subseteq F_m$ for every $m$ is an sc-operator (the continuity of $T_0 : E_m \to F_m$ is the closed graph theorem), and an sc-operator bijective on every level is an sc-isomorphism (the open mapping theorem). *Book:* Definition 1.3, p. 7. *Hidden hypotheses:* none. *Deviation (restated):* the level maps are data, with the text's predicate on the base map recovered as `Polyfold.IsScOperator`; the continuity of the inverse of a bijective sc-operator on each level, which the text's definition of an sc-isomorphism demands, is built into the formal notion, and derived from the open mapping theorem separately (`Polyfold.ScIso.ofBijective`); the continuity on the levels asked by Definition 1.3 is automatic for a bounded $T_0$ that maps $E_m$ into $F_m$ (Findings, Chapter 3, item 233).

**V1.3 (sc$^+$-operators; `Polyfold.ScPlus`, `Polyfold.ScBanach.shiftIncl`, `Polyfold.ScPlus.toScOperator`, `Polyfold.ScPlus.isCompactOperator`, `Polyfold.ScPlus.toScOperator_mem_range_toBase_succ`).**
An sc$^+$*-operator* $E \to F$ is an sc-operator $E \to F^1$, i.e. continuous linear $S_m : E_m \to F_{m+1}$ commuting with the inclusions. Composed with the inclusion $F^1 \to F$ (an sc-operator) it is an sc-operator $E \to F$ whose level-$m$ map $E_m \to F_m$ is compact for every $m$, and which raises the level: $x \in \iota_m(E_m)$ implies $S_0 x \in \iota_{m+1}(F_{m+1})$. *Book:* Definition 1.4, p. 7, and the remark that follows it: an sc-operator $S$ with $S(E_m) \subset F_{m+1}$ and $S : E \to F^1$ an sc-operator is sc-compact. *Hidden hypotheses:* none; the condition $S(E_m) \subset F_{m+1}$ is contained in "$S : E \to F^1$ is an sc-operator" and is dropped.

**V1.4 (sc-subspaces and sc-splittings; `Polyfold.ScBanach.levelSub`, `Polyfold.ScBanach.isClosed_levelSub`, `Polyfold.ScBanach.IsScSubspace`, `Polyfold.ScBanach.ofSubspaceScale`, `Polyfold.ScBanach.ofSubspace`, `Polyfold.ScBanach.IsScSplitting`, `Polyfold.ScBanach.IsScSplitting.closedComplemented`, `Polyfold.ScBanach.IsScSplitting.symm`).** For a subspace $A \subseteq E_0$ let $A_m := \iota_m^{-1}(A) \subseteq E_m$, closed if $A$ is. $A$ is an *sc-subspace* if $A$ is closed and $A_m$ is contained in the closure of $A_m \cap E_\infty$ for every $m$. A closed $A$ with the levels $A_m$ and the restricted inclusions is a scale (the restricted inclusions are compact), and an sc-subspace is in this way an sc-Banach space. $E = F \oplus G$ is an *sc-splitting* if $F$ and $G$ are sc-subspaces and $F_m$, $G_m$ are complementary subspaces of $E_m$ for every $m$; then $F_m$ is closed complemented in $E_m$, and the splitting is symmetric in $F$ and $G$. *Book:* Definition 1.5, p. 7: $F$ closed with $(F \cap E_m)$ an sc-structure; an sc-complement $G$ with $(G \cap E_m)$ an sc-structure and $E_m = F_m \oplus G_m$ a topological direct sum on every level. *Hidden hypotheses:* none. *Deviation (restated):* the closedness of $F_m$ in $E_m$ and the compactness of $F_{m+1} \to F_m$ are automatic and are not conditions; "topological direct sum" is read as

complementary closed subspaces, the continuity of the projections following from the closed graph theorem (`Polyfold.ScBanach.IsScSplitting.closedComplemented`).

**V1.5 (Proposition 1.6; `Polyfold.ScBanach.proposition_1_6_iff`, `Polyfold.ScBanach.proposition_1_6`).** Let $E$ be an sc-Banach space and $F \subseteq E_0$ a finite-dimensional subspace. $F$ is an sc-subspace if and only if $F \leq E_\infty$; and if $F \leq E_\infty$ there is a subspace $G \subseteq E_0$ such that $E = F \oplus G$ is an sc-splitting. *Book:* p. 7, the same. *Hidden hypotheses:* none. *Proof:* as in the text, with a continuous projection $E_0 \to F$ from Mathlib (Hahn–Banach), lifted to each level through $F_m \cong F$.

**V1.6 (Proposition 1.7; `Polyfold.ScBanach.quotientScale`, `Polyfold.ScBanach.quotient`, `Polyfold.ScBanach.quotientMap`, `Polyfold.ScBanach.proposition_1_7`).** Let $E$ be an sc-Banach space and $A \subseteq E_0$ a closed subspace. The quotient Banach spaces $E_m/A_m$, with the inclusions $E_{m+1}/A_{m+1} \to E_m/A_m$ induced by $\iota$ (injective and compact), and with the smooth points dense on every level, form an sc-Banach space $E/A$, and the quotient maps $E_m \to E_m/A_m$ form an sc-operator $E \to E/A$. As a statement: there exist an sc-Banach space $Q$ and an sc-operator $q : E \to Q$ with every $q_m$ surjective and $\ker q_m = A_m$. *Book:* p. 8, for an sc-subspace $A$: $E/A$ with the filtration $E_m/A_m$ is an sc-Banach space, where $E_m/A_m$ is identified with $\{(x + A) \cap E_m : x \in E\}$. *Hidden hypotheses:* none. *Deviation (stronger):* only closedness of $A$ is used, not that $A$ is an sc-subspace. The text's identification of $E_m/A_m$ with a subset of $E/A$ is the injectivity of the induced inclusion, which holds because $A_m$ is the full preimage of $A$. The text's description of $(E/A)_\infty$ as the classes of points of $E_\infty$ is V1.33; only the trivial inclusion is used here.

**V1.7 (sc-Fredholm operators and the index; `Polyfold.ScFredholmSplitting`, `Polyfold.IsScFredholm`, `Polyfold.ScFredholmSplitting.index`, `Polyfold.scIndex`, `Polyfold.ScFredholmSplitting.index_eq`, `Polyfold.ScFredholmSplitting.K_le_smooth`, `Polyfold.ScFredholmSplitting.ker_lvl`, `Polyfold.ScFredholmSplitting.range_lvl`, `Polyfold.ScFredholmSplitting.isFredholm`).** An *sc-Fredholm splitting* of an sc-operator $T : E \to F$ is given by subspaces $K, X \subseteq E_0$ and $C, Y \subseteq F_0$ such that $E = K \oplus X$ and $F = C \oplus Y$ are sc-splittings, $K = \ker T_0$, $Y = \operatorname{range} T_0$, $K$ and $C$ are finite-dimensional, and for every $m$ the map $T_m$ is a bijection of $X_m$ onto $Y_m$. $T$ is *sc-Fredholm* if a splitting exists. The index $\dim K - \dim C$ of a splitting equals $\dim \ker T_0 - \dim(F_0/\operatorname{range} T_0)$, hence does not depend on the splitting. Moreover $K \leq E_\infty$; $\ker T_m = K_m$ and $\operatorname{range} T_m = Y_m$; and every $T_m : E_m \to F_m$ is a Fredholm operator in Mathlib's sense (finite-dimensional kernel, closed range of finite codimension, strict, with closed complemented kernel). *Book:* Definition 1.8, p. 9, with the remarks that $K \subset E_\infty$, $T(X_m) = Y_m$, $E_m = K \oplus X_m$ and "$F = C \oplus T(E_m)$" (read $F_m$), and $\operatorname{ind}(T) = \dim K - \dim C$. *Hidden hypotheses:* none. *Deviation (restated):* condition (3), "$T : X \to Y$ is an sc-isomorphism", is taken as the bijectivity of $T_m : X_m \to Y_m$ for every $m$; the continuity of the inverses is automatic by the open mapping theorem on the closed subspaces $X_m$, $Y_m$. The independence of the index from the splitting and the level-wise Fredholm property are not stated in the text.

**V1.8 (Proposition 1.9; `Polyfold.proposition_1_9`, `Polyfold.ScOperator.IsRegularizing`, `Polyfold.IsScFredholm.isRegularizing`, `Polyfold.isScFredholm_iff`).** Let $T : E \to F$ be an sc-operator with an sc-Fredholm splitting, $e \in E_0$ and $m \in \mathbb{N}$. If $T_0 e \in \iota_m(F_m)$ then $e \in \iota_m(E_m)$. As a property of sc-operators (`Polyfold.ScOperator.IsRegularizing`): every sc-Fredholm operator is regularizing. Conversely, an sc-operator $T$ with $T_0$ Fredholm that is regularizing is sc-Fredholm (V1.9); so $T$ is sc-Fredholm if and only if $T_0 : E_0 \to F_0$ is Fredholm and $T$ is regularizing. *Book:* p. 9, the same ("an sc-Fredholm operator is regularizing"). *Hidden hypotheses:* none; the step "$c = 0$" uses $C \cap Y = 0$ on level 0, as in the text. The converse is not stated in the text; it is the tool for the text's unargued "is sc-Fredholm" for operators obtained by

conjugation and restriction in Chapter 3 (V3.8, V3.46, V3.96; Findings, Chapter 3, item 233).

**V1.9 (Proposition 1.10; `Polyfold.proposition_1_10`, `Polyfold.ScPlus.regularizing_add`, `Polyfold.isScFredholm_of_isRegularizing`, `Polyfold.isScFredholm_of_isRegularizing_index`).** Let $E$, $F$ be sc-Banach spaces, $T : E \to F$ an sc-Fredholm operator and $S : E \to F$ an $\mathrm{sc}^+$-operator. Then $T + S$ (with $S$ read as an sc-operator $E \to F$) is sc-Fredholm. Along the way, $T + S$ is regularizing in the sense of V1.8. *Book:* p. 10, the same; the proof is in §1.4.1, pp. 14–15. *Hidden hypotheses:* none. *Proof:* the text's, with three additions. The opening claim that $T + S : E_m \to F_m$ is Fredholm is the classical compact-perturbation theorem, proved here since Mathlib lacks it (V1.24–V1.25). The step "$y = (T + S)e$, so by Proposition 1.9 $e \in E_m$" needs a bootstrap, since $Se$ is only one level up: $Te = y - Se \in F_1$ gives $e \in E_1$, then $Se \in F_2$, and so on $m$ times; the same bootstrap gives $K_m = K_{m+1}$. The complement $C$ is the range of a linear right inverse of $F_0 \to F_0/Y$ with values in $F_\infty$ rather than the span of a lifted basis, and condition (3) of Definition 1.8 for $T + S$, which the text does not verify, is proved as the bijectivity of $(T + S)_m : X_m \to Y_m$. The text does not assert that the index is preserved; it is, and this is V1.30. The construction uses only two properties of $T + S$, that it is Fredholm on level 0 and regularizing; with these as hypotheses it is proved as a criterion: an sc-operator that is Fredholm on level 0 and regularizing is sc-Fredholm, and every level map is then Fredholm of the level-0 index (not stated in the text).

**V1.10 (partial quadrants and tuples; `Polyfold.quadrant`, `Polyfold.quadrant_convex`, `Polyfold.quadrant_isClosed`, `Polyfold.lvlSet_quadrant`, `Polyfold.IsPartialQuadrant`, `Polyfold.IsPartialQuadrant.convex`, `Polyfold.IsPartialQuadrant.isClosed`, `Polyfold.IsPartialQuadran` `Polyfold.isPartialQuadrant_univ`, `Polyfold.IsRelOpenInQuadrant`).** In the sc-Banach space $\mathbb{R}^n \oplus W$ (with $\mathbb{R}^n$ constant) the *quadrant* on level $m$ is $\{(a, w) : a_i \geq 0 \text{ for all } i\}$; it is closed and convex with non-empty interior, and the level sets of the level-0 quadrant are the level quadrants. $C \subseteq E_0$ is a *partial quadrant* if there are $n$, an sc-Banach space $W$ and an sc-isomorphism $T : E \to \mathbb{R}^n \oplus W$ with $C = T_0^{-1}(\text{quadrant})$. Then every $C_m$ is closed, convex, with non-empty interior in $E_m$, and $C_m$ has unique differentiability (in Mathlib's sense) at each of its points; $E_0$ itself is a partial quadrant ($n = 0$). $(U, C, E)$ is a *tuple*, i.e. $U$ is a *relatively open subset of the partial quadrant* $C$, if $C$ is a partial quadrant and $U = O \cap C$ with $O \subseteq E_0$ open. *Book:* Definition 1.11, p. 10, and the sentence after it. *Hidden hypotheses:* none; "closed convex" is derived from the model. *Deviation:* none in the definition. The text's tacit use of $C_k$ as a partial quadrant in $E^k$ is V1.36 (for $k = 1$, and by iteration).

**V1.11 ($\mathrm{sc}^0$ maps; `Polyfold.levelMap`, `Polyfold.IsSc0`, `Polyfold.IsSc0.levelMap_mapsTo`, `Polyfold.IsSc0.continuousOn_base`, `Polyfold.IsSc0.congr`, `Polyfold.IsSc0.mono`, `Polyfold.IsSc0.comp`, `Polyfold.isSc0_id`, `Polyfold.ScOperator.isSc0`).** For $f : E_0 \to F_0$ the *level map* $f_m : E_m \to F_m$ sends $x$ to the unique $y \in F_m$ with $\iota_m y = f(\iota_m x)$, when such $y$ exists (and to a junk value otherwise). $f$ is $\mathrm{sc}^0$ *on* $U \subseteq E_0$ if for every $m$ and $x \in U_m$ the point $f(\iota_m x)$ lies in $\iota_m(F_m)$, and $f_m$ is continuous on $U_m$. Then $f$ is continuous on $U$; if $f(U) \subseteq V$ then $f_m(U_m) \subseteq V_m$; the property depends only on the values of $f$ on $U$ and passes to subsets of $U$; the composition of an $\mathrm{sc}^0$ map on $U$ with values in $V$ and an $\mathrm{sc}^0$ map on $V$ is $\mathrm{sc}^0$ on $U$; the identity and every sc-operator are $\mathrm{sc}^0$, the latter with level maps $T_m$. *Book:* Definition 1.12, p. 10: $f(U_m) \subset V_m$ (meaning $U'_m$) and $f : U_m \to U'_m$ continuous for all $m$. *Hidden hypotheses:* none. *Deviation (restated):* maps are total functions on $E_0$, and the target tuple enters only through $f(U) \subseteq U'$.

**V1.12 (the tangent; `Polyfold.ScBanach.tangent`, `Polyfold.ScBanach.tangentSet`, `Polyfold.ScBanach.lvlSet` `Polyfold.ScBanach.tangentSet_mono`, `Polyfold.IsScDomain.tangentSet`, `Polyfold.ScBanach.shift_tangen`**

The *tangent* of an sc-Banach space $E$ is $TE := E^1 \oplus E$, with levels $E_{m+1} \times E_m$. For $U \subseteq E_0$, $TU \subseteq (TE)_0 = E_1 \times E_0$ is $\{(x,h) : x \in U_1\}$, with level-$m$ part $U_{m+1} \times E_m$; $U \subseteq V$ implies $TU \subseteq TV$. If $U$ is an sc-domain (V1.27), $TU$ is an sc-domain in $TE$; and the level-1 part of $TU$ is $T(U_1)$ in $T(E^1) = (TE)^1$. *Book:* Definition 1.13, p. 10: $T(U,C,E) = (U^1 \oplus E, C^1 \oplus E, E^1 \oplus E)$, "again a tuple". *Hidden hypotheses:* none. *Deviation:* none; that $TC$ is a partial quadrant in $TE$, so that $T(U,C,E)$ is again a tuple, is V1.36, while the proofs of §§1.2–1.3 propagate the sc-domain property (V1.27) instead.

**V1.13 (sc¹ maps and the tangent map; `Polyfold.tangentMapWith`, `Polyfold.IsSc1With`, `Polyfold.IsSc1`, `Polyfold.scDeriv`, `Polyfold.tangentMap`, `Polyfold.IsSc1With.eq_of_uniqueDiffWithinAt`, `Polyfold.IsSc1.scDeriv_eqOn`, `Polyfold.IsSc1.tangentMap_eqOn`, `Polyfold.IsSc1.congr`, `Polyfold.IsSc1With.continuousOn_deriv`, `Polyfold.IsSc1With.exists_delta_deriv`, `Polyfold.exists_isSc1_not_continuousOn_deriv`, `Polyfold.ScOneExample.isSc1With`, `Polyfold.ScOneExample.not_continuousAt_D`).** Let $f : E_0 \to F_0$, $U \subseteq E_0$ and $D : E_1 \to \mathcal{L}(E_0,F_0)$ a family of bounded operators. The tangent map $T_D f : (TE)_0 \to (TF)_0$ is $(x,h) \mapsto (f_1(x), D(x)h)$. $f$ is sc$^1$ *on $U$ with derivatives $D$* if (1) $f$ is sc$^0$ on $U$; (2) for every $x \in U_1$, the map $f \circ \iota_1 : E_1 \to F_0$ has derivative $D(x) \circ \iota_1$ at $x$ within $U_1$; (3) $T_D f$ is sc$^0$ on $TU$. $f$ is sc$^1$ *on $U$* if some $D$ makes it so; `Polyfold.scDeriv` is a chosen such $D$ (junk if none exists) and `Polyfold.tangentMap` the tangent map built from it. If $D$ and $D'$ both satisfy (2) at a point $x \in U_1$ where $U_1$ has unique differentiability, then $D(x) = D'(x)$, because $E_1$ has dense image in $E_0$; hence on an sc-domain the chosen derivative agrees on $U_1$ with any admissible one, and the tangent map on $TU$. The property depends only on the values of $f$ on $U$. Level 0 of condition (3) is the continuity of $(x,h) \mapsto D(x)h$ on $U_1 \times E_0$, and it implies: for $x \in U_1$, a compact $K \subseteq E_0$ and $c > 0$ there is $\delta > 0$ with $\|D(x')w - D(x)w\| \le c$ for all $x' \in U_1$ with $\|x' - x\|_1 < \delta$ and all $w \in K$ (continuity of $x \mapsto D(x)$ in the compact-open topology). It need not be norm continuous: on $E = \mathbb{R} \oplus X$, with $X$ the sc-Banach space whose level $m$ consists of the real sequences with $(k+1)^m x_k \to 0$, normed by $\sup_k (k+1)^m |x_k|$ (every level a copy of $c_0$, the inclusions $\mathrm{diag}(1/(k+1))$), the map $f(t,x)_k = \cos(kt)x_k$ is sc$^1$ with $Df(t,x)(\tau,\xi)_k = -k\sin(kt)x_k\tau + \cos(kt)\xi_k$, and for every derivative family $D$ of $f$ the map $x \mapsto D(x)$ is not continuous on $E_1$ into $\mathcal{L}(E_0,X_0)$ with the operator norm: $(\pi/(n+1),0) \to 0$ in $E_1$, but $\|Df(\pi/(n+1),0) - Df(0,0)\| \ge 2$. *Book:* Definition 1.14, pp. 10–11, with conditions (1)–(3) as stated, and the remarks that $x \mapsto Df(x)$ is continuous for the compact-open but not in general for the norm topology, and that sc¹ maps between finite-dimensional spaces are the $C^1$ maps. *Hidden hypotheses:* the uniqueness of $Df(x)$, without which "the tangent map" is undefined; it holds because $E_1$ is dense in $E_0$ and $U_1$ has unique differentiability (true for relatively open subsets of partial quadrants, V1.10). *Deviation (hypotheses explicit):* the derivative family is an explicit parameter of `Polyfold.IsSc1With`, with `Polyfold.IsSc1` the text's notion. The compact-open remark is V1.34, its negative half the example above, which the text does not give (Findings, Chapter 1, follow-up item 1), and the finite-dimensional remark is V1.35 (on relatively open subsets of quadrants, with boundary; the open case is also in `Polyfold/Test.lean`, `isSc1_iff_contDiffOn_of_finiteDimensional`).

**V1.14 (sc^k and sc-smooth maps; `Polyfold.IsSck`, `Polyfold.IsScSmooth`, `Polyfold.IsSck.of_succ`, `Polyfold.IsSck.of_le`, `Polyfold.isScSmooth_iff`, `Polyfold.IsSck.congr`).** By recursion: $f$ is sc$^0$ on $U$ in the sense of V1.11; $f$ is sc$^{k+1}$ on $U$ if it is sc$^1$ on $U$ and its tangent map (built from the chosen derivative) is sc$^k$ on $TU \subseteq (TE)_0$ as a map into $(TF)_0$. $f$ is *sc-smooth* on $U$ if it is sc$^k$ for every $k$. Then sc$^{k+1}$ implies sc$^k$, and sc$^l$ for $l \le k$; $f$ is sc-smooth iff it is sc$^1$ with sc-smooth tangent map; on an sc-domain, sc$^k$ depends only on the values on $U$. *Book:* p. 11, the inductive

definition after Definition 1.14 (numbered 1.15 here, as in the Lean files; the preprint's Remark 1.15 on partial cones is not formalized). *Hidden hypotheses:* none beyond V1.13.

**V1.15 (Proposition 1.16; `Polyfold.IsSc1Levels`, `Polyfold.IsSc1Levels.continuousOn_clm`, `Polyfold.IsSc1Levels.contDiffOn`, `Polyfold.IsSc1Levels.toBase_apply`, `Polyfold.IsSc1Levels.isSc1Wit` `Polyfold.IsSc1With.isSc1Levels`, `Polyfold.proposition_1_16`, `Polyfold.IsSc1With.contDiffOn_base`, `Polyfold.IsSc1With.scOperator_deriv`).** Let $U$ be an sc-domain in $E$ and $f$ sc$^0$ on $U$. Then $f$ is sc$^1$ on $U$ if and only if there is a family $D_m(x) \in \mathcal{L}(E_m, F_m)$, $m \in \mathbb{N}$, $x \in U_{m+1}$, such that for every $m$: (a) $f_m \circ \iota : U_{m+1} \to F_m$ has derivative $D_m(x) \circ \iota$ at every $x \in U_{m+1}$ within $U_{m+1}$; (b) $(x, h) \mapsto D_m(x)h$ is continuous on $U_{m+1} \times E_m$. For any such family, $x \mapsto D_m(x) \circ \iota$ is continuous $U_{m+1} \to \mathcal{L}(E_{m+1}, F_m)$ in the operator norm, so $f : U_{m+1} \to F_m$ is $C^1$ within $U_{m+1}$; and $\iota_m D_m(x) = D_0(\iota x)\iota_m$, so all $D_m(x)$ are restrictions of the single operator $Df(x) := D_0(x)$. In the direction "sc¹ □ levels" the family is the level operators of V1.26; in particular $f \circ \iota_1 : U_1 \to F_0$ is $C^1$. If $f$ is sc$^1$ with derivatives $Df$ and $x \in U_1$ is a smooth point, $Df(x) : E_0 \to F_0$ is an sc-operator. *Book:* p. 12: an sc$^0$ map on a relatively open subset of a partial quadrant is sc$^1$ iff (1) for $m \geq 1$, $f : U_m \to F_{m-1}$ is $C^1$ with $df : U_m \to \mathcal{L}(E_m, F_{m-1})$ continuous and (2) $df(x)$ extends to a bounded $Df(x) : E_{m-1} \to F_{m-1}$ with $(x, h) \mapsto Df(x)h$ continuous on $U_m \oplus E_{m-1}$; in particular $Df(x)$ is an sc-operator at smooth points. *Hidden hypotheses:* the compatibility of the extensions $Df(x)$ across levels, which the text's notation assumes and the direction (□ ) needs (Lemma 3.H of the exposition); it is proved. *Deviation (restated):* the extensions form an explicit family; the $C^1$ property and the norm continuity of $df$ are consequences of (a)–(b) rather than part of the condition; the hypothesis on $U$ is the weaker sc-domain condition.

**V1.16 (Proposition 1.17; `Polyfold.IsSc0.shift`, `Polyfold.IsSc1With.shift`, `Polyfold.IsSck.shift`, `Polyfold.IsScSmooth.shift`, `Polyfold.IsScDomain.shift`).** Let $U$ be an sc-domain in $E$ and $f$ sc$^k$ on $U$. Then the level-1 map $f_1 : E_1 \to F_1$ is sc$^k$ on $U_1$ as a map between the shifted sc-Banach spaces $E^1$ and $F^1$ (whose level sets of $U_1$ are the $U_{m+1}$). For $k = 0$ no hypothesis on $U$ is needed; for an sc$^1$ map with derivatives $Df$, the derivatives of $f_1$ are the level-1 operators $Df(x) : E_1 \to F_1$, $x \in U_2$; sc-smooth maps lift to sc-smooth maps; and $U_1$ is an sc-domain in $E^1$. *Book:* p. 12: if $f : U \to V$ is sc$^k$ between relatively open subsets of partial quadrants then $f : U^1 \to V^1$ is sc$^k$. *Hidden hypotheses:* none. *Deviation:* none; the target $V$ plays no role and the hypothesis on $U$ is weaker.

**V1.17 (Proposition 1.18; `Polyfold.IsSck.contDiffOn_levels'`, `Polyfold.IsSck.contDiffOn_levels`, `Polyfold.IsSck.contDiffOn_levels_of_le`, `Polyfold.IsSck.contDiffOn_base`, `Polyfold.IsScSmooth.contD`** Let $U$ be an sc-domain in $E$ and $f$ sc$^k$ on $U$. Then for every $m$, the map $f_m \circ \iota : U_{m+k} \to F_m$ is $C^k$ within $U_{m+k}$; for $l \leq k$, $f_m \circ \iota : U_{m+l} \to F_m$ is $C^l$; in particular $f \circ \iota_k : U_k \to F_0$ is $C^k$. If $f$ is sc-smooth, $f : U_{m+k} \to F_m$ is $C^k$ for all $m$ and $k$. *Book:* p. 13, the same for $f : U \to V$ between relatively open subsets of partial quadrants. *Hidden hypotheses:* the meaning of "$C^k$" on a relatively open subset of a quadrant, never defined in the text: here Mathlib's $C^k$ within the set, with unique derivatives on these sets. *Deviation:* none in the statement. *Proof:* a direct induction on $k$ over all sc-domains, with a compatible level-lowering operator $\rho : E_{m+k} \to E_m$ as parameter, and the compact-open lemma (V1.26) applied to the second component of the level tangent map; the 2010 paper's multilinear maps $\Gamma(x)$ and its differentiation under the integral sign are not used.

**V1.18 (Proposition 1.19; `Polyfold.proposition_1_19_text`, `Polyfold.IsScSmooth.of_contDiffOn_levels_te` `Polyfold.proposition_1_19_zero`, `Polyfold.proposition_1_19`, `Polyfold.IsScSmooth.of_contDiffOn_leve`** Let $U$ be an sc-domain in $E$ and $f$ sc$^0$ on $U$. If for every $m \geq 0$ and $0 \leq l \leq k$ the map

$f_m \circ \iota : U_{m+l} \to F_m$ (the level-$m$ map restricted along the inclusion $E_{m+l} \to E_m$) is $C^{l+1}$ within $U_{m+l}$, then $f$ is $\mathrm{sc}^{k+1}$ on $U$; if this holds for every $l$, then $f$ is sc-smooth. Also proved, under the stronger hypothesis that every level map $f_m : U_m \to F_m$ is $C^{k+1}$ within $U_m$: $f$ is $\mathrm{sc}^{k+1}$; for $k = 0$, where the two hypotheses agree, $f$ is $\mathrm{sc}^1$ with $Df(x)$ the derivative of $f : U_0 \to F_0$ within $U$ at $\iota_1 x$; and if $f$ maps $U_m$ into $F_m$ for every $m$ and every $f_m$ is $C^\infty$, then $f$ is sc-smooth. *Book:* p. 13: if $f : U_{m+l} \to F_m$ is $C^{l+1}$ for all $m \geq 0$ and $0 \leq l \leq k$, then $f$ is $\mathrm{sc}^{k+1}$. *Hidden hypotheses:* none ("$f$ induces a map $U_{m+l} \to F_m$" is the $\mathrm{sc}^0$ hypothesis). *Deviation:* none; the sc-smooth statement is a direct consequence. *Proof:* the 2010 paper's (Proposition 2.4): $f$ is the elementary map with no vectors, and every elementary map with $j$ vectors and $j + s \leq k + 1$ is $\mathrm{sc}^s$ (V1.38), with the derivatives $d^j f(x)$ supplied by the derivative tower of $f$ (V1.37). The induction cannot be run on the hypothesis itself: if $f$ satisfies it with $k$, then $Tf$ in general does not satisfy it with $k-1$, since its second component $(x, h) \mapsto Df(x)h : U_{m+l+1} \times E_{m+l} \to F_m$ is only $C^l$. Under the stronger hypothesis the level maps of $Tf$ are $C^k$ and the induction is immediate.

**V1.19 (Corollary 1.20; `Polyfold.corollary_1_20_text`, `Polyfold.IsScSmooth.of_contDiffOn_levels_finite` `Polyfold.IsScSmooth.of_contDiffOn_finiteDimensional`, `Polyfold.corollary_1_20_zero`, `Polyfold.corollary_1_20`, `Polyfold.corollary_1_20_base`).** Let $U$ be an sc-domain in $E$, $V$ a finite-dimensional normed space with its constant sc-structure and $f : E_0 \to V$. If for some $k$ and all $0 \leq l \leq k$ the map $f \circ \iota_l : U_l \to V$ is $C^{l+1}$ within $U_l$, then $f$ is $\mathrm{sc}^{k+1}$ on $U$. If $f \circ \iota_l$ is $C^{l+1}$ within $U_l$ for every $l$, then $f$ is sc-smooth; if $f$ is $C^\infty$ within $U$, then $f$ is sc-smooth. Also proved, under the stronger hypothesis: if $f \circ \iota_m$ is $C^{k+1}$ within $U_m$ for every $m$, in particular if $f$ is $C^{k+1}$ within $U$, then $f$ is $\mathrm{sc}^{k+1}$; for $k = 0$ this is the text's hypothesis. *Book:* p. 13: if $f : U_l \to \mathbb{R}^N$ is $C^{l+1}$ for $0 \leq l \leq k$ then $f$ is $\mathrm{sc}^{k+1}$. *Hidden hypotheses:* none; no $\mathrm{sc}^0$ hypothesis is needed, since $f \circ \iota_m$ is the continuous $f \circ \iota_0$ composed with an inclusion. *Deviation:* none ($\mathbb{R}^N$ is any finite-dimensional space); the two sc-smooth statements are the exposition's additions. *Proof:* Proposition 1.19 (V1.18) for the constant structure, the hypothesis on $U_{m+l} \to V$ being the restriction of the one on $U_l$.

**V1.20 (Theorem 1.21, the chain rule; `Polyfold.IsSc1With.hasFDerivWithinAt_comp`, `Polyfold.IsSc1With.comp`, `Polyfold.theorem_1_21'`, `Polyfold.theorem_1_21`, `Polyfold.IsSck.comp`, `Polyfold.IsSck.comp_of_isRelOpenInQuadrant`, `Polyfold.IsScSmooth.comp`).** Let $U \subseteq C \subseteq E$ and $V \subseteq D \subseteq F$ be relatively open subsets of partial quadrants, $G$ an sc-Banach space, $f : E_0 \to F_0$ $\mathrm{sc}^1$ on $U$ with $f(U) \subseteq V$, and $g : F_0 \to G_0$ $\mathrm{sc}^1$ on $V$. Then $g \circ f$ is $\mathrm{sc}^1$ on $U$, and $T(g \circ f) = Tg \circ Tf$ on $TU$. More precisely: if $g$ is $\mathrm{sc}^1$ on $V$ with derivatives $Dg$, $f$ is $\mathrm{sc}^1$ on $U$ with derivatives $Df$, $f(U) \subseteq V$, and $V_1$ is the intersection of an open and a convex set, then $g \circ f$ is $\mathrm{sc}^1$ on $U$ with derivatives $x \mapsto Dg(f_1(x)) \circ Df(x)$, with no hypothesis on $U$; and $g \circ f \circ \iota_1$ has this derivative at every $x \in U_1$ within $U_1$. Iterated: if $U, V$ are sc-domains (in particular relatively open subsets of partial quadrants), $f$ $\mathrm{sc}^k$ on $U$ with $f(U) \subseteq V$ and $g$ $\mathrm{sc}^k$ on $V$, then $g \circ f$ is $\mathrm{sc}^k$ on $U$; sc-smooth maps compose to sc-smooth maps. *Book:* p. 13, with a third tuple $W \subset Q \subset G$ for the target of $g$; the proof is §1.4.2, pp. 15–16. *Hidden hypotheses:* the segment from $f(x)$ to $f(x+h)$ lies in $V_1$ for $|h|_1$ small, which uses the convexity of $D_1$, the openness of $V_1$ in $D_1$ and the continuity of $f : U_1 \to F_1$; the uniqueness of $D(g \circ f)(x)$ behind "$T(g \circ f) = Tg \circ Tf$". *Deviation:* none in the statement; $W \subset Q \subset G$ plays no role. *Proof:* the text's two integrals are replaced by the mean value inequality for $t \mapsto g(y_t) - t\, Dg(f(x))(f(x+h) - f(x))$ on $[0, 1]$; the first integral is not needed since $Dg(f(x))$ is bounded on $F_0$; the uniform convergence on the compact closure of $\{Df(x)h/|h|_1\}$ is Mathlib's `IsCompact.eventually_forall_of_forall_eventually` rather than a subsequence argument. The text's appeal to Proposition 1.18 is not used.

**V1.21 (Definition 1.23; `Polyfold.quadrantDeg`, `Polyfold.degIndexWith`, `Polyfold.degIndex`, `Polyfold.degIndex_eq`, `Polyfold.degIndex_quadrant`).** For $p = (a, w) \in \mathbb{R}^n \oplus W$ let $\#(p)$

be the number of $i$ with $a_i = 0$; for an sc-isomorphism $T : E \to \mathbb{R}^n \oplus W$ presenting a partial quadrant $C$ let $d_T(x) := \#(T_0 x)$. The *degeneracy index* $d_C : E_0 \to \mathbb{N}$ is $d_T$ for a chosen presentation $T$ of $C$ (and 0 if $C$ is not a partial quadrant); for every presentation $T$ and every $x \in C$, $d_C(x) = d_T(x)$. On the quadrant itself $d(p) = \#(p)$. *Book:* p. 14: $d_C(x) = \#\{i : a_i = 0\}$ where $T(x) = (a_1, \dots, a_k, w)$; interior points, boundary points and corners. *Hidden hypotheses:* none. *Deviation:* none; $d_C$ is total on $E_0$ with junk values off $C$, and the classification into interior, boundary and corner points is terminology.

**V1.22 (Lemma 1.24; `Polyfold.quadrantDeg_eq_of_equiv`, `Polyfold.lemma_1_24`, `Polyfold.lemma_1_24_dim`).** Let $T : E \to \mathbb{R}^n \oplus W$ and $T' : E \to \mathbb{R}^{n'} \oplus W'$ be sc-isomorphisms with $C = T_0^{-1}(Q_n) = T_0'^{-1}(Q_{n'})$. Then $d_T(x) = d_{T'}(x)$ for every $x \in C$, and $n = n'$. Core: if $S : \mathbb{R}^n \times W_0 \to \mathbb{R}^{n'} \times W_0'$ is a continuous linear equivalence with $q \in Q_n \iff Sq \in Q_{n'}$, then $\#(p) = \#(Sp)$ for every $p \in Q_n$. *Book:* p. 14, "$d_C$ does not depend on the choice of $T$", proof on p. 17. *Hidden hypotheses:* none; the proof uses only linear algebra on level 0. *Proof:* the text's, except that $k = k'$ is not proved first but derived from the lemma at $x = 0$; the codimension of $E_x$ is $\dim(V/E_x) = \#I(x)$ with $E_x = \ker((b, v) \mapsto (b_i)_{i \in I(x)})$, and "$S(E_x) \subset E'_y$ hence equality" is completed by the same inclusion for $S^{-1}$.

**V1.23 (Theorem 1.25, boundary recognition; `Polyfold.theorem_1_25`, `Polyfold.theorem_1_25_lvl1`, `Polyfold.degIndexWith_le_of_germ`, `Polyfold.eventually_degIndexWith_le`, `Polyfold.exists_tendsto_lv` `Polyfold.IsSc1With.mapsTo_minimalSubspace`).** Let $U \subseteq C \subseteq E$ and $U' \subseteq C' \subseteq E'$ be relatively open subsets of partial quadrants, $f : E_0 \to E'_0$ $\mathrm{sc}^1$ on $U$ with $f(U) \subseteq U'$, $g : E'_0 \to E_0$ $\mathrm{sc}^1$ on $U'$ with $g(U') \subseteq U$, $g(f(x)) = x$ for $x \in U$ and $f(g(y)) = y$ for $y \in U'$. Then $d_C(x) = d_{C'}(f(x))$ for every $x \in U$. Ingredients: $d_{C'}$ is upper semicontinuous ($d_{C'}(y') \le d_{C'}(y)$ for $y'$ near $y$); every $x \in C$ is a limit of points of $C$ on level 1 with the same index; and for $x \in U_1$ the derivative $Df(x)$ maps $E_x := \{v : (T_0 v)_i = 0 \text{ whenever } (T_0 x)_i = 0\}$ into $E'_{f(x)}$, from which $d_C(x) \le d_{C'}(f(x))$ follows by comparing codimensions through the chain rule. *Book:* p. 14: for tuples $(U, C, E)$, $(U', C', E')$ and a germ $f : (U, x) \to (U', x')$ of a local sc-diffeomorphism, $d_C(x) = d_{C'}(f(x))$; no proof, a reference to "[20], Theorem 1.19". *Hidden hypotheses:* the notion "germ of a local sc-diffeomorphism", undefined in Chapter 1, read as an $\mathrm{sc}^1$ map $f : U \to U'$ with $\mathrm{sc}^1$ inverse $g : U' \to U$ on relatively open neighbourhoods $U$, $U'$ of $x$, $f(x)$ (which may be shrunk at will). *Deviation (stronger):* only $\mathrm{sc}^1$ is required of $f$ and $g$, where the text has sc-smooth. *Proof:* the argument of §2.3 (Lemma 2.29, Theorem 2.31, Corollary 2.32) for the identity retraction, with level-1 points in place of smooth points; the Lean file is `Polyfold/Chapter2/DegeneracyIndex.lean`.

**V1.24 (Fredholm operators between Banach spaces, Riesz–Schauder; `Polyfold.isStrictMap_of_isClosed_ran` `Polyfold.isFredholm_of_banach`, `Polyfold.isCompactOperator_of_finiteDimensional_range`, `Polyfold.finiteDimensional_ker_one_add`, `Polyfold.isClosed_range_one_add`, `Polyfold.coFG_range_one_` `Polyfold.isFredholm_one_add`).** Let $E$, $F$ be real Banach spaces. A continuous linear $u : E \to F$ with closed range is a strict map; if moreover $\ker u$ is finite-dimensional and range $u$ has finite codimension, $u$ is a Fredholm operator in Mathlib's sense. A continuous linear map of finite rank is a compact operator. For a compact operator $c : E \to E$: $\ker(1 + c)$ is finite-dimensional, $\mathrm{range}(1 + c)$ is closed and of finite codimension, and $1 + c$ is Fredholm. *Book:* not stated; used in §1.4.1, p. 14 ("$T + S : E_m \to F_m$ is Fredholm", "its image $Y$ is closed"). *Deviation (not in the book):* infrastructure Mathlib v4.33.1 lacks.

**V1.25 (compact perturbations of Fredholm operators; `Polyfold.isFredholm_add_isCompactOperator`).** Let $E$, $F$ be real Banach spaces, $u : E \to F$ Fredholm and $k : E \to F$ compact. Then $u + k$ is

Fredholm. *Book:* not stated; the first line of §1.4.1. *Deviation (not in the book):* infrastructure; the invariance of the index, which the exposition records (Kato IV.5.26) and the text does not, is V1.28.

**V1.26 (the compact-open lemma and the level operators; `Polyfold.exists_delta_of_continuousOn`, `Polyfold.continuousOn_clm_comp_of_isCompactOperator`, `Polyfold.contDiffOn_clm_comp_of_isCompactOp` `Polyfold.levelDeriv`, `Polyfold.IsSc1With.exists_levelDeriv`, `Polyfold.IsSc1With.toBase_levelDeriv`, `Polyfold.IsSc1With.levelMap_tangentMapWith`, `Polyfold.IsSc1With.continuousOn_levelDeriv`, `Polyfold.IsSc1With.continuousOn_levelDeriv_clm`, `Polyfold.IsSc1With.hasFDerivWithinAt_level`, `Polyfold.IsSc1With.contDiffOn_level`, `Polyfold.IsSc1With.levelDeriv_eq`).** *Compact-open lemma.* Let $X$, $Y$, $Y'$, $Z$ be real normed spaces, $\iota : Y' \to Y$ a compact operator and $A : X \to \mathcal{L}(Y, Z)$. If $(x, y) \mapsto A(x)y$ is continuous on $s \times Y$, then $x \mapsto A(x) \circ \iota$ is continuous on $s$ into $\mathcal{L}(Y', Z)$ with the operator norm; if $s$ is the intersection of an open and a convex set with unique differentiability at every point and $(x, y) \mapsto A(x)y$ is $C^k$ within $s \times Y$, then $x \mapsto A(x) \circ \iota$ is $C^k$ within $s$. *Level operators.* Let $f$ be $\mathrm{sc}^1$ on $U$ with derivatives $Df$, $m \in \mathbb{N}$ and $x \in U_{m+1}$. There is a unique continuous linear $Df(x) : E_m \to F_m$ (written `levelDeriv`) with $\iota_m \circ Df(x) = Df(\iota x) \circ \iota_m$, namely the second component of the level-$m$ tangent map, which is $(x, h) \mapsto (f_{m+1}(x), Df(x)h)$ on $U_{m+1} \times E_m$; $(x, h) \mapsto Df(x)h$ is continuous on $U_{m+1} \times E_m$; $x \mapsto Df(x) \circ \iota$ is continuous $U_{m+1} \to \mathcal{L}(E_{m+1}, F_m)$. If $U$ is an sc-domain, $f_m \circ \iota : U_{m+1} \to F_m$ has derivative $Df(x) \circ \iota$ at $x$ within $U_{m+1}$, hence is $C^1$ within $U_{m+1}$; and if $f_m : U_m \to F_m$ has derivative $L$ at $\iota x$ within $U_m$, then $L = Df(x)$. *Book:* not stated as such; the compactness argument of the 2010 paper's Propositions 2.1–2.3 (quoted in §1.2) and its integral identity $f(x + h) - f(x) = \int_0^1 Df(x + \tau h)h \, d\tau$. *Deviation (not in the book):* the $C^k$ case of the lemma, proved by induction with the mean value inequality, replaces the multilinear maps $\Gamma(x)$ and the differentiation under the integral sign of the 2010 paper; the integral identity is formed as a Bochner integral in $F_m$ and identified with $f(x + h) - f(x)$ by the fundamental theorem of calculus for the $C^1$ map $f : U_1 \to F_0$.

**V1.27 (sc-domains; `Polyfold.IsScDomain`, `Polyfold.IsRelOpenInQuadrant.isScDomain`, `Polyfold.IsScDomain.tangentSet`, `Polyfold.IsScDomain.shift`).** $U \subseteq E_0$ is an *sc-domain* if for every $m$ the level set $U_m$ is the intersection of an open and a convex subset of $E_m$, and $U_m$ has unique differentiability at each of its points. A relatively open subset of a partial quadrant is an sc-domain; if $U$ is an sc-domain, so are $TU$ in $TE$ and $U_1$ in $E^1$. *Book:* not stated; the text works with tuples $(U, C, E)$ and asserts that $T(U, C, E)$ is a tuple. *Deviation (not in the book):* the level-wise regularity that the proofs of §§1.2–1.3 use, stable under the tangent construction and the level shift, stated in place of the tuple.

**V1.28 (the Fredholm index and its invariance; `Polyfold.linIndex`, `Polyfold.HasFiniteIndex`, `Polyfold.linIndex_comp`, `Polyfold.linIndex_eq_zero_of_finiteRange_sub_id`, `Polyfold.HasFiniteIndex.` `Polyfold.HasFiniteIndex.of_comp_of_comp`, `Polyfold.fredholmIndex`, `Polyfold.IsFredholm.hasFiniteInd` `Polyfold.fredholmIndex_add_of_norm_lt`, `Polyfold.fredholmIndex_add_isCompactOperator`).** For a linear map $u : E \to F$ over a field let $\mathrm{ind}(u) := \dim \ker u - \dim(F / \operatorname{range} u) \in \mathbb{Z}$, meaningful when both dimensions are finite ("finite index"). If $u$ and $v : F \to G$ have finite index, so has $v \circ u$, and $\mathrm{ind}(v \circ u) = \mathrm{ind}(u) + \mathrm{ind}(v)$. If $a$ is bijective and $r$ has finite rank, $a + r$ has finite index 0 (in particular $1 + r$). If $v \circ u$ and $u \circ v$ have finite index, so has $u$. For real Banach spaces $E$, $F$: a Fredholm operator $u : E \to F$ has finite index, and $\mathrm{ind}(u)$ is its *Fredholm index*. If $v : F \to E$ is a quasi-inverse of $u$ ($vu - 1$ and $uv - 1$ have finite rank) and $\|v\| \, \|s\| < 1$, then $u + s$ has finite index and $\mathrm{ind}(u + s) = \mathrm{ind}(u)$. If $u$ is Fredholm and $k : E \to F$ is compact, then $\mathrm{ind}(u + k) = \mathrm{ind}(u)$. *Book:* not stated; the classical fact behind the exposition's (F2) (Kato IV.5.26), which the text's §1.4.1 would need to conclude that

the index is preserved. *Deviation (not in the book):* infrastructure Mathlib v4.33.1 lacks (it has Fredholm operators but no index). *Proof:* additivity by rank–nullity along the exact sequence $0 \to \ker u \to \ker vu \to \operatorname{range} u \cap \ker v \to F/\operatorname{range} u \to G/\operatorname{range} vu \to G/\operatorname{range} v \to 0$; small perturbations through $v(u+s) = (1+vs) + (vu-1)$ and $(u+s)v = (1+sv) + (uv-1)$, where $1+vs$, $1+sv$ are invertible by the Neumann series; compact perturbations by the local constancy of $t \mapsto \operatorname{ind}(u+tk)$ on $\mathbb{R}$ (each $u + t_0 k$ is Fredholm by V1.25), not by Kato's argument.

**V1.29 (Lemma 2.D, the level operators of an sc-Fredholm operator;** `Polyfold.ScFredholmSplitting.C_le_smoo` `Polyfold.ScBanach.finrank_levelSub_of_le_smooth`, `Polyfold.ScFredholmSplitting.fredholmIndex_lvl`, `Polyfold.IsScFredholm.fredholmIndex_lvl`, `Polyfold.scIndex_eq_fredholmIndex`**).** Let $T : E \to F$ be an sc-operator with an sc-Fredholm splitting $(K, X, C, Y)$ (V1.7). Then $C \le F_\infty$ (and $K \le E_\infty$, V1.7); a subspace $A \le E_\infty$ has $\dim A_m = \dim A$ for every $m$; and for every $m$ the level map $T_m : E_m \to F_m$ is a Fredholm operator with $\operatorname{ind}(T_m) = \dim K - \dim C$. Hence for an sc-Fredholm $T$, $\operatorname{ind}(T_m) = \operatorname{ind}(T) := \dim\ker T_0 - \dim(F_0/\operatorname{range} T_0)$ for every $m$; the latter is the Fredholm index of $T_0$. *Book:* p. 9, after Definition 1.8: $K \subset E_\infty$, $T(X_m) = Y_m$, $E_m = K \oplus X_m$, "$F = C \oplus T(E_m)$"; the level-wise index is not stated. *Hidden hypotheses:* none. *Deviation (stronger):* the Fredholm property and the index on every level are added (the exposition's Lemma 2.D). *Proof:* $\ker T_m = K_m \cong K$ and $F_m/\operatorname{range} T_m = F_m/Y_m \cong C_m \cong C$, the isomorphisms $A_m \cong A$ coming from $A \le E_\infty$.

**V1.30 (Proposition 1.10, the index;** `Polyfold.proposition_1_10_index`, `Polyfold.proposition_1_10_index_l`
Let $T : E \to F$ be sc-Fredholm and $S : E \to F$ an sc$^+$-operator. Then $\operatorname{ind}(T+S) = \operatorname{ind}(T)$, the index being $\dim\ker(\cdot)_0 - \dim\operatorname{coker}(\cdot)_0$; and for every $m$ the level map $(T+S)_m : E_m \to F_m$ is Fredholm of index $\operatorname{ind}(T)$. *Book:* p. 10, Proposition 1.10 states only that $T+S$ is sc-Fredholm (V1.9). *Hidden hypotheses:* none. *Deviation (stronger):* the index statement is the exposition's addition. *Proof:* on level 0, $T_0$ is Fredholm (V1.7) and $S_0$ compact (V1.3), so V1.28 applies; on level $m$, V1.29 for the sc-Fredholm operator $T+S$ (V1.9).

**V1.31 (finite-dimensional spaces;** `Polyfold.ScBanach.smooth_eq_top_of_finiteDimensional`, `Polyfold.ScBanach.toBase_bijective_of_finiteDimensional`, `Polyfold.ScBanach.isoConstOfFiniteDimer` `Polyfold.ScScale.finiteDimensional_of_surjective_incl`**).** Let $E$ be an sc-Banach space with $E_0$ finite-dimensional. Then every point of $E_0$ is smooth, every $\iota_m : E_m \to E_0$ is bijective, and the maps $\iota_m$ form an sc-isomorphism of $E$ onto $E_0$ with its constant structure. For a scale $E$ (no density assumed) and any $m$: if the inclusion $E_{m+1} \to E_m$ is surjective, then $E_m$ is finite-dimensional. *Book:* p. 6, after Definition 1.1: a finite-dimensional Banach space has precisely one sc-structure, the constant one; on an infinite-dimensional space the constant structure is not an sc-structure, because the inclusions are not compact. *Hidden hypotheses:* none. *Deviation (restated):* "precisely one" is read, for the indexed families of V1.1, as "sc-isomorphic to the constant structure by the inclusions" (so the level norms are all equivalent), and "the constant structure fails (1)" as "an onto inclusion forces finite dimension". *Proof:* $E_\infty$ is a dense, hence (finite dimension) closed, hence full subspace of $E_0$; an onto compact inclusion is an isomorphism (open mapping theorem), so the identity of $E_m$ is compact.

**V1.32 (sc-splittings are sc-projections;** `Polyfold.ScBanach.IsScSplitting.isTopCompl`, `Polyfold.ScBanach.IsScSplitting.scProj`, `Polyfold.ScBanach.IsScSplitting.scProj_idem`, `Polyfold.ScBanach.IsScSplitting.range_scProj`, `Polyfold.ScBanach.IsScSplitting.ker_scProj`, `Polyfold.ScOperator.idem_lvl`, `Polyfold.ScOperator.levelSub_range_eq`, `Polyfold.ScBanach.isScSplitt` `Polyfold.ScBanach.isScSplitting_iff_exists_scProj`**).** Let $A, B \subseteq E_0$. If $E = A \oplus B$ is an sc-splitting, then on every level $A_m$, $B_m$ are topological complements, and the continuous projections of $E_m$ onto $A_m$ along $B_m$ form an sc-operator $P : E \to E$ with $P_m^2 = P_m$, $P_m(E_m) = A_m$

and $\ker P_m = B_m$. Conversely, if $P : E \to E$ is an sc-operator with $P_0^2 = P_0$, then $P_m^2 = P_m$ and $(P_0(E_0))_m = P_m(E_m)$ for every $m$, and $E = P_0(E_0) \oplus \ker P_0$ is an sc-splitting. Hence $E = A \oplus B$ is an sc-splitting iff some sc-operator $P$ with $P_0^2 = P_0$ has range $A$ and kernel $B$. *Book:* pp. 7–8: the definition of an sc-splitting, and in the proof of Proposition 1.6 the step "since $P \circ P = P$, it is a sc-projection" with $G := (1 - P)(E)$. *Hidden hypotheses:* none. *Deviation (hypotheses explicit):* the equivalence the text uses is stated and proved (the exposition's Lemma 2.B); $\ker P_0$ is the text's $(1 - P)(E)$. The formal Proposition 1.6 (V1.5) does not depend on it.

**V1.33 (the smooth points of a quotient; `Polyfold.ScBanach.exists_smooth_mem_coset`, `Polyfold.ScBanach.mk_mem_quotient_smooth_iff`).** Let $A$ be an sc-subspace of $E$ and $x \in E_0$. If for every $j$ there is $y_j \in E_j$ with $\iota_j y_j - x \in A$, then there is $y \in E_\infty$ with $y - x \in A$. Equivalently: the class of $x$ is a smooth point of the quotient $E/A$ (V1.6) if and only if $x + A$ contains a smooth point of $E$. *Book:* p. 9, in the proof of Proposition 1.7: "$(E/A)_\infty$ consists of all elements of the form $x + A_\infty$ with $x \in E_\infty$". *Hidden hypotheses:* that $A$ is an sc-subspace (the density of $A_\infty$ in every $A_j$) is used; the text has it. *Deviation:* none. *Proof:* a Mittag-Leffler construction (the exposition's Lemma 2.C), with $\prod_{i<j} \max(1, \|E_{i+1} \to E_i\|)$ bounding all the inclusions $E_j \to E_i$, $i \le j$.

**V1.34 (continuity of $Df$ for the compact-open topology; `Polyfold.continuousOn_apply_iff_uniformOnCompac` `Polyfold.IsSc1With.deriv_uniformOnCompact`, `Polyfold.exists_isSc1_not_continuousOn_deriv`).** Let $X$ be a metric space, $Y$, $Z$ real normed spaces, $s \subseteq X$ and $A : X \to \mathcal{L}(Y, Z)$. Then $(x, h) \mapsto A(x)h$ is continuous on $s \times Y$ if and only if for every $x \in s$, every compact $K \subseteq Y$ and every $\varepsilon > 0$ there is a neighbourhood of $x$ in $s$ on which $\|A(x')w - A(x)w\| < \varepsilon$ for all $w \in K$. Applied to an sc$^1$ map with derivatives $Df$: $x \mapsto Df(x)$ is continuous on $U_1$ for the topology of uniform convergence on compact subsets of $E_0$. There is an sc$^1$ map on a relatively open subset of a partial quadrant for which no derivative family is continuous on $U_1$ into $\mathcal{L}(E_0, F_0)$ with the operator norm (the example of V1.13). *Book:* p. 11: $x \mapsto Df(x)$ is in general not continuous for the operator norm, but it is continuous for the compact-open topology. *Hidden hypotheses:* none. *Deviation (restated):* the compact-open topology on $\mathcal{L}(Y, Z)$ is written out as uniform convergence on compact sets, and the remark becomes an equivalence with level 0 of condition (3) of Definition 1.14 (the exposition's Lemma 3.F). The negative half of the remark is proved by a sequence-space example (V1.13), not by the shift map of the exposition's §6, which needs weighted Sobolev spaces.

**V1.35 (finite dimensions: sc$^1$ is $C^1$, also at the boundary; `Polyfold.ScBanach.lvlSet_const`, `Polyfold.isSc1_iff_contDiffOn_const`, `Polyfold.isSc1_iff_fderivWithin_const`, `Polyfold.isScSmooth_i` `Polyfold.isSc1_iff_contDiffOn_of_isRelOpenInQuadrant`, `Polyfold.isScSmooth_iff_contDiffOn_of_isRe`).** Let $V$, $W$ be finite-dimensional real normed spaces with their constant sc-structures, $f : V \to W$ and $U \subseteq V$ an sc-domain (V1.27), for instance a relatively open subset of a partial quadrant of $V$. Then $f$ is sc$^1$ on $U$ if and only if $f$ is $C^1$ within $U$; equivalently, $f$ is differentiable within $U$ at every point of $U$ and the derivative within $U$, $x \mapsto \mathrm{fderivWithin}\ f\ U\ x$, is continuous on $U$. And $f$ is sc-smooth on $U$ if and only if it is $C^\infty$ within $U$. *Book:* p. 11: "the sc$^1$-maps between finite dimensional Banach spaces are the familiar $C^1$-maps". *Hidden hypotheses:* the meaning of $C^1$ at boundary points of a quadrant, never defined in the text; here the derivative within $U$, which at a boundary point is the one-sided derivative along $U$ and is unique because the quadrant has non-empty interior. *Deviation (hypotheses explicit):* none beyond this; by V1.31 every sc-structure on a finite-dimensional space is the constant one up to sc-isomorphism. *Proof:* Propositions 1.16 and 1.19 at $k = 0$ (V1.15, V1.18) for the levels of the constant structure, which all equal $U$.

**V1.36 (level shifts and tangents of tuples; `Polyfold.ScIso.shift`, `Polyfold.IsPartialQuadrant.shift`, `Polyfold.IsRelOpenInQuadrant.shift`, `Polyfold.tangentQuadrantIso`, `Polyfold.IsPartialQuadrant.tange` `Polyfold.IsRelOpenInQuadrant.tangentSet`, `Polyfold.ScBanach.shift_tangentSet`).** Let $C$ be a partial quadrant in $E$, presented by an sc-isomorphism $T : E \to \mathbb{R}^n \oplus W$. Then $C_1$ is a partial quadrant in $E^1$, presented by the level shift $T^1 : E^1 \to \mathbb{R}^n \oplus W^1$; and $TC = \{(x,h) \in E_1 \times E_0 : x \in C_1\}$ is a partial quadrant in $TE$, presented by $(x,h) \mapsto (a,(w,T_0h))$ where $T_1x = (a,w)$, an sc-isomorphism $TE \to \mathbb{R}^n \oplus (W^1 \oplus (\mathbb{R}^n \oplus W))$. If $U$ is relatively open in $C$, then $U_1$ is relatively open in $C_1$ and $TU$ in $TC$; and the level-1 part of $TU$ is $T(U_1)$. *Book:* p. 10, after Definition 1.13: "$T(U,C,E)$ is a tuple consisting again of a relatively open subset $TU$ in the partial quadrant $TC$"; the corresponding fact for $C^1 \subset E^1$ is used without being stated. *Hidden hypotheses:* none. *Deviation:* none; $C^k$ for $k \geq 2$ is the iterate. These are the exposition's Lemmas 3.A(2) and 3.B, proved in the Chapter 2 files (`Polyfold/Chapter2/LevelPolyfold.lean`, `Polyfold/Chapter2/TangentBundle.lean`).

**V1.37 (the derivative tower; `Polyfold.IsDerivTower`, `Polyfold.levelDerivTower`, `Polyfold.baseDerivTower`, `Polyfold.contDiffOn_levels_min`, `Polyfold.levelDerivTower_toBase`, `Polyfold.baseDerivTower_toBase`, `Polyfold.isDerivTower_of_contDiffOn_levels`, `Polyfold.IsDerivTower.eq_of_toBase`).** A derivative tower of order $k+1$ for $f : E_0 \to F_0$ on $U$ consists of maps $\Phi_i : E_0 \times E_0^i \to F_0$ with $\Phi_0(x) = f(x)$ and, for $x \in U_{m+l}$, continuous $i$-linear maps $\Psi_{i,m,l}(x) : E_{m+l}^i \to F_m$, for $i \leq l+1$ and $i \leq k+1$, such that (a) $\iota_m \Psi_{i,m,l}(x)(h_1,\dots,h_i) = \Phi_i(\iota_{m+l}x)(\iota_{m+l}h_1,\dots,\iota_{m+l}h_i)$; (b) $\Psi_{i,m,l}$ is continuous on $U_{m+l}$ into the space of continuous $i$-linear maps; (c) for $i \leq l$ and $i \leq k$, $\Psi_{i,m,l}$ is differentiable within $U_{m+l}$ at every point, with derivative $\xi \mapsto \Psi_{i+1,m,l}(x)(\xi,\cdot)$. If $U$ is an sc-domain, $f$ is sc$^0$ on $U$ and $f_m \circ \iota : U_{m+l} \to F_m$ is $C^{l+1}$ within $U_{m+l}$ for all $m$ and $l \leq k$, then the iterated derivatives within the level sets, $\Psi_{i,m,l} := d^i(f_m \circ \iota)$ on $U_{m+l}$ and $\Phi_i$ the $i$-th derivative of $f : U_{i-1} \to F_0$ at lifts to $E_{i-1}$, form a derivative tower of order $k+1$. In a tower, two values $\Psi_{i,m,l}(x)(h)$, $\Psi_{i,m,l'}(x')(h')$ with the same images in $E_0$ are equal. *Book:* not stated; the 2010 paper's proof of Proposition 2.4 (pp. 51–53) writes one symbol $d^jf(x)$ for the derivatives on all level pairs. *Hidden hypotheses:* the compatibility (a) is the identification the text makes tacitly. *Deviation (not in the book):* a tool of the formalization. *Proof:* (a) from the chain rule for iterated derivatives within a set composed on the left and on the right with continuous linear maps (the inclusions), and the uniqueness of derivatives on the level sets; for $l > k$ the hypothesis is used with order $k+1$, restricted from $U_{m+k}$.

**V1.38 (elementary maps and the induction $(\mathrm{S}_k)$; `Polyfold.ElemData`, `Polyfold.ElemData.Good`, `Polyfold.ElemData.dom`, `Polyfold.ElemData.map`, `Polyfold.ElemData.map_toBase`, `Polyfold.ElemData.leve` `Polyfold.ElemData.isSc0`, `Polyfold.ElemData.deriv`, `Polyfold.ElemData.hasFDerivWithinAt_level`, `Polyfold.ElemData.continuousOn_deriv`, `Polyfold.ElemData.isSc1Levels`, `Polyfold.ElemData.isSc1With`, `Polyfold.ElemData.shiftData`, `Polyfold.ElemData.derivData`, `Polyfold.ElemData.slotData`, `Polyfold.ElemData.tangentMap_eqOn`, `Polyfold.ElemData.isSck`, `Polyfold.ElemData.base`, `Polyfold.ScBanach.shiftAdd`, `Polyfold.ScBanach.shiftAddZeroIso`, `Polyfold.ScBanach.shiftAddShiftIsc`** Fix a derivative tower of order $k+1$ for $f$ on $U$ (V1.37), and write $E^p$ for the sc-Banach space with levels $(E^p)_l = E_{p+l}$ ($E \cong E^0$ and $(E^p)^1 \cong E^{p+1}$ as sc-Banach spaces). Elementary data on an sc-Banach space $X$ are an sc-operator $\pi_0 : X \to E^p$, sc-operators $\pi_a : X \to E^{n_a}$ ($1 \leq a \leq i$) and a number $m$; the elementary map is $g(z) = \Phi_i(\pi_0 z)(\pi_1 z,\dots,\pi_i z)$, a map $X \to F^m$, on $W := \pi_0^{-1}(U_p)$. Assume $m+i \leq p$, $m+i \leq n_a+1$ for all $a$, and $i \leq k$. Then $g$ is sc$^0$ on $W$, its

level-$l$ map being $\Psi_i$ on level $(m+l, i-1)$ at the slots. If $W$ is an sc-domain, $g$ is $\mathrm{sc}^1$ with

$$Dg(z)\zeta = d^{i+1}f(\pi_0 z)(\pi_0\zeta, \pi_1 z, \dots, \pi_i z) + \sum_{a=1}^{i} d^i f(\pi_0 z)(\pi_1 z, \dots, \pi_a\zeta, \dots, \pi_i z),$$

and, if moreover $i+1 \le k$, its tangent map on $TW$ is $(g', g_0 + \sum_a g_a)$, through $(F^m)^1 \cong F^{m+1}$ in the first component, with $g'$ the elementary map of the data $(\pi_0, \pi_a)$ taken on the first factor of $TX = X^1 \oplus X$ (levels $p+1$, $n_a+1$, $m+1$), $g_0$ the one with $i+1$ vectors ($\pi_0$ on the second factor, $\pi_a$ on the first) and target shift $m$, and $g_a$ the one with $\pi_a$ moved to the second factor; all satisfy the three conditions. Finally, if $W$ is an sc-domain and $i+s \le k+1$, then $g$ is $\mathrm{sc}^s$ on $W$. *Book:* 2010 paper, Proposition 2.4, pp. 51–53, the inductive statement $(\mathrm{S}_k)$: the components of $T^{k+1}f$ are linear combinations of $d^j f(x)(h_1, \dots, h_j)$ on $U^{k+1} \oplus E^{n_1} \oplus \cdots \oplus E^{n_j} \to F^l$ with $j + l - 1 \le n_i \le k$, each $\mathrm{sc}^1$ with the derivative (11), and those of $T^{k+2}f$ are again of this form. *Hidden hypotheses:* the base-point condition $j + l \le k+1$, which the text derives from its bounds. *Deviation (not in the book):* the induction runs on $s$ over all elementary maps on all $X$, not on the description of $T^{k+1}f$; the upper bound $n_i \le k$ is not needed, and its consequences $j+l \le p$ and $j \le k$ are conditions of their own. *Proof:* $\mathrm{sc}^1$ by the level criterion of Proposition 1.16 (V1.15): $Dg$ exists one level down because $d^{i+1}f$ and $d^i f$ do, by the derivative of the multilinear evaluation (V1.39); $\mathrm{sc}^{s+1}$ from the tangent map, the pairing, the sum rule (V1.39) and the induction hypothesis.

**V1.39 (sums of $\mathrm{sc}^k$ maps and the derivative of multilinear evaluation; `Polyfold.IsSck.zero`, `Polyfold.IsSck.add`, `Polyfold.IsSck.sum`, `Polyfold.hasFDerivAt_cmm_eval`).** Let $W$ be an sc-domain in $X$. The constant map $0$ is $\mathrm{sc}^k$ on $W$; if $g_1$, $g_2$ are $\mathrm{sc}^k$ on $W$, so is $g_1 + g_2$; a finite sum of $\mathrm{sc}^k$ maps is $\mathrm{sc}^k$. For continuous multilinear maps $A : \prod_a M_a \to Z$ between normed spaces, the evaluation $(A, h) \mapsto A(h)$ is differentiable at every $(A, h)$ with derivative $(B, \eta) \mapsto B(h) + \sum_a A(h_1, \dots, \eta_a, \dots)$. *Book:* not stated; the 2010 paper's proof of Proposition 2.4 uses both (the terms of $\pi \circ T^{k+1}f$ treated one at a time, and the derivative (11)). *Hidden hypotheses:* none. *Deviation (not in the book):* tools; only sums, not scalar multiples, are needed. *Proof:* $g_1 + g_2$ is the sc-smooth addition $Y \oplus Y \to Y$ composed with the pairing $(g_1, g_2)$ (`Polyfold.IsSck.scPair`), by the chain rule (V1.20); the sum by induction.

**What is not verified in this chapter**

- pp. 6–7, Example 1.2: the weighted Sobolev spaces $H^{m,\delta_m}(\mathbb{R} \times S^1)$ as an sc-structure on $L^2$. Mathlib has no weighted Sobolev spaces and no Rellich–Kondrachov theorem; the exposition proves it (Proposition 1.C). Planned, with the shift map of the 2010 paper (exposition §6), as the project's step H2.
- p. 11, Remark 1.15: partial cones. Not formalized, as in the exposition.
- p. 11, after Definition 1.14: the text's own instance of a derivative that is not norm continuous, the shift map of the exposition's §6, needs the weighted Sobolev spaces of Example 1.2. The remark itself is verified, the negative half by a sequence-space example (V1.13, V1.34), the positive half by V1.34.

### 3. Chapter 2 (HWZ §§2.1–2.6): verified except Proposition 2.2 and Lemma 2.29 as stated; Remark 2.28's negative claim is false

Chapter 2 of Part I (preprint pp. 18–60) builds differential geometry on the images of sc-smooth retractions. §2.1 defines retractions and retracts, proves that the tangent $TO = Tr(TU)$ does not depend on the retraction (Proposition 2.4), defines sc-smooth maps between retracts and proves the chain rule (Definitions 2.1–2.7, Proposition 2.6, Theorem 2.8). §2.2 glues retracts into M-polyfolds, gives them a filtration $X_m$ and a tangent bundle $TX$, proves metrizability (Theorem 2.16, Proposition 2.19) and treats sub-M-polyfolds (Definitions 2.11–2.20, Proposition 2.21). §2.3 transports the degeneracy index to M-polyfolds and compares it with the tangent space (Definitions 2.22–2.27, Lemmas 2.23, 2.26, 2.29, Propositions 2.24, 2.30, Theorem 2.31, Corollary 2.32). §2.4 introduces tame retracts and the faces of tame M-polyfolds (Definitions 2.33–2.39, Propositions 2.35–2.46, with the products of M-polyfolds that Proposition 2.46 uses), §2.5 strong bundles (Definitions 2.47–2.53, Proposition 2.54), and the appendix §2.6 proves Proposition 2.2, Theorem 2.16 and Proposition 2.19 with Smirnov's theorem (Theorem 2.55), Lemma 2.56 and Proposition 2.57. Verified: every definition except 2.9 and 2.38 (which are given only in the forms the theory uses) and every numbered result of §§2.1–2.3 and §2.5, with these exceptions: Proposition 2.2 (the classical submanifold statement, motivation only), Lemma 2.29 in its path formulation (its content, the sign arguments, is verified and Proposition 2.30 is proved from it), the openness of $p : TX \to X_1$ in Proposition 2.19(2). The second statement of Theorem 2.31 (equality makes $C_xO$ a partial quadrant) is verified without its hypothesis, and **Remark 2.28's claim that $C_xO$ is in general not a partial quadrant in $T_xO$ is false**: it is one at every smooth point of every sc-smooth retract (V2.33), by Flor's theorem on nonnegative idempotent matrices, proved in the needed case. Every numbered result of §2.4 is verified: Propositions 2.35, 2.36 (with "$C_xO$ is a partial quadrant", which holds for every retract, V2.33), Lemma 2.40, Corollary 2.41, Propositions 2.42, 2.43 (the retraction onto a face, and the face as a tame sub-M-polyfold with $d_F = d_O - 1$), 2.44 (both sentences, the intersections of local faces included), Theorem 2.45 and Proposition 2.46 (fibred products, with the products of M-polyfolds it uses). In §2.5 the M-polyfold $Y[1]$, the sc-smoothness of $P[1]$ and the chart-independence of $Y_{m,k}$ are verified, and the chart-wise definition of sc$^+$-sections is shown to be the text's; equivalence of atlases, of M-polyfolds and of strong bundles, is an equivalence relation (V2.11, V2.57), strong bundle charts restrict to open subsets of the base (V2.57), and the charts of a strong bundle depend only on its class of atlases (V2.58). **Theorem 2.16 holds with the text's hypotheses**: the formal proof adds none, avoids Lemma 2.56 and Proposition 2.57 (which need a regularity of $X_m$ that the text does not check) and proves Smirnov's theorem, which Mathlib lacks, in a pull-back form. Several statements are proved in a stronger form: Propositions 2.30, 2.35 and Theorem 2.31 at points on level 1 and for sc$^1$ germs, Proposition 2.35 from condition (2) of tameness alone, Corollary 2.32 for every relatively open subset of a partial quadrant, and Proposition 2.36's $d_O = d_C$ in the form "a tame chart realizes $d_X$" for every M-polyfold.

The setting. Sc-Banach spaces and maps are those of Chapter 1. A *retraction* is a total map $r : E_0 \to E_0$ with an explicit domain $U$; an *M-polyfold* is a Hausdorff paracompact space with one sc-smooth atlas representing its class, and every notion is defined through the maximal atlas of all charts compatible with it. Sc-smoothness between retracts quantifies over all retractions onto the source (equivalent to the text's "one"). Where the text computes in the model $C = [0,\infty)^k \oplus W$, the formal statements use a presentation $T : E \cong \mathbb{R}^n \oplus W$ of $C$ and the coordinate functionals $\lambda_i = (T\,\cdot)_i$; the subspaces $E_x$, the cones $C_x$ and tameness are defined through $T$. The tangent bundle $TX$ is a quotient of tuples $(x, \varphi, h)$ with its topology and charts; its fibres $T_xX$ are sets, identified with $T_{\varphi(x)}O$ by the charts, with no linear structure of their own. Theorem 1.25 of Chapter

1 (V1.23) is proved in `Polyfold/Chapter2/DegeneracyIndex.lean` by the argument of §2.3.

**V2.1 (sc-smooth retractions; `Polyfold.ScRetraction`, `Polyfold.ScRetraction.retract`, `Polyfold.ScRetraction.mem_retract_iff`, `Polyfold.ScRetraction.apply_eq_self_of_mem_retract`, `Polyfold.ScRetraction.lvlSet_retract_subset`, `Polyfold.ScRetraction.scDeriv_comp_self`).** Let $C$ be a partial quadrant in $E$. An *sc-smooth retraction* on $C$ consists of a set $U \subseteq E_0$ relatively open in $C$ and a map $r : E_0 \to E_0$ with $r(U) \subseteq U$, $r$ sc-smooth on $U$ and $r(r(x)) = r(x)$ for $x \in U$. Its *retract* is $O = r(U)$; $O \subseteq U$, $x \in O$ iff $x \in U$ and $r(x) = x$, and $O_m \subseteq U_m$. For $x \in O_1$, $Dr(x) \circ Dr(x) = Dr(x)$. *Book:* Definition 2.1, p. 18, and the remark after Definition 2.3, p. 19. *Hidden hypotheses:* none. *Deviation:* none; the values of $r$ off $U$ play no role.

**V2.2 (sc-smooth retracts; `Polyfold.IsScRetract`, `Polyfold.ScRetraction.isScRetract`, `Polyfold.ScRetraction.id`, `Polyfold.IsRelOpenInQuadrant.isScRetract`, `Polyfold.LocalModel`).** $(O, C, E)$ is an *sc-smooth retract* if $O = r(U)$ for an sc-smooth retraction on $C$. A relatively open $U \subseteq C$ is a retract ($r = \mathrm{id}$). A *local model* is a triple $(E, C, O)$ with $C$ a partial quadrant and $O$ a retract in $C$. *Book:* Definition 2.3, p. 18. *Hidden hypotheses:* none.

**V2.3 (Proposition 2.4; `Polyfold.proposition_2_4`, `Polyfold.ScRetraction.comp_eq_self_of_retract_subse`, `Polyfold.ScRetraction.tangent_subset_of_retract_subset`, `Polyfold.ScRetraction.tangent`, `Polyfold.ScRetraction.mem_tangent_iff`).** For sc-smooth retractions $r$ on $U$ and $r'$ on $U'$, both relatively open in $C$, with $r(U) = r'(U')$: $Tr(TU) = Tr'(TU')$. If only $r(U) \subseteq r'(U')$, then $r' \circ r = r$ on $U$ and $Tr(TU) \subseteq Tr'(TU')$. A point $p$ lies in $Tr(TU)$ iff $p \in TU$ and $Tr(p) = p$. *Book:* Proposition 2.4, p. 19. *Hidden hypotheses:* the composition $r' \circ r$ is defined because $O \subseteq U'$. *Proof:* as in the text, by Theorem 1.21 applied to $r' \circ r = r$.

**V2.4 (Definition 2.5, the tangent of a retract; `Polyfold.retractTangent`, `Polyfold.retractTangent_eq`, `Polyfold.ScRetraction.mem_tangent_iff_exists`, `Polyfold.ScRetraction.mem_tangent_iff_mem_range`, `Polyfold.tangentSpace`, `Polyfold.tangentSpace_eq_range`, `Polyfold.tangentSpace_eq_coe_submodule`).** For a retract $O$, $TO \subseteq E_1 \times E_0$ is $Tr(TU)$ for any retraction $r$ onto $O$ (V2.3); $TO = \{(r(x), Dr(x)h) : x \in U_1, h \in E_0\}$, and for $x \in O_1$, $(x, h) \in TO$ iff $h$ is in the range of $Dr(x)$. The tangent space $T_xO = \{h : (x, h) \in TO\}$ is this range, a closed subspace of $E_0$. *Book:* Definition 2.5, p. 19, and Definition 2.17, p. 25 (the fibre $T_xO$). *Hidden hypotheses:* none. *Deviation (restated):* $TO$ is defined for every subset $O$ by a choice of retraction, and is empty if $O$ is not a retract.

**V2.5 (Proposition 2.6; `Polyfold.proposition_2_6_1`, `Polyfold.proposition_2_6_2`, `Polyfold.proposition_2`, `Polyfold.ScRetraction.comp_comp_eqOn`).** Let $r$ on $U$ and $s$ on $V$ be retractions onto the same $O$ and $f : E_0 \to E_0'$. (1) If $f \circ r$ is sc-smooth on $U$, then $f \circ s$ is sc-smooth on $V$. (2) Then $T(f \circ r) = T(f \circ s)$ on $Tr(TU)$. (3) If $f \circ r$ is sc-smooth and $f$ maps $O$ into the retract $O' = r'(U')$, then $T(f \circ r)$ maps $Tr(TU)$ into $Tr'(TU')$. *Book:* Proposition 2.6, p. 20. *Hidden hypotheses:* in (3), $f(O) \subseteq O'$ (contained in "$f : O \to O'$"). *Proof:* as in the text, with $f \circ r \circ s = f \circ s$ and $\rho \circ f \circ r = f \circ r$ and Theorem 1.21.

**V2.6 (Definition 2.7, sc-smooth maps between retracts; `Polyfold.IsScSmoothRetractMap`, `Polyfold.isScSmoothRetractMap_iff_exists`, `Polyfold.retractTangentMap`, `Polyfold.retractTangentMap_`, `Polyfold.IsScSmoothRetractMap.tangentMap_mapsTo`, `Polyfold.retractTangentMap_apply`, `Polyfold.IsScSmoothRetractMap.continuousOn`, `Polyfold.IsScSmoothRetractMap.mem_range_toBase`).** $f$ is *sc-smooth from* $(O, C, E)$ *to* $(O', C', E')$ if $f(O) \subseteq O'$ and $f \circ r$ is sc-smooth on $U$ for every sc-smooth retraction $r$ on $C$ with $r(U) = O$; for a retract $O$ this holds iff $f(O) \subseteq O'$ and $f \circ r$ is sc-smooth for one such $r$. The tangent map $Tf$ is $T(f \circ r)$ for a chosen $r$; on $TO$ it equals $T(f \circ r)$ for every $r$ onto $O$, maps $TO$ into $TO'$ when $O, O'$ are retracts, and

$Tf(x,h) = (f(x), D(f \circ r)(x)h)$. Such an $f$ is continuous on $O$ and maps points of $O$ on level $m$ to points on level $m$. *Book:* Definition 2.7, pp. 20–21. *Hidden hypotheses:* none. *Deviation (restated):* the definition quantifies over all retractions and contains $f(O) \subseteq O'$; the text's form is `Polyfold.isScSmoothRetractMap_iff_exists`.

**V2.7 (Theorem 2.8, the chain rule for retracts; `Polyfold.theorem_2_8`, `Polyfold.IsScSmoothRetractMap.comp`, `Polyfold.retractTangentMap_comp`, `Polyfold.isScSmoothRetractMap_id`, `Polyfold.ScRetraction.isScSmoo` `Polyfold.IsScSmoothRetractMap.congr`).** Let $(O,C,E)$ and $(O',C',E')$ be retracts, $f$ sc-smooth from $O$ to $O'$ and $g$ sc-smooth from $O'$ to $(O'',C'',E'')$. Then $g \circ f$ is sc-smooth from $O$ to $O''$ and $T(g \circ f) = Tg \circ Tf$ on $TO$. The identity is sc-smooth, a retraction is sc-smooth as a map $(U,C,E) \to (O,C,E)$, and sc-smoothness depends only on the values on $O$. *Book:* Theorem 2.8, p. 21, which writes $(O'',C',E'')$ for $(O'',C'',E'')$. *Hidden hypotheses:* $r' \circ f \circ r = f \circ r$, which needs $f(O) \subseteq O'$. *Proof:* the text gives none ("follows from Theorem 1.21"); $(g \circ f) \circ r = (g \circ r') \circ (f \circ r)$ on $U$ and Theorem 1.21.

**V2.8 (the tangent functor on retracts; `Polyfold.IsPartialQuadrant.tangentSet`, `Polyfold.IsRelOpenInQuadra` `Polyfold.ScRetraction.tangentRetraction`, `Polyfold.ScRetraction.tangentRetraction_retract`, `Polyfold.IsScRetract.tangentSet`, `Polyfold.LocalModel.tangent`, `Polyfold.IsScSmoothRetractMap.tange` `Polyfold.retractTangentMap_id`, `Polyfold.retractTangent_inter_isOpen`, `Polyfold.retractTangentMap_i`).** $TC$ is a partial quadrant in $TE$ and $TU$ is relatively open in $TC$; $Tr$ is an sc-smooth retraction on $TU$ with retract $TO$, so $(TO,TC,TE)$ is a retract. If $f$ is sc-smooth between the retracts $O$ and $O'$, then $Tf$ is sc-smooth from $(TO,TC,TE)$ to $(TO',TC',TE')$; $T$ of a map equal to the identity on $O$ is the identity on $TO$. For $W \subseteq E_0$ open: $T(O \cap W) = TO \cap (W_1 \times E_0)$, and $Tf$ computed on $(O \cap W, C, E)$ equals $Tf$ computed on $(O,C,E)$ there. *Book:* p. 21, after Theorem 2.8 ($T$ a functor $\mathcal{R} \to \mathcal{R}$), and p. 22, after Definition 2.9 ($Tr$ a bundle retraction). *Hidden hypotheses:* the locality of $Tf$ under restriction to open subsets, used in Definitions 2.17–2.18. *Deviation (restated):* the bundle structure (the category $\mathcal{BR}$) is not formalized.

**V2.9 (Proposition 2.10; `Polyfold.proposition_2_10_1`, `Polyfold.proposition_2_10_2`, `Polyfold.ScRetraction.restrict`, `Polyfold.ScRetraction.restrict_retract`, `Polyfold.IsScSmoothRetrac` `Polyfold.IsScSmoothRetractMap.of_locally`, `Polyfold.LocalModel.restrict`).** (1) If $(O,C,E)$ is a retract and $W \subseteq E_0$ is open, $(O \cap W, C, E)$ is a retract (with $r$ restricted to $U \cap r^{-1}(W)$). (2) If $(V,C,E)$ is a retract and $s$ is sc-smooth from $(V,C,E)$ to itself with $s \circ s = s$ on $V$, then $(s(V),C,E)$ is a retract, with the retraction $s \circ r$. Sc-smooth maps between retracts restrict to $O \cap W$, and are sc-smooth as soon as every point of $O$ has an open neighbourhood $W$ with $f$ sc-smooth on $O \cap W$. *Book:* Proposition 2.10, p. 23. *Hidden hypotheses:* the restriction of an sc-smooth map to a smaller relatively open set is sc-smooth (V2.63). *Deviation (restated):* an open subset of $O$ is written $O \cap W$.

**V2.10 (Definition 2.11, charts; `Polyfold.MChart`, `Polyfold.MChart.overlap`, `Polyfold.MChart.isScRetract_ov` `Polyfold.MChart.transition`, `Polyfold.MChart.Compatible`, `Polyfold.MChart.compatible_self`, `Polyfold.MChart.Compatible.restrict`, `Polyfold.MChart.compatible_of_locally`).** A *chart* on a space $X$ is a local model $(O,C,E)$ and a partial equivalence $\varphi$ between $X$ and $E_0$ with open source $V$ and target $O$, continuous on $V$, with inverse continuous on $O$. The overlap $\varphi(V \cap V')$ is a retract. Two charts are *compatible* if $\psi \circ \varphi^{-1}$ is sc-smooth from $(\varphi(V \cap V'),C,E)$ to $(\psi(V \cap V'),C',E')$ and $\varphi \circ \psi^{-1}$ back. A chart is compatible with itself, compatibility passes to restrictions, and two charts each compatible with every chart of a family covering $V \cap V'$ are compatible. *Book:* Definition 2.11, pp. 23–24. *Hidden hypotheses:* none. *Deviation (restated):* the chart is a partial equivalence; the last statement, the locality of compatibility, is not in the text and is what makes the maximal atlas (V2.12) work.

**V2.11 (Definition 2.12, atlases; `Polyfold.MAtlas`, `Polyfold.MAtlas.Equiv`, `Polyfold.MAtlas.equiv_iff`, `Polyfold.MAtlas.Equiv.refl`, `Polyfold.MAtlas.Equiv.symm`, `Polyfold.MAtlas.Equiv.trans`, `Polyfold.MAtlas.equivalence`).** An *atlas* is a set of pairwise compatible charts whose sources cover $X$; two atlases are *equivalent* if their union is an atlas, equivalently if every chart of one is compatible with every chart of the other; the relation is an equivalence relation. *Book:* Definition 2.12, p. 24. *Hidden hypotheses:* none; the transitivity of the equivalence, which "an equivalence class of atlases" (Definition 2.13) presupposes, is not discussed in the text and is proved from the locality of compatibility (Findings, Chapter 2, item 41).

**V2.12 (Definition 2.13, M-polyfolds; `Polyfold.MPolyfold`, `Polyfold.MPolyfold.IsChart`, `Polyfold.MPolyfold.IsChart.compatible`, `Polyfold.MPolyfold.IsChart.restrict`, `Polyfold.MPolyfold.is` `Polyfold.MPolyfold.exists_isChart_mem_source`, `Polyfold.LocalModel.toMPolyfold`).** An *M-polyfold* structure on a Hausdorff paracompact space $X$ is an atlas. The *charts of the structure* are all charts compatible with every chart of this atlas; any two are compatible, restrictions of charts are charts, every point lies in the source of one, and equivalent atlases give the same charts. A nonempty retract $O$ is an M-polyfold with the chart id. *Book:* Definition 2.13, p. 24. *Hidden hypotheses:* charts compatible with an atlas are compatible with each other (the text does not discuss it; it needs V2.10). *Deviation (restated):* one atlas represents its equivalence class.

**V2.13 (Definition 2.14, sc-smooth maps between M-polyfolds; `Polyfold.MPolyfold.IsScSmoothOn`, `Polyfold.MPolyfold.IsScSmoothMap`, `Polyfold.MPolyfold.IsScSmoothOn.of_exists_charts`, `Polyfold.MPolyfold.isScSmoothMap_id`, `Polyfold.MPolyfold.IsScSmoothOn.comp`, `Polyfold.MPolyfold.IsS`** $f : X \to Y$ is *sc-smooth on the open set* $U$ if it is continuous on $U$ and, for all charts $\varphi$ of $X$, $\psi$ of $Y$ and every open $V \subseteq U \cap \operatorname{dom}\varphi$ with $f(V) \subseteq \operatorname{dom}\psi$, the map $\psi \circ f \circ \varphi^{-1}$ is sc-smooth from $(\varphi(V), C, E)$ to $(O', C', E')$; $f$ is sc-smooth if it is sc-smooth on $X$. It suffices that every $x \in U$ has one such pair of charts and one such $V \ni x$. The identity is sc-smooth; if $f$ is sc-smooth on $U$ with $f(U) \subseteq U'$ and $g$ on $U'$, then $g \circ f$ is sc-smooth on $U$. *Book:* Definition 2.14, p. 24, and "the definition does not depend on the choice of charts". *Hidden hypotheses:* $f$ continuous (for charts with $f(V) \subseteq V'$ to exist around every point). *Deviation (hypotheses explicit):* continuity is part of the definition; maps on open subsets are included.

**V2.14 (the filtration $X_m$; `Polyfold.MPolyfold.levelSet`, `Polyfold.MPolyfold.mem_levelSet_iff`, `Polyfold.MPolyfold.levelSet_zero`, `Polyfold.MPolyfold.levelSet_succ_subset`, `Polyfold.MPolyfold.smo` `Polyfold.MPolyfold.levelTopology`, `Polyfold.MPolyfold.Level`, `Polyfold.MPolyfold.continuous_levelVa` `Polyfold.MPolyfold.isEmbedding_levelLift`).** $X_m$ is the set of points $x$ with $\varphi(x) \in \iota_m(E_m)$ for some chart $\varphi$ of the structure around $x$; then this holds for every chart around $x$. $X_0 = X$, $X_{m+1} \subseteq X_m$, $X_\infty = \bigcap_m X_m$. $X_m$ carries the topology generated by the sets $\{x \in V : \varphi(x) \in W\}$, $\varphi$ a chart, $W \subseteq E_m$ open (in terms of the lift of $\varphi(x)$ to $E_m$). The inclusion $X_m \to X$ is continuous, $X_m$ is Hausdorff (`Polyfold.MPolyfold.instT2SpaceLevel`), and on $V_m = V \cap X_m$ the lift of $\varphi$ to $E_m$ is a topological embedding. *Book:* p. 25, the paragraph after Definition 2.14. *Hidden hypotheses:* that the level is independent of the chart needs that transition maps are $\mathrm{sc}^0$ on the overlap retracts (the text: "of course"). *Deviation (restated):* the topology is the one generated by the text's family; that the family is a basis is not proved.

**V2.15 (Lemma 2.15; `Polyfold.lemma_2_15`, `Polyfold.MPolyfold.levelIncl`).** The inclusion $X_{m+1} \to X_m$ is continuous for every $m$. *Book:* Lemma 2.15, p. 25. *Hidden hypotheses:* none.

**V2.16 (the M-polyfold $X^1$; `Polyfold.ScRetraction.shift`, `Polyfold.ScRetraction.shift_retract`, `Polyfold.IsScRetract.shift`, `Polyfold.LocalModel.shift`, `Polyfold.IsScSmoothRetractMap.shift`, `Polyfold.MPolyfold.levelChart`, `Polyfold.MPolyfold.levelChart_compatible`, `Polyfold.MPolyfold.level`**

`Polyfold.MPolyfold.shift`, `Polyfold.MPolyfold.shift_isChart`, `Polyfold.MPolyfold.mem_shift_levelSet`
A retraction $r$ on $U$ induces an sc-smooth retraction on $U_1 \subseteq C_1 \subseteq E^1$ with retract $O_1$, so $(O_1, C_1, E^1)$ is a retract; an sc-smooth map between retracts induces an sc-smooth map between their level-1 parts (Proposition 1.17 for retracts). For every chart $\varphi$ of $X$, $(V_1, \varphi, (O_1, C_1, E^1))$ is a chart of $X_1$; any two are compatible, they form an atlas, and $X_1$ with it is an M-polyfold $X^1$. *Book:* p. 25 ("$(V_m, \varphi, (O_m, C_m, E_m))$ is a chart on $X_m$ … the collection of such charts is an atlas on $X_m$"). *Hidden hypotheses:* the paracompactness of $X_1$ (Theorem 2.16), an instance argument supplied by V2.17. The levels of $X^1$ are those of $X$ shifted by one, as sets: $(X^1)_m = X_{m+1}$. *Deviation (narrower):* $m = 1$ only; for $m \geq 2$ the iterate $(X^1)^1 \cdots$ has the points of $X_m$, but that its topology and charts are those of $X_m$ is not formalized (gap G.6; the set identity was added in a fourth addendum, Findings, Chapter 5, item 142).

**V2.17 (Theorem 2.16;** `Polyfold.theorem_2_16`, `Polyfold.MPolyfold.metrizableSpace`, `Polyfold.MPolyfold.metrizableSpace_level`, `Polyfold.MPolyfold.paracompactSpace_level`, `Polyfold.MPolyfold.SmoothLevel`, `Polyfold.MPolyfold.smoothLevel_topology_eq`, `Polyfold.MPolyfold.me` `Polyfold.MPolyfold.paracompactSpace_subtype`**).** Let $X$ be an M-polyfold. Then $X$ is metrizable; for every $m$, the level $X_m$ with the topology of V2.14 is metrizable and paracompact; and $X_\infty$, with the initial topology of the inclusions $X_\infty \to X_m$ ($m \geq 0$), is metrizable. Every subset of $X$ is paracompact. *Book:* Theorem 2.16, p. 25, proof pp. 57–58. *Hidden hypotheses:* the topology of $X_\infty$, which the text does not define. *Deviation (hypotheses explicit):* **no hypothesis is added.** *Proof:* not the text's. The text needs $X_m$ regular for Proposition 2.57 and does not check it. Here a partition of unity of the paracompact Hausdorff $X$ subordinate to the chart domains is pulled back to $X_m$ along the continuous inclusion; with the metrizable chart domains $V_m \cong O_m$ it gives an explicit metric (V2.62). Paracompactness of $X_m$ follows by Stone's theorem; $X_\infty$ embeds in $\prod_m X_m$.

**V2.18 (Definition 2.17, tangent vectors in charts;** `Polyfold.MPolyfold.TangentPre`, `Polyfold.MPolyfold.Tange` `Polyfold.MPolyfold.TangentPre.transport_self`, `Polyfold.MPolyfold.TangentPre.transport_mem`, `Polyfold.MPolyfold.TangentPre.transport_transport`, `Polyfold.MPolyfold.TangentPre.Rel`, `Polyfold.MPolyfold.TangentPre.Rel.symm`, `Polyfold.MPolyfold.TangentPre.Rel.trans`, `Polyfold.MPolyfold.TangentPre.setoid`**).** A tangent tuple is $(x, \varphi, h)$ with $x \in X_1$, $\varphi$ a chart of the structure around $x$ and $(\varphi(x), h) \in TO$. Its transport to a chart $\psi$ around $x$ is $T(\psi \circ \varphi^{-1})(\varphi(x), h)$, computed on the overlap retract; it lies in $TO'$ over $\psi(x)$, transport to $\varphi$ is the identity, and transport to $\psi$ followed by transport to $\chi$ is transport to $\chi$. Two tuples are equivalent if their points agree and the transport of the first to the chart of the second is the second; this is an equivalence relation. *Book:* Definition 2.17, pp. 25–26. *Hidden hypotheses:* that the relation is an equivalence relation (the text does not check it); it needs Theorem 2.8 on the triple overlap and the locality of $Tf$ (V2.8). *Deviation (restated):* none in substance.

**V2.19 (Definition 2.18, the tangent space $TX$;** `Polyfold.MPolyfold.TangentSpace`, `Polyfold.MPolyfold.tange` `Polyfold.MPolyfold.tangentFiber`, `Polyfold.MPolyfold.tangentChartSet`, `Polyfold.MPolyfold.tangentCh` `Polyfold.MPolyfold.tangentChartMap_mk_self`, `Polyfold.MPolyfold.tangentChartMap_injOn`, `Polyfold.MPolyfold.tangentChartMap_mk_ofVec`, `Polyfold.MPolyfold.tangentChartMap_mk_ofVec_of_mem`, `Polyfold.MPolyfold.tangentBasis`**).** $TX$ is the set of equivalence classes of tangent tuples, with $p : TX \to X_1$, $[(x, \varphi, h)] \mapsto x$, and $T_xX = p^{-1}(x)$. For a chart $\varphi$, $TV = p^{-1}(V_1)$ and $T\varphi : TV \to TO$ sends a class to its transport to $\varphi$; $T\varphi[(x, \varphi, h)] = (\varphi(x), h)$, $T\varphi$ is injective on $TV$ with inverse $a \mapsto [(\varphi^{-1}(a_1), \varphi, a_2)]$ on $TO$, and $T\psi \circ (T\varphi)^{-1} = T(\psi \circ \varphi^{-1})$ on $T(\varphi(V \cap V'))$. $TX$ carries the topology generated by the sets $\{\alpha \in TV : T\varphi(\alpha) \in W\}$, $W$ open. *Book:* Definition 2.18 and the paragraphs after it, pp. 26–27. *Hidden hypotheses:* none. *Deviation*

*(narrower):* the vector space, Banach and sc-Banach structures of $T_xX$ are not formalized.

**V2.20 (Proposition 2.19;** `Polyfold.MPolyfold.continuous_tangentProj`, `Polyfold.MPolyfold.continuous_le` `Polyfold.MPolyfold.isOpen_tangentChartSet`, `Polyfold.MPolyfold.metrizableSpace_tangentSpace`, `Polyfold.MPolyfold.paracompactSpace_tangentSpace`, `Polyfold.proposition_2_19_3`**).** (1) $TX$ is Hausdorff (the instance `Polyfold.MPolyfold.instT2SpaceTangentSpace`). (2) $p : TX \to X_1$ is continuous, and so is $TX \to X$; each $TV$ is open. (3) $TX$ is metrizable and paracompact. *Book:* Proposition 2.19, p. 27, proof pp. 58–60. *Hidden hypotheses:* none for (3). *Deviation (narrower):* the openness of $p$ in (2) is not proved; the basis property in (1) is not needed. *Proof:* (3) as for Theorem 2.16, pulling a partition of unity of $X$ back along $TX \to X_1 \to X$; the charts $TV \cong TO$ are metrizable.

**V2.21 ($TX$ as an M-polyfold;** `Polyfold.MPolyfold.tangentChart`, `Polyfold.MPolyfold.tangentChart_overla` `Polyfold.MPolyfold.tangentChart_transition`, `Polyfold.MPolyfold.tangentChart_compatible`, `Polyfold.MPolyfold.tangentAtlas`, `Polyfold.MPolyfold.tangentPolyfold`, `Polyfold.MPolyfold.isScSmoot` `Polyfold.MPolyfold.tangentProj_isScSmooth`**).** For a chart $\varphi$ of $X$, $(TV, T\varphi, (TO, TC, TE))$ is a chart of $TX$, with $T\varphi(TV \cap TV') = T(\varphi(V \cap V'))$ and transition map $T(\varphi' \circ \varphi^{-1})$ (formula (5)); these charts are pairwise compatible, form an atlas, and make $TX$ an M-polyfold. The projection $TO \to O_1$ is sc-smooth from $(TO, TC, TE)$ to $(O_1, C_1, E^1)$, and $p : TX \to X^1$ is sc-smooth. *Book:* p. 27, after Proposition 2.19. *Hidden hypotheses:* the paracompactness of $TX$ and $X_1$ (instance arguments, supplied by V2.20 and V2.17). *Deviation (narrower):* "sc-smooth M-polyfold bundle" is read as "$p$ is sc-smooth"; the fibrewise linear structure is not formalized.

**V2.22 (tangent maps of sc-smooth maps between M-polyfolds;** `Polyfold.MPolyfold.tangentMap`, `Polyfold.MPolyfold.tangentMap_mk`, `Polyfold.MPolyfold.levelVal_tangentProj_tangentMap`, `Polyfold.MPolyfold.tangentMap_mapsTo_fiber`, `Polyfold.MPolyfold.tangentChartMap_tangentMap`, `Polyfold.MPolyfold.tangentMap_id`, `Polyfold.MPolyfold.tangentMap_comp`, `Polyfold.MPolyfold.continuc` `Polyfold.MPolyfold.tangentMap_isScSmooth`, `Polyfold.MPolyfold.IsScSmoothMap.mapsTo_levelSet`**).** An sc-smooth $f : X \to Y$ maps $X_m$ into $Y_m$. Its tangent map $Tf : TX \to TY$, $[(x, \varphi, h)] \mapsto [(f(x), \psi, T(\psi \circ f \circ \varphi^{-1})(\varphi(x), h)_2)]$, is independent of the chart $\psi$ around $f(x)$ and of the representative; $p \circ Tf = f \circ p$, $Tf(T_xX) \subseteq T_{f(x)}Y$, $T\psi \circ Tf = T(\psi \circ f \circ \varphi^{-1}) \circ T\varphi$ in charts, $T(\mathrm{id}) = \mathrm{id}$, $T(g \circ f) = Tg \circ Tf$, and $Tf$ is continuous and sc-smooth $TX \to TY$. *Book:* not stated; the text uses $Tf$ for M-polyfolds from §2.3 on (Proposition 2.21(4), Definition 2.38). *Deviation (not in the book):* the construction the text leaves implicit.

**V2.23 (Definition 2.20, sub-M-polyfolds;** `Polyfold.MPolyfold.IsSubMPolyfold`, `Polyfold.MPolyfold.isSubMP` $A \subseteq X$ is a *sub-M-polyfold* if every $a \in A$ has an open $V \ni a$ and a map $r$ with $r(V) \subseteq V$, $r$ sc-smooth on $V$, $r \circ r = r$ on $V$ and $r(V) = A \cap V$. *Book:* Definition 2.20, p. 28. *Hidden hypotheses:* none.

**V2.24 (Proposition 2.21 (1), (2);** `Polyfold.MPolyfold.SubData`, `Polyfold.MPolyfold.SubData.shrink`, `Polyfold.MPolyfold.SubData.image_shrink`, `Polyfold.MPolyfold.SubData.rho_isScSmooth`, `Polyfold.MPolyfold.SubData.isScRetract_image`, `Polyfold.MPolyfold.subChart`, `Polyfold.MPolyfold.sub` `Polyfold.MPolyfold.subAtlas`, `Polyfold.MPolyfold.subPolyfold`, `Polyfold.MPolyfold.subPolyfold_inclu` `Polyfold.MPolyfold.subPolyfold_retraction_isScSmooth`, `Polyfold.IsScSmoothRetractMap.mono_source`, `Polyfold.MPolyfold.paracompactSpace_of_isClosed`**).** Let $A$ be a sub-M-polyfold with the subspace topology. For data $(V, r)$ at a point and a chart $\varphi$, the set $W = (V \cap \mathrm{dom}\,\varphi) \cap r^{-1}(V \cap \mathrm{dom}\,\varphi)$ is open with $r(W) \subseteq W$ and $r(W) = W \cap A$; $\rho = \varphi \circ r \circ \varphi^{-1}$ is sc-smooth and idempotent on $\varphi(W)$, so $\varphi(W \cap A)$ is a retract and $\varphi$ restricted to $W \cap A$ is a chart of $A$ with model $(\varphi(W \cap A), C, E)$. Any two such charts are compatible; they form an atlas, and $A$ is an M-

polyfold. (1) The inclusion $A \to X$ is sc-smooth. (2) For every data $(V, r)$, the map $i^{-1} \circ r$ is sc-smooth on $V$. A closed subset of $X$ is paracompact. *Book:* Proposition 2.21 (1), (2), pp. 28–30. *Hidden hypotheses:* the topology of $A$ (the subspace topology; then (1)'s homeomorphism is by construction) and the paracompactness of $A$, an instance argument of the structure, supplied for every $A$ by `Polyfold.MPolyfold.paracompactSpace_subtype` (V2.17). *Deviation (hypotheses explicit):* none otherwise.

**V2.25 (Proposition 2.21 (3), (4), in a chart; `Polyfold.ScRetraction.compRetraction`, `Polyfold.ScRetraction.compRetraction_retract`, `Polyfold.proposition_2_21_3`, `Polyfold.proposition_2`**
Let $r$ be a retraction onto $O$, $s$ sc-smooth from $(O, C, E)$ to itself with $s \circ s = s$, $t = s \circ r$ (a retraction onto $s(O)$) and $a \in s(O)_1$. If $a$ is smooth: $T_aO = Dr(a)E_0$, $T_as(O) = Dt(a)E_0 \subseteq T_aO$, the subspaces $Dt(a)E_0$ and $(Dr(a) - Dt(a))E_0$ are sc-subspaces of $E$, and on every level $m$ their level-$m$ parts are disjoint with sum the level-$m$ part of $T_aO$. For every $a \in s(O)_1$: $s$ is sc-smooth from $O$ to $s(O)$ and $Ts$ maps $T_aO$ onto $T_as(O)$. *Book:* Proposition 2.21 (3), (4), p. 30, proved in coordinates with "$T_aO = Ds(a)Dr(a)E \oplus (I - Ds(a))Dr(a)E$". *Hidden hypotheses:* $Ds(a)$ means $Dt(a)$; the splitting is an sc-splitting because $Dr(a)$, $Dt(a)$ are commuting sc-projections with $Dr(a)Dt(a) = Dt(a) = Dt(a)Dr(a)$. *Deviation (narrower):* in a chart only, as the text proves it; the transport to $T_aX$ is not formalized (no sc-structure on the fibres of $TX$).

**V2.26 (Definition 2.22, the degeneracy index; `Polyfold.MPolyfold.chartDegIndex`, `Polyfold.MPolyfold.degI`
`Polyfold.MPolyfold.degIndex_le`, `Polyfold.MPolyfold.exists_isChart_degIndex_eq`,
`Polyfold.MPolyfold.eventually_chartDegIndex_le`).** For a chart $\varphi$ around $x$, $d(x, \varphi) = d_C(\varphi(x))$; $d_X(x)$ is the least $d(x, \varphi)$ over the charts of the structure around $x$. The minimum is attained, $d_X(x) \le d(x, \varphi)$ for every such chart, and $d(\cdot, \varphi)$ is upper semicontinuous on the source of $\varphi$. *Book:* Definition 2.22, p. 31 ("smooth charts"). *Hidden hypotheses:* none. *Deviation (restated):* "smooth charts" are the charts of the maximal atlas.

**V2.27 (Lemma 2.23; `Polyfold.lemma_2_23`, `Polyfold.eventually_degIndexWith_le`, `Polyfold.MPolyfold.isO`
`Polyfold.MPolyfold.isClosed_setOf_le_degIndex`).** For every $x \in X$, $d_X(y) \le d_X(x)$ for all $y$ near $x$; hence $\{d_X < k\}$ is open and $\{d_X \ge k\}$ closed. The degeneracy index $d_C$ of a partial quadrant is upper semicontinuous. *Book:* Lemma 2.23, p. 31 ("evident"). *Hidden hypotheses:* none.

**V2.28 (Proposition 2.24; `Polyfold.MPolyfold.IsScDiffeoOn`, `Polyfold.proposition_2_24`,
`Polyfold.MPolyfold.transportChart`, `Polyfold.MPolyfold.transportChart_isChart`,
`Polyfold.MPolyfold.degIndex_le_of_isScDiffeoOn`).** Let $U \subseteq X$, $V \subseteq Y$ be open, $f$ sc-smooth on $U$ with $f(U) \subseteq V$ and $g$ sc-smooth on $V$ with $g(V) \subseteq U$, inverse to each other. Then $d_X(x) = d_Y(f(x))$ for every $x \in U$. For a chart $\psi$ of $Y$, the map $\psi \circ f$ on $U \cap f^{-1}(\operatorname{dom} \psi)$ is a chart of the structure of $X$. *Book:* Proposition 2.24, p. 31 ("from the definitions one deduces immediately"). *Hidden hypotheses:* none. *Deviation (restated):* a germ is represented by an sc-diffeomorphism of open sets.

**V2.29 (Definition 2.25, the boundary; `Polyfold.MPolyfold.boundary`, `Polyfold.MPolyfold.IsWithoutBoundar`
`Polyfold.MPolyfold.isWithoutBoundary_iff`, `Polyfold.MPolyfold.isClosed_boundary`,
`Polyfold.MPolyfold.isOpen_interior`, `Polyfold.MPolyfold.corners`, `Polyfold.MPolyfold.isClosed_corne`
`Polyfold.MPolyfold.IsScDiffeoOn.mem_boundary_iff`).** $\partial X = \{d_X \ge 1\}$; $X$ is without boundary iff $d_X \equiv 0$ iff $\partial X = \emptyset$. $\partial X$ and $\{d_X \ge 2\}$ are closed, $\{d_X = 0\}$ is open, and sc-diffeomorphisms of open sets preserve $\partial X$. *Book:* Definition 2.25, p. 31. *Hidden hypotheses:* none.

**V2.30 (Lemma 2.26; `Polyfold.lemma_2_26`).** For a sub-M-polyfold $A$ with the structure of V2.24, $d_A(a) \le d_X(a)$ for every $a \in A$. *Book:* Lemma 2.26, pp. 31–32. *Hidden hypotheses:* the paracom-

pactness of $A$, as in V2.24.

**V2.31 (the subspaces $E_x$ and the cones $C_x$; `Polyfold.minimalSubspaceWith`, `Polyfold.mem_minimalSubspaceWi` `Polyfold.isClosed_minimalSubspaceWith`, `Polyfold.finrank_quotient_minimalSubspaceWith`, `Polyfold.partialQuadrantAtWith`, `Polyfold.minimalSubspaceWith_subset_partialQuadrantAtWith`, `Polyfold.isClosed_partialQuadrantAtWith`, `Polyfold.exists_tendsto_lvl1_degIndexWith`, `Polyfold.exists_tendsto_smooth_degIndexWith`, `Polyfold.ScBanach.subset_closure_ker_inter_range_to` `Polyfold.partialQuadrantAtWith_subset_closure`).** Let $T : E \cong \mathbb{R}^n \oplus W$ present $C$. $E_x = \{v : (Tv)_i = 0 \text{ whenever } (Tx)_i = 0\}$ is a closed subspace of $E_0$ with $\dim(E_0/E_x) = d_C(x)$; $C_x = \{v : (Tv)_i \geq 0 \text{ whenever } (Tx)_i = 0\}$ is closed and convex and contains $E_x$. Every $x \in C$ is a limit of points of level 1 (and of smooth points) with the same index and in $C$. The level-1 points of the kernel of any continuous linear map into a finite-dimensional space are dense in it, and the level-1 points of $C_x$ are dense in $C_x$. *Book:* p. 32 ($E_i$, $E_I$, $I(x)$, $E_x$, $H_i$, $C_x$ in the model $\mathbb{R}^k \oplus W$). *Hidden hypotheses:* the densities, which the text uses without proof ("the closure of $(T_x^R O)_\infty$ is $T_x^R O$"). *Deviation (restated):* defined through a presentation of $C$.

**V2.32 (Definition 2.27; `Polyfold.reducedTangentSpaceWith`, `Polyfold.partialConeWith`, `Polyfold.quotDim`, `Polyfold.quotDim_inf_minimalSubspace_le`, `Polyfold.quotDim_reducedTangentSpaceWi`** For a retraction $r$, a presentation $T$ and $x \in O_1$: $T_x^R O = Dr(x)E_0 \cap E_x$ and $C_x O = Dr(x)E_0 \cap C_x$. For subspaces $A, B$, $\dim(A/(A \cap B))$ is the natural number $\mathrm{quotDim}(A, B)$; $\dim(T_x O / T_x^R O) \leq d_C(x)$. *Book:* Definition 2.27, p. 33. *Hidden hypotheses:* none. *Deviation (stronger):* defined at every $x \in O_1$, not only at smooth points.

**V2.33 (Remark 2.28; `Polyfold.reducedTangentSpaceWith_subset_partialConeWith`, `Polyfold.remark_2_28_` `Polyfold.remark_2_28_partialConeWith`, `Polyfold.flor_nonnegRange_simplicial`, `Polyfold.remark_2_28_` `Polyfold.ScRetraction.isScSubspace_reducedTangentSpaceWith`, `Polyfold.partialConeWith_eq_image_sc`** $T_x^R O \subseteq C_x O \subseteq T_x O$. Moreover, for every sc-smooth retraction $r$ in $C = T^{-1}([0,\infty)^n \oplus W)$ and every smooth $x \in O$ (no tameness), $T_x O$ and $T_x^R O$ are sc-subspaces of $E$ and $C_x O$ is a partial quadrant in the sc-Banach space $T_x O$ (with the sc-structure induced from $E$), of dimension $r = \operatorname{rank} M = \dim(T_x O / T_x^R O) \leq d_C(x)$, where $M$ is the matrix of $Dr(x)$ on $E/E_x \cong \mathbb{R}^{I(x)}$. The core is Flor's theorem in the needed case: for a linear $M$ on $\mathbb{R}^\iota$ with $M^2 = M$ mapping the orthant into itself, the cone $\operatorname{range} M \cap [0,\infty)^\iota$ is the set of nonnegative combinations of linearly independent columns of $M$. *Book:* Remark 2.28, p. 33: the inclusions, and "naively one might expect that $C_x O$ is a partial quadrant in $T_x O$. However, this is in general not the case", without an example. *Hidden hypotheses:* none. *Deviation (stronger):* **the remark's negative claim does not hold in our formal reading** (checked in Lean); the remark may mean the narrower point below. *Proof:* $Dr(x)$ maps $C_x$ and $E_x$ into themselves (the sign arguments of Lemma 2.29), so $C_x O = Dr(x)(C_x)$; modulo $T_x^R O$ it is the column cone of the nonnegative idempotent matrix $M$, simplicial by Flor's theorem (minimal column supports, generation by descent on the support, independence through a private index of each minimal support); the sc-isomorphism $T_x O \cong \mathbb{R}^r \oplus T_x^R O$ of the proof of Theorem 2.31 (V2.42) carries $C_x O$ onto $[0,\infty)^r \oplus T_x^R O$. What can fail for a non-tame retract is only $r = d_C(x)$ (Theorem 2.31), which the text proves for tame retracts (Proposition 2.36, V2.42) (Findings, Chapter 2, item 50; round-trip item V2.33).

**V2.34 (the sign arguments; `Polyfold.IsSc1With.deriv_mem_minimalSubspace_of_lvl1`, `Polyfold.IsSc1With.mapsTo_minimalSubspace`, `Polyfold.IsSc1With.deriv_mem_partialQuadrant_of_lvl1`, `Polyfold.IsSc1With.mapsTo_partialQuadrant`, `Polyfold.eventually_add_smul_mem_lvlSet`, `Polyfold.eventually_add_smul_mem_lvlSet_of_cone`, `Polyfold.ScRetraction.scDeriv_mapsTo_minimalSub`**

Let $f$ be sc¹ on $U$, relatively open in $C$ (presented by $T$), with values in $C'$ (presented by $T'$), and $x \in U_1$. For $v \in E_1$ with $\iota_1 v \in E_x$: $x + tv \in U_1$ for $|t|$ small and $Df(x)v \in E'_{f(x)}$; hence $Df(x)$ maps $E_x$ into $E'_{f(x)}$. For $v \in E_1$ with $\iota_1 v \in C_x$: $x + tv \in U_1$ for small $t \geq 0$ and $Df(x)v \in C'_{f(x)}$; hence $Df(x)$ maps $C_x$ into $C'_{f(x)}$. For a retraction $r$ and $x \in O_1$, $Dr(x)$ maps $E_x$ into $E_x$. *Book:* the proofs of Lemma 2.29, pp. 33–35, and Proposition 2.35, p. 40. *Deviation (not in the book):* the derivative-level content of Lemma 2.29, isolated; Lemma 2.29 itself, about sc-smooth paths, is not formalized. *Proof:* the function $t \mapsto (T'f(x+tv))_j$ is $\geq 0$ near 0 and vanishes at 0; its derivative at 0 vanishes (two-sided) or is $\geq 0$ (one-sided).

**V2.35 (Proposition 2.30; `Polyfold.ScRetraction.IsDiffeoGerm`, `Polyfold.IsScSmoothRetractMap.isDiffeoG` `Polyfold.ScRetraction.IsDiffeoGerm.deriv_inv_deriv`, `Polyfold.ScRetraction.IsDiffeoGerm.deriv_der` `Polyfold.proposition_2_30_reduced`, `Polyfold.proposition_2_30_cone`).** A *germ of sc-diffeomorphisms* between the retracts $O = r(U)$ and $O' = r'(U')$ is a pair $F, G$ with $F \circ r$, $G \circ r'$ sc¹, $F(O) \subseteq O'$, $G(O') \subseteq O$, $G \circ F =$ id on $O$ and $F \circ G =$ id on $O'$; an sc-smooth map between retracts with an sc-smooth inverse is one. For $x \in O_1$ and $y = (F \circ r)_1(x)$: $D(G \circ r')(y) \circ D(F \circ r)(x) = Dr(x)$, $D(F \circ r)(x) \circ D(G \circ r')(y) = Dr'(y)$, and, with $Tf(x) = D(F \circ r)(x)$, $Tf(x)(T^R_x O) = T^R_y O'$ and $Tf(x)(C_x O) = C_y O'$. *Book:* Proposition 2.30, p. 35 (smooth $x$, sc-smooth germs; deduced from Lemma 2.29). *Hidden hypotheses:* none. *Deviation (stronger):* level-1 points and sc¹ germs. *Proof:* from the sign arguments (V2.34) and the two derivative identities, not from Lemma 2.29.

**V2.36 (a retract as an M-polyfold; `Polyfold.LocalModel.selfChart`, `Polyfold.LocalModel.toMPolyfold`, `Polyfold.IsRelOpenInQuadrant.toMPolyfold`, `Polyfold.IsRelOpenInQuadrant.selfChart_isChart`, `Polyfold.MChart.Compatible.isDiffeoGerm`).** A nonempty retract $O$, and a nonempty relatively open $U \subseteq C$, is an M-polyfold on the subtype with the single chart id. Two compatible charts give a germ of sc-diffeomorphisms (in the sense of V2.35) between the overlap retracts, with the restricted retractions. *Book:* p. 35, before Theorem 2.31. *Deviation (restated):* the text's $d_O(x) = \min d_{C'}(\varphi(x))$ over germs into retracts is Definition 2.22 for this structure.

**V2.37 (Theorem 2.31, first statement; `Polyfold.theorem_2_31`, `Polyfold.theorem_2_31_retract`, `Polyfold.ScRetraction.IsDiffeoGerm.quotDim_le`, `Polyfold.MPolyfold.quotDim_le_chartDegIndex`, `Polyfold.ScRetraction.scDeriv_restrict`, `Polyfold.degIndexWith_le_of_germ`, `Polyfold.remark_2_28_is`** Let $X$ be an M-polyfold, $\varphi$ a chart of the structure around $x$ with $\varphi(x) = \iota_1 z$ on level 1, $r$ a retraction onto the model $O$ and $T$ a presentation of $C$. Then $\dim(T_z O / T^R_z O) \leq d_X(x)$. For a retract $O$ itself as an M-polyfold and $x \in O_1$: $\dim(T_x O / T^R_x O) \leq d_O(x)$. Along a germ of sc-diffeomorphisms $F$ between retracts, $\dim(T_x O / T^R_x O) \leq d_{C'}(F(x))$. *Book:* Theorem 2.31, p. 35 (a retract, a smooth $x$). *Hidden hypotheses:* none. *Deviation (stronger):* every M-polyfold, every chart, points on level 1. The second statement of the theorem (equality makes $C_x O$ a partial quadrant) holds without its hypothesis: $C_x O$ is a partial quadrant at every smooth point of every retract (V2.33). *Proof:* for every chart $\psi$ around $x$, $D(\psi \circ \varphi^{-1})$ induces $T_z O / T^R_z O \cong T_y O' / T^R_y O'$, which embeds in $E'/E'_y$ of dimension $d_{C'}(\psi(x))$; the text instead reduces to a chart realizing $d_O(x)$.

**V2.38 (Corollary 2.32; `Polyfold.corollary_2_32`, `Polyfold.IsRelOpenInQuadrant.degIndex_le`, `Polyfold.IsRelOpenInQuadrant.degIndex_le_degIndex_of_lvl1`, `Polyfold.degIndexWith_le_of_germ`).** For a nonempty relatively open subset $U$ of a partial quadrant $C$, viewed as the M-polyfold $X_U$, $d_{X_U}(x) = d_C(x)$ for every $x \in U$. *Book:* Corollary 2.32, p. 37 ($U = C$). *Hidden hypotheses:* the semicontinuity of $d_{C'}$ and Proposition 2.24 at the approximating points. *Deviation (stronger):* every relatively open $U$; level-1 approximants suffice. Theorem 1.25 (V1.23, `Polyfold.theorem_1_25`)

is proved in the same file by the same argument.

**V2.39 (Definition 2.33; `Polyfold.partialQuadrantAtWith`, `Polyfold.minimalSubspaceWith`, `Polyfold.minimalSubspaceWith_subset_partialQuadrantAtWith`, `Polyfold.convex_partialQuadrantAtWith`, `Polyfold.finrank_quotient_minimalSubspaceWith`, `Polyfold.minimalSubspaceWith_le`, `Polyfold.minimalSubspaceWith_eq`).** For a presentation $T$ of $C$ and $e \in C$: the partial quadrant $C_e$ and the minimal subspace $E_e$ of V2.31, with $E_e \subseteq C_e$, $C_e$ convex and $\dim(E_0/E_e) = d_C(e)$. For two presentations $T, T'$ of the same $C$, the two subspaces $E_e$ agree. *Book:* Definition 2.33, p. 39, through the extended faces. *Hidden hypotheses:* none. *Deviation (restated):* through a presentation, of which $E_e$ is independent (the argument of Lemma 1.24); the intrinsic description by extended faces (p. 39) is not formalized.

**V2.40 (Definition 2.34, tame retractions; `Polyfold.ScRetraction.IsTameWith`, `Polyfold.ScRetraction.IsTame`, `Polyfold.IsTameRetract`, `Polyfold.IsTameRetract.isScRetract`, `Polyfold.ScRetraction.id_isTame`, `Polyfold.ScRetraction.IsTameWith.isTame`, `Polyfold.ScRetraction.IsTameWith.restrict`).** A retraction $r$ is *tame for* $T$ if (1) $d_C(r(x)) = d_C(x)$ for $x \in U$ and (2) for every smooth $x \in O_1$ there is a subspace $A \leq E_x$ such that $E = Dr(x)E_0 \oplus A$ is an sc-splitting. $r$ is *tame* if it is tame for every presentation of $C$; tameness for one presentation implies it. A *tame retract* is the retract of a tame retraction. Identities of relatively open subsets are tame, and the restriction of a tame retraction to $U \cap r^{-1}(V)$, $V$ open, is tame. *Book:* Definition 2.34, pp. 39–40. *Hidden hypotheses:* that the definition does not depend on the presentation, which the text uses tacitly. *Deviation (restated):* none in substance.

**V2.41 (Proposition 2.35; `Polyfold.proposition_2_35`, `Polyfold.ScRetraction.scDeriv_mapsTo_minimalSubs`, `Polyfold.ScRetraction.isScOperator_scDeriv`, `Polyfold.ScRetraction.isScSplitting_range_scDeriv`, `Polyfold.coord_scDeriv_eq`).** Let $r$ be tame for $T$ and $x \in O_1$ smooth. Then $(1 - Dr(x))E_0 \leq E_x$, and $E = Dr(x)E_0 \oplus (1 - Dr(x))E_0$ is an sc-splitting; equivalently $\lambda_j(Dr(x)v) = \lambda_j(v)$ for $j \in I(x)$ and all $v$. *Book:* Proposition 2.35, p. 41, with the discussion on p. 40. *Hidden hypotheses:* none. *Deviation (stronger):* only condition (2) of tameness is used; $Dr(x)E_x \subseteq E_x$ holds for every retraction (V2.34), where the text uses condition (1).

**V2.42 (Proposition 2.36; `Polyfold.proposition_2_36`, `Polyfold.proposition_2_36_degIndex`, `Polyfold.quotDim_reducedTangentSpaceWith_eq`, `Polyfold.MPolyfold.degIndex_eq_chartDegIndex_of_isT`, `Polyfold.MPolyfold.degIndex_eq_chartDegIndex_of_isTameRetract`, `Polyfold.LocalModel.degIndex_toMP`, `Polyfold.quotDim_eq_finrank_quotient_of_isCompl`, `Polyfold.proposition_2_36_partialCone`, `Polyfold.splitIso`, `Polyfold.isPartialQuadrant_ofSubspace`).** Let $r$ be tame for $T$. (13) At a smooth $x \in O_1$, $\dim(T_xO/T_x^RO) = d_O(x) = d_C(x)$, where $d_O$ is the index of the M-polyfold $O$; moreover $T_xO$ and $T_x^RO$ are sc-subspaces of $E$, and $C_xO = T_xO \cap C_x$ is a partial quadrant in the sc-Banach space $T_xO$ (with the sc-structure induced from $E$). (14) $d_O(x) = d_C(x)$ for every $x \in O$. More generally, a chart $\varphi$ of any M-polyfold whose model is a tame retract satisfies $d_X(y) = d_C(\varphi(y))$ for every $y$ in its source. For sc-subspaces $B \leq A$ with $B = A \cap \bigcap_j \ker \ell_j$ and smooth $e_j \in A$ with $\ell_i(e_j) = \delta_{ij}$, the map $v \mapsto ((\ell_j v)_j, v - \sum_j \ell_j(v)e_j)$ is an sc-isomorphism $A \cong \mathbb{R}^d \oplus B$, and $\{v \in A : \ell_j(v) \geq 0 \ \forall j\}$ is a partial quadrant in $A$. *Book:* Proposition 2.36, p. 41. *Hidden hypotheses:* none. *Deviation (stronger):* the chart form, which Proposition 2.42 needs; $T_x^RO$ is an sc-subspace, where the text says a Banach space. *Proof:* (13), (14) as in the text: $E_x = T_x^RO \oplus A$ gives $\dim(T_xO/T_x^RO) = d_C(x)$, Theorem 2.31 gives $\leq d_X$; a general point is approximated by smooth points $r(T^{-1}(a, w_j))$ of the same index, with Lemma 2.23. The partial quadrant not from the equality case of Theorem 2.31, as in the text, but directly (it holds without tameness, V2.33; what tameness adds is the dimension $d_O(x)$ of the quadrant): the smooth

vectors $\hat{e}_j = Dr(x)T^{-1}(e_{i_j}, 0)$ satisfy $\lambda_{i_k}(\hat{e}_j) = \delta_{jk}$ by Proposition 2.35, and the splitting applies to $A = T_xO$, $B = T^R_xO$; $T^R_xO$ is an sc-subspace because the smooth points $Dr(x)T^{-1}(a, w')$, $w' \in W_\infty$, lie in it ($Dr(x)E_x \subseteq E_x$) and approximate $u$, $Tu = (a, w)$, on every level.

**V2.43 (Definition 2.37; `Polyfold.MPolyfold.IsTame`, `Polyfold.MPolyfold.isChart_of_mem_equiv`, `Polyfold.MPolyfold.IsTame.exists_chart`).** $X$ is *tame* if some atlas equivalent to its atlas consists of charts with tame models; then every point lies in a chart of the structure whose model is a tame retract. *Book:* Definition 2.37, p. 42. *Hidden hypotheses:* none.

**V2.44 (Definition 2.39; `Polyfold.MPolyfold.IsFace`, `Polyfold.MPolyfold.IsFaceStructured`, `Polyfold.MPolyfold.IsFace.isClosed`, `Polyfold.MPolyfold.IsFace.subset_boundary`).** A *face* is the closure of the connected component in $\{d_X = 1\}$ of a point with $d_X = 1$; $X$ is *face-structured* if every $x$ lies in exactly $d_X(x)$ faces. Faces are closed and contained in $\partial X$. *Book:* Definition 2.39, p. 43. *Hidden hypotheses:* none.

**V2.45 (the local geometry of faces; `Polyfold.subset_coordHyper_of_isPreconnected`, `Polyfold.IsSc1With.apply_deriv_eq_zero_of_ray`, `Polyfold.ray_zeroSet`, `Polyfold.local_face_structur` `Polyfold.eventually_zeroSet_subset`, `Polyfold.convex_ball_inter_zeroSet_eq`, `Polyfold.ScRetraction.` `Polyfold.coordHyper`).** With $\lambda_i = (T\,\cdot)_i$, $I(z) = \{i : \lambda_i(z) = 0\}$ and $f_i = \{\lambda_i = 0\}$: (a) a preconnected subset of $\{z \in C : d_C(z) = 1\}$ containing a point of $f_i$ lies in $f_i$; (b) if $r$ is tame for $T$, $x \in O_1$ smooth, $i \in I(x)$, $\lambda_i(y) = 0$ and $\lambda_l(y) > 0$ for the other $l \in I(x)$, then for all small $\tau > 0$, $x + \tau y \in U$, $I(x + \tau y) = \{i\}$ and $I(r(x + \tau y)) = \{i\}$; (c) for $e \in O$ with $\lambda_i(e) = 0$ there are $\varepsilon > 0$ and $p \in S = \{z \in O : d_C(z) = 1\}$ with $\lambda_i(p) = 0$ such that $B = B_\varepsilon(e) \cap C \subseteq U$, $I(z) \subseteq I(e)$ and $I(r(z)) \subseteq I(e)$ on $B$, $r$ maps the points of $B$ with $I = \{i\}$ into the component of $S$ through $p$, and maps $f_i \cap B$ into its closure. *Book:* the proof of Lemma 2.40, p. 43 (a sketch). *Deviation (not in the book):* the exposition's Lemmas 4.A–4.C; (b) uses tameness condition (2) through Proposition 2.35.

**V2.46 (Lemma 2.40; `Polyfold.ScRetraction.IsRetractFace`, `Polyfold.lemma_2_40`, `Polyfold.lemma_2_40_in` `Polyfold.closure_connectedComponentIn_faceStratum`, `Polyfold.connectedComponentIn_coordHyper_eq`, `Polyfold.LocalModel.isFace_toMPolyfold_iff`, `Polyfold.subset_of_relClopen_of_isPreconnected`).** Let $r$ be tame for $T$. A *face of the retract* is $\overline{\widehat{F}} \cap O$ for a connected component $\widehat{F}$ of $S$. Then $G$ is a face of the retract iff $G$ is the connected component of $f_i \cap O$ through some $e \in O$ with $\lambda_i(e) = 0$; and for $x \in O$ with $\lambda_i(x) = 0$, equal components of $f_i \cap O$ and $f_j \cap O$ through $x$ force $i = j$. The faces of the M-polyfold $O$ are exactly the faces of the retract. *Book:* Lemma 2.40, p. 43. *Hidden hypotheses:* the face preservation of V2.45, which the text's sketch does not address. *Deviation (hypotheses explicit):* extended faces are the coordinate hyperplanes of a presentation.

**V2.47 (Corollary 2.41; `Polyfold.corollary_2_41`, `Polyfold.corollary_2_41_retract`, `Polyfold.LocalModel.isFaceStructured_toMPolyfold`).** For a nonempty tame retract $O$, the M-polyfold $O$ is face-structured: every $x \in O$ lies in exactly $d_O(x)$ faces. *Book:* Corollary 2.41, p. 44. *Hidden hypotheses:* none.

**V2.48 (Proposition 2.42; `Polyfold.proposition_2_42`, `Polyfold.MPolyfold.faces_encard_le`, `Polyfold.MPolyfold.IsTame.exists_chart`).** Let $X$ be tame and $x \in X$. There is a chart $\varphi : V \to O$ of the structure around $x$ such that for all $y \in V$, $d_X(y) = d_C(\varphi(y))$ and $d_O(\varphi(y)) = d_X(y)$, and the M-polyfold $O$ is face-structured. Globally, $x$ lies in at most $d_X(x)$ faces of $X$. *Book:* Proposition 2.42, p. 44. *Hidden hypotheses:* "$d_V = d_X$" needs that a tame chart realizes $d_X$ (V2.42). *Deviation (restated):* the neighbourhood $V$ is represented by the M-polyfold $O$.

**V2.49 (Proposition 2.43;** `Polyfold.proposition_2_43`, `Polyfold.killCoord`, `Polyfold.killCoord_mem`, `Polyfold.killCoord_eq_self`, `Polyfold.LocalModel.isSubMPolyfold_of_isFace`, `Polyfold.faceIncl`, `Polyfold.faceProj`, `Polyfold.degIndexWith_eq_quadrantDeg_faceProj_add_one`, `Polyfold.proposition_2_`
`Polyfold.MPolyfold.faceChart`, `Polyfold.MPolyfold.faceChart_compatible_subChart`, `Polyfold.LocalModel.proposition_2_43_isTame`**).** Let $r$ be tame for $T$ and $F$ a face of the retract. (a) $F \subseteq f_i$ for some $i$, and there is an sc-smooth retraction $s$ on a set $U^*$ with $F \subseteq U^* \subseteq U$, $U^*$ relatively open in $C$, with retract $F$ and $s = r \circ P_i$ on $U^*$, where $P_i$ is the sc-projection onto $f_i$ killing the $i$-th coordinate; every face of the M-polyfold $O$ is a sub-M-polyfold of $O$. (b) With the sc-operators $\iota : \mathbb{R}^k \oplus W \to E$ (insert a zero $i$-th coordinate, onto $f_i$) and $\pi : E \to \mathbb{R}^k \oplus W$ (delete it), $\pi \circ \iota = 1$, $\iota \circ \pi = 1$ on $f_i$ and $d_C(y) = d_{C'}(\pi y) + 1$ on $f_i$ ($C' = [0,\infty)^k \oplus W$); $r' = \pi \circ r \circ \iota$ is a tame retraction of $C'$ onto $\pi(F)$, and $d_{C'}(\pi y) + 1 = d_C(y)$ for $y \in F$. (c) For a face $F$ of a nonempty tame retract $O$, with its sub-M-polyfold structure, $F$ is a tame M-polyfold and $d_F(x) + 1 = d_O(x)$ for $x \in F$; the chart $\pi : F \to \pi(F)$ is compatible with the sub-charts of Proposition 2.21. *Book:* Proposition 2.43, pp. 44–45. *Hidden hypotheses:* the face preservation of V2.45; the paracompactness of $F$ (faces are closed). *Deviation (restated):* the tame model lives on $\mathbb{R}^k \oplus W$ through $\iota$, $\pi$ instead of the sc-subspace $f_1$; tameness (2) of $r'$ is proved with the complement $(1 - Dr'(x))E'$ by Proposition 2.35, not by intersecting with $f_1$. $U^*$ is a union of balls intersected with their $P_i$-preimages, since $P_i$ need not be norm-decreasing. *No hypothesis is added.*

**V2.50 (Proposition 2.44;** `Polyfold.MPolyfold.proposition_2_44`, `Polyfold.proposition_2_42`, `Polyfold.LocalModel.proposition_2_43_isTame`, `Polyfold.MPolyfold.proposition_2_44_corner`, `Polyfold.corner_retraction`, `Polyfold.corner_tameModel`, `Polyfold.cornerIncl`, `Polyfold.cornerProj`, `Polyfold.cornerKill`, `Polyfold.degIndexWith_eq_quadrantDeg_cornerProj_add_card`**).** Let $X$ be tame and $x \in X$. There is a chart $\varphi : V \to O$ of the structure around $x$, with a tame retraction $r$ onto $O$ (tame for a presentation $T$), such that $\varphi(x)$ lies in exactly $d_X(x)$ faces of the M-polyfold $O$ (the local faces), and every face $F$ of $O$ is a sub-M-polyfold of $O$ which, with the induced structure, is tame with $d_F(\varphi(x)) + 1 = d_X(x)$ when $\varphi(x) \in F$. For every $\sigma \subseteq I(\varphi(x))$ (a set of $\#\sigma$ face germs at $x$) there are pairwise distinct faces $F_i \ni \varphi(x)$ of the retract, $F_i \subseteq f_i$ ($i \in \sigma$), and an open $B \ni \varphi(x)$ such that $A = \{y \in V : \varphi(y) \in B,\ \varphi(y) \in F_i\ \forall i \in \sigma\}$ is a sub-M-polyfold of $X$, tame with its induced structure, with $d_A(y) + \#\sigma = d_X(y)$ for all $y \in A$. The ingredients: the corner $f_\sigma = \bigcap_{i\in\sigma} f_i$ is modelled on $\mathbb{R}^{n-\#\sigma} \oplus W$ through sc-operators $\iota_\sigma$, $\pi_\sigma$, with $d_C(y) = d_{C'}(\pi_\sigma y) + \#\sigma$ on $f_\sigma$; near a point $o \in O$ with $\sigma \subseteq I(o)$, $r \circ P_\sigma$ ($P_\sigma = \iota_\sigma \pi_\sigma$) is an sc-smooth retraction onto $O \cap V \cap f_\sigma = O \cap V \cap \bigcap_{i\in\sigma} F_i$; and $\pi_\sigma \circ r \circ \iota_\sigma$ is a tame retraction of $[0,\infty)^{n-\#\sigma} \oplus W$ onto its image under $\pi_\sigma$. *Book:* Proposition 2.44, p. 46 (stated without proof). *Hidden hypotheses:* none. *Deviation (restated):* local faces are the faces of the M-polyfold $O \cong V$, face germs at $x$ are the faces $F_i$, $i \in I(\varphi(x))$, "the germ $F_x^\sigma$" is represented by the set $A$, and "codimension $\#\sigma$" is the number of quadrant coordinates removed in the tame model; the index identity holds on all of $A$. *Proof:* not by the exposition's induction on $\#\sigma$ but at once: Lemma 4.C for each $i \in \sigma$ shows that $r$ maps points of $C \cap f_\sigma$ near $\varphi(x)$ into $f_\sigma$; then as in Proposition 2.43 and Theorem 2.45.

**V2.51 (Theorem 2.45;** `Polyfold.MPolyfold.theorem_2_45_interior`, `Polyfold.MPolyfold.isSubMPolyfold_o`
`Polyfold.MPolyfold.theorem_2_45_isSubMPolyfold_interior`, `Polyfold.MPolyfold.exists_localFace_of_`
`Polyfold.MPolyfold.theorem_2_45_isSubMPolyfold_face`, `Polyfold.MPolyfold.theorem_2_45_isTame_face`
`Polyfold.MPolyfold.isTame_of_forall_exists_chart`**).** Let $X$ be tame, $G$ the connected component of $\{d_X = 1\}$ through a point and $F = \overline{G}$ the face. (a) $F \cap \{d_X = 1\} = G$, and $G$ is a sub-M-polyfold of $X$. (b) If $X$ is face-structured, every face $F$ agrees near each of its points, in a

tame chart, with a face of the model retract; $F$ is a sub-M-polyfold of $X$, tame with its induced structure, and $d_X(x) = d_F(x) + 1$ for $x \in F$. (A subset $A$ is a sub-M-polyfold as soon as near each point $\varphi(A)$ agrees on an open set with an sc-smooth retract contained in $O$; $X$ is tame as soon as every point lies in a chart of the structure with a tame model.) *Book:* Theorem 2.45, p. 46 (stated without proof). *Hidden hypotheses:* the meaning of "the interior of a face", here $F \cap \{d_X = 1\}$. *Deviation (hypotheses explicit):* the counting argument behind "each global face through $x$ contains exactly one local face" (onto by Lemma 4.C, injective by face-structuredness) is made explicit. The theorem's face counts are V2.48 and the definition of face-structured (V2.44).

**V2.52 (products of M-polyfolds, exposition Lemma 4.D;** `Polyfold.IsPartialQuadrant.prod`, `Polyfold.prod_eq_prodPresentation`, `Polyfold.degIndex_prod`, `Polyfold.minimalSubspaceWith_prodPrese` `Polyfold.ScRetraction.prod`, `Polyfold.IsScRetract.prod`, `Polyfold.ScRetraction.IsTameWith.prod`, `Polyfold.MChart.prod`, `Polyfold.MChart.Compatible.prod`, `Polyfold.MPolyfold.paracompactSpace_prod`, `Polyfold.MPolyfold.prod`, `Polyfold.MPolyfold.degIndex_prod_le`, `Polyfold.MPolyfold.isTame_prod`, `Polyfold.MPolyfold.isScSmoothMap_fst`, `Polyfold.MPolyfold.isScSmoothMap_snd`, `Polyfold.MPolyfold.Is` For partial quadrants $C \subseteq E$, $C' \subseteq E'$, $C \times C'$ is a partial quadrant of $E \oplus E'$, presented by $E \oplus E' \cong \mathbb{R}^{n+n'} \oplus (W \oplus W')$, with $d_{C\times C'}(x, x') = d_C(x) + d_{C'}(x')$ and minimal subspace $E_x \oplus E'_{x'}$; products of sc-smooth (tame) retractions are sc-smooth (tame) retractions. For M-polyfolds $X$, $Y$ the space $X \times Y$ is paracompact and the product charts are an atlas of an M-polyfold structure; $d_{X\times Y}(x, y) \le d_X(x) + d_Y(y)$, and if $X$, $Y$ are tame, $X \times Y$ is tame with equality. The projections are sc-smooth and the pairing of sc-smooth maps is sc-smooth. *Book:* used without statement in the proof of Proposition 2.46, p. 47 ("$d_{X\times Y} = d_X + d_Y$"). *Hidden hypotheses:* the additivity needs tameness; in general only the inequality holds. Paracompactness of $X \times Y$ is Theorem 2.16. *Deviation (extra):* not stated in the text; the exposition's Lemma 4.D.

**V2.53 (Proposition 2.46;** `Polyfold.MPolyfold.IsLocalScDiffeo`, `Polyfold.MPolyfold.fibredProduct`, `Polyfold.MPolyfold.fibredSubDataL`, `Polyfold.MPolyfold.isSubMPolyfold_fibredProduct`, `Polyfold.MPolyfold.isScDiffeoOn_snd_fibredProduct`, `Polyfold.MPolyfold.isScDiffeoOn_fst_fibredPro` `Polyfold.MPolyfold.degIndex_fibredProduct_eq_snd`, `Polyfold.MPolyfold.degIndex_fibredProduct_eq_f` `Polyfold.MPolyfold.isTame_fibredProduct`, `Polyfold.MPolyfold.proposition_2_46`, `Polyfold.MPolyfold.proposition_2_46_of_isLocalScDiffeo`**).** Let $f : X \to Z$ be a local sc-diffeomorphism (every point has an open neighbourhood that $f$ maps sc-diffeomorphically onto an open subset of $Z$) and $g : Y \to Z$ sc-smooth. Then $X_f \times_g Y = \{(x, y) : f(x) = g(y)\}$ is closed in $X \times Y$, hence paracompact, and a sub-M-polyfold of $X \times Y$, near each point the retract of $R(x', y') = ((f|_U)^{-1}(g(y')), y')$; $d_{X_f\times_g Y}(x, y) = d_Y(y)$; if $Y$ is tame, so is the fibred product. If $g$ is also a local sc-diffeomorphism, then $d_{X_f\times_g Y}(x, y) = d_X(x) = d_Y(y)$, and if moreover $X$ and $Y$ are tame, $2\, d_{X_f\times_g Y}(x, y) = d_{X\times Y}(x, y)$. *Book:* Proposition 2.46, pp. 46–47. *Hidden hypotheses:* the meaning of "local sc-diffeomorphism"; the paracompactness of the fibred product. The text's $\pi_2 : X_f \times_g Y \to X$ has target $Y$. *Deviation (stronger):* $d = d_Y$, and $d = d_X = d_Y$ in the last statement, hold without tameness (Proposition 2.24 along $\pi_2$, $\pi_1$); tameness is used only for $d_{X\times Y} = d_X + d_Y$ (V2.52). The inverse of $\pi_2$ near $(x, y)$ is $y' \mapsto ((f|_U)^{-1}(g(y')), y')$; the exposition's factorization through the constant-in-$x$ map $y' \mapsto (x, y')$ is not sc-smooth unless $x$ is a smooth point. *No hypothesis is added.*

**V2.54 (Definition 2.47;** `Polyfold.doubleLevel`, `Polyfold.shiftFiber`, `Polyfold.IsStrongBundleMap`, `Polyfold.IsStrongBundleRetraction`, `Polyfold.IsPartialQuadrant.prod_univ`, `Polyfold.IsRelOpenInQuad` `Polyfold.IsStrongBundleRetraction.baseRetraction`, `Polyfold.IsStrongBundleRetraction.fiber_idem`, `Polyfold.IsStrongBundleRetraction.toScRetraction`, `Polyfold.IsStrongBundleRetraction.toScRetracti`

$C \times F_0$ is a partial quadrant in $E \oplus F$ and $U \times F_0$ is relatively open in it. A *strong bundle map* on $U \lhd F$ is $\Phi : E_0 \times F_0 \to E_0' \times F_0'$ mapping $U \times F_0$ into $U' \times F_0'$, whose first component does not depend on $h$ and second is linear in $h$ for $u \in U$, mapping $U_m \oplus F_k$ into $U_m' \oplus F_k'$ for $k \le m+1$, sc-smooth on $U \times F_0$ for $E \oplus F$, and whose induced map on $U \times F_1$ (fibre component lifted to $F_1'$) is sc-smooth for $E \oplus F^1$. A *strong bundle retraction* is one with $R \circ R = R$ on $U \times F_0$. Then $r(u) = R(u,0)_1$ is an sc-smooth retraction on $U$, $\Gamma(u,\cdot)$ is a projection for $u \in O$, and $R$ is an sc-smooth retraction on $(U \lhd F)[0]$ and $[1]$. *Book:* Definition 2.47 and the remarks after it, pp. 48–49. *Hidden hypotheses:* $U \lhd F$ relatively open in a partial quadrant (for "sc-smooth" to make sense). *Deviation (restated):* strong bundle isomorphisms are not defined separately.

**V2.55 (Definition 2.48 and $K[i]$; `Polyfold.IsStrongBundleRetract`, `Polyfold.strongLevel`, `Polyfold.strongLevelShift`, `Polyfold.strongShiftOne`, `Polyfold.IsStrongBundleRetraction.mem_image_i` `Polyfold.IsStrongBundleRetraction.toScRetractionOne_retract`, `Polyfold.IsStrongBundleRetract.isSc` `Polyfold.IsStrongBundleRetract.isScRetract_one`, `Polyfold.IsStrongBundleRetract.isScSmoothRetract` `Polyfold.IsStrongBundleRetract.isScSmoothRetractMap_fst_one`, `Polyfold.IsStrongBundleRetract.isSc`).** $K$ is a *local strong bundle retract* over $C$ if $K = R(U \times F_0)$ for a strong bundle retraction on a relatively open $U \subseteq C$; then $K = \{(u,h) : u \in O, \Gamma(u,h) = h\}$. $K_{m,k}$ and $K[1] = \{(u,h) \in E_0 \times F_1 : (u, \iota_1 h) \in K\}$ are defined level by level; $K$ is a retract in $(C \times F_0, E \oplus F)$, $K[1]$ (the image of $R$ on $(U \lhd F)[1]$) a retract in $(C \times F_0, E \oplus F^1)$, and the projections $K[i] \to O$ and the inclusion $K[1] \to K[0]$ are sc-smooth. *Book:* Definition 2.48 and pp. 49–50. *Hidden hypotheses:* that $R$ preserves the double filtration at $(0,1)$, for $K[1] = R((U \lhd F)[1])$. *Deviation (restated):* none in substance.

**V2.56 (Definition 2.49; `Polyfold.IsScSmoothSection`, `Polyfold.IsScPlusSection`, `Polyfold.IsScPlusSectio` `Polyfold.IsScPlusSection.isScSmoothSection`).** An *sc-smooth section* of $p : K \to O$ is $f$ with $f(x)_1 = x$ on $O$, sc-smooth from $(O, C, E)$ to $(K, C \times F_0, E \oplus F)$; an *sc⁺-section* is $f : E_0 \to E_0 \times F_1$ with $f(x)_1 = x$ on $O$, sc-smooth into $(K[1], C \times F_0, E \oplus F^1)$. An sc⁺-section is an sc-smooth section. *Book:* Definition 2.49, p. 50 ($K(0)$, $K(1)$ for $K[0]$, $K[1]$). *Deviation (restated):* the category $\mathcal{SBR}$ is not formalized.

**V2.57 (Definition 2.50; `Polyfold.IsBanachable`, `Polyfold.StrongBundleChart`, `Polyfold.StrongBundleChart` `Polyfold.StrongBundleChart.continuous_fiber`, `Polyfold.StrongBundleChart.Compatible`, `Polyfold.StrongBundleChart.Compatible.symm`, `Polyfold.StrongBundleAtlas`, `Polyfold.StrongBundleAtla` `Polyfold.StrongBundleChart.compatible_of_locally`, `Polyfold.StrongBundleAtlas.equiv_iff`, `Polyfold.StrongBundleAtlas.Equiv.trans`, `Polyfold.StrongBundleAtlas.equivalence`, `Polyfold.IsStrongBundleRetraction.restrict`, `Polyfold.StrongBundleChart.restrict`, `Polyfold.StrongBundleChart.Compatible.restrict`).** For $P : Y \to X$ with linear structures on the fibres: a *strong bundle chart* is a chart $\varphi$ of the M-polyfold $X$, an sc-Banach space $F$, a strong bundle retraction $R$ on a relatively open $U \subseteq C$ with base retract $O$, and a homeomorphism $\Phi : P^{-1}(V) \to K = R(U \times F_0)$ with $\Phi(y)_1 = \varphi(P(y))$ and linear on each fibre (hence continuous on fibres). Two charts are *compatible* if $\Phi' \circ \Phi^{-1}$ and $\Phi \circ \Phi'^{-1}$ are sc-smooth between the overlaps on $[0]$ and, through the lift to $F_1$, on $[1]$, and carry fibre components in $F_1$ to fibre components in $F_1'$. Atlases and their equivalence as in V2.11; two atlases are equivalent iff every chart of one is compatible with every chart of the other, and the relation is an equivalence relation (it rests on the locality of compatibility: near a point of the overlap a transition factors through a chart of the middle atlas, on $[0]$ and on $[1]$). A strong bundle chart restricts to every open $s \subseteq V$ (retraction domain $U \cap r^{-1}(W)$ for $\varphi(s) = O \cap W$, $K_s = \{(u,h) \in K \mid u \in \varphi(s)\}$, $\Phi$ unchanged), and restrictions of compatible charts are compatible. A space is *Banachable* if it is linearly homeomorphic to a Banach space. *Book:* Definition 2.50, pp. 51–52. *Hidden hypotheses:* $\varphi$ a

chart of $X$ (the text: a homeomorphism onto $O$); the $[1]$-transition condition. *Deviation (hypotheses explicit):* both stated. The transitivity of the equivalence, which "an equivalence class of strong bundle atlases" (Definition 2.51) presupposes, and the restriction of charts, which the text uses silently (Lemma 5.2), are proved (Findings, Chapter 2, items 40, 42); a restriction whose retraction domain lies inside a given open $W \ni 0$ of $E$ is not, since $U \cap W$ need not be invariant under $r$.

**V2.58 (Definition 2.51, strong bundles, $Y[0]$, $Y[1]$, $P[0]$, $P[1]$; `Polyfold.StrongBundle`, `Polyfold.StrongBundleChart.toMChart`, `Polyfold.StrongBundleChart.Compatible.toMChart`, `Polyfold.StrongBundle.atlas0`, `Polyfold.StrongBundle.toMPolyfold0`, `Polyfold.StrongBundle.isScSmoot` `Polyfold.StrongBundle.doubleLevel`, `Polyfold.StrongBundle.mem_doubleLevel_iff`, `Polyfold.StrongBundle.oneChart`, `Polyfold.StrongBundle.oneChart_compatible`, `Polyfold.StrongBundle.` `Polyfold.StrongBundle.toMPolyfold1`, `Polyfold.StrongBundle.isScSmoothMap_proj1`, `Polyfold.StrongBundle.maximalAtlas`, `Polyfold.StrongBundle.maximalAtlas_equiv`, `Polyfold.StrongBundle.isChart_iff_exists_equiv`, `Polyfold.StrongBundle.isChart_congr`, `Polyfold.StrongBundle.IsChart.compatible`, `Polyfold.StrongBundle.IsChart.restrict`).** A *strong bundle* is a continuous surjection $P : Y \to X$, $Y$ Hausdorff paracompact, with Banachable fibres and a strong bundle atlas. The charts $\Phi$ form an sc-smooth atlas of $Y$ (the M-polyfold $Y[0]$), and $P : Y[0] \to X$ is sc-smooth. $Y_{m,k}$ ($k \le m+1$) is the set of $y$ with $\Phi(y) \in E_m \times F_k$ in the charts of the atlas around $y$; one chart around $y$ suffices. The charts $\Phi[1] : y \mapsto (\Phi(y)_1, \Phi(y)_2$ lifted to $F_1$), defined on $Y_{0,1}$ with the topology they generate and onto the retracts $K[1]$, are pairwise compatible, their transition maps being the $[1]$-transitions of Definition 2.50, and make $Y[1]$ an M-polyfold; $P[1] : Y[1] \to X$ is sc-smooth. The charts of the strong bundle (compatible with the atlas) are those of the maximal atlas, which is equivalent to the given one: a chart is one iff it belongs to some atlas of the class, any two are compatible, restrictions of charts are charts, and equivalent atlases have the same charts. *Book:* Definition 2.51 and the remarks after it, p. 52. *Hidden hypotheses:* the paracompactness of $Y[1]$ (proved, V2.64); for the chart-independence of $Y_{m,m+1}$ and of $Y_{0,1}$, the $F_1$-condition on the $[1]$-transition (V2.57). *Deviation (restated):* $Y[1]$ carries the topology generated by the charts $\Phi[1]$, as $X_m$ in §2.2; the charts are those induced by the given atlas. The notions defined through the charts compatible with the atlas depend only on the class; those defined with the charts of the given atlas itself ($Y_{m,k}$, $Y[1]$ and its atlas, sc$^+$-sections) are not shown invariant (Findings, Chapter 2, item 43).

**V2.59 (Definition 2.52; `Polyfold.IsSection`, `Polyfold.StrongBundle.IsScSmoothSection`, `Polyfold.StrongBundle.localRepOne`, `Polyfold.StrongBundle.IsScPlusSection`).** A *section* is $f$ with $P \circ f = \mathrm{id}$; it is *sc-smooth* if it is sc-smooth $X \to Y[0]$, and an *sc$^+$-section* if in every chart of the atlas its fibre values lie in $F_1$ and its local representative, lifted to $E_0 \times F_1$, is sc-smooth from $O$ into $K[1]$. *Book:* Definition 2.52, p. 53. *Deviation (restated):* sc$^+$ is defined chart by chart; this is equivalent to the text's definition: $f$ is an sc$^+$-section iff $P \circ f = \mathrm{id}$ and $f = \iota \circ g$ for an sc-smooth $g : X \to Y[1]$, $\iota : Y[1] \to Y$ the inclusion (`Polyfold.StrongBundle.isScPlusSection_iff`).

**V2.60 (Definition 2.53; `Polyfold.pullbackSpace`, `Polyfold.pullbackProj`, `Polyfold.pullbackProj'`, `Polyfold.pullback_comm`, `Polyfold.continuous_pullbackProj`, `Polyfold.isClosed_pullbackSpace`).** $f^*Y = \{(z, y) : P(y) = f(z)\}$ with the projections $P_f$, $P'$; $P \circ P' = f \circ P_f$; $P_f$ is continuous, and $f^*Y$ is closed in $Z \times Y$ when $X$ is Hausdorff and $P$, $f$ are continuous. *Book:* Definition 2.53, p. 53. *Hidden hypotheses:* none.

**V2.61 (Proposition 2.54; `Polyfold.proposition_2_54`, `Polyfold.pullbackAtlas`, `Polyfold.PullbackChartDat` `Polyfold.PullbackChartData.toChart`, `Polyfold.PullbackChartData.transition_eqOn`, `Polyfold.PullbackChartData.transition_isScSmooth`, `Polyfold.PullbackChartData.transitionOne_isScS` `Polyfold.PullbackChartData.compatible_toChart`, `Polyfold.paracompactSpace_pullbackSpace`,**

`Polyfold.pullbackFiberCLE`, `Polyfold.pullbackRetraction`, `Polyfold.isStrongBundleRetraction_pullbac`
`Polyfold.pullback_image`, `Polyfold.isStrongBundleRetract_pullback`). Let $P : Y \to X$ be a strong bundle and $f : Z \to X$ sc-smooth. Then $f^*Y$ is paracompact, and $P_f : f^*Y \to Z$ is a strong bundle whose atlas consists of the pulled-back charts: for a strong bundle chart $\Phi$ over $\varphi$ and a chart $\psi$ of $Z$ with $f(\operatorname{dom}\psi) \subseteq \operatorname{dom}\varphi$ and $g = \varphi \circ f \circ \psi^{-1}$, the chart $\Phi_f(z, y) = (\psi(z), \operatorname{pr}_F\Phi(y))$ onto $K_g = \{(w, h) : w \in O_Z, (g(w), h) \in K\}$, the image of the strong bundle retraction $R_g(w, h) = (r_Z(w), \Gamma(g(r_Z(w)), h))$. Its fibres are $P_f^{-1}(z) \cong P^{-1}(f(z))$ (a linear homeomorphism). Transition maps of pulled-back charts are $(w, h) \mapsto (\psi_2\psi_1^{-1}(w), \operatorname{pr}_F(\Phi_2\Phi_1^{-1})(g_1(w), h))$ and are sc-smooth on $[0]$ and $[1]$. *Book:* Proposition 2.54, p. 53 ("the easy proof is left to the reader"). *Hidden hypotheses:* $f^*Y$ paracompact (required by Definition 2.51, not mentioned); proved from Theorem 2.16 for $Z$ and $Y[0]$. *Deviation (hypotheses explicit):* "natural structure" is the atlas of all pulled-back charts. That the projection $P' : f^*Y \to Y$ is a strong bundle map (sc-smooth on $[0]$ and $[1]$), which "natural" presumably also says, is not formalized (Findings, Chapter 2, item 51).

**V2.62 (Theorem 2.55, Smirnov's theorem;** `Polyfold.metrizableSpace_of_locally`, `Polyfold.metrizableSpac`
`Polyfold.metrizableSpace_of_pullback`, `Polyfold.cone_triangle`, `Polyfold.truncDist`,
`Polyfold.paracompactSpace_of_metrizableSpace`). A paracompact Hausdorff space in which every point has a metrizable open neighbourhood is metrizable. More generally, a space $Y$ with open metrizable subspaces $V_i$ and a locally finite family of continuous $\rho_i \geq 0$ with $\operatorname{supp}\rho_i \subseteq V_i$, not all vanishing at any point, is metrizable, with the metric $\sum_i |\rho_i(x) - \rho_i(y)| + \min(\rho_i(x), \rho_i(y)) \min(d_i(x, y), 1)$; and if $\pi : Y \to X$ is continuous into a paracompact Hausdorff $X$ whose points have open neighbourhoods $U$ with $\pi^{-1}(U)$ metrizable, $Y$ is metrizable. Metrizable spaces are paracompact. *Book:* Theorem 2.55, p. 56, quoted from Smirnov. *Deviation (stronger):* proved, in the pull-back form used for $X_m$ and $TX$.

**V2.63 (restriction and locality of sc^k maps;** `Polyfold.IsSck.of_subset`, `Polyfold.IsScSmooth.of_subset`,
`Polyfold.IsSc1.tangentMap_eqOn_of_subset`, `Polyfold.IsSc1.of_locally`, `Polyfold.IsSck.of_locally`,
`Polyfold.IsScSmooth.of_locally`, `Polyfold.IsScSmooth.congr`). On sc-domains: an sc$^k$ map restricts to sc$^k$ on every smaller sc-domain, with the same tangent map there; a map that is sc$^1$ (resp. sc$^k$, sc-smooth) on $U \cap W$ for an open $W$ around each point of $U$ is sc$^1$ (resp. sc$^k$, sc-smooth) on $U$; sc-smoothness depends only on the values on $U$. *Book:* used without statement in §2.1–§2.2 (Propositions 2.4, 2.10, Definitions 2.13–2.14). *Deviation (not in the book):* infrastructure the text uses silently.

**V2.64 (the space $Y[1]$ and the inclusion $Y[1] \to Y[0]$;** `Polyfold.StrongBundle.oneSet`,
`Polyfold.StrongBundle.mem_oneSet_iff`, `Polyfold.StrongBundle.One`, `Polyfold.StrongBundle.continuous`
`Polyfold.StrongBundle.metrizableSpace_one`, `Polyfold.StrongBundle.paracompactSpace_one`,
`Polyfold.StrongBundle.isScSmoothMap_oneVal`). For a strong bundle $P : Y \to X$, $Y_{0,1}$ is the set of points whose fibre component lies in $F_1$ in every chart of the atlas around them, and one such chart suffices. With the topology generated by the charts $\Phi[1]$ (V2.58), the inclusion $Y[1] \to Y$ is continuous, so $Y[1]$ is Hausdorff; $Y[1]$ is metrizable, hence paracompact; and the inclusion $Y[1] \to Y[0]$ is sc-smooth (in the charts $\Phi[1]$, $\Phi$ it is $(u, h) \mapsto (u, \iota h)$). *Book:* not stated; the text takes $Y[1]$ to be an M-polyfold on $Y_{0,1}$ (p. 52). *Hidden hypotheses:* paracompactness of $Y[1]$, required by Definition 2.13. *Deviation (not in the book):* proved; metrizability by the pull-back form of Smirnov's theorem over the paracompact base (V2.62), as in Theorem 2.16.

**What is not verified in this chapter**

- p. 18, Proposition 2.2 (classical $C^\infty$ retractions have submanifold images): motivation only; not attempted (the exposition proves it, §8.1).
- p. 22, Definition 2.9 (sc-smooth bundle retractions, the categories $\mathcal{R}$, $\mathcal{BR}$): not formalized as a definition; its instances ($Tr$, strong bundle retractions) are.
- p. 25, $X_m$ as an M-polyfold for $m \geq 2$: only $X^1$ is constructed (V2.16); the iterate $(X^1)^1 \cdots$ has the points of $X_m$, its topology is not compared with that of $X_m$; Theorem 2.16 is proved for every $m$.
- pp. 26–30, the vector space, Banach and sc-Banach structures of $T_xX$, and Proposition 2.21 (3),
  (4) in $T_aX$ rather than in a chart: not formalized.
- p. 27, Proposition 2.19 (2), openness of $p : TX \to X_1$: not proved (not used).
- pp. 33–35, Lemma 2.29 in its path formulation, and $T_xO = C_xO - C_xO$: not formalized; the sign arguments are V2.34.
- p. 39, the intrinsic description of extended faces ($k$ components of $\{d_C = 1\}$, minimal subspaces, independence of the presentation): the formalization works in a presentation.
- p. 42, Definition 2.38 ($T^R_xX$, $C_xX$ in $T_xX$): only the local form, its invariance and the partial quadrant $C_oO$ (in a tame chart V2.42, in any chart V2.33); not the transport to $T_xX$ or $d_{C_xX}(0_x) = d_X(x)$.
- pp. 48–51, strong bundle isomorphisms, the category $\mathcal{SBR}$ and the functors forget$[i]$: not formalized.
- p. 53, Proposition 2.54: that the projection $P' : f^*Y \to Y$ of the pull-back is a strong bundle map; the pulled-back atlas is V2.61.
- p. 56, Lemma 2.56 and Proposition 2.57: not needed, since the proofs of Theorem 2.16 and Proposition 2.19 (3) avoid them.

---

## 4. Chapter 3 (HWZ §§3.1–3.8): partly verified

Chapter 3 of Part I (preprint pp. 61–146) develops sc-Fredholm theory. §3.1 defines germs, sections of strong bundles, fillers, basic germs, sc-Fredholm germs and sections, proves local compactness (Theorem 3.12) and the stability under sc$^+$-perturbations (Theorem 3.15), and states the implicit function theorems (Theorems 3.19, 3.23), with good position (Definition 3.21, Proposition 3.22). §3.2 introduces the tangent germ property and shows that it makes a subset a sub-M-polyfold (Definition 3.25, Proposition 3.26, Theorem 3.27). §3.3 proves the germ implicit function theorem (Definition 3.28, Theorems 3.29–3.35). §3.4 proves that basic germs are stable under sc$^+$-perturbations (Theorem 3.38). §3.5 proves local regularity and compactness (Theorem 3.40) and local solvability (Corollary 3.41, Theorem 3.43, Corollary 3.47). §3.6 states and proves the implicit function theorem for sc-Fredholm sections (Proposition 3.52, Definition 3.53, Theorem 3.54, Lemma 3.55) and its global forms (Theorems 3.56, 3.57). §3.7 gives a criterion for being conjugate to a basic germ (Theorem 3.58, Lemmas 3.59–3.61). The appendix §3.8 proves Proposition 3.22, Lemma 3.46, Lemma 3.55, and the inverse and implicit function theorems in partial quadrants (Theorems 3.72, 3.78).

What is verified. **The germ implicit function theorem (Theorem 3.33) holds with the text's hypotheses**, and so do Corollary 3.34 and Theorem 3.35: the solution germ is sc^k on the germ of neighbourhoods shrunk by $2^k - 1$ levels, and sc-smooth on one diagonal germ, which shrinks fast with the level (V3.31, V3.32). **The implicit function theorems also hold with the text's hy-**

**potheses, local and global**: Theorems 3.54, 3.23 and 3.19 (V3.94, V3.95), Theorem 3.57, Corollary 3.24 and Theorem 3.56 (V3.87, V3.70, V3.86), once three steps of the text's proof are replaced. The surjectivity of $f'(y)$ at the zeros $y$ near a given zero, which the text derives from Corollary 3.41 and which does not follow from it (V3.82; the same step stands in the book's arXiv version), is proved by a new argument, an implicit function theorem at the nearby zero for a perturbed equation (V3.93); the regularity of the solution germ away from the centre is proved by re-centring (V3.73, V3.74); and the relative openness of the set $\Sigma$ in the proof of Theorem 3.27 by the choice of the good complement (V3.75, V3.77). With them: the dimension of the solution set is the Fredholm index (V3.96, V3.97), the index is locally constant along the solution set (V3.98; also without good position, with the text's hypotheses: tame germ data and no bump functions), and so, with the text's hypotheses, is the index of Definition 6.1 (read with the text's locally defined sc$^+$-sections) along sc-smooth paths $\mathbb{R} \to X$, the text's Proposition 6.2 for paths on $\mathbb{R}$ (V3.98), ker $f'(y)$ is the tangent space of the solution set in a chart (V3.99), and good position is the surjectivity of $f'(y)$ at zeros with $d_X(y) = 0$, so that Theorems 3.19 and 3.56 hold with the text's hypothesis "$f'(x)$ surjective" (V3.100, V3.101). The forms first proved under an added filler condition at the nearby zeros (V3.68, V3.69, V3.82, V3.85) are kept; that condition follows from good position (V3.102), and with Definition 3.4 as stated the text's argument gives the kernel half of Theorem 3.54 (1) (V3.81) but not the surjectivity. All these statements are the repaired ones. They were first proved under the hypothesis that the base retraction of every strong bundle chart compatible with the atlas is tame, which no strong bundle over a nonempty base satisfies (V3.104, where only the half of the argument that concerns retractions is machine-checked), so those first statements were vacuous; they now assume what the text's Definitions 3.50, 3.51 and 3.53 ask, germ data and good position in a chart whose base retraction is tame, and conclude good position in such charts (V3.103, V3.107). Also verified: every definition of §§3.1–3.6 (Definitions 3.1–3.4, 3.6, 3.7, 3.11, 3.14, 3.21, 3.25, 3.28, 3.37, 3.48, 3.50, 3.51, 3.53; since the switch of Definition 3.11 to the text's germ reading, the fillings, the local sc$^+$-sections and every notion built on them are germs whose levels may shrink with $m$, as in the text, V3.7, V3.9, V3.13, and so, since a fourth addendum, is Definition 3.3, V3.5); Proposition 3.5, also in sc form (V3.8); Lemma 3.9 as stated (V3.11); Proposition 3.10, both statements, the second on every level (V3.12); Theorems 3.12 (with fewer hypotheses), 3.15 (without tameness), Corollary 3.13; Theorem 3.18 (without tameness), with the linearization of the perturbed section (V3.91, V3.92); Lemmas 3.16, 3.17; Proposition 3.22 (without the extreme-ray lemmas); Proposition 3.26; Theorem 3.27 under two added regularity conditions, each with a counterexample to the step of the text's proof it replaces (for solution sets both are theorems); Theorems 3.29, 3.30, Lemmas 3.31, 3.32 (as one induction), Theorem 3.38 (with a linear coordinate change), Corollary 3.39, Theorem 3.40 (with compact closures), Corollary 3.41, Lemmas 3.42, 3.44–3.46; Theorem 3.43 and Corollary 3.47 (their clause (1), "the solution map is an sc-smooth germ", in the level-wise sense: $C^k$ into level $k$ on a neighbourhood depending on $k$); Remark 3.49; Proposition 3.52 in charts, as an sc-Fredholm operator with the index $n - N$ of the conjugating basic germ on every level, in every chart (V3.46, V3.96); Lemma 3.81 as stated (V3.66); Lemma 3.55 (with the added hypothesis that the good complement is closed); Theorem 3.54 as described above, with its graph description; Theorem 3.58 in the coordinates of its proof; Lemmas 3.59 and 3.60 (in an abstract form, with their hypotheses derived in Theorem 3.58) and Lemma 3.61; and the appendix except Lemmas 3.62–3.64, 3.66–3.68, 3.76, which the formal proofs avoid. Not verified: Proposition 3.52 on $T_xX$ itself, which carries no linear structure here; ker $f'(y) = T_yS$ with $T_yS$ a subspace of $T_yX$ (it is verified in a chart); Proposition 6.2 for paths sc-smooth on $[0,1]$ only (the project has no M-polyfold $[0,1]$); the passage from an atlas of tame charts to tame germ data (Definition 3.11 $\Rightarrow$ Definition 3.50), which is open. Remarks 3.8 and 3.20 compare the definitions with the earlier paper [21] and make no claim to verify. Items V3.68–V3.90 were added after the first edition of this section

(the Lean formalized after it); the earlier items whose content changed with them (V3.13, V3.23, V3.24, V3.34, V3.38, V3.42, V3.46–V3.54, V3.59, V3.62) are updated in place. Items V3.91–V3.102 were added in a second addendum (Theorem 3.18, Theorem 3.54 (1) with the text's hypotheses, the index of the solution set, the kernel as tangent space, good position at interior zeros); the items whose content changed with them (V3.46, V3.47, V3.68, V3.69, V3.81–V3.87) are updated in place. Items V3.103–V3.107 were added after the repair of the tameness hypothesis (Findings, Chapter 3, items 190–199: the tame germ notions, the non-tame retraction behind the vacuity, Theorems 3.15 and 3.18 for tame germ data, good position in any chart of the germ data); the items whose statements changed with it (V3.45, V3.47, V3.48, V3.68–V3.70, V3.80–V3.82, V3.84–V3.87, V3.94, V3.95, V3.97–V3.102) are updated in place. A third addendum (sc-Fredholm theory on all levels, the one-germ form of Theorem 3.35, Lemmas 3.9 and 3.81 as stated, good position with the text's sc-complement, the index without good position, and the difference between the text's germs and the formal definitions on the whole chart) updated V3.5, V3.7–V3.9, V3.11–V3.13, V3.16, V3.19, V3.20, V3.22, V3.28, V3.31, V3.32, V3.43, V3.45–V3.47, V3.66, V3.68, V3.70, V3.86, V3.87, V3.94–V3.98 and V3.103 in place. The switch of Definition 3.11 and of everything built on it to the text's germ reading (phase 2: the canonical notions and theorems are the germ reading, the project's earlier whole-chart ones are kept as `…WholeChart`, `…_wholeChart`; Findings, Chapter 3, items 330–334) updated V3.5, V3.7–V3.9, V3.13, V3.16, V3.17, V3.19, V3.43, V3.45–V3.47, V3.68, V3.78, V3.79, V3.81, V3.82, V3.91, V3.93, V3.96, V3.98, V3.102, V3.103, V3.105 and V3.107 in place. A fourth addendum (2026-11-02: Definition 3.3 in the germ reading, Definition 6.1 and Proposition 6.2, germ data and the charts of a tame atlas; Findings, Chapter 3, items 340–355) updated V3.5, V3.13, V3.98 and V3.103 in place. A fifth addendum (2026-11-02: the restriction of a strong bundle to an open set, the index along the zero set and Proposition 6.2 without bump functions, Definition 6.1 with the text's local sections; Findings, Chapter 3, items 349, 360–367) updated V3.68 and V3.98 in place.

The setting. Germs are fixed sequences of sets (Definition 3.1), and the calculus of germs — restriction, chain rule, pairing, Propositions 1.16–1.18 for germs — is proved, not assumed. The finite-dimensional parameter spaces $\mathbb{R}^n$, $\mathbb{R}^N$ are arbitrary finite-dimensional normed spaces $P$, $Q$, and the parameter set $\widetilde{C}$ a convex set containing $0$ with nonempty interior, which is closed or a partial quadrant where a theorem needs it. A "strong bundle chart" in Definitions 3.3, 3.6, 3.11 is a chart compatible with the atlas. The fillings and the local sc$^+$-sections of Definition 3.11 are, as in the text, germs on sc-germs of neighbourhoods whose levels may shrink with $m$ (the text's germ reading, since phase 2; V3.7, V3.9, V3.13); the project's earlier notions, with these data on the whole $U$ of the chart (levels $U \cap E_m$), are stricter and are kept under the names `…WholeChart`. The sc-smooth section germs of Definition 3.3, which no theorem assumes, are germs in the same sense since a fourth addendum (V3.5; Findings, Chapter 3, items 220–221, 330). Tameness of the strong bundle is nowhere used in §3.1. It enters §3.6 as the text's Definitions 3.50, 3.51 and 3.53 put it: the germ data of $f$ at a point, and its good position, lie in a strong bundle chart whose base retraction is tame (V3.103; "tame germ data" and "good position in a tame chart" below). It is used, for the chart at the zero $x_0$, through Proposition 2.35, $(1 - Dr(0))E \subseteq E_0$, and, for the tangent germ property of the solution set, through the tameness of the model of the chart $\psi \circ \varphi$ (V3.71). The reading first formalized, that the base retractions of all charts compatible with the atlas are tame, cannot be met (V3.104, the retraction-level half machine-checked). Classical regularity in the appendix is Mathlib's $C^k$ within the set, which on relatively open subsets of partial quadrants is the text's Definition 3.71. Findings referred to below are those of `Exposition/Findings.md`, "Chapter 3".

**V3.1 (Definition 3.1, sc-germs of neighbourhoods; `Polyfold.ScNbhdGerm`, `Polyfold.ScNbhdGerm.tangent`,**

`Polyfold.ScNbhdGerm.ofSet`, `Polyfold.IsQuadrantLike`, `Polyfold.IsPartialQuadrant.isQuadrantLike`). An sc-germ of neighbourhoods around 0 in $C \subseteq E_0$ is a family $U_m \subseteq E_m$, each of the form $O \cap C_m$ with $O$ open in $E_m$, containing 0, with $\iota U_{m+1} \subseteq U_m$. Its tangent germ has levels $U_{m+1} \times E_m$. A relatively open $U \ni 0$ in $C$ gives the germ $U_m = U \cap E_m$. $C$ is quadrant-like if $0 \in C$ and every $C_m$ is convex with nonempty interior; partial quadrants are. *Book:* Definition 3.1, p. 62. *Hidden hypotheses:* none. *Deviation (restated):* a germ is a fixed sequence, not a class up to shrinking; the text's "the size of the sets $U_m$ does not matter" is not built in, and the theorems say on which germ they hold. $(U_{m+1})_m$ is a germ only when all levels coincide (finite-dimensional $E$).

**V3.2 (Definition 3.2, sc^k-germs;** `Polyfold.IsSc0Germ`, `Polyfold.IsSc1GermWith`, `Polyfold.IsSc1Germ`, `Polyfold.IsSckGerm`, `Polyfold.IsScSmoothGerm`, `Polyfold.IsSck.isSckGerm`**).** $f$ is an sc⁰-germ on $(U_m)$ if $f(U_m) \subseteq F_m$ and $f : U_m \to F_m$ is continuous for every $m$; sc¹ if moreover, for a family of bounded $Df(x) : E_0 \to F_0$, $x \in U_1$, $f$ is differentiable within $U_1$ at $x$ (increments in $E_1$, remainder in $F_0$) with derivative $Df(x)$, and the tangent map $(x, h) \mapsto (f(x), Df(x)h)$ is an sc⁰-germ on the tangent germ; sc^{k+1} if sc¹ with sc^k tangent map; sc-smooth if sc^k for all $k$. An sc^k map on a relatively open $U \ni 0$ is an sc^k-germ on $(U \cap E_m)$. *Book:* Definition 3.2, pp. 62–63. *Hidden hypotheses:* for quadrant-like $C$ the derivative is unique on $U_1$. *Deviation:* none.

**V3.3 (the calculus of germs;** `Polyfold.IsSckGerm.congr`, `Polyfold.IsSckGerm.mono`, `Polyfold.ScNbhdGerm.MapsToGerm`, `Polyfold.IsSckGerm.comp`, `Polyfold.IsSc1Germ.comp`, `Polyfold.ScOperator.isScSmoothGerm`, `Polyfold.IsSckGerm.pair`, `Polyfold.ScNbhdGerm.preimage`**).** For quadrant-like sets: sc^k-germs depend only on their values on $U_0$ and restrict to smaller germs; if $f$ is an sc^k-germ on $(U_m)$, $g$ an sc^k-germ on $(V_m)$ and $f(U_m) \subseteq V_m$ for every $m$, then $g \circ f$ is an sc^k-germ, with $T(g \circ f) = Tg \circ Tf$ on the tangent germ (k = 1); sc-operators are sc-smooth germs on every germ; pairings of sc^k-germs are sc^k-germs; for an sc⁰-germ $f$ with $f(0) = 0$ and a germ $G'$ around 0 there is a germ inside $(U_m)$ on which $f$ maps into $G'$ level by level. *Book:* not stated; used throughout §§3.3–3.6. *Hidden hypotheses:* the level-wise inclusion $f(U_m) \subseteq V_m$ for compositions. *Deviation (not in the book).*

**V3.4 (Propositions 1.16–1.18 for germs;** `Polyfold.IsSc1GermWith.exists_levelDeriv`, `Polyfold.IsSc1GermWith.continuousOn_levelDeriv_clm`, `Polyfold.IsSc1GermWith.hasFDerivWithinAt_lev` `Polyfold.ScNbhdGerm.shift`, `Polyfold.IsSckGerm.shift`, `Polyfold.IsSckGerm.contDiffOn_levels`**).** For an sc¹-germ with derivative family $D$ and $x \in U_{m+1}$, $D(x)$ restricts to bounded operators $E_m \to F_m$, continuous in $x$ in the operator norm on $U_{m+1}$, and $f : U_{m+1} \to F_m$ is differentiable within $U_{m+1}$ with these derivatives. $(U_{m+1})_m$ is a germ in $E^1$ around $0 \in C^1$, and an sc^k-germ on $(U_m)$ is an sc^k-germ $(U_{m+1})_m \to F^1$. An sc^k-germ is $C^k$ from $U_{m+k}$ to $F_m$. *Book:* not stated for germs; the text applies Propositions 1.16–1.18 to germs (Lemma 3.9: "recalling Proposition 1.17", which is 1.16; Lemma 3.31; Corollary 3.34). *Hidden hypotheses:* none. *Deviation (not in the book):* the germ versions, with the proofs of Chapter 1.

**V3.5 (Definition 3.3, sc-smooth section germs;** `Polyfold.StrongBundle.IsScSmoothSectionGerm`, `Polyfold.StrongBundle.IsChart`, `Polyfold.StrongBundle.IsScSmoothSection.isScSmoothSectionGerm`, `Polyfold.StrongBundle.isChart_congr`**).** $(f, x_0)$ is an sc-smooth section germ of the strong bundle $P : Y \to X$ if $x_0$ is smooth and in some strong bundle chart compatible with the atlas, centred at $x_0$, the principal part $\mathbf{f}$ of $f$ satisfies: $\mathbf{f} \circ r$ is an sc-smooth germ on some sc-germ of neighbourhoods $G$ of 0 with $G_0 \subseteq U$. An sc-smooth section is an sc-smooth section germ at every smooth point that has a centred chart (with the germ $U \cap E_m$). *Book:* Definition 3.3, p. 63. *Hidden hypotheses:* "a strong bundle chart" means a chart compatible with the atlas (a

chart of the atlas need not be centred); the notion does not depend on the atlas within its class (`Polyfold.StrongBundle.isChart_congr`, V2.58). *Deviation (hypotheses explicit):* the chart, as just said. Since a fourth addendum the germ is the text's: $\mathbf{f} \circ r$ is an sc-smooth germ on some sc-germ of neighbourhoods $G$ with $G_0 \subseteq U$, whose levels may shrink with $m$, as for Definitions 3.4, 3.6 and 3.11 since phase 2. Until then it was required on the whole $U$ of the chart, which is stricter (a function $C^k$ on $(-1/k, 1/k)$ for every $k$ is an sc-smooth germ in the constant scale $\mathbb{R}$ and sc-smooth on no neighbourhood of 0, `Polyfold.exists_isScSmoothGerm_not_isScSmooth`), and the item was "narrower". No theorem assumes the definition (Findings, Chapter 3, items 220–221, 260).

**V3.6 (the section algebra and the sum rule; `Polyfold.IsScSmooth.add`, `Polyfold.sectionAdd`, `Polyfold.StrongBundleChart.principal_sectionAdd`, `Polyfold.StrongBundleChart.isZeroAt_iff`, `Polyfold.StrongBundleChart.add_mem_K`, `Polyfold.StrongBundle.IsScSmoothSection.isScSmoothRetractM` `Polyfold.StrongBundle.IsScPlusSection.isScSmoothRetractMap_one`).** Sums, differences and negatives of sc-smooth maps on relatively open subsets of partial quadrants are sc-smooth. The fibrewise sum $f + s$ of sections is a section, with principal part $\mathbf{f} + \mathbf{s}$ in every chart; $f(x) = 0$ iff $\mathbf{f}(\varphi(x)) = 0$; the fibres of $K$ over $O$ are linear. In every strong bundle chart compatible with the atlas, the representative $z \mapsto (z, \mathbf{f}(z))$ of an sc-smooth section is sc-smooth $O \to K$, and the representative $z \mapsto (z, \mathbf{s}_1(z))$ of an sc⁺-section is sc-smooth $O \to K[1]$. *Book:* not stated; used in Theorems 3.12, 3.15. *Hidden hypotheses:* the compatibility on $[1]$ of Definition 2.50 (its first use). *Deviation (not in the book).*

**V3.7 (Definition 3.4, fillers; `Polyfold.IsFillingGerm`, `Polyfold.fillDefect`, `Polyfold.fillDefect_isScSmoot` `Polyfold.IsFillerOf`, `Polyfold.StrongRetr.r`, `Polyfold.StrongRetr.`, `Polyfold.StrongRetr.CLM`).** For a strong bundle retraction $R(u,h) = (r(u), \rho(u)h)$ on $U \lhd F$ with $O = r(U)$, $0 \in U$: $g$ is a filling of $f$ if $g$ is an sc-smooth germ on an sc-germ of neighbourhoods $G$ with $G_0 \subseteq U$ and (1) $f = g$ on $O$ near 0; (2) a point $y \in U$ near 0 with $g(y) = \rho(r(y))g(y)$ lies in $O$; (3) $y \mapsto [1 - \rho(r(y))]g(y)$ is an sc¹-germ on $G$ with a derivative family $Dk$, and $Dk(0)$ restricts to a topological linear isomorphism $\ker Dr(0) \to \ker \rho(0)$. ($\rho(u)$ is a bounded operator, $\rho(x)$ a projection for $x \in O$.) *Book:* Definition 3.4, p. 64. *Hidden hypotheses:* the map in (3) is differentiable at 0 (true for every sc-smooth germ $g$, and $Dk(0)$ is unique). *Deviation (hypotheses explicit):* stated for the principal parts; tameness unused. Since the switch of Definition 3.11 to the text's germ reading (phase 2) $g$ and the map in (3) are germs whose levels may shrink with $m$; the stricter whole-chart filler (`Polyfold.IsFillerOf`, the map in (3) sc¹ on all of $U$, the sc-smoothness of $g$ on $U$ added in Definition 3.6) is kept for the whole-chart notions (Findings, Chapter 3, items 220–221, 261, 330). Formerly "narrower", for the whole-chart reading.

**V3.8 (Proposition 3.5 and the matrix of $Dg(0)$; `Polyfold.IsFillerOf.proposition_3_5`, `Polyfold.proposition_3_5`, `Polyfold.fillerRestrict`, `Polyfold.fillerRestrict_kerEquiv`, `Polyfold.fillerRestrict_cokerEquiv`, `Polyfold.StrongRetr.scDeriv_fiber_zero`, `Polyfold.IsFillerOf.d` `Polyfold.IsFillerOf.scDeriv_section_comp_r`, `Polyfold.IsScFredholm.restrict`, `Polyfold.IsFillerOf.p` `Polyfold.IsFillerOf.filler_condition_three_sc`, `Polyfold.IsFillerOf.proposition_3_5_sc_converse`, `Polyfold.IsFillerOf.proposition_3_5_sc_iff`).** Let $g$ be sc-smooth on $U$ and fill the section $f$ of $K \to O$, with $g(0) = 0$. Then $D(\rho(r(\cdot))g)(0) = \rho(0)Dg(0)$, $(1 - \rho(0))Dg(0)Dr(0) = 0$, $D(f \circ r)(0) = Dg(0)Dr(0)$; and $Tf(0) := Dg(0)$ restricted to $Dr(0)E \to \rho(0)F$ is surjective iff $Dg(0)$ is, Fredholm iff $Dg(0)$ is, and has the same index ($\dim\ker - \dim\operatorname{coker}$). The linear algebra holds for any idempotents $p$, $q$ and $L$ with $(1-q)Lp = 0$ and $(1-q)L : \ker p \to \ker q$ a topological isomorphism. In sc form: if $Dg(0)$ is an sc-Fredholm sc-operator, then $Dr(0)E$, $\rho(0)F$, $\ker Dr(0)$ and $\ker \rho(0)$ are sc-subspaces, $Tf(0) : Dr(0)E \to \rho(0)F$ is sc-Fredholm with the index of $Dg(0)$

on every level, and $(1-\rho(0))Dg(0) : \ker Dr(0) \to \ker \rho(0)$ is an sc-isomorphism (condition (3) on every level); for $Dg(0)$ an sc-operator, $Dg(0)$ is sc-Fredholm iff $Tf(0)$ is sc-Fredholm and condition (3) holds on every level. *Book:* the paragraph after Definition 3.4 and Proposition 3.5, pp. 64–65. *Hidden hypotheses:* $(u,h) \mapsto \rho(u)h$ differentiable at $(0,0)$; $g(0)=0$. *Deviation (stronger):* the index equality without any finiteness assumption, via isomorphisms of kernels and cokernels instead of the text's factorization. The text's target "$(1-D\rho(0))F$" is $(1-\rho(0))F$. The sc form needs condition (3) only on level 0, as the text states it (the higher levels of $Tf(0)$ come from the regularity of $Dg(0)$); its converse needs condition (3) on every level and fails with level 0 alone (an example with the shift on a weighted $\ell^2$, not formalized; Findings, Chapter 3, items 234, 237, 238). The same holds for a filling germ (V3.7), with $Dg(0)$ the value at 0 of any derivative family of $g$ (`Polyfold.IsFillingGerm.proposition_3_5`, `Polyfold.IsFillingGerm.proposition_3_5_sc`, `Polyfold.IsFillingGerm.proposition_3_5_sc_iff`; Findings, Chapter 3, items 272, 289, 314).

**V3.9 (Definition 3.6, filled versions; `Polyfold.IsFillingGerm`, `Polyfold.IsFilledVersion`, `Polyfold.IsFilledVersion.isFillingGerm`, `Polyfold.IsFillingGerm.isFilledVersion`).** A filling of the principal part **f** (V3.7) is an sc-smooth germ $g$ on an sc-germ of neighbourhoods inside $U$, whose levels may shrink with $m$, filling **f** near 0; the centred chart is that of Definition 3.11 (V3.13). *Book:* Definition 3.6, p. 66. *Hidden hypotheses:* none. *Deviation (restated):* the datum in a given chart, for the principal parts; the chart is quantified in Definition 3.11. The project's earlier filled version (`Polyfold.IsFilledVersion`), $g$ sc-smooth on all of $U$, is stricter for germs whose levels shrink with $m$ (round-trip items V3.9, V3.43); it is exactly a filling sc-smooth on all of $U$ (`IsFilledVersion.isFillingGerm`, `IsFillingGerm.isFilledVersion`; Findings, Chapter 3, items 220–221, 260–262, 330). Formerly "narrower", for the whole-chart reading.

**V3.10 (Definition 3.7, basic germs; `Polyfold.IsBasicGerm`, `Polyfold.IsSc0ContractionGerm`).** For a finite-dimensional $P$, a set $\widetilde{C} \subseteq P$, a finite-dimensional $Q$ and a germ $G$ around 0 in $\widetilde{C} \oplus W$: $f$ is a basic germ with contraction map $B$ if $f$ is an sc-smooth germ on $G$ with $f(0)=0$, the $W$-component of $f(a,w)$ is $w - B(a,w)$ on $U_0$, and $B$ is an sc$^0$-contraction germ (V3.25). *Book:* Definition 3.7, p. 66 (Remark 3.8: historical). *Hidden hypotheses:* the points of the contraction estimate lie in $U_m$. *Deviation (restated):* $\mathbb{R}^n$, $\mathbb{R}^N$ replaced by finite-dimensional spaces.

**V3.11 (Lemma 3.9; `Polyfold.ContractionGerm.norm_levelDeriv_snd_le`, `Polyfold.IsSc0ContractionGerm.d` `Polyfold.IsSc0ContractionGerm.exists_norm_deriv_snd_le`, `Polyfold.IsSc0ContractionGerm.exists_nor`** For an sc$^0$-contraction germ $B$ that is an sc$^1$-germ: (a) in the working form, $\|D_2B(a,w)\|_{L(W_m)} \le \frac{1}{2}$ at points $(a,w) \in U_{m+1}$ with $a \in N(m+1) \cap \widetilde{C}$, $w \in O(m+1)$, $\iota w \in O(m)$; (b) $D_2B(0,0)=0$; (c) for every $\varepsilon > 0$ there is $r>0$ with $|D_2B(x)\zeta|_0 \le \varepsilon|\zeta|_0$ for $x \in U_1$, $|x|_1 < r$, $\zeta \in W_0$; (d) as stated, for $\widetilde{C}$ quadrant-like: for every $\varepsilon>0$ and $m$ there is $r>0$ with $|D_2B(x)\zeta|_m \le \varepsilon|\zeta|_m$ for $\zeta \in W_m$ at every $x \in U_{m+1}$ with $|x|_m < r$. *Book:* Lemma 3.9, p. 67 (every $\varepsilon$, every level, $(a,w) \in E_{m+1}$ small in $E_m$; $B$ sc-smooth). *Hidden hypotheses:* differentiability of $B : U_{m+1} \to W_m$ (Proposition 1.16 for germs, V3.4). *Deviation (stronger):* sc$^1$ suffices, and the general statement is proved as stated (d); smallness on level $m+1$ is not needed, since the contraction estimate of Definition 3.28 is applied at points small on level 0 (Findings, Chapter 3, item 215). Formerly "narrower", when only (a)–(c) were proved.

**V3.12 (Proposition 3.10; `Polyfold.proposition_3_10`, `Polyfold.proposition_3_10_scPlus`, `Polyfold.IsSc1GermWith.isScOperator_deriv_zero`, `Polyfold.ScBanach.scIndex_projWOp`, `Polyfold.StabilityData.exists_isFredholm_Dfs'`, `Polyfold.proposition_3_10_level`, `Polyfold.proposition_3_10_full`, `Polyfold.IsBasicGerm.levelDeriv_add_eq`, `Polyfold.IsBasicGerm.snd_`** Let $\widetilde{C}$ be a partial quadrant, $h$ a basic germ, $s$ an sc$^1$-germ into $(Q \oplus W)^1$ and $f = h + \iota s$ an sc$^1$-germ

with derivative family $Df$. Then $Df(0)$ is an sc-operator, sc-Fredholm of index $\dim P - \dim Q$: it is $T_0 + S$ with $T_0(\delta a, \delta w) = (0, \delta w)$ and $S$ an sc$^+$-operator. For the data of Theorem 3.38, $D(h+s)(x) : E_0 \to F_0$ is Fredholm of index $\dim P - \dim Q$ for every $x \in U_1$ small on level 1. On every level (the second statement): for every $m$ there is $\delta > 0$ such that at every $x \in U_{m+1}$ with $|x|_m < \delta$ the level-$m$ operator $Df(x)_m : P \oplus W_m \to Q \oplus W_m$ is Fredholm of index $\dim P - \dim Q$ ($\widetilde{C}$ quadrant-like); $f(0) = 0$ is not used. *Book:* Proposition 3.10, p. 68. *Hidden hypotheses:* the finite-rank terms are sc$^+$ because $Dh(0)$ is an sc-operator at the smooth point 0. *Deviation (stronger):* both statements as in the text, with $s$ and $f$ only sc$^1$ and $f(0) = 0$ unused. The second statement by the text's proof on level $m$: $Df(x)_m = T_0 + S(x) + K(x)$ with $\|S(x)\| \le \varepsilon$ by Lemma 3.9 on level $m$ (V3.11) and $K(x)$ compact; the $W$-component of $Df(x)_m$, which the text uses silently, follows from the uniqueness of derivatives within $U_1$ (Findings, Chapter 3, item 230). Formerly "narrower", the second statement on level 0 only.

**V3.13 (Definition 3.11, sc-Fredholm germs;** `Polyfold.StrongBundle.IsScFredholmGerm`, `Polyfold.IsFillingGerm`, `Polyfold.IsScPlusGerm`, `Polyfold.IsConjugateToBasicGerm`, `Polyfold.IsStrongBundleIso`, `Polyfold.StrongBundle.IsScFredholmGermWholeChart`, `Polyfold.StrongBundle.IsScFredholmGermWholeChart.isScFredholmGerm`, `Polyfold.StrongBundle.isScFre`**)** $(f, x_0)$ is an sc-Fredholm germ if $x_0$ is smooth and in a strong bundle chart of $P$ centred at $x_0$ the principal part has a filling $g$ (V3.9: an sc-smooth germ on a germ of neighbourhoods inside $U$), there is an sc$^+$-germ $s$ (`IsScPlusGerm`: $s : E_0 \to F_1$ an sc-smooth germ into $F^1$ on a germ inside $U$) with $\iota s(0) = g(0)$, and $k = g - \iota s$ is conjugated to a basic germ: strong bundle maps $\Psi : U \lhd F \to U' \lhd F'$ and $\Psi'$ back, inverse to each other, with base map $\psi$, $\psi(0) = 0$, $U'$ relatively open in $\widetilde{C} \oplus W$ ($\widetilde{C}$ a partial quadrant of a finite-dimensional space), a basic germ $h$ on a germ inside $U'$, and $(\Psi(u, k(u)))_2 = h(\psi(u))$ for $u \in U$ near 0. *Book:* Definition 3.11, p. 69. *Hidden hypotheses:* none. *Deviation (restated):* the identity is required for $u \in U$; tameness unused; $C' = T\psi(0)C$ not imposed (it is proved, V3.48). The witnesses of the conjugation are also available as data (`Polyfold.ConjugationData`, V3.46). Since the switch to the text's germ reading (phase 2) the filling and the sc$^+$-section are germs whose levels may shrink with $m$; $\Psi$ on all of $U$ is no restriction (on a relatively open $U_0 \subseteq U$ it is the same after restricting the chart on level 0). The project's earlier notion, with $g$, $s$ sc-smooth on the whole $U$ (`IsScFredholmGermWholeChart`), implies this one and is exactly this one with $g$, $s$ sc-smooth on $U$; whether the two agree for sections sc-smooth on $X$ is open, and no theorem depends on it (Findings, Chapter 3, items 220–221, 261–264, 330). The existential over charts is essential: a filling in a chart gives a topological isomorphism $\ker Dr(0) \cong \ker \rho(0)$, a condition on the chart (`Polyfold.IsFillingGerm.nonempty_kerEquiv`), so a filling in one chart gives none in another unless these normal spaces match; a compatible tame chart of the trivial bundle carries no filling of any section (`Polyfold.TrivialBundle.not_isFillingGerm_evenBundleChart`; Findings, Chapter 3, items 350–352).

**V3.14 (Theorem 3.12;** `Polyfold.theorem_3_12`, `Polyfold.theorem_3_12_local_subset`, `Polyfold.MPolyfold.continuousOn_levelChartSymm`**).** Let $f$ be a section of a strong bundle $P : Y \to X$ ($P \circ f = \mathrm{id}$) with $(f, x_0)$ an sc-Fredholm germ and $f(x_0) = 0$. There are open sets $O(0) \supseteq O(1) \supseteq \cdots$ of $X$ containing $x_0$ such that for every $i$ the closure in $X$ of $\{x \in O(i) \mid f(x) = 0\}$ is the image of a compact subset of $X_i$, with the level topology, under the inclusion $X_i \to X$. *Book:* Theorem 3.12, p. 70 (sc-smooth section of a tame strong bundle). *Hidden hypotheses:* none beyond those used. *Deviation (stronger):* only $P \circ f = \mathrm{id}$ of the section is used; no tameness. *Proof:* the local form (Theorem 3.40 for any subset of the solution set), transported through chart, filler and conjugation, with the inverse chart $O_i \to X_i$ continuous for the level

topology.

**V3.15 (Corollary 3.13; `Polyfold.corollary_3_13`, `Polyfold.corollary_3_13_section`, `Polyfold.tendsto_of_isCompact_of_tendsto_map`).** (a) Under the hypotheses of V3.14, if $f(x_k) = 0$ and $x_k \to x_0$ in $X$, then for every $m$ there are $y_k \in X_m$ with $\iota y_k = x_k$ for $k$ large and $y_k \to y_0$ in $X_m$, $\iota y_0 = x_0$. (b) If $f$ is an sc-Fredholm section and $x_k \to x$ are zeros, then $x$ is a zero and a smooth point, and the same holds at $x$. *Book:* Corollary 3.13, p. 70. *Hidden hypotheses:* the germ property at the limit. *Deviation (restated):* the text's arbitrary limit $x$ is covered for $x = x_0$ and, for sc-Fredholm sections, for every $x$.

**V3.16 (Definition 3.14, sc-Fredholm sections; `Polyfold.StrongBundle.IsScFredholmSection`, `Polyfold.StrongBundle.IsScFredholmSectionWholeChart`, `Polyfold.StrongBundle.IsScFredholmSectionWh`).** $f$ is an sc-Fredholm section of the strong bundle $P$ if it is sc-smooth, regularizing ($x \in X_m$, $f(x) \in Y_{m,m+1}$ imply $x \in X_{m+1}$) and an sc-Fredholm germ (V3.13) at every smooth point. *Book:* Definition 3.14, p. 70. *Hidden hypotheses:* none. *Deviation:* none; the bundle need not be tame. Clause (3) is the text's germ reading since phase 2; the project's earlier whole-chart notion (`IsScFredholmSectionWholeChart`), stricter, implies it (Findings, Chapter 3, items 220–221, 330). Formerly "restated", for the whole-chart reading.

**V3.17 (Theorem 3.15; `Polyfold.theorem_3_15`, `Polyfold.StrongBundle.IsScSmoothSection.sectionAdd`, `Polyfold.StrongBundle.regularizing_sectionAdd`, `Polyfold.StrongBundle.IsScFredholmGerm.sectionAdd`).** If $f$ is an sc-Fredholm section and $s$ an sc$^+$-section of a strong bundle, the fibrewise sum $f + s$ is an sc-Fredholm section. *Book:* Theorem 3.15, p. 71 (tame bundle). *Hidden hypotheses:* the sum rule; the [1]-compatibility of charts (V3.6). *Deviation (stronger):* no tameness. In the germ reading of Definition 3.11 (V3.13), with the text's proof: $\tilde{t} = \hat{s} \circ r$ is sc-smooth on the whole $U$, and a germ plus a whole-chart map is a germ on the same germ (V3.19; Findings, Chapter 3, item 300); the whole-chart form is `Polyfold.theorem_3_15_wholeChart`.

**V3.18 (Lemma 3.16; `Polyfold.lemma_3_16`).** For a section $s$ of $K \to O$: $s(r(x)) = s(x)$ for $x \in O$, and $\rho(r(u))s(r(u)) = s(r(u))$, $R(r(u), s(r(u))) = (r(u), s(r(u)))$ for $u \in U$. *Book:* Lemma 3.16, p. 72. *Hidden hypotheses:* $s$ takes values in $K$. *Deviation (restated):* the sc$^+$-regularity of $s \circ r$ is proved inside Theorem 3.15 (V3.17).

**V3.19 (Lemma 3.17; `Polyfold.IsFillingGerm.add_section_germ`, `Polyfold.IsFillingGerm.add_section`, `Polyfold.lemma_3_17`).** If $g$ is a filling of $\mathbf{f}$ (V3.7) and $s$ a section of $K \to O$, then $g + s \circ r$ is a filling of $\mathbf{f} + s$, with the same derivative family in (3): for $s$ sc-smooth on $U$, on the same germ; for $s$ an sc-smooth germ on $G_s$, on $G \cap r^{-1}(G_s)$. *Book:* Lemma 3.17, p. 72. *Hidden hypotheses:* none. *Deviation (stronger):* the regularity of $g + s \circ r$ is part of the conclusion, and of $s$ only the sc-smoothness is used, not that it is sc$^+$. This is the text's statement for fillings that are germs, proved in the project since the switch to the germ reading (phase 2); the whole-chart `Polyfold.lemma_3_17` (fillers in the sense of `IsFillerOf`, any section $s$, the regularity left to Theorem 3.15) neither implies it nor follows from it (round-trip item V3.19; Findings, Chapter 3, items 220–221, 273). Formerly "restated".

**V3.20 (Definition 3.21, good position; `Polyfold.IsGoodComplement`, `Polyfold.IsInGoodPosition`, `Polyfold.IsScGoodComplement`, `Polyfold.IsInGoodPositionSc`, `Polyfold.IsInGoodPosition.isInGoodPosit`, `Polyfold.isScGoodComplement_iff`, `Polyfold.isInGoodPositionSc_iff`).** A subspace $P$ is a good complement of $N$ with respect to $C$ if $E_0 = N \oplus P$ (algebraically) and for some $\varepsilon > 0$, $n \in N$, $p \in P$, $|p|_0 \le \varepsilon |n|_0$ imply ($n \in C \iff n + p \in C$). $N$ is in good position to $C$ if $N \cap C$ has nonempty interior in $N$ and $N$ has a good complement. Definition 3.21 as written

(`IsScGoodComplement`: $E = N \oplus P$ an sc-splitting with the constant $\varepsilon$; `IsInGoodPositionSc`) implies this; on a partial quadrant a smooth finite-dimensional $N$ in good position is in good position with an sc-complement, also inside any sc-subspace $T \supseteq N$, and for smooth $N$ the text's good complements are the project's closed ones. *Book:* Definition 3.21, pp. 76–77 (an sc-complement, or a topological one). *Hidden hypotheses:* none. *Deviation (stronger):* only the algebraic splitting is required, as a hypothesis. As a conclusion the algebraic form is weaker (an algebraic good complement need not be closed, round-trip item V3.20), so the text's form is formalized as well and proved where the theorems conclude good position (V3.47, V3.94; Findings, Chapter 3, items 210, 211).

**V3.21 (Proposition 3.22; `Polyfold.proposition_3_22`, `Polyfold.exists_quadrantNormalForm`).** If the finite-dimensional subspace $N$ is in good position to the partial quadrant $C$, then $N \cap C$ is a partial quadrant in $N$. More precisely, for $C = \{\ell_i \geq 0\}$ with continuous functionals jointly onto $\mathbb{R}^n$, there is a linear isomorphism $e : N \to \mathbb{R}^k \times N_0$ carrying $N \cap C$ onto $[0,\infty)^k \times N_0$, with $k = \dim N - \dim(N \cap W)$ and the degeneracy counts preserved off $N \cap W$. *Book:* Proposition 3.22, p. 77, proof pp. 120–126. *Hidden hypotheses:* none. *Deviation:* none in the statement. *Proof:* a segment argument and Lemma 3.65 instead of the extreme-ray Lemmas 3.62–3.64, 3.66–3.68.

**V3.22 (Definition 3.25, the tangent germ property; `Polyfold.MPolyfold.TangentGermChart`, `Polyfold.MPolyfold.HasTangentGermProperty`, `Polyfold.IsScInftyGermAt`).** $M \subseteq X$ has the $n$-dimensional tangent germ property if $M \subseteq X_\infty$ and every $x \in M$ has: a chart $\varphi : (U, x) \to (O, 0)$ of $X$ onto a tame retract $(O, C, E)$; an $n$-dimensional smooth $N \subseteq E$ in good position to $C$ with a good complement $Y$ that is **closed**; a relatively open $V \ni 0$ in $N \cap C$; and a continuous $\delta : V \to E$ with values in $Y$, such that $\varphi(M \cap U) = \{v + \delta(v) \mid v \in V\}$, $\delta(0) = 0$, $D\delta(0) = 0$, and for every $m$, $k$ there is a relatively open $V_{m,k} \ni 0$ in $N \cap C$ with $\delta(V_{m,k}) \subseteq E_m$ and $\delta : V_{m,k} \to E_m$ of class $C^k$. *Book:* Definition 3.25, p. 79. *Hidden hypotheses:* $Y$ closed (the text's sc-complement is). *Deviation (narrower):* the added closedness; "sc^∞-germ" read with $k$-dependent neighbourhoods, which the fixed-germ and the set-sense readings imply (`Polyfold.IsScSmoothGerm.isScInftyGermAt`, `Polyfold.IsScSmooth.isScInftyGermAt`). The germ implicit function theorem gives the fixed-germ reading as well (V3.31, V3.32); Theorem 3.43 (1) is proved in the $k$-dependent one (V3.38).

**V3.23 (Proposition 3.26; `Polyfold.proposition_3_26`, `Polyfold.proposition_3_26_image`, `Polyfold.MPolyfold.TangentGermChart.transport`).** Let $\psi : X \to X'$ be an sc-diffeomorphism of M-polyfolds (sc-smooth on all of $X$, with an sc-smooth inverse $\psi'$ on all of $X'$) and $\psi(M) = M'$. Then $M'$ has the $n$-dimensional tangent germ property (V3.22) if and only if $M$ has it; the data at $\psi(x)$ are the chart $\varphi \circ \psi^{-1}$ with the same $N$, $Y$, $V$, $\delta$. *Book:* Proposition 3.26, p. 80. *Hidden hypotheses:* none. *Deviation:* none (the tameness of $X$ is not used). The text's "$X' = \psi(X)$ is tame" is not needed here; that tameness is carried by sc-diffeomorphisms is V3.71.

**V3.24 (Theorem 3.27; `Polyfold.theorem_3_27`, `Polyfold.theorem_3_27_isSubMPolyfold`, `Polyfold.MPolyfold.TangentGermChart.IsRegular`, `Polyfold.MPolyfold.theorem_3_27_isChart`, `Polyfold.MPolyfold.theorem_3_27_isTame`, `Polyfold.MPolyfold.theorem_3_27_isManifoldWithCorners`, `Polyfold.MPolyfold.theorem_3_27_degIndex`).** Let $M \neq \emptyset$ be a subset of an M-polyfold $X$ such that every point of $M$ has data of Definition 3.25 (V3.22) that are *regular*: (a) $\delta$ is sc-smooth on $V$ as a map on a set, and (b) the projection $\pi$ along $Y$ maps $O$ into $C$. Then $M$ is a sub-M-polyfold of $X$; its induced M-polyfold structure (Proposition 2.21) is tame and has an equivalent atlas of charts on relatively open subsets of $n$-dimensional partial quadrants; each map $y \mapsto \pi\varphi(y)$ on $M \cap \varphi^{-1}(\Sigma \cap O)$, modelled on $(V, N \cap C, N)$, is a chart of the induced structure containing $x$; and

$d_M(y) = d_{N\cap C}(\pi\varphi(y))$ there. *Book:* Theorem 3.27, p. 81, proof pp. 81–84. *Hidden hypotheses:* (a) and (b); the text's proof asserts both and each fails in an example (a smooth curve with a vertical tangent written over the horizontal axis; $C = [0,\infty)^3$, $N = \mathbb{R}(1,1,1)$, $Y = \{x+y-z=0\}$). *Deviation (narrower):* the two added conditions; $X$ need not be tame; the relation $d_{N\cap C} = d_X$ via Lemma 3.70 is not transported to a general presentation of $C$. For the solution set of an sc-Fredholm section both conditions are theorems (V3.74, V3.77, V3.85); for any set with the tangent germ property, (a) follows from (b) after shrinking the data (V3.74). With a dimension that may vary from point to point: V3.72.

**V3.25 (Definition 3.28, sc⁰-contraction germs; `Polyfold.IsSc0ContractionGerm`, `Polyfold.ContractionGerm`, `Polyfold.IsSc0ContractionGerm.toContractionGerm`).** For a germ $G$ around 0 in $\widetilde{C}\oplus W$: $B$ is an sc⁰-contraction germ if $B$ is an sc⁰-germ, $B(0,0) = 0$, and for every $\varepsilon > 0$ and $m$ there is $r > 0$ such that the points $(a,w)$ with $a \in \widetilde{C}$, $|a| < r$, $|w|_m < r$ lie in $U_m$ and $|B(a,w) - B(a,w')|_m \le \varepsilon|w-w'|_m$ for such $(a,w)$, $(a,w')$. Every such $B$ has a working form: open $N(0) \supseteq N(1) \supseteq \cdots \ni 0$ in $P$ and $0 \in K(m) \subseteq O(m) \subseteq W_m$ ($K$ closed, $O$ open, $\iota K(m+1) \subseteq K(m)$) with $\{a\}\times O(m) \subseteq U_m$, $B_m(a,\cdot)$ a $\frac{1}{2}$-contraction on $O(m)$ mapping $K(m)$ into itself, for $a \in N(m)\cap\widetilde{C}$. *Book:* Definition 3.28, p. 85. *Hidden hypotheses:* the points lie in $U_m$. *Deviation (restated):* explicit regions, needed for Lemma 3.31.

**V3.26 (Theorem 3.29; `Polyfold.theorem_3_29`, `Polyfold.theorem_3_29_of_isSc0ContractionGerm`, `Polyfold.exists_fixedPoint_continuousOn`).** For a contraction germ in working form with quadrant-like $\widetilde{C}$: the level-0 fixed point $\delta(a)$ of $B(a,\cdot)$ is an sc⁰-germ on $V_m = N(m)\cap\widetilde{C}$ with $\delta(0) = 0$, $(a,\delta_m(a)) \in U_m$ for $a \in V_m$, $B(a,\delta(a)) = \delta(a)$ on $V_0$, and every $w \in O(0)$ with $B(a,w) = w$, $a \in V_0$, equals $\delta(a)$. From Definition 3.28 directly: the same, with uniqueness among small solutions. The parametrized Banach fixed point theorem: for $\frac{1}{2}$-contractions $g_l$ of a nonempty closed $K$ in a complete metric space, continuous in $l$ pointwise, the fixed point is continuous in $l$ and unique in $K$. *Book:* Theorem 3.29, p. 86. *Hidden hypotheses:* the uniqueness region. *Deviation (hypotheses explicit).*

**V3.27 (Theorem 3.30; `Polyfold.theorem_3_30`, `Polyfold.ContractionGerm.isSc1GermWith_sol`).** If $B$ is an sc¹-germ, the solution germ $\delta$ is a solution germ and an sc¹-germ on $V_m = N(m+1)\cap\widetilde{C}$ (the germ shifted by one level), with $D\delta(a) = (1 - D_2B(a,\delta(a)))^{-1}D_1B(a,\delta(a))$. *Book:* Theorem 3.30, p. 86, quoted from [21] without proof. *Hidden hypotheses:* none. *Deviation (restated):* sc¹ on the shifted germ; a germ only because $\widetilde{C}$ is finite-dimensional. *Proof:* supplied; it needs $(a_0,\delta(a_0)) \in U_2$, hence the shift.

**V3.28 (Lemma 3.31; `Polyfold.ContractionGerm.lemma_3_31`, `Polyfold.ContractionGerm.next`, `Polyfold.ContractionGerm.isSc0ContractionGerm_next`).** If $B$ is sc^{k+1} with derivative family $D$ and $\delta$ is sc^j on $(N(m+s)\cap\widetilde{C})_m$, then $f^{(1)}(v,b,u,w) = (u - B(v,u), w - DB(v,\delta(v))(b,w))$, in working form with parameter set $\widetilde{C}\times P$, fibre $TW$ and regions $N(m+2+s)\times P$, $O(m+1)\times W_m$, $K(m+1)\times W_m$, is a contraction germ of class sc^{min(k, j)} whose solution germ is $(v,b)\mapsto(\delta(v), D\delta(v)b)$ on $(N(2+s)\cap\widetilde{C})\times P$. If the working form also satisfies Definition 3.28 and $B$ is sc¹, then $f^{(1)}$ satisfies Definition 3.28 on its germ, for every $s$ (an $\varepsilon$-contraction near 0 for every $\varepsilon > 0$ on every level), by Lemma 3.9 on level $m$ (V3.11). *Book:* Lemma 3.31, pp. 86–87. *Hidden hypotheses:* uniqueness of the fixed point only in a region unbounded in the tangent directions; the shifted solution germ (Proposition 1.17 for germs). *Deviation (restated):* the working form records a $\frac{1}{2}$-contraction on explicit regions, all the proof of Theorem 3.33 uses; the text's "$f^{(1)}$ is an sc⁰-contraction germ" in the sense of Definition 3.28 is proved separately

(Findings, Chapter 3, item 252).

**V3.29 (Lemma 3.32, as an induction; `Polyfold.ContractionGerm.sol_isSckGerm`, `Polyfold.ContractionGerm.`** For every contraction germ over any finite-dimensional parameter space: if $B$ is sc^k and $j \le k$, then $\delta$ is an sc^j-germ on $(N(m+s)\cap\widetilde{C})_m$ for some $s$; the step from $j$ to $j+1$ turns shifts $s$ (for $\delta$) and $s'$ (for $T\delta$) into $s+s'+1$. *Book:* Lemma 3.32, p. 87 (the iterated germs $f^{(j)}$ with solution $T^j\delta$). *Hidden hypotheses:* none. *Deviation (restated):* the $f^{(j)}$ are not constructed; one induction quantified over all contraction germs.

**V3.30 (Theorem 3.33, the germ implicit function theorem; `Polyfold.theorem_3_33`, `Polyfold.theorem_3_33_sm` `Polyfold.theorem_3_33_of_isSc0ContractionGerm`).** If $B$ is an sc^k-germ, there is $s$ such that the solution germ $\delta$ is a solution germ and an sc^k-germ on $(N(m+s)\cap\widetilde{C})_m$; the proof gives $s = 2^k - 1$. If $B$ is sc-smooth, this holds for every $k$ (with $s$ depending on $k$). From Definition 3.28 with balls: there is an sc^k solution germ. *Book:* Theorem 3.33, p. 88. *Hidden hypotheses:* **none; the theorem holds with the text's hypotheses**. *Deviation (restated):* the germ on which $\delta$ is sc^k shrinks with $k$; in the text's up-to-shrinking reading this is the theorem.

**V3.31 (Corollary 3.34; `Polyfold.corollary_3_34`, `Polyfold.corollary_3_34_isScSmoothGerm`, `Polyfold.isSckGerm_of_levels`).** If $B$ is an sc-smooth germ, then for all $m$, $k$ there is a relatively open $V \ni 0$ in $\widetilde{C}$ with $\delta(V) \subseteq W_m$ and $\delta : V \to W_m$ of class $C^k$. The last sentence on one germ: $\delta$ is an sc-smooth germ (sc^k for every $k$ on the same sequence of sets) on a diagonal germ $V_m = N(m+t_m)\cap\widetilde{C}$, $t_m = \sum_{j\le m} s_j$, and a solution germ there. *Book:* Corollary 3.34, p. 88. *Hidden hypotheses:* Proposition 1.18 for germs. *Deviation:* none; its last sentence ("$\delta$ is sc-smooth at 0") holds on one germ: sc^j involves only the levels $\ge j$ of a germ (`Polyfold.isSckGerm_of_levels`), so the $k$-dependent germs of Theorem 3.33 can be replaced by one diagonal germ, which shrinks fast with the level (Findings, Chapter 3, item 250).

**V3.32 (Theorem 3.35; `Polyfold.theorem_3_35`, `Polyfold.theorem_3_35_smooth`, `Polyfold.theorem_3_35_isSc` `Polyfold.theorem_3_35_of_isSc0ContractionGerm`, `Polyfold.isSckGerm_contractionMap_iff`).** For $\widetilde{C}$ a partial quadrant in a finite-dimensional $P$: if $B$ is sc^k, $\delta$ is sc^k on $(N(m+s)\cap\widetilde{C})_m$ for some $s$; if $B$ is sc-smooth, then for every $k$ there is such an $s$, and $\delta$ is an sc-smooth germ on one germ $N(m+t_m)\cap\widetilde{C}$. In the text's form: for an sc$^0$-contraction germ $f = w - B(a,w)$ of Definition 3.28 on a quadrant-like $\widetilde{C}$ and any representative $(V,\delta)$ of its solution germ, if $f$ is sc^k then $\delta$ is sc^k on a germ $V' \subseteq V$, and if $f$ is sc-smooth then $\delta$ is an sc-smooth germ on one germ $V' \subseteq V$. *Book:* Theorem 3.35, p. 89. *Hidden hypotheses:* none. *Deviation (restated):* the conclusion holds on a smaller germ $V' \subseteq V$, as for Theorem 3.33 (V3.30); the hypotheses are on $f$ ($f$ is sc^k iff $B$ is); $\mathbb{R}^n$ is generalized to a quadrant-like $\widetilde{C}$ in a finite-dimensional $P$ (Findings, Chapter 3, items 250, 251). Before the third addendum "sc-smooth" held only for every $k$ on a germ depending on $k$.

**V3.33 (Definition 3.37, the basic class; `Polyfold.IsBasicGerm`).** As V3.10. *Book:* Definition 3.37, p. 90. *Hidden hypotheses:* none. *Deviation:* none.

**V3.34 (Theorem 3.38; `Polyfold.theorem_3_38`, `Polyfold.StabilityData`, `Polyfold.StabilityData.exists_of` `Polyfold.StabilityData.pushMap`, `Polyfold.StabilityData.pushB_isSc0ContractionGerm`, `Polyfold.scPlusRemainder_contraction`, `Polyfold.StabilityData.pushMap_eq_zero_iff`, `Polyfold.StabilityData.finrank_Cs_eq_finrank_Zs`).** Let $f$ be a basic germ on $G$ with contraction $B$, $s$ an sc-smooth germ into $W^1$ with $s(0) = 0$ (the $W$-component of an sc$^+$-germ), and $W = C \oplus X = R \oplus Z$ an sc-Fredholm splitting of $1 + D_2 s(0)$ (it exists, Proposition 1.10).

With $P_1 : W \to X$, $P_2 : W \to R$ the projections and $L = (1 + D_2 s(0))|_X$, the linear maps $\psi(a, w) = ((a, (1 - P_1)w), P_1 w)$ and $\Psi(q, w) = ((q, (1 - P_2)w), L^{-1} P_2 w)$ carry $f + \iota s$ to the germ $\Psi \circ (f + \iota s) \circ \psi^{-1}$ on the index-raised spaces $(P \oplus C) \oplus X^1 \to (Q \oplus Z) \oplus X^1$, with germ $\psi(U_{m+1})$, which is a basic germ with contraction map $L^{-1} P_2 (B - S)$, $S = s - D_2 s(0)$; its zeros correspond to those of $f + \iota s$; and $\dim C = \dim Z$. *Book:* Theorem 3.38, pp. 90–93. *Hidden hypotheses:* the operator-norm continuity of the level derivatives of $s$ (for "$S$ is an arbitrarily small contraction"). *Deviation (restated):* a linear coordinate change; only the $W$-component of $s$; the text's $\tau : Z \to C$ not applied. The coordinate change is a strong bundle isomorphism of the index-raised structures (`Polyfold.StabilityData.linIso_isStrongBundleIso`, V3.76).

**V3.35 (Theorem 3.40; `Polyfold.theorem_3_40_compactness`, `Polyfold.theorem_3_40_regularity`, `Polyfold.IsSc0ContractionGerm.exists_lift`, `Polyfold.theorem_3_12_local`).** Let $B$ be an sc$^0$-contraction germ on $G$ around 0 in $\widetilde{C} \oplus W$ with $\widetilde{C}$ **closed**, $\mathbf{s}$ an sc$^0$-germ into $W^1$ with $\mathbf{s}(0) = 0$, and $S$ any set of solutions in $U_0$ of $w = B(a, w) - \iota \mathbf{s}(a, w)$. There are open $O(0) \supseteq O(1) \supseteq \cdots \ni 0$ in $E_0$ and compact $K_i \subseteq E_i$ with $\mathrm{cl}_{E_0}(S \cap O(i)) = \iota_i(K_i)$; in particular these closures lie in $E_i$. For a basic germ $h$ and an sc$^0$-germ $s$ into $(Q \oplus W)^1$ with $s(0) = 0$, the closures of $\{h + \iota s = 0\} \cap O(i)$ are compact subsets of $E_i$. *Book:* Theorem 3.40, p. 94, proof pp. 95–98. *Hidden hypotheses:* $\widetilde{C}$ closed (automatic for a partial quadrant). *Deviation (stronger):* compact closures; $B$ only sc$^0$ with the contraction estimate, $\mathbf{s}$ only sc$^0$ into $W^1$.

**V3.36 (Corollary 3.41; `Polyfold.StabilityData.corollary_3_41`, `Polyfold.StabilityData.exists_isFredho` `Polyfold.StabilityData.exists_ker_Dfs'_inf_eq_bot`, `Polyfold.isFredholm_add_of_norm_lt`, `Polyfold.HasFiniteIndex.surjective_of_finrank_ker_le`).** For the data of V3.34 with $D(f + \iota s)(0)$ surjective: there is $r > 0$ such that every zero $x$ of $f + \iota s$ in $U_0$ with $|x|_0 < r$ is $\iota(x_1)$ for some $x_1 \in U_1$, and $D(f + \iota s)(x_1) : E_0 \to F_0$ is a surjective Fredholm operator of index $\dim P - \dim Q$. *Book:* Corollary 3.41, p. 98. *Hidden hypotheses:* none. *Deviation (restated):* the $\mathbb{R}^N$-component of the sc$^+$-map is absorbed into the basic germ. *Proof:* the Fredholm property holds at every $x \in U_1$ small on level 1 (small plus compact perturbation of $D(f + \iota s)(0)$); surjectivity from $\dim \ker \leq \mathrm{ind}$.

**V3.37 (Lemma 3.42; `Polyfold.StabilityData.lemma_3_42`).** For the push-forward $g = \Phi_*(f + \iota s)$ of V3.34 (on level 0): for every $\varepsilon > 0$ there is $r > 0$ such that every zero $x'$ of $g$ in $\psi(U_0)$ with $|x'|_0 < r$ has its $X$-component $\iota(w_1)$ with $w_1 \in X_1$, $|w_1|_1 < \varepsilon$. *Book:* Lemma 3.42, p. 101. *Hidden hypotheses:* none. *Deviation (restated):* the form Theorem 3.43 uses; $w' = \delta(a')$ is derived there.

**V3.38 (Theorem 3.43; `Polyfold.theorem_3_43_sc_smooth`, `Polyfold.theorem_3_43`, `Polyfold.ShiftedBasicDa` `Polyfold.ShiftedBasicData.Sol.exists_contDiffOn_level`, `Polyfold.exists_proj_mem_of_eq_zero`).** Let $\widetilde{C}$ be a partial quadrant, $g$ an sc-smooth germ on $G$ with $g(0) = 0$ whose index-raised germ is a basic germ, with $Dg(0)$ surjective, $K' = \ker Dg(0)$ in good position to $\widetilde{C} \oplus W$, and such that for every $\varepsilon > 0$ there is $r > 0$ with every zero $x$ of $g$ in $U_0$, $|x|_0 < r$, having $W$-component $\iota(w_1)$, $|w_1|_1 < \varepsilon$. Then there are a good complement $Y$ of $K'$, a relatively open $V \ni 0$ in $K' \cap (\widetilde{C} \oplus W)$ and $\tau : K' \to P \times W_0$ with values in $Y$ and $W$-components on level 1, $C^1$ on $V$, $\tau(0) = 0$, $D\tau(0) = 0$, such that $y + \tau(y) \in U_0$ and $g(y + \tau(y)) = 0$ for $y \in V$; for every $k \geq 1$ a relatively open $V_k \subseteq V$ containing 0 and a $C^k$ map $\tau_k : V_k \to E_k$ with $\iota_k \tau_k = \tau$ on $V_k$; and an open $U'' \ni 0$ in which every zero of $g$ in $U_0$ is $y + \tau(y)$ with $y \in V$. *Book:* Theorem 3.43, p. 102. *Hidden hypotheses:* the regularity of Lemma 3.42 (made a hypothesis); the projection of a solution to $\widetilde{K}$ lies in $\widetilde{C}$ (a gap in the text, proved); that the solutions given by Theorem 3.78 at the different regularities $k$ coincide near 0 (uniqueness clause). *Deviation (restated):* clause (1), "$\tau$ is an

sc-smooth germ", in the level-wise sense with $k$-dependent neighbourhoods. With $Y$ closed and its projection onto $K'$ mapping $\widetilde{C} \oplus W$ into itself: V3.77.

**V3.39 (Lemma 3.44; `Polyfold.ShiftedBasicData.Deriv.ker_Dg_eq`).** In the setting of V3.38, $\ker Dg(0) = \{(\alpha, D\delta(0)\alpha) \mid \alpha \in \ker DH(0)\}$, $H(a) = \mathrm{pr}_Q\, g(a, \delta(a))$. *Book:* Lemma 3.44, p. 103. *Hidden hypotheses:* $D_2B(0) = 0$ on level 0, by density from level 1. *Deviation:* none.

**V3.40 (Lemma 3.45; `Polyfold.IsGoodComplement.map`, `Polyfold.IsInGoodPosition.map`).** A continuous linear isomorphism $S : E_0 \to E_0'$ with $x \in C \iff Sx \in C'$ carries a good complement $P$ of $N$ to a good complement $S(P)$ of $S(N)$ (constant $\varepsilon/(\|S\|\|S^{-1}\| + 1)$), and good position to good position. *Book:* Lemma 3.45, p. 104 (for $T\varphi(0)$). *Hidden hypotheses:* none. *Deviation (stronger):* any such $S$.

**V3.41 (Lemma 3.46; `Polyfold.lemma_3_46`, `Polyfold.lemma_3_46_isInGoodPosition`).** If $K' = \{(\alpha, L\alpha) \mid \alpha \in \widetilde{K}\}$ has a good complement with respect to $\widetilde{C} \oplus W$, then $\widetilde{K}$ has a good complement $Z$ with respect to $\widetilde{C}$ such that $Z \times W$ is a good complement of $K'$; good position of $K'$ gives good position of $\widetilde{K}$. *Book:* Lemma 3.46, p. 104, proof pp. 129–132. *Hidden hypotheses:* none. *Deviation:* none (the proof avoids the intermediate complement $Y_0$).

**V3.42 (Corollary 3.47; `Polyfold.theorem_3_47_sc_smooth`, `Polyfold.theorem_3_47`).** For $\widetilde{C}$ a partial quadrant, a basic germ $f$ and the data of V3.34 with $D(f + \iota s)(0)$ surjective and kernel $K$ in good position to $\widetilde{C} \oplus W$: there are a good complement $Y$ of $K$, a relatively open $V \ni 0$ in $K \cap (\widetilde{C} \oplus W)$, and $\sigma$ with values in $Y$ and $W$-components on level 1, $C^1$ on $V$, $\sigma(0) = 0$, $D\sigma(0) = 0$, with $v + \sigma(v) \in U_0$ a zero for $v \in V$; for every $k \geq 1$ a $C^k$ lift $\sigma_k : V_k \to E_k$ of $\sigma$ on a relatively open $V_k \subseteq V$ containing 0; and an open $U'' \ni 0$ in which every zero in $U_0$ is of this form. *Book:* Corollary 3.47, p. 106. *Hidden hypotheses:* none. *Deviation (restated):* (1) level-wise, as in V3.38. *Proof:* Theorem 3.43 for the push-forward, transported back linearly; Lemma 3.70 and Theorem 3.72 are not needed. With $Y$ closed and its projection onto $K$ mapping $\widetilde{C} \oplus W$ into itself: V3.77.

**V3.43 (Definition 3.48; `Polyfold.IsFillingGerm`, `Polyfold.IsFillerOf`).** As V3.7. *Book:* Definition 3.48, p. 109. *Hidden hypotheses:* none. *Deviation:* none. Since the switch to the text's germ reading (phase 2) the filling is the germ notion of V3.7; the stricter whole-chart filler (`IsFillerOf`, with `IsFilledVersion`), for which "filled version = filling" on a given $U$ is false (round-trip item V3.43), is kept (Findings, Chapter 3, items 220–221, 330). Formerly "narrower", for the whole-chart reading.

**V3.44 (Remark 3.49; `Polyfold.IsFillerOf.remark_3_49`).** For a filler there is an open $W \ni 0$ with $f = g$ on $O \cap W$ and, for $y \in U \cap W$, $g(y) = \rho(r(y))g(y)$ only if $y \in O$. *Book:* Remark 3.49, p. 109. *Hidden hypotheses:* none. *Deviation:* none.

**V3.45 (Definitions 3.50, 3.51; `Polyfold.StrongBundle.IsScFredholmGerm`, `Polyfold.StrongBundle.IsScFredh`
`Polyfold.StrongBundle.IsScFredholmGermTame`, `Polyfold.StrongBundle.IsScFredholmSectionTame`,
`Polyfold.StrongBundle.IsScFredholmGermTameWholeChart`).** As V3.13 and V3.16, with the chart of the germ data tame: V3.103. *Book:* Definitions 3.50, 3.51, p. 110 (for a tame strong bundle; the chart of Definition 3.50 has a tame local strong bundle $K \to O$). *Hidden hypotheses:* none. *Deviation:* none. The forms with a tame chart (V3.103) are the ones the theorems of §3.6 use; since the switch to the text's germ reading (phase 2) they take the filling and the sc$^+$-section as germs, as the text does, and the earlier whole-chart forms (…`WholeChart`), which imply them (round-trip

item V3.45), are kept (Findings, Chapter 3, items 220–221, 330). The forms of V3.13 and V3.16 drop the tameness, which §3.1 does not need. Formerly "narrower", for the whole-chart reading.

**V3.46 (Proposition 3.52; `Polyfold.proposition_3_52_index`, `Polyfold.proposition_3_52_local_index`, `Polyfold.ConjugationData`, `Polyfold.ConjugationData.scDeriv_isFredholm_index`, `Polyfold.ConjugationData.index_eq`, `Polyfold.proposition_3_52`, `Polyfold.proposition_3_52_local`, `Polyfold.proposition_3_52_sc`, `Polyfold.proposition_3_52_local_sc`, `Polyfold.ConjugationData.scDeri` `Polyfold.StrongBundle.linAt_isScFredholm`).** Let $f$ be a section of a strong bundle with $(f, x_0)$ an sc-Fredholm germ and $f(x_0) = 0$. In the strong bundle chart of Definition 3.11 centred at $x_0$, with the filling germ $g$ (a derivative family $Dg$ on a germ inside $U$) and the local sc⁺-germ $s$ for which $g - s$ is conjugated to a basic germ, $Dg(0) : E_0 \to F_0$ and $Tf(x_0) := Dg(0)|_{Dr(0)E_0} : Dr(0)E_0 \to \rho(0)F_0$ are Fredholm operators, and for **every** conjugation of $g - s$ to a basic germ $h : O(\widetilde{C} \oplus W, 0) \to \mathbb{R}^N \oplus W$, $\widetilde{C} \subseteq \mathbb{R}^n$, both have index $n - N$ (`ConjugationData`: the witnesses of a conjugation, Definition 3.11, as data). In particular $n - N$ is the same for all conjugations of one germ: for $k$ sc-smooth with $k(0) = 0$ and any conjugation of $k$ to a basic germ, $Dk(0)$ is Fredholm of index $n - N$. They are sc-Fredholm operators: $Dg(0) : E \to F$ is an sc-operator, $T_0O = Dr(0)E$ and $K_0 = \rho(0)F$ are sc-subspaces, and $Tf(x_0) : T_0O \to K_0$ is sc-Fredholm, both of index $n - N$ on every level; for an sc-Fredholm section the same holds in every strong bundle chart around a zero $y$, at the point $z$ over $y$, for $\mathrm{Lin}(z) : T_zO \to K_z$. *Book:* Proposition 3.52, p. 110 ($Tf(x) : T_xX \to Y_x$ is sc-Fredholm); the index $n - N$ is used in the proof of Theorem 3.54. *Hidden hypotheses:* none. *Deviation (restated):* in charts ($T_xX$ carries no linear structure here, Chapter 2). The hypothesis is the text's since the switch to the germ reading (phase 2): the filling and the sc⁺-section are germs (V3.13), and $Dg(0)$ is the value at 0 of any derivative family of the filling germ; the forms for germ data on the whole chart are the `_wholeChart` declarations of the same names (Findings, Chapter 3, items 220–221, 289, 310, 331). Formerly "narrower", for that hypothesis. The sc-Fredholm property, first proved on level 0 only, follows the text's argument with two steps it leaves out: $D(g - s)(0) = \Gamma(0)^{-1} Dh(0) D\psi(0)$ is sc-Fredholm by the criterion of V1.8, and $T_0O$, $K_0$ are sc-subspaces (Definition 2.47 at the smooth point 0, the closed graph theorem); in other charts the levels are transported by the chart change at smooth points (Findings, Chapter 3, items 235, 239). The first version (`Polyfold.proposition_3_52`, kept for its consumers) states the index as "$n - N$ for some natural numbers $n$, $N$", which carries no information; `proposition_3_52_index` identifies $n$ and $N$ with the dimensions of the conjugating basic germ (Findings, Chapter 3, item 120). In every other strong bundle chart around $x_0$ the linearization is Fredholm of the same index (V3.96).

**V3.47 (Definition 3.53; `Polyfold.IsGoodPositionAtGerm`, `Polyfold.IsGoodPositionAt`, `Polyfold.IsInGoodPositionIn`, `Polyfold.StrongBundle.IsGoodPositionGerm`, `Polyfold.StrongBundle.IsGo` `Polyfold.IsScGoodComplementIn`, `Polyfold.IsInGoodPositionScIn`, `Polyfold.IsGoodPositionAtScGerm`, `Polyfold.IsGoodPositionAtSc`, `Polyfold.StrongBundle.IsGoodPositionGermSc`, `Polyfold.StrongBundle.Is` `Polyfold.StrongBundle.isGoodPositionGermSc_iff`).** In a chart centred at the zero, with the filling germ $g$ and $Dg(0)$ the value at 0 of a derivative family of $g$ (the same for all): the germ is in good position if $Tf(0) = Dg(0)|_{T_0O} : T_0O \to K_0$ is onto and $\ker Dg(0)$ is in good position to $T_0O \cap C$ inside $T_0O$ (interior of $\ker Dg(0) \cap C$ in $\ker Dg(0)$ nonempty, a good complement inside $T_0O$). For a section, $(f, x_0)$ is in good position if it has the data of Definition 3.11 at $x_0$ with the germ in good position in that chart. With the text's sc-complement (`IsGoodPositionAtSc`; for a section `IsGoodPositionGermSc`, in a tame chart) the kernel has a good sc-complement inside $T_0O$ ($T_m = N_m \oplus P_m$ on every level); at a zero of an sc-Fredholm section with tame germ data

the two forms agree. *Book:* Definition 3.53, p. 111. *Hidden hypotheses:* none. *Deviation (restated):* in a chart rather than in $T_xX$; no case distinction $d_X(x) \geq 1$, which is recovered as a theorem for tame germ data: at a zero with $d_X = 0$ good position in a tame chart is the surjectivity of the linearization (V3.100). At a zero the linearization and the two conditions do not depend on the chart (V3.78); Definition 3.53 at a zero from a chart not centred there is V3.79, and in any chart of the germ data V3.107. The form in a tame chart, which the theorems of §3.6 assume and conclude, is V3.103. The germ data are the text's since the switch to the germ reading (phase 2, V3.13); the whole-chart forms are `IsGoodPositionGermWholeChart`, `IsGoodPositionGermScWholeChart` (Findings, Chapter 3, items 220–221, 280, 320, 330); that "an sc-splitting of $T_0O$", written in $E$, is an sc-splitting of the sc-Banach space $T_0O$ is not formalized (item 214).

**V3.48 (Theorem 3.54, the graph description of the solution set; `Polyfold.theorem_3_54_graph`, `Polyfold.theorem_3_54_linear`, `Polyfold.theorem_3_54_pushforward`, `Polyfold.scDeriv_mem_iff_of_inve`**
Let $f$ be a section of a strong bundle, with $f(x_0) = 0$ and $(f, x_0)$ in good position in a tame chart (V3.103). Then there are: a tame strong bundle chart $\varphi$ centred at $x_0$ (compatible with the atlas, with tame base retraction); an sc-smooth $\psi$ on its $U$ with $\psi(0) = 0$ into a relatively open $U'$ of $\widetilde{C} \oplus W$ ($\widetilde{C}$ a partial quadrant), with an sc-smooth inverse $\psi'$ on $U'$; the data of Theorem 3.38 and a basic germ $h'$ with $D(h' + \iota t)(0)$ surjective and kernel $K$ in good position, $K$ a smooth subspace; a closed good complement $Y$ of $K$; a relatively open $V' \ni 0$ in $K \cap (\widetilde{C} \oplus W)$; $\sigma$ with values in $Y$ and $W$-components on level 1, $C^1$ on $V'$, $\sigma(0) = 0$, $D\sigma(0) = 0$, such that every $v + \sigma(v)$, $v \in V'$, is a zero of $h' + \iota t$ in its germ, and for every $k \geq 1$ a $C^k$-lift $\sigma_k : V_k \to E'_k$ of $\sigma$ on a relatively open $V_k \ni 0$, $V_k \subseteq V'$ (Theorem 3.47 (1)); and an open $V_x \ni x_0$ in the chart domain on which $\psi\varphi$ is injective, $f(y) = 0 \iff (h' + \iota t)(\psi\varphi(y)) = 0$, and $\psi\varphi(\{y \in V_x \mid f(y) = 0\}) = \{v + \sigma(v) \mid v \in V'\}$. On the way: in the chart, $Dg(0)$ is surjective with kernel in good position to $C$ in $E$, $x \in C \iff D\psi(0)x \in C'$, and $D(\Psi_* g)(0) = \Gamma(0)Dg(0)D\psi(0)^{-1}$ is surjective Fredholm with kernel in good position. *Book:* Theorem 3.54, p. 111, the step (3) of its proof, pp. 113–114. *Hidden hypotheses:* the tameness of the chart at $x_0$, used only as $(1 - Dr(0))E \subseteq E_0$; the neighbourhood $V$ made explicit; the kernel is smooth (the text: without comment). *Deviation (narrower):* only the graph description (3) of the proof; conclusions (1) and (2) of the theorem are V3.82 and V3.85, the choice of $Y$ with the projection property V3.77.

**V3.49 (Lemma 3.55; `Polyfold.lemma_3_55`, `Polyfold.lemma_3_55_core`).** In $E = \mathbb{R}^n \oplus W$, $C = [0, \infty)^n \oplus W$: let $N$ be finite-dimensional in good position with good complement $Y$, $Y$ **closed**, and $\tau : N \to E$ with values in $Y$, $C^1$ on $N \cap C$, $\tau(0) = 0$, $D\tau(0) = 0$. There is $\rho > 0$ such that for $v \in N \cap C$ with $|v| < \rho$: $v + \tau(v) \in C$, and $Y$ is a good complement of $N_v = \{n + D\tau(v)n \mid n \in N\}$ with respect to the partial quadrant $C_{v+\tau(v)}$ (so $E_0 = N_v \oplus Y$, both closed, with the good-complement constant independent of $v$). The core holds for any $C$ cut out by coordinate functionals. *Book:* Lemma 3.55, p. 114, proof pp. 132–135. *Hidden hypotheses:* $Y$ closed (needed for $D\tau(v)n \in Y$). *Deviation (narrower):* the added closedness; $\tau$ given on all of $N \cap C$. The form used in Theorem 3.54 — for a relatively open $V \subseteq N \cap C$ and any presentation of $C$, with the good position of $N_v$ — is `Polyfold.lemma_3_55_relOpen` (V3.80).

**V3.50 (Theorem 3.56 from the tangent germ property; `Polyfold.theorem_3_56_of_tangentGerm`).** If $X$ has no boundary and the nonempty zero set $S$ of $f$ has at every point regular data of Definition 3.25 (V3.24), then $S$ is a sub-M-polyfold whose induced structure is an $n$-dimensional manifold with corners without boundary. *Book:* Theorem 3.56, p. 115. *Hidden hypotheses:* the regularity conditions of V3.24. *Deviation (narrower):* assumes the tangent germ property of $S$, which the text derives from Theorem 3.54. The theorem with the text's hypotheses is V3.86.

**V3.51 (Theorem 3.57 from the tangent germ property; `Polyfold.theorem_3_57_of_tangentGerm`).** If the nonempty zero set $S$ of $f$ has at every point regular data of Definition 3.25, then $S$ is a sub-M-polyfold whose induced structure is tame and an $n$-dimensional manifold with boundary with corners. *Book:* Theorem 3.57, p. 115. *Hidden hypotheses:* as V3.50. *Deviation (narrower):* as V3.50. The theorem with the text's hypotheses is V3.87.

**V3.52 (Lemma 3.59, abstract form; `Polyfold.lemma_3_59`, `Polyfold.FamilyCompactness`).** Let $b \mapsto L(b)$ be a family of sc-operators $X \to Y$, $b$ in a finite-dimensional space, with $(b,h) \mapsto L(b)h$ continuous on $O \times X_0$ for a set $O \ni 0$, $L(0)$ injective on $X_0$, and the compactness property on level 0: if $b_j \to b$ in $O$, $(h_j)$ is bounded in $X_0$ and $L(b_j)h_j \to 0$, then $(h_j)$ has a convergent subsequence. Then $L(b)$ is injective on $X_0$ for $b \in O$ near 0. *Book:* Lemma 3.59, p. 117 (for $L(b) = P_Y \circ Df(b)|_X$ in the setting of Theorem 3.58). *Hidden hypotheses:* none. *Deviation (restated):* for an abstract family; in the proof of Theorem 3.58 its hypotheses are derived from those of the theorem for $L(b) = P_Y \circ Df(b,0)|_X$ (`Polyfold.familyCompactness_derivFamOp`, `Polyfold.continuousOn_derivFamOp`, V3.89).

**V3.53 (Lemma 3.60, abstract form; `Polyfold.lemma_3_60`).** For a compact set $K$ of parameters, a family of sc-operators $L(b)$ with $(b,h) \mapsto L(b)_m h$ continuous on $K \times X_m$, every $L(b)_m$, $b \in K$, injective, and the compactness property on level $m$: there is $c_m > 0$ with $c_m|h|_m \le |L(b)h|_m$ for all $b \in K$, $h \in X_m$. *Book:* Lemma 3.60, p. 118. *Hidden hypotheses:* none. *Deviation (restated):* abstract form; the injectivity on level $m$ is a hypothesis, which in Theorem 3.58 comes from level 0 (`Polyfold.ScOperator.injective_lvl`, V3.89).

**V3.54 (Lemma 3.61; `Polyfold.lemma_3_61`).** Let $V$ be a germ of relatively open neighbourhoods of 0 in a partial quadrant $\widetilde{C}$ of a finite-dimensional $G$, and $L(v) : E \to F$ sc-isomorphisms such that $(v,h) \mapsto L(v)h$ is an sc-smooth germ on $V \oplus E$ (levels $V_m \times E_m$) and for every $m$ there is $c_m > 0$ with $c_m|h|_m \le |L(v)h|_m$ for $v \in V_0$, $h \in E_m$. Then $(v,k) \mapsto L(v)^{-1}k$ is an sc-smooth germ on $V \oplus F$. *Book:* Lemma 3.61, p. 118, quoted from [25] without proof. *Hidden hypotheses:* none (no operator-norm continuity of $v \mapsto L(v)$). *Deviation (restated):* for germs. *Proof:* supplied; the differentiability uses the compactness of $F_1 \to F_0$. Used in Theorem 3.58 in set form (`Polyfold.famInv_isScSmooth_of`, V3.88).

**V3.55 (Lemma 3.65; `Polyfold.lemma_3_65`).** For functionals $\ell_i$ cutting out $C$, a good complement $P$ of $N$ (constant $\varepsilon$) satisfies $\ell_i(p) = 0$ for $p \in P$ and every $i$ vanishing at some nonzero point of $N \cap C$. *Book:* Lemma 3.65, p. 123. *Hidden hypotheses:* none. *Deviation:* none.

**V3.56 (Proposition 3.69; `Polyfold.proposition_3_69`).** For $N$ finite-dimensional in good position to $C = [0,\infty)^n \oplus W$ in $\mathbb{R}^n \oplus W$ and $x \in N \cap C$: $d_{N\cap C}(x) = d_C(x)$ if $x \notin N \cap W$, and $d_{N\cap C}(x) + \dim(N \cap W) = \dim N$ if $x \in N \cap W$. *Book:* Proposition 3.69, p. 126. *Hidden hypotheses:* none. *Deviation (restated):* (2) additively.

**V3.57 (Lemma 3.70; `Polyfold.lemma_3_70`).** For $N$ in good position to $C = [0,\infty)^n \oplus W$ with good complement $Y$ and $\tau : N \to E$ with values in $Y$, $C^1$ on $N \cap C$, $\tau(0) = 0$, $D\tau(0) = 0$: there is $\rho > 0$ such that for $v \in N \cap C$, $|v| < \rho$, $v + \tau(v) \in C$, and $d_{N\cap C}(v) = d_C(v + \tau(v))$ if $v \notin N \cap W$. *Book:* Lemma 3.70, p. 127. *Hidden hypotheses:* none. *Deviation (hypotheses explicit):* "close to 0" made explicit.

**V3.58 (Theorem 3.72; `Polyfold.theorem_3_72`).** For $C$ a partial quadrant in a finite-dimensional $V$, $U$ a relatively open neighbourhood of 0 in $C$ and $f$ of class $C^j$, $j \ge 1$, on $U$ with $f(U) \subseteq C$, $f(0) = 0$, $Df(0)C = C$ and $d_C(x) = d_C(f(x))$ on $U$: there are relatively open $U' \subseteq U$ and $V'$ containing 0 with $f(U') = V'$, $f$ injective on $U'$, and an inverse $g$ of class $C^j$ on $V'$. *Book:*

Theorem 3.72, p. 136 (smooth). *Hidden hypotheses:* none. *Deviation (stronger):* $C^j$; the degeneracy hypothesis used only as "interior iff interior".

**V3.59 (Proposition 3.73; `Polyfold.proposition_3_73`, `Polyfold.InvData`).** For $U$ relatively open and convex in $C$, $f : U \to C$ of class $C^j$, $j \geq 1$, $\|Df - 1\| \leq \frac{1}{2}$ on $U$, and $x$ interior iff $f(x)$ interior: $f(U)$ is relatively open in $C$, $f$ is injective on $U$, with an inverse of class $C^j$ on $f(U)$. *Book:* Proposition 3.73, p. 137. *Hidden hypotheses:* none. *Deviation (stronger):* $0 \in U$, $f(0) = 0$ not needed; (3) weakened. With the hypotheses its proof uses, without the partial quadrant: V3.90.

**V3.60 (Lemma 3.74; `Polyfold.InvData.injOn`, `Polyfold.InvData.ball_inter_subset_image`, `Polyfold.InvData.image_relOpen`).** Under the hypotheses of V3.59: $\frac{1}{2}|x-x'| \leq |f(x)-f(x')| \leq \frac{3}{2}|x - x'|$, $f$ is injective, $B(x', 3r) \cap C \subseteq U$ implies $B(f(x'), r) \cap C \subseteq f(U)$, and $f(U)$ is relatively open in $C$. *Book:* Lemma 3.74, p. 137. *Hidden hypotheses:* the interior starting point with $f(x_0) \in B(y', r)$. *Deviation:* none in the statement (openness of $f$ on subsets not stated separately).

**V3.61 (Corollary 3.75; `Polyfold.corollary_3_75`).** Under the hypotheses of V3.59, $f : U \to f(U)$ is a homeomorphism. *Book:* Corollary 3.75, p. 139. *Hidden hypotheses:* none. *Deviation:* none.

**V3.62 (Lemma 3.77; `Polyfold.InvData.hasFDerivWithinAt_inv`).** Under the hypotheses of V3.59 the inverse is differentiable within $f(U)$ with $Dg(y) = [Df(g(y))]^{-1}$. *Book:* Lemma 3.77, p. 140 (for any homeomorphism $f : U \to V$ with $V$ relatively open and $Df$ invertible). *Hidden hypotheses:* none. *Deviation (narrower):* under the hypotheses of Proposition 3.73, which imply the lemma's. With the hypotheses its proof uses, without the partial quadrant: V3.90.

**V3.63 (Theorem 3.78; `Polyfold.theorem_3_78`, `Polyfold.ContractionData.exists_solution`).** For a partial quadrant $C$ in a finite-dimensional $V$, $U$ a relatively open neighbourhood of 0 in $C$, $f : U \to F$ of class $C^j$, $j \geq 1$, $f(0) = 0$, $Df(0)$ onto, $K = \ker Df(0)$ in good position with good complement $Y$: there are a relatively open $V' \ni 0$ in $K \cap C$, $\tau : K \to Y$ of class $C^j$ on $V'$, $\varepsilon, \sigma > 0$ with (1) $\tau(0) = 0$, $D\tau(0) = 0$, $k + \tau(k) \in U$ and $f(k + \tau(k)) = 0$ for $k \in V'$; (2) $|\tau(k)| \leq \sigma$ for $k \in V'$, $|k| \leq \varepsilon$; (3) $k \in V'$, $|k| \leq \varepsilon$, $y \in Y$, $|y| \leq \sigma$, $k + y \in U$, $f(k + y) = 0$ imply $y = \tau(k)$. *Book:* Theorem 3.78, p. 141. *Hidden hypotheses:* $f(k + \tau(k)) = 0$ and $k + y \in U$, omitted in the text. *Deviation (stronger):* the two clauses; $F$ any normed space; only convexity and interior of $C$ used.

**V3.64 (Lemma 3.79; `Polyfold.ContractionData.exists_radii`, `Polyfold.Radii.eq_` ).** In the normalized data there are $\varepsilon, a, b > 0$, $\varepsilon a \leq b$, with the region $R = \{|k| \leq a, |y| \leq b\} \cap C \subseteq U$ and $\|DB\| \leq \varepsilon/(2(1 + \varepsilon))$ on $R$; a fixed point $y$ of $B(k, \cdot)$ with $(k, y) \in R$ is $\tau(k)$. *Book:* Lemma 3.79, p. 142. *Hidden hypotheses:* none. *Deviation:* none.

**V3.65 (Lemma 3.80; `Polyfold.Radii.mapsTo_ball`).** $B(k, \cdot)$ maps $\{|y| \leq \varepsilon|k|\}$ into itself for $k \in K \cap C$, $|k| < a$, and has a fixed point there. *Book:* Lemma 3.80, p. 143. *Hidden hypotheses:* the maximum is over $B \cap C$. *Deviation (hypotheses explicit).*

**V3.66 (Lemma 3.81; `Polyfold.Radii.norm_ _sub_le`, `Polyfold.Radii.norm_ _sub_le_global`, `Polyfold.Radii.norm_ _sub_le_eps`, `Polyfold.Radii.lipschitzOnWith_` ).** $|\tau(k)| \leq \varepsilon|k|/2$, $\tau$ is continuous, and $|\tau(k) - \tau(k_0)| \leq \frac{\varepsilon}{1+\varepsilon}|k - k_0| \leq \varepsilon|k - k_0|$ for all $k, k_0 \in V_0$ (an older form with the constant $1 + 2\varepsilon$ is kept for the proofs that use it). *Book:* Lemma 3.81, p. 144 (constant $\varepsilon$). *Hidden hypotheses:* none. *Deviation (stronger):* the sharper constant $\varepsilon/(1 + \varepsilon)$. The text's argument uses the point $(k, \tau(k_0))$, which lies in the region of the contraction when $|k_0| \leq 2|k|$; otherwise $|k| < 2|k_0|$ and the same argument with $k$ and $k_0$ exchanged applies, a case distinction the text omits (Findings, Chapter 3, item 216). Formerly "narrower", with the constant $1 + 2\varepsilon$.

**V3.67 (Lemma 3.82; `Polyfold.Radii.contDiffOn_` ).** $\tau$ is differentiable on $V_0 = \{k \in K \cap C : |k| < a\}$ with $D\tau(k) = (1 - D_2B(k,\tau(k)))^{-1}D_1B(k,\tau(k))$ and of class $C^j$ on $V_0$. *Book:* Lemma 3.82, p. 144 ("on a possibly smaller neighbourhood"). *Hidden hypotheses:* none. *Deviation (stronger):* no shrinking.

**V3.68 (Theorem 3.19, the implicit function theorem, interior case, first form; `Polyfold.theorem_3_19`, `Polyfold.theorem_3_19_linearization`, `Polyfold.MPolyfold.IsWithoutBoundary`, `Polyfold.theorem_3_19_` `Polyfold.StrongBundle.linAt_isScFredholm`).** Let $X$ be an M-polyfold with $d_X \equiv 0$, $f$ an sc-Fredholm section (V3.16) of a strong bundle $P : Y \to X$ with tame germ data (V3.103), $f(x_0) = 0$, and let $(f, x_0)$ be in good position in a tame chart (V3.47, V3.103) **with a filler that satisfies condition (3) of Definition 3.4 at the zeros of the filler near** $0$ (V3.82). Then there are an open $U \ni x_0$ and $d \in \mathbb{N}$ such that every zero $y \in U$ is a smooth point with $(f, y)$ in good position in a tame chart (in particular $f'(y)$ is onto), and $S(f,U) = \{y \in U \mid f(y) = 0\}$ is a sub-M-polyfold of $X$ whose induced structure (Proposition 2.21) is equivalent to that of a $d$-dimensional manifold with boundary with corners and has degeneracy index $0$ at every point. The first clause: for an sc-Fredholm section and a zero $x_0$ in good position, in the chart of Definition 3.11 centred at $x_0$, $Dg(0)$ and $f'(x_0) = Tf(x_0) = Dg(0)|_{T_0O} : T_0O \to K_0$ are sc-Fredholm operators with the same index $n - N$ on every level, and $Tf(x_0)$ is onto on every level (`Polyfold.theorem_3_19_linearization_sc`; Proposition 3.52 in sc form, V3.46); with no chart singled out, in every strong bundle chart around $x_0$ the linearization $\mathrm{Lin}(z) : T_zO \to K_z$ is an sc-Fredholm operator with the same index on every level (`Polyfold.StrongBundle.linAt_isScFredholm`, V3.96). *Book:* Theorem 3.19, p. 74 ("$f'(x_0)$ surjective"; the local solution set is a manifold of dimension the Fredholm index, $f'(y)$ onto and $\ker f'(y) = T_yS(f,U)$). *Hidden hypotheses:* tameness of the bundle, read as Definitions 3.50, 3.51 and 3.53 read it: germ data and good position in charts with tame base retraction (V3.103; the reading "every chart compatible with the atlas has a tame base retraction", under which this item was first proved, cannot be met, V3.104). *Deviation (narrower):* (i) the filler condition at the nearby zeros, through Theorem 3.54 (1) (V3.82), which was an added hypothesis when this form was proved and is now a theorem: it holds whenever $(f, x_0)$ is in good position in a tame chart (V3.102), so this form has the text's hypotheses after all, and `Polyfold.theorem_3_19'` states it without the condition (V3.95); (ii) the hypothesis "$f'(x_0)$ surjective" is read as good position (Definition 3.53 in the chart form), which at $d_X = 0$ is the same (V3.100; the form with "$f'(x_0)$ surjective" is V3.101); (iii) $d$ is existentially quantified here; "the dimension agrees with the Fredholm index" is V3.97; (iv) "$\ker f'(y) = T_yS$" in the chart is V3.99; (v) $f'(x_0)$ in charts (V3.46); its sc-Fredholm property, first proved only on level 0, holds on every level. *Proof:* Theorem 3.23 (V3.69) and Lemma 2.26 ($d_S \le d_X = 0$).

**V3.69 (Theorem 3.23, the implicit function theorem, boundary case; `Polyfold.theorem_3_23_of_nearby`, `Polyfold.theorem_3_23`, `Polyfold.MPolyfold.subPolyfold`, `Polyfold.MPolyfold.IsManifoldWithCorners`).** Let $f$ be an sc-Fredholm section of a strong bundle, $f(x) = 0$, $(f, x)$ in good position in a tame chart (V3.47, V3.103), and let every zero in an open $V_{x_0} \ni x$ be in good position in a tame chart. Then there are an open $U$ with $x \in U \subseteq V_{x_0}$ and $d \in \mathbb{N}$ such that every zero in $U$ is a smooth point and $S(f,U) = \{y \in U \mid f(y) = 0\}$ is a sub-M-polyfold of $X$ whose induced structure (Proposition 2.21) is tame and equivalent to that of a $d$-dimensional manifold with boundary with corners (an equivalent atlas of charts on relatively open subsets of $d$-dimensional partial quadrants). For $f$ with tame germ data (V3.103) and the filler condition of V3.82 at $x$ in place of the nearby good position (`Polyfold.theorem_3_23`): the same, and every zero in $U$ is in good position in a tame chart. *Book:* Theorem 3.23, p. 77 ($f'(x)$ surjective with kernel in good position; $S(f,U)$ consists of

smooth points and is a tame sub-M-polyfold with a uniquely determined structure of a manifold with boundary with corners, the induced one). *Hidden hypotheses:* tameness as in V3.68. *Deviation (narrower):* the good position of the nearby zeros (Theorem 3.54 (1)) is a hypothesis of the first form, the filler condition at the nearby zeros (V3.82) of the second; both are now theorems (V3.94, V3.102), and the form with the text's hypotheses is V3.95, with the dimension as the Fredholm index V3.97; $d$ is existentially quantified here; "uniquely determined" is the equivalence of atlases, not stated separately. *Proof:* regular data of Definition 3.25 at every point of $S(f, U)$, of the same dimension by the local constancy (V3.85, V3.74), then Theorem 3.27 (V3.24).

**V3.70 (Corollary 3.24;** `Polyfold.corollary_3_24`, `Polyfold.corollary_3_24_sc`, `Polyfold.forall_isGoodPos`**)** Let $f$ be an sc-Fredholm section of a strong bundle such that $(f, x)$ is in good position in a tame chart (V3.103) at every zero $x$, and let $S = \{x \in X \mid f(x) = 0\}$ be nonempty. Then $S$ is a sub-M-polyfold of $X$ whose induced structure is tame and equivalent to that of a manifold with boundary with corners of locally constant dimension (V3.72). The same with good position in the text's sense (sc-complement, V3.47) at every zero, which for an sc-Fredholm section with tame germ data is the same hypothesis (`corollary_3_24_sc`). *Book:* Corollary 3.24, p. 78. *Hidden hypotheses:* tameness as in V3.68; $S \neq \emptyset$ (for $S = \emptyset$ the formal charts have no values; the statement is then empty). *Deviation (hypotheses explicit):* **no hypothesis added**; the text names no dimension, the formal one is locally constant. It is Theorem 3.57 (V3.87) without its last clause.

**V3.71 (tameness is carried by sc-diffeomorphisms and by open subsets;** `Polyfold.IsScDiffeoQuadrant`, `Polyfold.IsScDiffeoQuadrant.transport`, `Polyfold.IsScDiffeoQuadrant.scDeriv_transport`, `Polyfold.IsScSubspace.map`, `Polyfold.IsScSplitting.map`, `Polyfold.IsScDiffeoQuadrant.isTame_transpo` `Polyfold.IsScDiffeoQuadrant.isTameRetract_image`, `Polyfold.ScRetraction.restrict_isTame`, `Polyfold.IsTameRetract.inter_isOpen`, `Polyfold.MChart.isTameRetract_image`**).** Let $\psi : U \to U'$ be an sc-diffeomorphism between relatively open subsets of partial quadrants ($\psi$ sc-smooth on $U$, $\psi'$ sc-smooth on $U'$, mapping $U$ and $U'$ into each other, inverse to each other) and $r : U \to U$ a tame sc-smooth retraction onto $O$ whose domain is $U$. Then $r' = \psi \circ r \circ \psi'$ is a tame retraction of $U'$ onto $\psi(O)$, so $(\psi(O), C', E')$ is a tame retract; on the way $d_{C'}(\psi(x)) = d_C(x)$ (Theorem 1.25), $Dr'(y) = D\psi(x)Dr(x)D\psi'(y)$ at the level-1 points $y = \psi(x)$ with $x \in O$, and an sc-operator with an sc-operator inverse carries sc-subspaces and sc-splittings. The restriction of a tame retraction to $U \cap r^{-1}(W)$, $W$ open, is tame; an open subset $O \cap W$ of a tame retract is a tame retract; the model $\varphi(s)$ of a chart restricted to an open $s$ of its domain is tame if the model of $\varphi$ is. *Book:* used without statement in the proof of Proposition 3.26 ("$X' = \psi(X)$ is tame", p. 80) and in the proof of Theorem 3.54 (the chart $\psi \circ \varphi$ of a tame M-polyfold, p. 114). *Hidden hypotheses:* condition (2) of Definition 2.34 is transported at smooth points, where $D\psi(x)$ is an sc-isomorphism, with $D\psi(x)E_x \subseteq E'_{\psi(x)}$ (the one-sided sign argument). *Deviation (not in the book).*

**V3.72 (Theorem 3.27 with locally constant dimension;** `Polyfold.theorem_3_27_loc`, `Polyfold.MPolyfold.IsManifoldWithCornersLoc`, `Polyfold.MPolyfold.theorem_3_27_loc_isSubMPolyfold`, `Polyfold.MPolyfold.theorem_3_27_loc_isTame`, `Polyfold.MPolyfold.theorem_3_27_loc_isManifoldWithCo` `Polyfold.MPolyfold.theorem_3_27_loc_atlas_equiv`**).** Let $M \neq \emptyset$ be a subset of an M-polyfold $X$ such that every point of $M$ has regular data of Definition 3.25 (V3.22, conditions (a), (b) of V3.24) of some dimension, which may depend on the point. Then $M$ is a sub-M-polyfold of $X$, its induced structure is tame, and the charts $y \mapsto \pi\varphi(y)$ of all these data form an atlas equivalent to the induced one, consisting of charts on relatively open subsets of partial quadrants of finite-dimensional spaces (`IsManifoldWithCornersLoc`: a manifold with boundary with corners of locally constant dimension); each of these charts is a chart of the induced structure containing

its point. *Book:* Theorem 3.27, p. 81 (for a fixed dimension $n$). *Hidden hypotheses:* (a), (b) of V3.24. *Deviation (narrower):* the conditions (a), (b); the dimension may vary from point to point (same proof as V3.24).

**V3.73 (the re-centring lemma; `Polyfold.recentre_contDiffOn_levels`, `Polyfold.finrank_le_of_interior_i` `Polyfold.exists_contDiffOn_inverse`).** Let $\chi_1$, $\chi_2$ be charts of an M-polyfold $X$ and $S \subseteq X$. Suppose that $\chi_1(S \cap V_{x_1}) = \{v + \sigma_1(v) \mid v \in V_1\}$ for an open $V_{x_1}$ in the domain of $\chi_1$, where $K_1$ is a finite-dimensional smooth subspace in good position to $C_1$ with a closed complement $Y_1$, $V_1$ is relatively open in $K_1 \cap C_1$, and $\sigma_1$ has values in $Y_1$ and is of class $C^1$ on all of $V_1$; that $\chi_2$ is centred at a point $y \in V_{x_1} \cap V_{x_2}$ and $\chi_2(S \cap V_{x_2})$ is the graph of $\sigma_2$ over a relatively open $V_2 \ni 0$ in $K_2 \cap C_2$ (the same conditions on $K_2$, $Y_2$), with $\sigma_2$ continuous, $\sigma_2(0) = 0$, $D\sigma_2(0) = 0$ and $C^k$-lifts $\sigma_{2,k} : V_{2,k} \to E_{2,k}$ on relatively open neighbourhoods of 0 for every $k \geq 1$; and that $\chi_1(y) = v_0 + \sigma_1(v_0)$, $v_0 \in V_1$. Then $\dim K_1 = \dim K_2$, and for all $m$, $k$ there is a relatively open $B' \ni v_0$ in $K_1 \cap C_1$, $B' \subseteq V_1$, on which $\sigma_1$ takes values in $E_{1,m}$ and is of class $C^k$ into $E_{1,m}$. On the way: a map differentiable on a set whose image has nonempty interior does not lower the dimension (Sard's lemma in fixed dimension); and a $C^k$ map between relatively open subsets of convex sets with nonempty interior in finite-dimensional spaces, with a continuous left inverse on its relatively open image and an invertible derivative at a point, has a $C^k$ inverse near the image of that point. *Book:* the step "$\delta$ is an sc$^\infty$-germ near every point $v_0 \in V$" of the proof of Theorem 3.27, p. 84. *Hidden hypotheses:* $\sigma_1$ differentiable at $v_0$ (here $C^1$ on $V_1$; without it the step fails, V3.24); $Y_1$, $Y_2$ closed; the lifts at 0 (Theorem 3.47 (1), V3.42). *Deviation (narrower):* for two graph descriptions of one set, with $\sigma_1$ of class $C^1$ on $V_1$. *Proof:* the linearization of $\beta = \pi_1 \circ \chi_1\chi_2^{-1} \circ (1 + \sigma_2)$ at 0 is injective by the chain rule for the transition and its inverse on level 1 and the tangency of the graph of $\sigma_2$ to the retract; it is onto by the dimension count, not by the symmetric chain rule (which would need $\sigma_1$ differentiable into level 1 at $v_0$); the inverse is $C^k$ by V3.90, not by Theorem 3.72 (which would need $d_C(x) = d_C(\beta(x))$).

**V3.74 (condition (a) of `IsRegular` by re-centring; `Polyfold.MPolyfold.TangentGermChart.exists_isRegular_` `Polyfold.MPolyfold.TangentGermChart.shrink`, `Polyfold.MPolyfold.TangentGermChart.exists_shrinkDat` `Polyfold.MPolyfold.TangentGermChart.shrink_isRegular`, `Polyfold.MPolyfold.TangentGermChart.exists`).** Let $D$ be data of Definition 3.25 (V3.22) of dimension $n$ for $M \subseteq X$ at $x$ with $\pi(O) \subseteq C$ (condition (b) of V3.24), and let every point of $M$ in an open $U \ni x$ have data of Definition 3.25 of some dimension. Then $M$ has at $x$ regular data (conditions (a) and (b) of V3.24) of dimension $n$ whose chart domain lies in $U$, and every datum of Definition 3.25 for $M$ at a point of $M$ in that domain has dimension $n$. (The proof shrinks $D$ — the chart restricted to an open subset, $V$ to a relatively open $V'$ inside the neighbourhood of 0 on which $\delta$ is $C^1$, the same $N$, $Y$, $\delta$ — and applies V3.73 at every $v_0 \in V'$; the statement exports only the existence of the regular data.) *Book:* the proof of Theorem 3.27, p. 84; the proof of Theorem 3.54, p. 115 ("proceeding as above"). *Hidden hypotheses:* the data at the nearby points. *Deviation (narrower):* condition (b) is a hypothesis. Read off the statement (not a separate Lean theorem): every point of a set with the $n$-dimensional tangent germ property has data, so *for such a set, data at $x$ that satisfy (b) can be shrunk to regular data*; condition (a) of V3.24 follows from (b). For the solution set of an sc-Fredholm section (b) holds by the choice of the good complement (V3.75, V3.84).

**V3.75 (good complements whose projection maps the partial quadrant into itself; `Polyfold.ProjectsInto`, `Polyfold.exists_goodComplement_projectsInto`, `Polyfold.exists_proj_goodComplement`, `Polyfold.exists_proj_quadCone`, `Polyfold.ProjectsInto.mapL`).** `ProjectsInto C N Y` means: $z \in C$, $k \in N$, $z - k \in Y$ imply $k \in C$; for $E_0 = N \oplus Y$ the projection onto $N$ along $Y$ maps $C$ into $N \cap C$. If the finite-dimensional subspace $N$ is in good position to the partial quadrant

$C$ of the sc-Banach space $E$, there is a continuous projection $\pi : E_0 \to N$ with $\pi(C) \subseteq N \cap C$ whose kernel $Y$ is a good complement of $N$; $Y$ is closed and `ProjectsInto C N Y` holds. The property is carried by continuous linear isomorphisms $S$ with $x \in C \iff Sx \in C'$. *Book:* not stated; the proof of Theorem 3.27 (p. 83) asserts that $\Sigma = \{v + y \in C \mid v \in V, y \in Y\}$ is relatively open in $C$, which fails for an arbitrary good complement (V3.24) and holds when $\pi(O) \subseteq C$. *Hidden hypotheses:* none. *Deviation (not in the book):* the construction keeps the coordinates of a set $S$ of indices; when no coordinate vanishes at a nonzero point of $N \cap C$ and $N \cap C \neq N$, one coordinate is kept (the natural choice $S = \Sigma$ fails there: for $C = [0,\infty)^2$, $N = \mathbb{R}(1,1)$, the projection along the good complement $\mathbb{R}(1,2)$ maps $(0,1)$ to $-(1,1)$).

**V3.76 (Corollary 3.39; `Polyfold.corollary_3_39`, `Polyfold.corollary_3_39_isConjugate`, `Polyfold.ConjugationData.compIso_eq_strongShift`, `Polyfold.IsStrongBundleIso.shift`, `Polyfold.IsStrongBundleIso.comp`, `Polyfold.IsStrongBundleIso.pushforward_comp`, `Polyfold.StabilityData.linIso_isStrongBundleIso`, `Polyfold.ScOperator.isStrongBundleIso_prodMap`).** Let $U$ be relatively open in a partial quadrant of $E$ with $0 \in U$, $g : U \to F$ sc-smooth and conjugated to a basic germ (V3.13), and $s : U \to F_1$ sc-smooth into $F^1$ with $s(0) = 0$. Then the level-1 map $(g + \iota s)^1 : U_1 \to F_1$ is conjugated to a basic germ on the shifted structures $E^1 \lhd F^1$. Explicitly, for a conjugation $\Phi$ of $g$ to the basic germ $h$, with $t = \Phi[1]_* s$ ($t(0) = 0$) and $h' = h + (t_{\mathbb{R}^N}, 0)$, there are the data of Theorem 3.38 (V3.34) for $t_W$ such that: $h'$ is a basic germ and $\Phi_*(g + s) = h + t$ near 0; the composite $\Gamma = \Lambda \circ \Phi^1$ of the index raise of $\Phi$ and the linear coordinate change $\Lambda$ of Theorem 3.38 is a strong bundle isomorphism of the shifted structures onto a relatively open subset of $(\widetilde{C} \oplus C) \oplus X^1$, with base map vanishing at 0 and $\Gamma_* = \Lambda_* \circ \Phi^1_*$; and $(\Gamma(u, (g+s)^1(u)))_2 = (\Lambda_*(h+t))^1(\gamma(u))$ for $u \in U_1$ near 0, a basic germ. The text's form "$\Gamma = \Psi \circ \Phi$, then raise the index" is the same map (`compIso_eq_strongShift`). On the way: a pair of sc-isomorphisms, the index raise of a strong bundle isomorphism and a composition of strong bundle isomorphisms are strong bundle isomorphisms, and push-forwards compose. *Book:* Corollary 3.39, p. 93. *Hidden hypotheses:* the index raise of a strong bundle isomorphism is one (used silently). *Deviation (restated):* "sc-germ" read as sc-smooth on $U$, $s$ as an sc$^+$-map on all of $U$ with $s(0) = 0$; the conclusion is a conjugation of $(g+s)^1$ on the shifted structures; the text's "strong bundle map $\Psi$" is an isomorphism; the $\mathbb{R}^N$-component of $t$ is absorbed into $h$; the text's $\tau : Z \to C$ is not applied.

**V3.77 (Theorems 3.43, 3.47 and 3.54 (3) with a closed good complement whose projection maps the quadrant into itself; `Polyfold.theorem_3_43_closed`, `Polyfold.theorem_3_43_proj`, `Polyfold.theorem_3_47_proj`, `Polyfold.theorem_3_47_closed`, `Polyfold.theorem_3_54_graph_proj`, `Polyfold.IsGoodComplement.prod_top_graph`, `Polyfold.ShiftedBasicData.exists_sol_of_goodComplement`).** The conclusions of V3.38, V3.42 and V3.48 hold with a good complement $Y$ that is closed and satisfies `ProjectsInto` (V3.75) for the partial quadrant ($\widetilde{C} \oplus W$, resp. $C' = \widetilde{C} \oplus W$): the projection onto the kernel along $Y$ maps the partial quadrant into the kernel's cone. In Theorem 3.43, $Y = Z \oplus W$ for a good complement $Z$ of $\widetilde{K}$ chosen by V3.75: every good complement $Z$ of $\widetilde{K}$ gives the good complement $Z \oplus W$ of $K' = \{(\alpha, L\alpha)\}$ (constant $\varepsilon/(\|L\| + 1)$), the data of the proof exist for a prescribed $Z$, and the projection property passes from $(\widetilde{K}, Z)$ to $(K', Z \oplus W)$; in Corollary 3.47 both properties are carried by the linear homeomorphism of Theorem 3.38; in Theorem 3.54 (3) the shrinking of $V'$ leaves $Y$ unchanged. *Book:* Theorem 3.43, p. 102; Corollary 3.47, p. 106; Theorem 3.54, step (3) of the proof, p. 114. *Hidden hypotheses:* none. *Deviation (stronger):* two additional conclusions; the closedness is the text's (its $Y'$ is an sc-complement), the projection property is not in the text and is what makes condition (b) of Theorem 3.27 hold for the solution set (V3.84).

**V3.78 (the linearization at a zero in charts, and its chart independence; `Polyfold.linAt`, `Polyfold.rho_scDeriv_comp_r`, `Polyfold.linAt_eq_scDeriv`, `Polyfold.IsFilledVersion.fillerRestrict_e` `Polyfold.linAt_eq_germDeriv`, `Polyfold.IsFillingGerm.fillerRestrict_eq_linAt`, `Polyfold.linAt_chart_change`, `Polyfold.StrongBundle.linAt_chart_independence`, `Polyfold.IsInGoodPositionIn.transport`).** In a strong bundle chart (retraction $r$ on $U$ onto $O$, fibre projections $\rho$, principal part **f**), $\mathrm{Lin}(z) = D(\mathbf{f}\circ r)(z)|_{T_zO}$, $T_zO = Dr(z)E$, at every level-1 point $z$ of the retract. At a zero its values lie in $K_z = \rho(z)F$; it equals $Dg(z)$ on $T_zO$ for every $g$ sc¹ on $U$ agreeing with **f** on $O$ near $z$, and for every sc¹-germ $g$ agreeing with **f** near $z$ with $z$ in its germ ($Dg(z)$ the value of its derivative family; Findings, Chapter 3, item 284); at a centred zero it is the $Tf(0)$ of Proposition 3.5 (V3.8), also for a filling germ. For strong bundle charts $c$, $c'$ whose transition is sc-smooth on $[0]$, a section with sc-smooth principal parts in both, and a zero $y$ in both domains with level-1 points $z$, $z'$ over $\varphi(y)$, $\varphi'(y)$: there are topological linear isomorphisms $A : T_zO \to T_{z'}O'$ and $\Gamma : K_z \to K'_{z'}$ with $\Gamma\circ\mathrm{Lin}(z) = \mathrm{Lin}'(z')\circ A$ and, on $T_zO$, $x \in C_z \iff Ax \in C'_{z'}$ (the partial quadrants at the two points). Hence for two strong bundle charts of $P$ and an sc-smooth section, $\mathrm{Lin}(z)$ and $\mathrm{Lin}'(z')$ are simultaneously onto, simultaneously Fredholm with the same index, and their kernels are simultaneously in good position to the partial quadrants inside $T_zO$, $T_{z'}O'$ (good position inside a subspace is carried by isomorphisms respecting the cones). *Book:* Definition 3.53 (p. 111) presupposes it; Definition 2.38 (p. 42) asserts the chart independence of $T^R_xX$, $C_xX$ by Proposition 2.30; nothing is said about $f'(x)$. *Hidden hypotheses:* the point is a zero (away from zeros the linearization is not chart independent). *Deviation (not in the book):* the chart form of what gap G.13 leaves open; of the compatibility of Definition 2.50 only the sc-smoothness of the transition on $[0]$ is used; level-1 points suffice; no tameness.

**V3.79 (Definition 3.53 at a zero, from a chart not centred there; `Polyfold.StrongBundle.isGoodPositionGermTa` `Polyfold.StrongBundle.isGoodPositionGerm_of_chart`, `Polyfold.StrongBundle.isGoodPositionGerm_of_l` `Polyfold.StrongBundle.isGoodPositionAtGerm_of_chart_dense`).** Let $f$ be an sc-smooth section, $y$ a zero with tame germ data (V3.103), $c$ a strong bundle chart of $P$ with $y$ in its domain (not necessarily centred), $z$ a level-1 point over $\varphi(y)$, and $g$ an sc¹-germ on a germ $G$ of $c$ with derivative family $Dg$ and $z \in G_1$, agreeing with **f** on $O$ near $\varphi(y)$. If the range of $Dg(z)|_{T_zO}$ contains $\rho(z)k$ for every smooth $k \in F_\infty$, and $\ker Dg(z)\cap T_zO$ is in good position to $C_{\varphi(y)}$ inside $T_zO$, then $(f, y)$ is in good position in a tame chart (V3.47, V3.103). Variants: with all of $K_z$ in the range, for an sc-Fredholm germ, in the chart centred at $y$ of Definition 3.11; with the hypotheses on $\mathrm{Lin}(z)$ (V3.78). *Book:* the proof of Theorem 3.54, p. 115 ("proceeding as above", at a nearby zero). *Hidden hypotheses:* the kernel hypothesis is on $\ker Dg(z)\cap T_zO$, not on $\ker Dg(z)$ (condition (3) of the filler holds at its centre only); $z$ lies in the germ of $g$. *Deviation (not in the book):* a dense part of $K_z$ in the range suffices, since the range of the linearization in the chart of the germ data is closed (Proposition 3.52, V3.46), which is why the dense form is stated for tame germ data. In the germ reading since phase 2; the whole-chart forms are `Polyfold.StrongBundle.isGoodPositionGerm_of_chart_dense`, `Polyfold.StrongBundle.isGoodPositionGerm_of_linAt` and the `_wholeChart` declarations (Findings, Chapter 3, items 285, 301, 313).

**V3.80 (property (4) of the proof of Theorem 3.54 in the chart at $x_0$: Corollary 3.41 and Lemma 3.55 on the graph; `Polyfold.theorem_3_54_nearby`, `Polyfold.lemma_3_55_relOpen`, `Polyfold.StabilityData.exists_surjective_Dfs'_graph`).** In the data of V3.48, for a section $f$ of a strong bundle with $f(x_0) = 0$ and $(f, x_0)$ in good position in a tame chart (V3.103), there are a presentation $T$ of $C' = \widetilde{C}\oplus W$ and an open $V_{x,1} \ni x_0$ such that for every zero $y \in V_{x,1}$, with $\psi\varphi(y) = v + \sigma(v)$, $v \in V'$: $v + \sigma(v)$ lies on level 1 and $D(h + t)$ there is a surjective Fredholm

operator of index $n - N$ ($n = \dim \mathbb{R}^n$, $N = \dim \mathbb{R}^N$ of the basic germ); $v + \sigma(v) \in C'$; and $Y$ is a good complement of $N_v = \{k + D\sigma(v)k \mid k \in K\}$ with respect to the partial quadrant $C'_{v+\sigma(v)}$, and $N_v$ is in good position to it. Lemma 3.55 is proved in the form used: for $N$ in good position to $C$ (any presentation) with closed good complement $Y$, $V$ relatively open in $N \cap C$, $\sigma : V \to Y$ of class $C^1$ on $V$ with $\sigma(0) = 0$, $D\sigma(0) = 0$, for $v \in V$ near 0: $v + \sigma(v) \in C$, $Y$ is a good complement of $N_v$ with respect to $C_{v+\sigma(v)}$, and $N_v$ is in good position to it. *Book:* the proof of Theorem 3.54, property (4), p. 114; Lemma 3.55, p. 114. *Hidden hypotheses:* $Y$ closed (V3.49); that $N_v \cap C_z$ has interior in $N_v$, which Lemma 3.55 does not state. *Deviation (narrower):* in the chart at $x_0$, for the tangent space $N_v$ of the graph; its identification with the kernel at $y$ is V3.81.

**V3.81 (Theorem 3.54 (1) at the nearby zeros, the kernel;** `Polyfold.theorem_3_54_nearby_kernel, Polyfold.theorem_3_54_chart_kernel, Polyfold.StabilityData.graphSpace_eq_ker_Dfs', Polyfold.IsStrongBundleIso.ker_scDeriv_eq_map, Polyfold.scDeriv_mem_partialQuadrantAtWith_iff, Polyfold.IsInGoodPosition.isInGoodPositionIn_of_le, Polyfold.IsStrongBundleIso.ker_deriv_eq_map_`
Let $f$ be a section of a strong bundle, $f(x_0) = 0$, $(f, x_0)$ in good position in a tame chart (V3.103). In a tame chart centred at $x_0$ (filling germ $g$ with a derivative family $Dg$, conjugation covering $\psi$ with inverse $\psi'$, the germ $h + t$ with kernel $K$, closed good complement, graph $\sigma : V' \to Y$) there is an open $V_{x,2} \ni x_0$ such that for every zero $y \in V_{x,2}$, with $z_0 = \varphi(y)$, $\psi(z_0) = v + \sigma(v)$ and level-1 points $z_{0,1}$ over $z_0$, which lies in the germ of $g$, $x_1$ over $v + \sigma(v)$: $\mathbf{f} = g$ on $O$ near $z_0$ and $g(z_0) = 0$; $D(h + t)(x_1)$ is a surjective Fredholm operator of index $n - N$; $N_v = \ker D(h + t)(x_1)$; $D(\Psi_* g)(x_1) = D(h + t)(x_1)$ and $\ker Dg(z_{0,1}) = D\psi'(x_1)(N_v)$; $Dg(z_{0,1})$ is onto; $\ker Dg(z_{0,1}) \subseteq T_{z_0}O$; and $\ker Dg(z_{0,1})$ is in good position to $C_{z_0}$ inside $T_{z_0}O$. *Book:* Theorem 3.54 (1), the kernel half, "from Corollary 3.41 and Lemma 3.55", p. 114. *Hidden hypotheses:* the $C^1$-lift of $\sigma$ into level 1 (Theorem 3.47 (1)) for $N_v \subseteq \ker D(h+t)$; $\ker Dg(z_0) \subseteq T_{z_0}O$ comes from the tangency of the solution set to $O$, not from the filler (whose condition (3) holds at 0 only). *Deviation (narrower):* in the chart at $x_0$; **no added hypothesis**; the surjectivity of $f'(y)$ is not part of it (V3.93, V3.94). In the germ reading $z_{0,1}$ lies in the germ of the filling because the basic germ is taken on the points that $\psi'$ maps into the germs of $g$ and $s$ (Findings, Chapter 3, items 276–278, 295).

**V3.82 (Theorem 3.54 (1) under the filler condition at the nearby zeros;** `Polyfold.theorem_3_54_one, Polyfold.StrongBundle.IsGoodPositionGermFilled, Polyfold.exists_range_deriv_eq_ _of_normal_injec`
`IsGoodPositionGermFilled` at $x_0$: $x_0$ is smooth and there are the data of Definition 3.53 at $x_0$ in a tame chart (V3.47, V3.103: a centred chart with tame base retraction, a filling germ $g$ with a derivative family $Dg$ on a germ $G$, a local sc$^+$-germ $s$ with $g - s$ conjugated to a basic germ, the germ in good position) and an open $N \ni 0$ such that at every level-1 point $z_1 \in G_1$ with $z_1 \in O \cap N$ and $g(z_1) = 0$, $Dr(z_1)b = 0$ and $\rho(z_1)Dg(z_1)b = Dg(z_1)b$ imply $b = 0$: condition (3) of Definition 3.4, the injectivity of $(1-\rho)Dg$ on $\ker Dr$, at the zeros of $g$ near 0. Theorem: if $f$ is an sc-Fredholm section with tame germ data (V3.103), $f(x_0) = 0$, and $(f, x_0)$ is in good position in this strengthened sense, there is an open $V_x \ni x_0$ such that $(f, y)$ is in good position in a tame chart (V3.47, V3.103) at every zero $y \in V_x$. The step it needs: at a zero $z$ of a filler with $Dg(z)$ onto and condition (3) at $z$, $\rho(z)F \subseteq Dg(z)(T_zO)$ (Proposition 3.5 (1) at $z$). *Book:* Theorem 3.54 (1), p. 111, proof p. 114. *Hidden hypotheses:* the filler condition at the nearby zeros, added when this form was proved; it is a consequence of good position (V3.102), so the form has the text's hypotheses after all. *Deviation (narrower):* the condition is a hypothesis of the statement. Definition 3.4 imposes (3) at 0 only (so does the 2017 version, Definition 3.13 there), and then the text's argument does not give the surjectivity of $f'(y)$: with respect to $E = T_{z_0}O \oplus \ker Dr(z_0)$, $F = K_{z_0} \oplus \ker \rho(z_0)$, $Dg(z_0)$ is block triangular and $Tf(z_0)$ is onto iff the block $(1 - \rho(z_0))Dg(z_0)|_{\ker Dr(z_0)}$ is injective,

which is the local constancy of the index along the solution set (Findings, Chapter 3, items 100, 157); both directions are formalized (V3.102). The surjectivity holds with Definition 3.4 as stated, by an implicit function theorem at the nearby zero (V3.93), and Theorem 3.54 (1) with the text's hypotheses is V3.94.

**V3.83 (the parameter extension of the proof of Proposition 6.2: the obstruction and the tilted kernel; `Polyfold.mem_lineality_of_goodComplement_prod`, `Polyfold.sup_lineality_eq_top_of_goodComplem` `Polyfold.isInGoodPosition_extendedKernel`).** (a) Let $C \subseteq V$ be closed under nonnegative scaling with $0 \in C$ and $N \subseteq V$ a subspace. For every good complement of $\mathbb{R} \times N$ with respect to $\mathbb{R} \times C$, the $V$-components of its vectors lie in the lineality space $C \cap (-C)$; so $\mathbb{R} \times N$ is in good position only if $N + (C \cap -C) = V$. (b) Let $K$, $Y'$ be subspaces of $\mathbb{R}^n \oplus W$ and $j : \Lambda \to \mathbb{R}^n \oplus W$ linear, $\Lambda$ finite-dimensional, with $K \cap j(\Lambda) = 0$, $(K + j(\Lambda)) \oplus Y' = \mathbb{R}^n \oplus W$, $Y'$ in the lineality space of $\widetilde{C} \oplus W$ ($v + y \in \widetilde{C} \oplus W \iff v \in \widetilde{C} \oplus W$ for $y \in Y'$), and $K + j(\Lambda)$ meeting the interior of $\widetilde{C} \oplus W$. Then $\{((\lambda, a), w) \mid (a, w) + j(\lambda) \in K\}$ is in good position to $(\Lambda \times \widetilde{C}) \oplus W$. *Book:* not in Chapter 3; the linear algebra of the perturbation argument of the text's Proposition 6.2, tried for the surjectivity of V3.82. *Hidden hypotheses:* none. *Deviation (not in the book):* the rest of the argument is not formalized; it would give the surjectivity only on a neighbourhood depending on $\Lambda$ (Findings, Chapter 3, item 106). The surjectivity is proved by another argument, with one perturbation direction at a time and the implicit function theorem at the nearby zero (V3.93).

**V3.84 (Theorem 3.54 (2) at the centre: the data of Definition 3.25 for the solution set at $x_0$; `Polyfold.theorem_3_54_tangentGermChart_proj`, `Polyfold.theorem_3_54_tangentGermChart`, `Polyfold.MChart.compDiffeo`, `Polyfold.MChart.compDiffeo_isChart`, `Polyfold.isScInftyGermAt_of_lifts` `Polyfold.StabilityData.ker_Dfs_le_smooth`, `Polyfold.MPolyfold.TangentGermChart. _mem_of_projectsI`**
Let $f$ be a section of a strong bundle, $f(x_0) = 0$, $(f, x_0)$ in good position in a tame chart (V3.103). Then there are an open $V_x \ni x_0$ and $d \in \mathbb{N}$ such that $\{y \in V_x \mid f(y) = 0\}$ has at $x_0$ data of Definition 3.25 of dimension $d$ (V3.22) with $\pi(O) \subseteq C$ (condition (b) of V3.24). The data: the chart $\psi \circ \varphi$ onto $\psi(O)$ (for a chart $\varphi$ of $X$ and an sc-diffeomorphism $\psi$ between relatively open subsets of partial quadrants whose domain contains the retract $O$ of $\varphi$, $\psi \circ \varphi$ is a chart of $X$ when $\psi(O)$ is a retract), whose model is a tame retract by V3.71; $N = \ker D(h + t)(0)$, a smooth subspace; the closed good complement with the projection property and the map $\sigma$ of V3.77; $\sigma$ is an sc$^\infty$-germ in the sense of V3.22 because it has $C^k$-lifts into level $k$ for every $k$. *Book:* the proof of Theorem 3.54, p. 115 ("$S$ possesses the $d$-dimensional tangent germ property", $d = \dim \ker f'(x)$). *Hidden hypotheses:* the kernel is a smooth subspace (the text takes it so without comment); $\psi(O)$ is tame (V3.71). *Deviation (narrower):* at $x_0$ only (the other points: V3.85); $d$ is existentially quantified here (the proof takes $d = \dim \ker D(h + t)(0)$); that $d$ is the Fredholm index of $f'(x_0)$ is V3.97.

**V3.85 (Theorem 3.54 (2), and Theorem 3.54; `Polyfold.theorem_3_54_two`, `Polyfold.theorem_3_54`, `Polyfold.theorem_3_54_tangentGermChart_solutionSet`).** Let $f$ be a section of a strong bundle, $f(x_0) = 0$, $(f, x_0)$ in good position in a tame chart (V3.103). (i) The solution set $S = \{f = 0\}$ has at $x_0$ data of Definition 3.25 with $\pi(O) \subseteq C$. (ii) If moreover every zero in an open $V_{x_0} \ni x_0$ is in good position in a tame chart, there are $d \in \mathbb{N}$ and regular data (V3.24) of dimension $d$ for $S$ at $x_0$ whose chart domain lies in $V_{x_0}$, and every datum of Definition 3.25 for $S$ at a point of $S$ in that domain has dimension $d$. (iii) Theorem 3.54, first form: if $f$ is an sc-Fredholm section with tame germ data and $(f, x_0)$ is in good position with the filler condition of V3.82 (in a tame chart), there is an open $V_x \ni x_0$ such that every zero in $V_x$ is in good position in a tame chart, and (ii) holds with $V_{x_0} = V_x$. *Book:* Theorem 3.54 (1), (2), p. 111. *Hidden hypotheses:* tameness as in V3.68. *Deviation (narrower):* (2) is proved from (1), with the nearby good position as a hypothesis;

(1) under the filler condition of V3.82, which follows from good position (V3.102). With the text's hypotheses: V3.94; with the dimension as the index and its local constancy: V3.97, V3.98. "$S$ is a sub-M-polyfold with the structure of a manifold with boundary with corners" follows from (ii) by Theorem 3.27 (V3.69). The regularity conditions (a), (b) of Theorem 3.27, hypotheses for general sets (V3.24), are theorems for the solution set (V3.74, V3.77).

**V3.86 (Theorem 3.56; `Polyfold.theorem_3_56, Polyfold.theorem_3_56_sc`).** Let $X$ be an M-polyfold with $d_X \equiv 0$, $f$ an sc-Fredholm section of a strong bundle, with $(f, x)$ in good position in a tame chart (V3.103) at every zero $x$, and $S = \{f = 0\}$ nonempty. Then $S$ is a sub-M-polyfold of $X$ whose induced structure is equivalent to that of a manifold with boundary with corners of locally constant dimension and has degeneracy index 0 at every point (a manifold without boundary). The same with good position in the text's sense (sc-complement, V3.47) at every zero, which for an sc-Fredholm section with tame germ data is the same hypothesis (`theorem_3_56_sc`). *Book:* Theorem 3.56, p. 115 ($f'(x)$ surjective at every zero). *Hidden hypotheses:* as V3.70. *Deviation (restated):* the hypothesis "$f'(x)$ surjective" is read as good position, which by the text's Definition 3.53 is surjectivity at points with $d_X(x) = 0$; in the chart form the kernel condition is stated for the chart's partial quadrant, and that it is void when $d_X(x) = 0$ is V3.100; the form with "$f'(x)$ surjective" is V3.101. No other hypothesis is added. The tameness of the induced structure is in V3.87 and V3.101.

**V3.87 (Theorem 3.57; `Polyfold.theorem_3_57, Polyfold.theorem_3_57_dim, Polyfold.solutionSetDim, Polyfold.theorem_3_57_sc`).** Let $f$ be an sc-Fredholm section of a strong bundle, with $(f, x)$ in good position in a tame chart (V3.103) at every zero $x$. If $S = \{f = 0\}$ is nonempty, $S$ is a sub-M-polyfold of $X$, its induced structure is tame and equivalent to that of a manifold with boundary with corners of locally constant dimension, and at every $x \in S$ there are regular data of Definition 3.25 of a dimension $d(x)$ such that every datum at a point of $S$ in their chart domain has dimension $d(x)$. Moreover $S$ is the disjoint union of the sets $S_n$ of points at which $S$ has data of dimension $n$, each $S_n$ is open and closed in $S$, and each nonempty $S_n$ is a sub-M-polyfold whose induced structure is tame and equivalent to that of an $n$-dimensional manifold with boundary with corners. The same with good position in the text's sense (sc-complement, V3.47) at every zero, which for an sc-Fredholm section with tame germ data is the same hypothesis (`theorem_3_57_sc`). *Book:* Theorem 3.57, p. 115. *Hidden hypotheses:* as V3.70. *Deviation (stronger):* **the text's hypotheses**; the local constancy of the dimension and the parts $S_n$ are not in the text. Two steps of the text's proof are replaced: the regularity of the solution germ away from the centre (V3.73, V3.74) and the relative openness of $\Sigma$ (V3.75, V3.77). The dimension $d(x)$ is the Fredholm index of $f'(y)$ at the zeros near $x$ (V3.97).

**V3.88 (Theorem 3.58, given the family of sc-isomorphisms; `Polyfold.theorem_3_58_of_family, Polyfold.DerivCondition, Polyfold.contrMap358_contraction, Polyfold.norm_sub_sub_le_of_hasFDerivW Polyfold.fiberIsoMap_isStrongBundleIso, Polyfold.famInv_isScSmooth_of`).** In the coordinates of the text's proof — $E = \mathbb{R}^n \oplus X$ ($\mathbb{R}^n$ any finite-dimensional space with a partial quadrant $\widetilde{C}$), $C = \widetilde{C} \oplus X$, $F = \mathbb{R}^N \oplus Y$ — let $U$ be relatively open in $C$ with $0 \in U$ and $f : U \to F$ sc-smooth; let $V_o \ni 0$ be open in $\mathbb{R}^n$ with $(b, 0) \in U$ for $b \in V_o \cap \widetilde{C}$, and $L(b) : X \to Y$ sc-isomorphisms with $L(b)h = (Df(b, 0)(0, h))_Y$ on level 0 and lower bounds $c_m|h|_m \le |L(b)h|_m$ for $b \in V_o \cap \widetilde{C}$, $h \in X_m$, every $m$; and let condition (3) hold (`DerivCondition`: for all $m$ and $\varepsilon > 0$ there is $r > 0$ with $|[D_2 f(b, 0) - D_2 f(b, x)]h|_m \le \varepsilon |h|_m$ for $b \in \widetilde{C}$, $|b| < r$, $x \in X_{m+1}$, $|x|_m < r$, $(b, 0), (b, x) \in U$, $h \in X_m$). Then on $U' = U \cap (V_o \oplus X)$: $k(b, x) = ((f(b, x) - f(0))_{\mathbb{R}^N}, L(b)^{-1}(f(b, x) - f(0))_Y)$ is a basic germ with contraction map

$H(b,x) = x - L(b)^{-1}(f(b,x) - f(0))_Y$; $\Phi((b,x),(c,y)) = ((b,x),(c,L(b)^{-1}y))$ is a strong bundle isomorphism $U' \lhd (\mathbb{R}^N \oplus Y) \to U' \lhd (\mathbb{R}^N \oplus X)$ covering the identity; and $f - f(0)$ is conjugated to a basic germ. *Book:* the second half of the proof of Theorem 3.58, pp. 118–120. *Hidden hypotheses:* the mean value inequality along a segment with the derivative bound in the weaker norm of $X_m$ (the text's integral; Findings, Chapter 3, item 73). *Deviation (restated):* in the coordinates of the proof; the sc-smoothness of $(b,y) \mapsto L(b)^{-1}y$ is Lemma 3.61 (V3.54) in set form.

**V3.89 (Theorem 3.58, the basic germ criterion; `Polyfold.theorem_3_58`, `Polyfold.CompactnessCondition`, `Polyfold.familyCompactness_derivFamOp`, `Polyfold.fredholmIndex_block`, `Polyfold.surjective_of_fredh` `Polyfold.derivFamOp`).** In the coordinates of V3.88, let $U$ be relatively open in $\widetilde{C} \oplus X$ with $0 \in U$ and $f : U \to \mathbb{R}^N \oplus Y$ sc-smooth with: (1) at every $(b,0) \in U$, $b \in \widetilde{C}$, $Df(b,0)$ is sc-Fredholm with the sc-index of $Df(0)$; (2) $h \mapsto (Df(0)(0,h))_Y$ is bijective on level 0, and the compactness condition holds (`CompactnessCondition`: there is $r > 0$ such that for every $m$, if $b_j \to b_0$ in $\widetilde{C}$ with $(b_j,0) \in U$ and $|b_0| < r$, $(\eta_j)$ is bounded in $X_m$ and $Df(b_j,0)(0,\eta_j) = y_j + z_j$ with $y_j \to 0$ in $F_m$ and $(z_j)$ bounded in $F_{m+1}$, then $(\eta_j)$ has a convergent subsequence in $X_m$); (3) `DerivCondition` (V3.88). Then there are an open $V_o \ni 0$ in $\mathbb{R}^n$ and sc-isomorphisms $L(b) : X \to Y$ with $L(b)h = (Df(b,0)(0,h))_Y$ for $b \in V_o \cap \widetilde{C}$, for which the conclusions of V3.88 hold; in particular $f - f(0)$ is conjugated near 0 to a basic germ. On the way: Lemmas 3.59, 3.60 for this family, with their hypotheses derived from (2) (V3.52, V3.53); the index of a block operator, $\operatorname{ind} T = \dim \mathbb{R}^n + \operatorname{ind} L - \dim \mathbb{R}^N$ for $T : \mathbb{R}^n \oplus X \to \mathbb{R}^N \oplus Y$ Fredholm with block $L$ of finite index, and an injective block is onto when $\operatorname{ind} T = \dim \mathbb{R}^n - \dim \mathbb{R}^N$. *Book:* Theorem 3.58, p. 116, proof pp. 116–120. *Hidden hypotheses:* the smallness of $b$ in (2) uniform in the level $m$. *Deviation (narrower):* in the coordinates of the proof; the reduction of a general sc-splitting $E = B \oplus X$, $F = A \oplus Y$ to them is not formalized; (1) is needed only at the points $(b,0)$; the text's "$B \supseteq \ker Df(0)$ and $Y = Df(0)X$" is replaced by the bijectivity it implies.

**V3.90 (Proposition 3.73 and Lemma 3.77 with the hypotheses their proofs use; `Polyfold.InvDataW`, `Polyfold.InvDataW.injOn`, `Polyfold.InvDataW.hasFDerivWithinAt_inv`, `Polyfold.InvDataW.contDiffOn_in`** Let $U$ be a convex subset of a finite-dimensional normed space $V$ with unique differentiability at its points, $f : V \to V$ of class $C^j$ ($j \geq 1$) on $U$ with $\|Df(x) - 1\| \leq \frac{1}{2}$ for $x \in U$, and let $f(U)$ have unique differentiability. Then $f$ is injective on $U$ ($\frac{1}{2}|x - x'| \leq |f(x) - f(x')|$), its inverse $g$ is differentiable within $f(U)$ with $Dg(y) = [Df(g(y))]^{-1}$, and $g$ is of class $C^j$ on $f(U)$. *Book:* Proposition 3.73, p. 137; Lemma 3.77, p. 140. *Hidden hypotheses:* none. *Deviation (restated):* no partial quadrant; the hypothesis "$x$ interior iff $f(x)$ interior" of V3.59 is replaced by the unique differentiability of $f(U)$, which is what the bootstrapping uses; the relative openness of $f(U)$ is not concluded. Used in the re-centring (V3.73), where $f(U)$ is relatively open in a cone for another reason.

**V3.91 (Theorem 3.18, parametrized perturbations; `Polyfold.theorem_3_18`, `Polyfold.theorem_3_18'`, `Polyfold.paramSection`, `Polyfold.paramSection_eq_sectionAdd`, `Polyfold.StrongBundle.pullback`, `Polyfold.StrongBundle.IsScFredholmSection.param`, `Polyfold.StrongBundle.isScPlusSection_paramComb` `Polyfold.IsBasicGerm.paramExtend`, `Polyfold.IsConjugateToBasicGerm.paramExtend`, `Polyfold.rnPolyfold`, `Polyfold.prod_rn_isTame`).** Let $P : Y \to X$ be a strong bundle over an M-polyfold ($X$, $Y$ Hausdorff and paracompact, as throughout), $f$ an sc-Fredholm section (V3.16) and $s_1, \dots, s_n$ sc$^+$-sections of $P$ (Definition 2.52). Let $\mathbb{R}^n$ carry its M-polyfold structure with the constant sc-structure and the charts $r \mapsto r - r_0$, $\mathbb{R}^n \times X$ the product M-polyfold, $\pi(r,x) = x$, and $\pi^*P : \pi^*Y \to \mathbb{R}^n \times X$ the pull-back strong bundle with the natural structure of

Proposition 2.54 (the atlas of all charts pulled back from charts of $P$ through charts of $\mathbb{R}^n \times X$). Then $F(r,x) = f(x) + \sum_i r_i s_i(x)$, computed in the fibre $P^{-1}(x)$ and read in the fibre of $\pi^*Y$ over $(r,x)$, is an sc-Fredholm section of $\pi^*P$. The paracompactness of $\mathbb{R}^n \times X$ and of $\pi^*Y$, instance arguments of `theorem_3_18`, is supplied by Theorem 2.16 in `theorem_3_18'`. On the way: $\tilde f(r,x) = f(x)$ is an sc-Fredholm section of $\pi^*P$ — in the product chart $(r \mapsto r - r_0) \times \varphi$, centred at $(r_0, x)$, the filling germ is $(a,u) \mapsto g(u)$ on $\mathbb{R}^n \times G$ (filler conditions (1)–(3) from those of $g$), the sc$^+$-germ $(a,u) \mapsto t_0(u)$, and the conjugation is the product of the given one with the identity of $\mathbb{R}^n$, onto the $\mathbb{R}^n$-extension $((a,\lambda),w) \mapsto h(a,w)$ of the basic germ, again a basic germ; $\tilde s(r,x) = \sum_i r_i s_i(x)$ is an sc$^+$-section (the product rule with a scalar factor, from Proposition 1.19); $F = \tilde f + \tilde s$ fibrewise, and Theorem 3.15 (V3.17) applies; $\mathbb{R}^n \times X$ is tame when $X$ is. *Book:* Theorem 3.18, p. 74 (for a tame strong bundle; "$\tilde f$ … is a sc-Fredholm section as is readily verified", p. 73). *Hidden hypotheses:* the translation charts of $\mathbb{R}^n$ (a chart centred at $(r_0,x)$ is needed for the germ there), the strong bundle structure of $\pi^*P$, the scalar product rule. *Deviation (stronger):* tameness is not used, as for Theorem 3.15.

**V3.92 (the linearization of the parametrized section; `Polyfold.linAt_paramSection_zero`, `Polyfold.linAt_paramSection`, `Polyfold.scDeriv_paramSection`, `Polyfold.PullbackChartData.ofProd`).** In the setting of V3.91, let $c$ be a strong bundle chart of $P$ compatible with the atlas, with 0 in its retract $O$, and consider the product chart of $\pi^*P$ through $(r \mapsto r - r_0) \times c$, with the retraction $1 \times r$. At its centre: $D(\hat F \circ (1\times r))(0)(\delta a, \delta u) = D(\hat G \circ r)(0)\delta u + \sum_i \delta a_i \hat s_i(0)$, $\hat G = \hat f + \sum_i r_{0,i}\hat s_i$; in the chart form of the linearization (V3.78), on $T_0(\mathbb{R}^n \times O) \subseteq \mathbb{R}^n \times T_0O$: $\mathrm{Lin}_F(0)(\delta a, \delta u) = \mathrm{Lin}_{f+\sum r_{0,i}s_i}(0)\delta u + \sum_i \delta a_i \hat s_i(0)$; and at $r_0 = 0$: $\mathrm{Lin}_F(0)(\delta r, \xi) = \mathrm{Lin}_f(0)\xi + \sum_i \delta r_i \hat s_i(0)$. *Book:* after Theorem 3.19, p. 75 ("the linearization of $\tilde f$ at $(0,x_0)$ is $(h,u) \mapsto f'(x_0)u + \sum h_i e_i$"). *Hidden hypotheses:* none (the formula does not use $F = 0$ at the point). *Deviation (restated):* in the chart form (Definition 3.53 is formalized in charts); at $(r_0,x)$ with $r_0 \neq 0$ the first term is the linearization of $f + \sum r_{0,i}s_i$; the text's "sc$^1$-sections $s_i$" are sc$^+$ here, as Theorem 3.18 needs.

**V3.93 (Theorem 3.54 (1), the surjectivity at the nearby zeros, in the chart at $x_0$; `Polyfold.theorem_3_54_chart_o`, `Polyfold.StrongRetr.mem_0_of_perturbed_zero`, `Polyfold.exists_inward_of_isInGoodPosition`, `Polyfold.inward_add_smul`, `Polyfold.exists_curve_of_inward`, `Polyfold.ShiftedBasicData.exists_radii`, `Polyfold.ShiftedBasicData.perturbContraction`, `Polyfold.ShiftedBasicData.perturbContraction_isSc1`, `Polyfold.xiMap`, `Polyfold.exists_xi_radii_bounds`, `Polyfold.ShiftedBasicData.surjective_DH_and_ker_`, `Polyfold.ShiftedBasicData.exists_perturbed_curve`).** In the data of V3.48 at a zero $x_0$ in good position — the strong bundle chart centred at $x_0$ with retraction $(r,\rho)$ on $U$ onto $O$, the filling germ $g$, the conjugation $\Psi$ covering $\psi$ of $g - s$ to a basic germ, $h' + \iota t$ with kernel $K = \ker D(h' + \iota t)(0)$ in good position, the closed good complement, the $C^1$ map $\sigma$ on $V'$ with its lifts, and an open $N_2 \ni 0$ on which $\psi$ maps the zeros of $g$ in $O$ onto the graph of $\sigma$ — assume condition (2) of Definition 3.4 on $N_2$ and $K \neq 0$. Then there is an open $N_4 \subseteq N_2$ containing 0 such that for every zero $z_0 \in O \cap N_4$ of $g$, every level-1 point over $z_0$ and every smooth $k \in F_\infty$, $\rho(z_0)k = Dg(z_0)x$ for some $x \in T_{z_0}O = Dr(z_0)E$. On the way: (a) a zero of $g + \mu\rho(r(\cdot))k$ at which condition (2) holds lies in $O$; (b) a nonzero finite-dimensional subspace in good position to the partial quadrant $C_z$ at $z \in C$ contains a vector $v$ with $z + t(v+\zeta) \in C$ for $t \in [0,t_0]$, $|\zeta| < \eta$, and such inward vectors stay inward after adding any vector and rescaling; (c) for $\widetilde{C}$ convex and $H$ of class $C^1$ on $\Omega \cap (\mathbb{R} \times \widetilde{C})$ with $H(0,a_0) = 0$, $D_aH(0,a_0)$ onto and an inward $\alpha_0$ with $DH(0,a_0)(1,\alpha_0) = 0$, there is a curve $a(\mu)$, $0 \le \mu < \mu_0$, $a(0) = a_0$, with right derivative $\alpha_0$ at 0 and $H(\mu, a(\mu)) = 0$; (d) for the normal form of Theorem 3.38 (a basic germ $(G_1, w - B)$ on the shifted structure) there are $\frac14$-contraction regions of $B$ below any prescribed radii, and

$B - \mu P\hat{e}$ is a contraction germ in the working form on $(-\mu_m, \mu_m) \times \{|a| < \nu_m\}$ with the same regions in $a$ for every perturbation $\hat{e}$ with given bounds, its solution germ sc[1] (Theorem 3.30); (e) the push-forward $\hat{e}_k = \Xi(\cdot, k)$ of $u \mapsto \rho(r(u))k$ to the normal form, with $\Xi$ sc-smooth into the index-raised target and linear in $k$, is bounded with its Lipschitz constant on the contraction regions by $c_m = 2|k|_{m+2}(M_m + 1)/\delta_m$, $\delta_m$, $M_m$ independent of $k$; (f) at a zero $(a_0, w_0)$ of the normal form with $|a_0| < \nu_3$, $|w_0|_1 < \rho_0$ and a kernel of dimension $n - N$, $D_a H(0, a_0)$ is onto for $H$ the $\mathbb{R}^N$-component of $(G + \mu\hat{e})(a, \delta_\mu(\mu, a))$, the kernel is the graph of $D_a\delta_\mu(0, a_0)$ over $\ker D_a H(0, a_0)$, and there is a curve of zeros of $G + \mu\hat{e}$ through it, right differentiable at 0 into level 2. *Book:* the proof of Theorem 3.54, property (4), p. 114 (from Corollary 3.41 and Lemma 3.55). *Hidden hypotheses:* none added. *Deviation (not in the book):* a new argument (Findings, Chapter 3, items 180–187): the text's argument gives the surjectivity of the filled $Dg(z_0)$ only (V3.82). The neighbourhood $N_4$ is determined by $g$, $r$, $\rho$, $\Psi$ and the normal form, not by $k$; Corollary 3.39 is not used. In the germ reading the curve of zeros lies in the germ of the filling, so $g$ is differentiated within its germ (Findings, Chapter 3, items 276–278, 298).

**V3.94 (Theorem 3.54 (1), and Theorem 3.54, with the text's hypotheses; `Polyfold.theorem_3_54_one'`, `Polyfold.theorem_3_54'`, `Polyfold.theorem_3_54_one_sc_of_tame`, `Polyfold.theorem_3_54_one_sc`, `Polyfold.theorem_3_54_sc`).** Let $f$ be an sc-Fredholm section of a strong bundle with tame germ data (V3.103), $f(x_0) = 0$, and $(f, x_0)$ in good position in a tame chart (V3.47, V3.103). Then there is an open $V_x \ni x_0$ such that $(f, y)$ is in good position in a tame chart at every zero $y \in V_x$; in particular $f'(y)$ is onto (`theorem_3_54_one'`). Moreover (`theorem_3_54'`) there are $d \in \mathbb{N}$ and regular data of Definition 3.25 (V3.24) of dimension $d$ for $S = \{f = 0\}$ at $x_0$ whose chart domain lies in $V_x$, and every datum of Definition 3.25 for $S$ at a point of $S$ in that domain has dimension $d$. With the text's notion of good position (sc-complement, V3.47) in the conclusion, from either form at $x_0$: every zero $y \in V_x$ is in good position with an sc-complement. *Book:* Theorem 3.54 (1), (2), p. 111. *Hidden hypotheses:* tameness as in V3.68. *Deviation (restated):* no hypothesis added; (2) as regular tangent germ data with locally constant dimension (the manifold structure of the local solution set is V3.95); the proof of the surjectivity in (1) is V3.93, through Definition 3.53 from a chart not centred at $y$ (V3.79), read in the tame chart of the germ data at $y$ (V3.107); for $\dim K = 0$ the only zero near $x_0$ is $x_0$. The project's good position asks only for an algebraic good complement (V3.20), less than the text's as a conclusion; the sc form follows at each zero (Findings, Chapter 3, items 210, 213).

**V3.95 (Theorems 3.23 and 3.19 with the text's hypotheses; `Polyfold.theorem_3_23'`, `Polyfold.theorem_3_19'`, `Polyfold.theorem_3_23_sc`).** Let $f$ be an sc-Fredholm section of a strong bundle with tame germ data (V3.103), $f(x) = 0$ and $(f, x)$ in good position in a tame chart. Then there are an open $U \ni x$ and $d \in \mathbb{N}$ such that every zero in $U$ is in good position in a tame chart and a smooth point, and $S(f, U) = \{y \in U \mid f(y) = 0\}$ is a sub-M-polyfold whose induced structure is tame and equivalent to that of a $d$-dimensional manifold with boundary with corners (`theorem_3_23'`). If moreover $d_X \equiv 0$, the induced structure also has degeneracy index 0 everywhere (`theorem_3_19'`). Theorem 3.23 with the text's notion of good position (sc-complement, V3.47) at $x$ and at every zero in $U$: `theorem_3_23_sc`. *Book:* Theorem 3.23, p. 77; Theorem 3.19, p. 74. *Hidden hypotheses:* tameness as in V3.68. *Deviation (hypotheses explicit):* no hypothesis added. For Theorem 3.19 the hypothesis "$f'(x_0)$ surjective" is read as good position, the same at $d_X = 0$ (V3.100, V3.101); "the dimension agrees with the Fredholm index" is V3.97, "$\ker f'(y) = T_y S$" V3.99; "uniquely determined" is the equivalence of atlases.

**V3.96 (the index of the linearization at a zero, in every chart; `Polyfold.StrongBundle.IsSurjLinOfIndex`, `Polyfold.StrongBundle.linAt_index_of_conjugationData`, `Polyfold.StrongBundle.isSurjLinOfIndex_of_`**

`Polyfold.StrongBundle.IsGoodPositionGerm.exists_isSurjLinOfIndex`, `Polyfold.theorem_3_19_lineariz`
`Polyfold.StrongBundle.linAt_isScFredholm`, `Polyfold.StrongBundle.linAt_isScFredholm_of_isSurjLin`
`Polyfold.proposition_3_52_sc`, `Polyfold.theorem_3_19_linearization_sc`). `IsSurjLinOfIndex f y d`: in every strong bundle chart compatible with the atlas whose domain contains $y$, at the level-1 point $z$ over $\varphi(y)$, the linearization $\mathrm{Lin}(z) : T_zO \to K_z$ (V3.78) is surjective and Fredholm with $\mathrm{ind}\,\mathrm{Lin}(z) = \dim\ker\mathrm{Lin}(z) = d$. Theorem: let $f$ be an sc-smooth section, $x_0$ a zero, $c$ the strong bundle chart of Definition 3.11 centred at $x_0$ with filling germ $g$, sc$^+$-germ $s$, $s(0) = g(0)$, and conjugation data of $g - s$ to a basic germ $h : O(\widetilde{C} \oplus W, 0) \to \mathbb{R}^N \oplus W$, $\widetilde{C} \subseteq \mathbb{R}^n$. Then in every strong bundle chart around $x_0$ the linearization at the point over $x_0$ is Fredholm of index $n - N$, and it is onto if the germ is in good position in $c$. Hence at a zero in good position `IsSurjLinOfIndex f x_0 d` holds with $d = n - N$ for every conjugation in the chart of Definition 3.53. For an sc-Fredholm section and a zero $x_0$ in good position, in the chart of Definition 3.53, $Dg(0)$ and $Tf(x_0) = Dg(0)|_{T_0O}$ are Fredholm, $Tf(x_0)$ is onto, and for every conjugation $\mathrm{ind}\,Dg(0) = \mathrm{ind}\,Tf(x_0) = \dim\ker Tf(x_0) = n - N$. In sc form: for an sc-Fredholm section, a zero $y$ and any strong bundle chart around $y$, at the point $z$ over $\varphi(y)$ (a smooth point), $T_zO = Dr(z)E$ and $K_z = \rho(z)F$ are sc-subspaces and $\mathrm{Lin}(z) : T_zO \to K_z$ is an sc-Fredholm operator of index $n - N$ on every level; under `IsSurjLinOfIndex f y d` it is onto on every level, of index $d$ on every level; in the chart of Definition 3.53, $Dg(0)$ and $f'(x_0)$ are sc-Fredholm and $f'(x_0)$ is onto on every level. *Book:* Proposition 3.52, p. 110, and the first clause of Theorem 3.19, p. 74. *Hidden hypotheses:* none. *Deviation (stronger):* in charts ($T_xX$ carries no linear structure here, Chapter 2); in every chart by the chart independence at zeros (V3.78), which the text does not state, realised on all levels by the chart change at smooth points (Findings, Chapter 3, items 236, 239); the index on every level, and the surjectivity on every level at a zero in good position, go beyond the text's statement. The hypothesis is the text's since the switch to the germ reading (phase 2): Proposition 3.52 enters at the centre of the chart of the germ datum, where $\mathrm{Lin}(0) = Dg(0)$ on $T_0O$ for the filling germ (Findings, Chapter 3, items 284, 310, 314, 330). `theorem_3_19_linearization_sc` gives the conclusions in the chart at $x_0$, and `linAt_isScFredholm_of_isSurjLinOfIndex` with `IsGoodPositionGerm.exists_isSurjLinOfIndex` in every chart. The sc-Fredholm property was first proved on level 0 only; formerly "narrower", for the whole-chart hypothesis.

**V3.97 (the dimension of the solution set is the Fredholm index;** `Polyfold.theorem_3_54_tangentGermChart_in`
`Polyfold.theorem_3_54_two_index`, `Polyfold.theorem_3_23_index`, `Polyfold.theorem_3_19_index`,
`Polyfold.theorem_3_57_index`, `Polyfold.theorem_3_54_graph_proj_conj`, `Polyfold.theorem_3_54_tangent`
`Polyfold.StrongBundle.linAt_isScFredholm_of_isSurjLinOfIndex`**).** Let $f$ be an sc-smooth section of a strong bundle, $f(x_0) = 0$, $(f, x_0)$ in good position in a tame chart (V3.103). (i) The data of Definition 3.25 at $x_0$ of V3.84 have a dimension $d$ with `IsSurjLinOfIndex f x_0 d` (V3.96): $d = n - N$ for the conjugation in the chart from which $\psi\varphi$ is built. (ii) The regular data of V3.85 (ii) have such a dimension. (iii) For $f$ sc-Fredholm with tame germ data there are an open $U \ni x_0$ and $d$ such that every zero $y \in U$ is in good position in a tame chart with `IsSurjLinOfIndex f y d`, the zeros in $U$ are smooth points, and $S(f, U)$ is a sub-M-polyfold whose induced structure is tame and equivalent to that of a $d$-dimensional manifold with boundary with corners (`theorem_3_23_index`); if $d_X \equiv 0$, also of degeneracy index 0 everywhere (`theorem_3_19_index`). (iv) Under the hypotheses of V3.87 (good position in a tame chart at every zero, $S \neq \emptyset$): $S$ is a sub-M-polyfold with tame induced structure, a manifold with boundary with corners of locally constant dimension, and at every $x \in S$ there are regular data of a dimension $d$ such that every datum at a point of $S$ in their chart domain has dimension $d$ and `IsSurjLinOfIndex f y d` holds at every zero $y$ of that domain (`theorem_3_57_index`). For an sc-Fredholm section, at a zero with `IsSurjLinOfIndex f y d`, $\mathrm{Lin}(z)$ is moreover sc-Fredholm and onto on every level, of index $d$

on every level, in every chart (V3.96). *Book:* Theorem 3.19, p. 74 ("whose dimension agrees with the Fredholm index"); the proof of Theorem 3.54, p. 115 ($d = \dim\ker f'(x)$). *Hidden hypotheses:* tameness as in V3.68. *Deviation (stronger):* the index at every zero near the point and in every chart; the text states no dimension in Theorems 3.23, 3.57.

**V3.98 (the Fredholm index along the solution set and along paths, Definition 6.1, Proposition 6.2;** `Polyfold.theorem_3_54_index_locally_constant`, `Polyfold.theorem_3_54_index`, `Polyfold.isSurjLinOfIndex_of_forall_dim`, `Polyfold.theorem_3_54_index_locally_constant_sc`, `Polyfold.StrongBundle.linAt_isScFredholm`, `Polyfold.index_locally_constant_tame`, `Polyfold.theorem_3_54_index_locally_constant_general_tame`, `Polyfold.index_locally_constant_of_ac` `Polyfold.theorem_3_54_index_locally_constant_general`, `Polyfold.StrongBundle.IsLinOfIndex`, `Polyfold.StrongBundle.restrictOpen`, `Polyfold.StrongBundle.IsIndexAtLocal`, `Polyfold.StrongBundle.I` `Polyfold.StrongBundle.exists_isIndexAtLocal`, `Polyfold.StrongBundle.IsIndexAt`, `Polyfold.StrongBundle.IsIndexAt.unique`, `Polyfold.StrongBundle.exists_isIndexAt`, `Polyfold.StrongBundle.isIndexAtLocal_iff_isIndexAt`, `Polyfold.exists_isScPlusSection_pullback_of_` `Polyfold.StrongBundle.isIndexAt_of_pullback_isLinOfIndex`, `Polyfold.proposition_6_2_local`, `Polyfold.proposition_6_2_local'`, `Polyfold.proposition_6_2`**).** Let $f$ be an sc-Fredholm section of a strong bundle with tame germ data (V3.103), $f(x_0) = 0$, $(f, x_0)$ in good position in a tame chart. Then there are an open $V_x \ni x_0$ and $d$ such that `IsSurjLinOfIndex f y d` holds at every zero $y \in V_x$: $f'(y)$ is onto with $\operatorname{ind} f'(y) = \dim\ker f'(y) = d = \operatorname{ind} f'(x_0)$ (`theorem_3_54_index_locally_constant`). Theorem 3.54 with the index (`theorem_3_54_index`): moreover every zero in $V_x$ is in good position in a tame chart, and $S$ has at $x_0$ regular data of dimension $d$ whose chart domain is $V_x$, every datum at a point of $S$ in $V_x$ having dimension $d$. On the way: at a zero in good position in a tame chart at which every datum of Definition 3.25 for $S$ has dimension $d$, `IsSurjLinOfIndex f y d` ($f$ sc-smooth). In sc form, at every zero in $V_x$ and in every chart, $\mathrm{Lin}(z)$ is sc-Fredholm, onto on every level, of index $d$ on every level (`theorem_3_54_index_locally_constant_sc`); at any zero of an sc-Fredholm section, in every chart, $\mathrm{Lin}(z)$ is sc-Fredholm with the same index on every level (`linAt_isScFredholm`). Without good position, with the text's hypotheses: if $f$ is an sc-Fredholm section with tame germ data and $f(x_0) = 0$, there are an open $V \ni x_0$ and $i \in \mathbb{Z}$ such that at every zero $y \in V$, in every strong bundle chart, $f'(y)$ is Fredholm of index $i$ (`IsLinOfIndex f y i`; `index_locally_constant_tame`) and $\mathrm{Lin}(z)$ is sc-Fredholm of index $i$ on every level (`theorem_3_54_index_locally_constant_general_tame`); no bump functions are assumed (the first forms, for $X$ admitting sc-smooth bump functions, Definition 5.9: `index_locally_constant_of_admitsBumpFunctions`, `theorem_3_54_index_locally_constant_general`). **Definition 6.1**, with the text's locally defined section: an sc-Fredholm section $f$ has index $i$ at a smooth point $x$ (`IsIndexAtLocal f x i`) if for some open $V \ni x$ and some sc$^+$-section $s$ of the restriction $P|V$ of $P$ to $V$ (`StrongBundle.restrictOpen`: the strong bundle over the M-polyfold $V$ whose atlas consists of the charts of $P$ with base domain inside $V$), $(f|_V + s)(x) = 0$ and $(f|_V + s)'(x)$ is Fredholm of index $i$ in every strong bundle chart of $P|V$; the index is unique, at a zero it is the index of $f'(x)$, and it exists at every smooth point. With an sc$^+$-section on all of $X$ in place of $s$ (`IsIndexAt f x i`) the index is unique as well and implies the local one; if $X$ admits sc-smooth bump functions it exists at every smooth point and the two readings agree (`isIndexAtLocal_iff_isIndexAt`). **Proposition 6.2**: if $f$ is an sc-Fredholm section with tame germ data, then for every sc-smooth path $\varphi : \mathbb{R} \to X$ the index $\operatorname{ind}(f, \varphi(t))$ (local reading) is the same for all $t$ (`proposition_6_2_local`), and any two values of the index at $\varphi(0)$ and $\varphi(1)$ agree (`proposition_6_2_local'`); if $X$ admits sc-smooth bump functions, the same holds for the global reading (`proposition_6_2`). *Book:* Definition 6.1, pp. 200–201; Proposition 6.2, p. 201 (over a tame

M-polyfold, the index is constant along sc-smooth paths $[0,1] \to X$ of smooth points; the proof: "we do not need a sc-smooth partition of unity", p. 202), whose proof uses the step of V3.82; in Chapter 3 the statement along the solution set is implicit in Theorem 3.54 (1). *Hidden hypotheses:* tameness as in V3.68; for Definition 6.1, the text's local sc$^+$-section $s$ with $s(x) = f(x)$ is a section of the restriction of the bundle to an open neighbourhood of $x$ ($-s$ here), a construction the text uses without comment. *Deviation (narrower):* the path is defined on $\mathbb{R}$, not on $[0,1]$ ($\mathbb{R} \times X$ is the product of Theorem 3.18 and has no boundary, and the text's tame M-polyfold $[0,1] \times X$ is not used): a path on $[0,1]$ that extends sc-smoothly to $\mathbb{R}$ is covered, a path sc-smooth on $[0,1]$ only is not, and the project has no M-polyfold $[0,1]$. This is the only narrowing left; the hypotheses on $f$ are the text's, in the reading of tameness of V3.103 (as in V3.94), and no bump functions are assumed. *Proof:* Proposition 6.2 is proved by the text's device, run on the restriction $P|V$ to the domain $V$ of a chart around $\varphi(t_0)$, where the text's sections need no cut-off — for each $t_0$ an sc$^+$-section $S$ of the pull-back to $\mathbb{R} \times V$ with $\tilde{f} + S = 0$ along the graph of the path near $t_0$ (the text's $R(x, f(\varphi(t)))$ in the chart), the local constancy of the index of $\tilde{f} + S$ along its zero set, the count $\operatorname{ind}(\tilde{f} + S)'(t,x) = \operatorname{ind}(f,x) + 1$ through the slice $S(t,\cdot)$, and the connectedness of $\mathbb{R}$; the earlier form for M-polyfolds with bump functions cuts the sections off instead (Findings, Chapter 3, items 340–349, 360–367). The local constancy along the zero set without good position is the same device: sc$^+$-sections $R(\cdot, e_i)$ on $P|V$ put the parametrized section of Theorem 3.18 into general position at $(0, x_0)$, its index is constant along its solution set, at $(0,y)$ it is $k + \operatorname{ind} f'(y)$, and the index at a zero is the same for $P$ and $P|V$, through the germ datum (items 240–242, 360–363). Definition 6.1 is well defined because two local sections are compared on the intersection of their domains; its independence of $s$ is proved from the germ data and Theorem 3.15, not by the text's appeal to Proposition 1.10; and it exists at every smooth point through the section $-R(\cdot, f(x))$ on a chart domain (items 340, 364). Under bump functions a local section is cut off to a global one with the same index (item 367). The candidate counterexample of items 188 and 243 is excluded by Theorem 3.54 (1). Without tame germ data the local constancy is open: it amounts to the index 0 of the block $(1 - \rho(z))Dg(z)|_{\ker Dr(z)}$ of the filling at the nearby zeros, which condition (3) of Definition 3.4 gives at $z = 0$ only (items 243, 343).

**V3.99 (**$\ker f'(y) = T_yS$ **in a chart;** `Polyfold.theorem_3_54_ker_eq_tangent`**).** Let $f$ be an sc-smooth section of a strong bundle, $f(x_0) = 0$, $(f, x_0)$ in good position in a tame chart (V3.103). Then there are a tame strong bundle chart $c$ centred at $x_0$ (as in Definition 3.53), the conjugating sc-diffeomorphism $\psi : U \to U' \subseteq C'$ (inverse $\psi'$, $\psi(0) = 0$), the finite-dimensional $K = \ker D(h+t)(0)$, the solution graph $\sigma : V' \to E'$ over a relatively open $V' \ni 0$ of $K \cap C'$ ($C^1$, $\sigma(0) = 0$, $D\sigma(0) = 0$), an open $V_x \ni x_0$ with $\psi\varphi(\{y \in V_x \mid f(y) = 0\}) = \{v + \sigma(v) \mid v \in V'\}$, and an open $V_{x,2} \subseteq V_x$ containing $x_0$, such that for every zero $y \in V_{x,2}$, with $\psi(\varphi(y)) = v + \sigma(v)$, $v \in V'$, and every level-1 point $z$ over $\varphi(y)$: $D\psi(z)$ maps the kernel of $\operatorname{Lin}(z) : T_zO \to K_z$ onto $N_v = \{n + D\sigma(v)n \mid n \in K\}$, the tangent space at $v$ of the graph. *Book:* Theorem 3.19, p. 74 ("ker $f'(y) = T_yS(f,U)$ is the tangent space"). *Hidden hypotheses:* tameness as in V3.68. *Deviation (restated):* in the chart $\psi\varphi$ of the tangent germ data, the kernel of the linearization in the chart $c$; $T_yS$ as a subspace of $T_yX$ is not formalized (Chapter 2, gap G.13).

**V3.100 (good position is surjectivity at a zero with** $d_X = 0$**;** `Polyfold.StrongBundle.IsLinSurjAt`**,** `Polyfold.isInGoodPositionIn_of_subset`**,** `Polyfold.StrongBundle.isGoodPositionGerm_of_linAt_surj_of`
`Polyfold.StrongBundle.isGoodPositionGerm_of_isLinSurjAt`**,** `Polyfold.StrongBundle.IsGoodPositionGer`
`Polyfold.StrongBundle.isGoodPositionGerm_iff_isLinSurjAt`**,** `Polyfold.StrongBundle.IsLinSurjAt.linA`
`IsLinSurjAt f y` ("$f'(y)$ is surjective"): in some strong bundle chart compatible with the atlas whose domain contains $y$, at a level-1 point $z$ over $\varphi(y)$, $\operatorname{Lin}(z)$ maps onto the fibre

$K_z = \{k \mid (\varphi(y), k) \in K\}$; then, at a zero, it does so in every such chart and at every level-1 point over $\varphi(y)$ (V3.78). Let $f$ be an sc-Fredholm section of a strong bundle with tame germ data (V3.103) and $y$ a zero with $d_X(y) = 0$. Then $(f, y)$ is in good position in a tame chart if and only if `IsLinSurjAt f y`. On the way: a subspace $N \subseteq T$ is in good position inside $T$ to every cone containing $T$; in a chart with tame base retraction around a zero $y$ with $d_X(y) = 0$ at which $(f, y)$ is an sc-Fredholm germ, if the linearization is onto there, $(f, y)$ is in good position, since $d_C(\varphi(y)) = d_X(y) = 0$ makes the partial quadrant at $\varphi(y)$ the whole space; the surjectivity is carried to the tame chart of the germ data at $y$, and the good position read there (V3.107). *Book:* the remark after Theorem 3.54, p. 111 (for $\partial X = \emptyset$ good position just means surjectivity), and Corollary 3.24 ("an empty condition if $d_X(x) = 0$", p. 79). *Hidden hypotheses:* the chart is tame, for $d_C = d_X$ (in a non-tame chart $d_C(\varphi(y))$ can exceed $d_X(y)$). *Deviation (hypotheses explicit).*

**V3.101 (Theorems 3.19 and 3.56 with "$f'(x)$ surjective", and Theorem 3.56 with tameness; `Polyfold.theorem_3_19_of_surjective`, `Polyfold.theorem_3_56_of_surjective`, `Polyfold.theorem_3_56_tame`).** Let $X$ be an M-polyfold with $d_X \equiv 0$ and $f$ an sc-Fredholm section of a strong bundle. (i) If $f$ has tame germ data (V3.103), $f(x_0) = 0$ and `IsLinSurjAt f x_0` (V3.100), the conclusions of `theorem_3_19_index` hold (V3.97 (iii)): an open $U \ni x_0$ and $d$ with every zero $y \in U$ in good position in a tame chart, `IsSurjLinOfIndex f y d`, the zeros in $U$ smooth, and $S(f, U)$ a sub-M-polyfold whose induced structure is tame, a $d$-dimensional manifold with degeneracy index 0 everywhere. (ii) If $(f, x)$ is in good position in a tame chart at every zero (`theorem_3_56_tame`), or $f$ has tame germ data and `IsLinSurjAt f x` at every zero (`theorem_3_56_of_surjective`), and $S \neq \emptyset$, then $S$ is a sub-M-polyfold whose induced structure is tame, a manifold of locally constant dimension with degeneracy index 0 everywhere, with regular data of Definition 3.25 at every zero of a dimension $d$ constant near it and equal to the Fredholm index at the zeros of the chart domain. *Book:* Theorem 3.19, p. 74; Theorem 3.56, p. 115. *Hidden hypotheses:* tameness as in V3.68; $S \neq \emptyset$ as in V3.70. *Deviation (stronger):* the text's hypotheses; the tameness of the induced structure and the index are added conclusions.

**V3.102 (the filler condition at the nearby zeros follows from good position; `Polyfold.normal_injective_of_ex` `Polyfold.exists_range_deriv_eq_ _iff_normal_injective_germ`, `Polyfold.isGoodPositionGermFilled_of`).** (a) Let $U$ be relatively open in a partial quadrant, $R$ a strong bundle retraction on $U \lhd F$, $g$ an sc$^1$-germ with derivative family $Dg$, and $z$ a level-1 point in its germ over a point $z_0$ of the retract $O$ with $\ker Dg(z) \subseteq T_{z_0}O$. If for every $k \in F$ there is $x \in T_{z_0}O$ with $Dg(z)x = \rho(z_0)k$, then $Dr(z)b = 0$ and $\rho(z_0)Dg(z)b = Dg(z)b$ imply $b = 0$: condition (3) of Definition 3.4 at $z_0$. (b) If moreover $\mathbf{f} \circ r$ is sc$^1$, $\mathbf{f}$ a section of $K$ on $O$ with $\mathbf{f}(z_0) = 0$, $\mathbf{f} = g$ on $O$ near $z_0$, and $Dg(z)$ is onto, the two conditions are equivalent. (c) For an sc-Fredholm section with tame germ data (V3.103) and a zero $x_0$ in good position in a tame chart, $(f, x_0)$ is in good position in a tame chart with the filler condition at the nearby zeros (`IsGoodPositionGermFilled`, V3.82). *Book:* not in the text (Definition 3.4 (3), p. 64, at 0 only). *Hidden hypotheses:* none. *Deviation (not in the book):* (c) uses V3.94 and the tangency $\ker Dg(z_0) \subseteq T_{z_0}O$ of V3.81; it shows that the added hypothesis of V3.68, V3.69, V3.82, V3.85 is satisfied whenever the text's hypotheses are.

**V3.103 (Definitions 3.50, 3.51 and 3.53 in a tame chart; `Polyfold.StrongBundle.IsTameChart`, `Polyfold.StrongBundle.IsScFredholmGermTame`, `Polyfold.StrongBundle.IsGoodPositionGermTame`, `Polyfold.StrongBundle.IsScFredholmSectionTame`, `Polyfold.StrongBundle.IsScFredholmGermTame.isScFr` `Polyfold.StrongBundle.IsGoodPositionGermTame.isGoodPositionGerm`, `Polyfold.StrongBundle.IsGoodPos` `Polyfold.StrongBundle.IsGoodPositionGermFilled.isGoodPositionGermTame`, `Polyfold.StrongBundle.isT` `Polyfold.ScRetraction.IsTame.congr`, `Polyfold.IsFillingGerm.nonempty_kerEquiv`, `Polyfold.TrivialBundle.tame_atlas_no_germData_in_atlas_chart`).** A strong bundle

chart of $P$ is *tame* (`IsTameChart`) if it is compatible with the atlas and its base retraction $r : U \to U$ is tame (Definition 2.34). `IsScFredholmGermTame f x` (Definition 3.50, "tame germ data at $x$"): $x$ is a smooth point and there are the data of Definition 3.11 (V3.13: a strong bundle chart centred at $x$, a filling germ $g$ of the principal part, a local sc$^+$-germ $s$ with $s(0) = g(0)$, and $g - s$ conjugated near $0$ to a basic germ; the text's germ reading) in a tame chart. `IsGoodPositionGermTame f x` ("good position in a tame chart"): the same data with the germ in good position in that chart (Definition 3.53 in the chart form of V3.47). `IsScFredholmSectionTame f` (Definition 3.51, "tame germ data"): $f$ is an sc-Fredholm section (V3.16) with tame germ data at every smooth point. Each of these implies the form without tameness (V3.13, V3.47), good position in a tame chart gives tame germ data, and the strengthened good position of V3.82 is stated in a tame chart and implies good position in it. A retraction agreeing on its domain with a tame retraction is tame; hence a compatible chart whose base retraction is the identity of $U$ (the retract $O = U$ relatively open in the partial quadrant) is tame, so the hypotheses can be met. *Book:* Definition 3.50, p. 110 (a strong bundle chart around $x$ "where $K \to O$ is a tame strong local bundle", as in Definition 3.6, p. 66); Definition 3.51, p. 110, and Definition 3.53, p. 111 (for a tame strong bundle); a strong bundle retraction is tame when its $r$ is (p. 49); the proof of Proposition 3.52 starts from "a tame strong bundle chart" (p. 110). *Hidden hypotheses:* none. *Deviation (hypotheses explicit):* these are the hypotheses of the implicit function theorems of §3.6 (V3.48, V3.68–V3.70, V3.80–V3.82, V3.84–V3.87, V3.94–V3.102), in place of the reading of "tame strong bundle" as the tameness of the base retraction of every compatible chart, under which they were first proved and which cannot be met (V3.104). Whether germ data in some chart (V3.13) can be moved to some compatible tame chart when the atlas consists of tame charts is open; it cannot be done by re-centring a chart of the tame atlas, since whether a filling exists in a chart depends on the chart: the trivial bundle with a one-chart tame atlas has sc-Fredholm sections with tame germ data in compatible charts and no germ data in the chart of the atlas (Findings, Chapter 3, items 350–353). The text builds the tame chart into Definition 3.50, and so do the theorems. Since the switch to the text's germ reading (phase 2) the three notions carry the text's germ data (V3.13, V3.45); the earlier whole-chart forms are `IsScFredholmGermTameWholeChart`, `IsGoodPositionGermTameWholeChart`, `IsScFredholmSectionTameWholeChart`, and the non-vacuity witnesses satisfy both readings (Findings, Chapter 3, items 220–221, 304, 330). For Definition 3.53 the comparison is that of V3.47. Formerly "restated", for the whole-chart reading (round-trip item V3.103).

**V3.104 (a retraction onto a product with a point that is not tame: the earlier tameness hypothesis is unsatisfiable; `Polyfold.ScRetraction.halfLineConst_prod_not_isTame`, `Polyfold.ScRetraction.halfLineConst`, `Polyfold.ScRetraction.halfLineConst_retract`, `Polyfold.halfLineSc`).** Let $W$ be an sc-Banach space and $c$ the constant retraction $(a, w) \mapsto (1, 0)$ of the partial quadrant $[0, \infty) \oplus W$ of $\mathbb{R} \oplus W$ (sc-smooth, $(1, 0)$ being a smooth point), with retract $\{(1, 0)\}$. For every sc-smooth retraction $r$ of a nonempty relatively open subset $U$ of a partial quadrant $C$, the product retraction $c \times r$ of $([0, \infty) \oplus W) \times U$, whose retract is $\{(1, 0)\} \times r(U)$, is not tame: at $((0, 0), u)$ with $u \in r(U)$ the degeneracy index is $1 + d_C(u)$, at its image $((1, 0), u)$ it is $d_C(u)$, against condition (1) of Definition 2.34. *Book:* not in the text (Definition 2.34, pp. 39–40; Definition 2.50, pp. 51–52). *Hidden hypotheses:* none. *Deviation (not in the book):* the retraction-level half of the argument (Findings, Chapter 3, item 190) that the hypothesis under which the theorems of §3.6 were first proved, "the base retraction of every strong bundle chart compatible with the atlas is tame", fails for every strong bundle over a nonempty base: from a compatible chart $(\varphi, U, R, \Phi)$, the chart $(1, \varphi)$ onto $\{1\} \times O$ with base retraction $(a, u) \mapsto (1, r(u))$ on $[0, \infty) \times U$ (the retraction above, with $W = 0$) has the transition maps of the first chart composed with affine maps, hence is compatible with every chart the first one is. That chart and its compatibility are

not formalized (see "What is not verified"); so the earlier statements, vacuous by this argument, are replaced by those with tame germ data (V3.103).

**V3.105 (Theorem 3.15 for tame germ data;** `Polyfold.theorem_3_15_tame`, `Polyfold.StrongBundle.IsScFredho`
If $f$ is an sc-Fredholm section with tame germ data (V3.103) and $s$ an sc$^+$-section, then $f+s$ is an sc-Fredholm section with tame germ data: at every smooth point the germ data of $f+s$ are taken in the tame chart of those of $f$ (filling germ $g+\widehat{s}\circ r$, sc$^+$-germ $\widehat{s}_1\circ r+t_0$, the same conjugation, as in V3.17). *Book:* Theorem 3.15, p. 71, recalled after Definition 3.51, p. 110, for the tame strong bundle of §3.6. *Hidden hypotheses:* none. *Deviation (hypotheses explicit):* Theorem 3.15 for the tame reading of Definitions 3.50, 3.51; the proof of V3.17 does not change the chart.

**V3.106 (Theorem 3.18 for tame germ data;** `Polyfold.theorem_3_18_tame`, `Polyfold.PullbackChartData.ofPro`
`Polyfold.StrongBundle.IsScFredholmGermTame.param`, `Polyfold.StrongBundle.IsScFredholmSectionTame.`
In the setting of V3.91, if $f$ is an sc-Fredholm section with tame germ data (V3.103) and $s_1,\dots,s_n$ are sc$^+$-sections, then $F(r,x)=f(x)+\sum_i r_i s_i(x)$ is an sc-Fredholm section of $\pi^*P$ with tame germ data. On the way: the product chart $(r\mapsto r-r_0)\times c$ of a strong bundle chart $c$ of $P$ (V3.92) has the base retraction $1\times r$ on $\mathbb{R}^n\times U$, which is tame when $r$ is ($\mathbb{R}^n$ carries the identity retraction of the whole space, so $d(a,u)=d_C(u)$, and the minimal subspaces and tangent spaces are products); so $\widetilde{f}(r,x)=f(x)$ has germ data at every smooth $(r_0,x)$ in the tame product chart, and V3.105 adds $\widetilde{s}$. *Book:* Theorem 3.18, p. 74 (for a tame strong bundle). *Hidden hypotheses:* none. *Deviation (hypotheses explicit):* the tame form of V3.91, which is what the implicit function theorems of §3.6 need when applied to $F$; the tameness of the base $\mathbb{R}^n\times X$ (`prod_rn_isTame`, V3.91) is a different statement.

**V3.107 (good position in any chart of the germ data;** `Polyfold.StrongBundle.isGoodPositionAtGerm_of_chart`
`Polyfold.StrongBundle.isGoodPositionAtGerm_of_chart_dense`, `Polyfold.StrongBundle.IsGoodPosition`
(a) Let $f$ be an sc-smooth section, $y$ a zero, $c$ a strong bundle chart of $P$ around $y$ (not necessarily centred), $z$ a level-1 point over $\varphi(y)$, and $g$ an sc$^1$-germ agreeing with $\mathbf{f}$ on $O$ near $\varphi(y)$, with $z$ in its germ, such that $Dg(z)$ maps $T_zO$ onto $K_z$ and $\ker Dg(z)\cap T_zO$ is in good position to $C_{\varphi(y)}$ inside $T_zO$. Then for **every** datum of Definition 3.11 at $y$ (a strong bundle chart $c'$ compatible with the atlas and centred at $y$, a filling germ $g'$ with a derivative family $Dg'$) the germ is in good position in $c'$ at $Dg'(0)$; with only $\rho(z)k$ in the range for every smooth $k\in F_\infty$ (the hypotheses of V3.79), the same when $Dg'(0)$ is Fredholm. (b) If $(f,y)$, $f(y)=0$, is in good position in some chart (V3.47) and has tame germ data at $y$ (V3.103), it is in good position in that tame chart. *Book:* Definition 3.53, p. 111, presupposes that good position does not depend on the chart; the proof of Theorem 3.54, p. 115 ("proceeding as above"). *Hidden hypotheses:* none. *Deviation (not in the book):* V3.79 gives the good position in the chart that Definition 3.11 provides at $y$; (a) gives it in every chart carrying germ data at $y$. (b) applies (a) at $z=0$ in the chart of the given good position, centred at $y$, where 0 lies in every germ, $Tf(0)$ maps onto $K_0$, $\ker Dg(0)\subseteq T_0O$ and $C_0=C$; in the germ reading no Fredholm input is needed (Findings, Chapter 3, item 301). This is how the good position of the nearby zeros in Theorem 3.54 (1) (V3.94) and the good position from surjectivity (V3.100) are obtained in tame charts. The whole-chart form of (a) is `Polyfold.StrongBundle.isGoodPositionAt_of_chart_dense`.

**What is not verified in this chapter**

- pp. 111–114, Theorem 3.54 (1): nothing of the statement; the proof is not the text's (V3.93), and the first forms under the filler condition at the nearby zeros (V3.68, V3.69, V3.82, V3.85) are kept beside the forms with the text's hypotheses (V3.94, V3.95). Of the perturbation argu-

ment of the text's Proposition 6.2, used for the surjectivity at the nearby zeros, only the linear algebra is formalized (V3.83); for the index it is formalized (V3.98).

- p. 201 (Chapter 6), Proposition 6.2 for paths sc-smooth on $[0,1]$ only: it is verified with the text's hypotheses (tame germ data, no bump functions, Definition 6.1 with local sections) for paths on $\mathbb{R}$ (V3.98); a path on $[0,1]$ that does not extend sc-smoothly to $\mathbb{R}$ is not covered, the project having no M-polyfold $[0,1]$. Without tame germ data the local constancy of the index at the zeros is open (Findings, Chapter 3, items 243, 343).
- p. 74, Theorem 3.19: "ker $f'(y) = T_yS(f,U)$" with $T_yS$ a subspace of $T_yX$; it is verified in the chart of the tangent germ data (V3.99).
- p. 89, Remark 3.36 in quantitative form: its content is V3.26 and V3.31.
- p. 110, Proposition 3.52 on $T_xX \to Y_x$ itself: verified in charts, as an sc-Fredholm operator $T_zO \to K_z$ in every chart (V3.46, V3.96); $T_xX$ carries no linear structure here (Chapter 2).
- pp. 116–120, Theorem 3.58 for a general sc-splitting $E = B \oplus X$, $F = A \oplus Y$: verified in the coordinates of the text's proof (V3.89).
- pp. 121–125, Lemmas 3.62–3.64, 3.66–3.68 and p. 139, Lemma 3.76: superseded by the formal proofs of Proposition 3.22 and Proposition 3.73, not formalized.
- p. 135, Definition 3.71: Mathlib's $C^k$ within the set is used, equal to it on these sets.
- Lemma 3.70 in a general presentation of $C$ (hence $d_M = d_X$ in Theorem 3.27); Lemma 3.55 is verified in the model, in coordinates, and for a general presentation of $C$ (V3.49, V3.80).
- Theorem 3.27 for a set whose data do not satisfy condition (b) of V3.24 (it would need a re-graphing along another complement); for solution sets (b) holds by the choice of the complement (V3.77, V3.84).
- The linearization $f'(x) : T_xX \to Y_x$ and good position in $T_xX$ (Definition 2.38, gap G.13 of Chapter 2): Definition 3.53 is verified in charts, with its chart independence at zeros (V3.78).
- The chart-level half of the argument that the earlier hypothesis "the base retraction of every compatible strong bundle chart is tame" cannot be met (Findings, Chapter 3, items 190, 199): that the strong bundle chart $(1,\varphi)$ with base retraction $(a,u) \mapsto (1, r(u))$ on $[0,\infty) \times U$ is a chart compatible with every chart $(\varphi, U, R, \Phi)$ is (sc-calculus for affine maps, not done). Only the non-tameness of its base retraction is verified (V3.104); the theorems no longer use the hypothesis.
- Definition 3.11 $\Rightarrow$ Definition 3.50 (Findings, Chapter 3, items 199, 353): whether in a strong bundle whose atlas consists of tame charts every sc-Fredholm germ has germ data in some compatible tame chart is open. Re-centring a tame chart of the atlas does not do it: a filling needs $\ker Dr(0) \cong \ker \rho(0)$ in its chart, and a one-chart tame atlas of the trivial bundle carries no germ data at all (V3.103). The text builds the tame chart into Definition 3.50, and the theorems of §3.6 assume tame germ data.
- §3.1: Definitions 3.4, 3.6, 3.11, 3.14 (and 3.48, 3.50, 3.51, 3.53) are in the text's germ reading since phase 2 (V3.7, V3.9, V3.13, V3.16, V3.43, V3.45, V3.47), and Definition 3.3 since the fourth addendum (V3.5). Whether the text's germ reading and the project's earlier whole-chart reading of Definition 3.11 agree for sections sc-smooth on $X$ is open; no theorem depends on it (Findings, Chapter 3, items 220–221, 264, 330).

---

## 5. Chapter 4 (HWZ §§4.1–4.4): verified except Lemma 4.10 and §4.4

Chapter 4 of Part I (preprint pp. 147–168) asks when an M-polyfold is an ordinary manifold with boundary with corners. §4.1 defines a compatible manifold structure with corners (Definition 4.1),

states the characterization (Theorem 4.2: $X_0 = X_\infty$, $1_0^1 : X \to X^1$ sc-smooth, and centred tame charts whose tangent space at the centre is finite-dimensional and in good position), and prepares its proof: Remark 4.3, sc$^+$-retractions and sc$^+$-retracts (Definitions 4.4, 4.6, Lemmas 4.5, 4.7, Proposition 4.8), and the description of an sc$^+$-retract near a point as a graph over its tangent space (Proposition 4.9, with Lemmas 4.10–4.12). §4.2 defines smooth finite-dimensional submanifolds (Definition 4.13, Remark 4.14, Proposition 4.15, an open question), good position and tameness of submanifolds (Definitions 4.16, 4.17), proves that a submanifold in good position is a tame manifold with corners (Theorem 4.18, Lemma 4.19, Remark 4.20), defines general position (Definition 4.21) and describes a submanifold near a point of general position (Theorem 4.22). §4.3 proves, with Sard's theorem, that the fibres of a family $A \subseteq \mathbb{R}^m \times X$ over almost every parameter are compact manifolds with corners in general position (Theorem 4.23, Remark 4.24). §4.4 sketches sc-differential forms, de Rham cohomology and Stokes' theorem (Definitions 4.25, 4.27, Proposition 4.26, Theorem 4.28).

What is verified. **Theorems 4.2, 4.18, 4.22 and 4.23 hold with the text's hypotheses**, and Theorem 4.2 holds without its tameness hypothesis (V4.5). Every numbered statement of §§4.1–4.3 is verified except Lemma 4.10, which is no longer used: Definition 4.1 (V4.1–V4.4), Theorem 4.2 (V4.5, V4.27, V4.28), Remark 4.3, with an example for "$X^1 \neq X$ is possible", which the text does not give (V4.6), Definitions 4.4, 4.6 and Lemmas 4.5, 4.7 (V4.9–V4.12), Proposition 4.8 with its chart form (V4.13, V4.14), Proposition 4.9 with Lemmas 4.11 and 4.12 (V4.20–V4.25), Definition 4.13 (V4.29), Remark 4.14 in a chart (V4.31, V4.32), Proposition 4.15 (V4.33–V4.36), Definitions 4.16, 4.17, 4.21 in charts with their chart independence (V4.38–V4.40, V4.44, V4.45), Theorem 4.18, Lemma 4.19, Remark 4.20 (V4.41–V4.43), Theorem 4.22 (1)–(4) (V4.49–V4.52), Theorem 4.23 (V4.54, V4.63, V4.64) and Remark 4.24 (V4.55). Four places of the text needed more than a reading. (a) Proposition 4.9 is proved with two hypotheses the text leaves implicit — the good complement $Y$ is closed, and $p$ is the projection along $Y$, not $Dt(0)$ — and by a direct argument, because the text's route through Lemma 4.10 and Corollary 3.47 yields the graph map only as a germ over a complement of its own choosing (V4.20); where Proposition 4.9 is applied the closed complement is chosen, so Theorems 4.2, 4.18, 4.22, 4.23 need nothing extra. (b) One step of Lemma 4.19 is false ("$z \in Z \subset C$" does not give $n + y \in C$) and is repaired (V4.42). (c) Theorem 4.22 (4) has no proof in the text and is proved (V4.51, V4.52). (d) In Theorem 4.23 "regular value" is the stratified notion of the proof — every stratified regular value is a regular value, and with ordinary regular values the conclusion fails at the value — and the finite cover of the proof does not exist in general; a countable one is used (V4.53, V4.54, V4.64). Proved although the text does not construct or state it: charts and strong bundle charts centred at a smooth point (V4.15–V4.18), the sc$^+$-retraction onto the fibre $A_\lambda$ and its general position (V4.61, V4.62), Theorem 3.78 in $C^\infty$ form (V4.58), and Sard's theorem in all dimensions (V4.56, vendored from our Milnor project, `math_faithful`). Not verified: Lemma 4.10, §4.4, the open question after Proposition 4.15, and the chart independence of stratified regularity (see the list at the end of this section). A fourth addendum (2026-11-02; Findings, items 140–146) added the example of Remark 4.3 to V4.6.

The setting. An sc$^+$-map is a map whose values lie in $F_1$ and whose lift into $F^1$ is sc-smooth (V4.7); the text uses the word for maps and defines it only for retractions. $T_xX$ carries no linear structure in the formalization (gap G.7 of Chapter 2), so every condition the text states in $T_aX$ — the tangent space of a submanifold, good and general position, stratified regularity — is stated in a chart and, except for stratified regularity, proved independent of the chart. A chart "with $\psi(x) = 0$" is a chart re-centred at the smooth point $x$, modelled in the partial quadrant $C_{\varphi(x)}$ of Definition 2.33 (V4.17). The map $1_0^1 : X \to X^1$ is defined on $X_1$, with an arbitrary value off $X_1$ that plays no

role. Tameness of $X$ enters §4.2 only through Lemma 4.19 (Theorem 4.18, Remark 4.20), the tame centred chart of Theorem 4.22 and Proposition 2.36 (general position implies good position, V4.48). Findings referred to below are those of `Exposition/Findings.md`, "Chapter 4".

**V4.1 (Definition 4.1, corner charts and corner structures; `Polyfold.MChart.IsCornerChart`, `Polyfold.MPolyfold.HasCornerStructure`, `Polyfold.MPolyfold.hasCornerStructure_iff`, `Polyfold.MPolyfold.HasCornerStructure.isTame`, `Polyfold.MChart.IsCornerChart.isTameRetract`, `Polyfold.MChart.IsCornerChart.isScPlusRetract`).** A chart of $X$ is a *corner chart* if its local model is $(U, C, V)$ with $V$ a finite-dimensional real normed space carrying the constant sc-structure, $C$ a partial quadrant in $V$, $U$ relatively open in $C$, and the identity retraction. $X$ has a *corner structure* if it has an atlas of corner charts equivalent to its atlas; equivalently, every point of $X$ lies in a corner chart that is a chart of the M-polyfold structure. The dimension of $V$ may vary from chart to chart. The model of a corner chart is a tame retract and an sc$^+$-retract, and an M-polyfold with a corner structure is tame. *Book:* Definition 4.1, p. 147. *Hidden hypotheses:* "sc-smoothly compatible charts" with "$\varphi$ an sc-diffeomorphism" is read as compatibility with the atlas of $X$ (equivalence of atlases, as for tameness). *Deviation (restated):* the classical manifold structure with corners is represented by the atlas of corner charts; local constancy of the dimension is neither stated by the text nor formalized.

**V4.2 (Definition 4.1, the transition maps are $C^\infty$; `Polyfold.MChart.IsCornerChart.contDiffOn_transition`, `Polyfold.MChart.isRelOpenInQuadrant_overlap`).** For compatible corner charts $\chi$, $\chi'$ the overlap $\chi(V \cap V')$ is relatively open in the partial quadrant of $\chi$, and the transition map $\chi' \circ \chi^{-1}$ is of class $C^\infty$ on it (derivatives within the set, one-sided at the boundary). *Book:* Definition 4.1, p. 147, the paragraph after the definition. *Hidden hypotheses:* none. *Deviation:* "the atlas defines the structure of a smooth manifold with boundary with corners" is not formalized beyond this (Mathlib's manifolds with corners have a single model space).

**V4.3 (finite-dimensional sc-Banach spaces; `Polyfold.ScBanach.finiteDimensional_lvl`, `Polyfold.ScBanach.toBase_surjective`, `Polyfold.ScBanach.scIsoConst`, `Polyfold.ScBanach.mem_smooth_c`**
If $E_0$ is finite-dimensional, every level $E_m$ is finite-dimensional, every inclusion $E_m \to E_0$ is onto, $E$ is sc-isomorphic to $E_0$ with the constant structure (the level maps being the inclusions), and every point of $E_0$ is smooth. *Book:* §1.1, p. 6, after Definition 1.1 ("a finite-dimensional Banach space has precisely one sc-structure"), used in Definition 4.1. *Hidden hypotheses:* the density of $E_\infty$ in every level (Definition 1.1 (2)) is what makes the inclusions onto; compact injective inclusions alone do not ($E_0 = \mathbb{R}$, $E_m = 0$ for $m \geq 1$). *Deviation (restated):* "one sc-structure" as an sc-isomorphism onto the constant structure.

**V4.4 (Definition 4.1 with arbitrary finite-dimensional models, and the manifolds with corners of Chapter 3; `Polyfold.MChart.exists_isCornerChart_of_finiteDimensional`, `Polyfold.IsRelOpenInQuadrant.isScDiffeoQuadrant_toConst`, `Polyfold.MPolyfold.hasCornerStructure_i`, `Polyfold.MPolyfold.isManifoldWithCorners_iff`, `Polyfold.MPolyfold.IsManifoldWithCorners.hasCorner`**
A chart of $X$ onto a relatively open subset of a partial quadrant in an sc-Banach space with finite-dimensional $E_0$ is, with the same map and domain, a corner chart for the constant structure on $E_0$, of the same dimension. Hence $X$ has a corner structure iff it has an equivalent atlas of charts onto relatively open subsets of partial quadrants in sc-Banach spaces with finite-dimensional $E_0$ (the manifolds with corners of locally constant dimension of Theorem 3.27). The manifolds with corners of dimension $n$ of Theorem 3.27 are exactly the corner structures all of whose corner charts have dimension $n$. *Book:* Definition 4.1, p. 147 ("or in a finite-dimensional vector space"); the relation to Chapter 3 is not stated. *Hidden hypotheses:* none. *Deviation (not in the book).*

**V4.5 (Theorem 4.2, the characterization; `Polyfold.MPolyfold.theorem_4_2`, `Polyfold.MPolyfold.theorem_4_2`**
Let $X$ be an M-polyfold and $x_0 \in X_1$ the junk value of $1^1_0$ off $X_1$. The following are equivalent. (i) $X$ has a corner structure (V4.1). (ii) Every point of $X$ is smooth, $1^1_0 : X \to X^1$ is sc-smooth, and every $x \in X$ lies in a chart $\psi$ of $X$ with $\psi(x) = 0$ whose model $(O, C, E)$ is a tame retract and whose tangent space $N = T_0O$ (Definition 2.17) is finite-dimensional and in good position to $C$ (Definition 3.21). `theorem_4_2` is the text's form, with the hypothesis "$X$ tame". *Book:* Theorem 4.2, p. 147; proof pp. 156–157. *Hidden hypotheses:* the chart with $\psi(x) = 0$ is a re-centred chart (V4.17). *Deviation (stronger):* the tameness of $X$ is not used, and (i) implies it (V4.1). *Proof:* V4.27, V4.28.

**V4.6 (Remark 4.3; `Polyfold.ScBanach.liftOne`, `Polyfold.MPolyfold.toLevelOne`, `Polyfold.MPolyfold.remar`
`Polyfold.MPolyfold.remark_4_3_isScDiffeo`, `Polyfold.remark_4_3_const`, `Polyfold.remark_4_3_example`,
`Polyfold.remark_4_3_retract_example`, `Polyfold.IsLossFamily.isScSmooth`, `Polyfold.exists_retract_sm`
`Polyfold.MPolyfold.mem_shift_levelSet_iff`).** The inclusion $1^0_1 : X^1 \to X$ is always sc-smooth; in the charts $(V_1, \varphi, (O_1, C_1, E^1))$ and $\varphi$ it is the sc-operator $\iota : E^1 \to E$. If every point of $X$ lies in $X_1$ and $1^1_0 : X \to X^1$ is sc-smooth, then $1^1_0$ is an sc-diffeomorphism with inverse $1^0_1$. For a finite-dimensional $V$ with the constant structure, $V^1 = V$ as sc-Banach spaces, the lift $V \to V^1$ is the identity, and every sc-smooth retract in $V$ is an sc$^+$-retract. The levels of $X^1$ are those of $X$ shifted by one, as sets: $(X^1)_m = X_{m+1}$. **The example:** there is an M-polyfold all of whose points are smooth ($X_0 = X_\infty$) for which $1^1_0 : X \to X^1$ is not sc-smooth, so $X^1 \neq X$ as M-polyfolds. It is a retract $O \subseteq \mathbb{R} \oplus X$, $X$ the weighted sequence space of V1.13 ($|x|_m = \sup_k (k+1)^m |x_k|$), with a single chart: the image of $r(t, z) = (t, L(t)z)$, $L(t)$ the projection onto $\cos\theta\, e_{n_k} + \sin\theta\, e_{n_{k+1}}$ ($n_k = 2^k$), the angle $\theta$ turning from 0 to $\pi/2$ with flat ends while $t$ runs through $[1/(k+2), 1/(k+1)]$, and $L(t) = 0$ for $t \leq 0$. Its points are smooth, and the lift $O \to E_1$ is discontinuous at 0, so $O$ is not an sc$^+$-retract (Proposition 4.8). The sc-smoothness of $r$ comes from a criterion proved for it: for level operators $L(p)_m : Z_m \to W_m$ depending on a finite-dimensional parameter $p$, compatible with the inclusions, with $(p, z) \mapsto L(p)_m z$ continuous and $p \mapsto L(p)_m$ of class $C^\infty$ as a map into the operators $Z_{m+1} \to W_m$ (one level of loss), $(p, z) \mapsto L(p)z$ is sc-smooth. *Book:* Remark 4.3, p. 148. *Hidden hypotheses:* $1^1_0$ is defined on $X_1$ (the text: when $X_0 = X_\infty$). *Deviation (stronger):* the sc-diffeomorphism clause needs only $X = X_1$, not $X_0 = X_\infty$; "$X^1 \neq X$ is possible" has an example, which the text does not give, and in it $X$ and $X^1$ even have different topologies on the same set. "$X^1 \neq X$" is read as "$1^1_0$ is not sc-smooth", equivalently "the identity is not an sc-diffeomorphism". An example needs a retraction that loses a level in its first derivative, as the splicings of HWZ do (a retraction with $C^1$ level maps gives none: an argument, not formalized), so its sc-smoothness does not follow from Proposition 1.19. "$(X^1)_m = X_{m+1} = X_i$" misprints $X_0$; the topologies of the iterated levels are gap G.6 (Findings, items 5, 125, 140–146). Formerly "narrower", for the missing example (gap G.31, now closed).

**V4.7 (sc$^+$-maps; `Polyfold.IsScPlus`, `Polyfold.IsScPlus.mem_range_succ`, `Polyfold.IsScPlus.iff_exists`,
`Polyfold.IsScPlus.isScSmooth`, `Polyfold.IsScPlus.comp_isScSmooth`, `Polyfold.IsScSmooth.comp_isScPlu`
`Polyfold.ScPlus.isScPlus`).** A map $f$ on $U \subseteq E_0$ is *sc$^+$* if $f(U) \subseteq F_1$ and the lift $U \to F^1$, $x \mapsto f(x) \in F_1$, is sc-smooth. Then $f(U_m) \subseteq F_{m+1}$ for every $m$. On an sc-domain $U$: $f$ is sc$^+$ iff some sc-smooth $g : U \to F^1$ has $\iota \circ g = f$ on $U$; an sc$^+$-map is sc-smooth; $g \circ f$ is sc$^+$ if $g$ is sc$^+$ on an sc-domain $V$ and $f$ is sc-smooth with $f(U) \subseteq V$; $g \circ f$ is sc$^+$ if $f$ is sc$^+$ with $f(U) \subseteq V$ and $g$ is sc-smooth on $V$ (through Proposition 1.17 for $g$); an sc$^+$-operator is an sc$^+$-map. *Book:* not defined (Definition 4.4 is for retractions; the proof of Lemma 4.7 says "$r$ is a sc$^+$-map"). *Hidden hypotheses:* $U$, $V$ sc-domains for the composition rules. *Deviation (not in the book).*

**V4.8 (the derivative of an sc⁺-map; `Polyfold.IsScPlus.exists_scPlus_deriv`, `Polyfold.IsScPlus.isCompact`, `Polyfold.finiteDimensional_range_of_isCompactOperator_of_idem`).** Let $f$ be sc⁺ on an sc-domain $U$ and $x \in U_1$ a smooth point. There is an sc⁺-operator $S : E \to F$ whose level-0 operator is $Df(x)$; $Df(x) : E_0 \to F_0$ is compact, and so is every level operator of $S$. A compact continuous linear projection $P = P \circ P$ of a normed space has finite-dimensional range. *Book:* proof of Lemma 4.7 (ii), p. 149 ("a sc⁺-operator, since $r$ is a sc⁺-map … must be finite-dimensional"). *Hidden hypotheses:* $x$ smooth; the range of a compact projection is finite-dimensional by Riesz's theorem. *Deviation (hypotheses explicit).*

**V4.9 (Definition 4.4, sc⁺-retractions; `Polyfold.ScRetraction.IsScPlus`, `Polyfold.ScRetraction.IsScPlus.mem`, `Polyfold.MPolyfold.IsScPlusRetractionOn`).** An sc-smooth retraction $r : U \to U$ is an *sc⁺-retraction* if it is an sc⁺-map on $U$ (V4.7); then $r(U_m) \subseteq U_{m+1}$ for every $m$. On an open $V$ of an M-polyfold: $s$ is an sc⁺-retraction on $V$ if $s$ is sc-smooth on $V$, $s(V) \subseteq V$, $s \circ s = s$ on $V$, $s(V) \subseteq X_1$, and $1^1_0 \circ s : V \to X^1$ is sc-smooth (for every junk value). *Book:* Definition 4.4, p. 148. *Hidden hypotheses:* in the M-polyfold form, $s(V) \subseteq X_1$, so that "$r : V \to V^1$" makes sense. *Deviation (restated):* $r(U_m) \subseteq U_{m+1}$ is derived, not assumed.

**V4.10 (Lemma 4.5; `Polyfold.lemma_4_5`).** If $t$ is an sc⁺-retraction and $s$ an sc-smooth retraction (possibly of another relatively open subset of $C$) with the same retract, then $s$ is an sc⁺-retraction. *Book:* Lemma 4.5, p. 148. *Hidden hypotheses:* none. *Proof:* the text's, $s = t \circ s$ and sc⁺ □ sc-smooth is sc⁺.

**V4.11 (Definition 4.6, sc⁺-retracts; `Polyfold.IsScPlusRetract`, `Polyfold.IsScPlusRetract.isScPlus`, `Polyfold.isScPlusRetract_iff_forall`).** $(O, C, E)$ is an *sc⁺-retract* if $O$ is the retract of an sc⁺-retraction of a relatively open $U \subseteq C$; then every sc-smooth retraction onto $O$ is sc⁺, and $O$ is an sc⁺-retract iff it is a retract all of whose retractions are sc⁺. *Book:* Definition 4.6 and the remark after it, pp. 148–149. *Hidden hypotheses:* none.

**V4.12 (Lemma 4.7; `Polyfold.lemma_4_7_i`, `Polyfold.lemma_4_7_ii`, `Polyfold.lemma_4_7_ii_tangentSpace`).** Let $r$ be an sc⁺-retraction onto $O$. (i) $O \subseteq E_\infty$. (ii) For every level-1 point $o$ with $\iota(o) \in O$ the range of $Dr(o)$ is finite-dimensional; every $o \in O$ has a smooth lift $o_1 \in E_1$, and $T_{o_1}O = Dr(o_1)E_0$ is a finite-dimensional linear subspace. *Book:* Lemma 4.7, p. 149. *Hidden hypotheses:* none. *Deviation (hypotheses explicit):* the tangent space of Definition 2.17 is indexed by level-1 points; the step "$Dr(o)$ is an sc⁺-operator" is V4.8.

**V4.13 (Proposition 4.8; `Polyfold.proposition_4_8`).** An sc-smooth retract $(O, C, E)$ is an sc⁺-retract iff $O \subseteq E_\infty$ and the lift $1^1_0$ is sc-smooth from $(O, C, E)$ to $(O_1, C_1, E^1)$ in the sense of Definition 2.7. *Book:* Proposition 4.8, p. 149. *Hidden hypotheses:* none. *Deviation:* none (the proof ends "the lemma is proved").

**V4.14 (Proposition 4.8 in charts; `Polyfold.MPolyfold.levelChart_toLevelOne`, `Polyfold.MPolyfold.proposit`, `Polyfold.MPolyfold.isScSmoothMap_toLevelOne_iff`, `Polyfold.MPolyfold.isScSmoothMap_toLevelOne_of_`).** In the charts $\varphi$ of $X$ and $(V_1, \varphi, (O_1, C_1, E^1))$ of $X^1$ the map $1^1_0$ is the lift $O \to O_1$. For a chart $\varphi : V \to O$ of $X$: the model is an sc⁺-retract iff $V \subseteq X_\infty$ and $1^1_0$ is sc-smooth on $V$. Hence "every point smooth and $1^1_0$ sc-smooth" holds iff every chart of $X$ is modelled on an sc⁺-retract, iff every point lies in some chart so modelled. *Book:* not stated; used in the proofs of Theorem 4.2 and Proposition 4.15 (4). *Hidden hypotheses:* none. *Deviation (not in the book).*

**V4.15 (the partial quadrant $C_e$ and the neighbourhood $G(e)$; `Polyfold.recentreNbhd`, `Polyfold.recentreIso`, `Polyfold.isPartialQuadrant_partialQuadrantAtWith`, `Polyfold.image_sub_partia`, `Polyfold.recentreNbhd_inter_eq`, `Polyfold.degIndex_sub_recentre`, `Polyfold.minimalSubspaceWith_rece`**

`Polyfold.partialQuadrantAtWith_recentreIso`). Let $T$ present $C$ with coordinates $\lambda_1, \dots, \lambda_n$ and let $e \in E_0$. $G(e) = \{v : \lambda_i(v) > 0 \text{ whenever } \lambda_i(e) \neq 0\}$ is open and, for $e \in C$, contains $e$. $C_e$ is a partial quadrant, presented by reordering the coordinates as $\mathbb{R}^{I(e)} \oplus (\mathbb{R}^{\text{rest}} \oplus W)$; $C_e - e = C_e$; $G(e) \cap C = G(e) \cap C_e$; and for $v \in G(e)$ the vanishing coordinates of $v - e$ for this presentation are those of $v$, so that $d_{C_e}(v-e) = d_C(v)$ ($v \in C$) and the minimal subspace and the partial quadrant of $v - e$ are $E_v$ and $C_v$. *Book:* Definition 2.33, p. 39, asserts without proof that $C_e$ is a partial quadrant; the rest is not in the text. *Hidden hypotheses:* none. *Deviation (not in the book).*

**V4.16 (translations and re-centred retracts;** `Polyfold.isScSmooth_add_const`, `Polyfold.isScSmooth_sub_cons`
`Polyfold.relOpen_translate`, `Polyfold.isScDiffeoQuadrant_translate`, `Polyfold.ScRetraction.restrict`
`Polyfold.ScRetraction.restrictIn_isTame`, `Polyfold.isScRetract_recentre`, `Polyfold.isTameRetract_re`
Translation by a smooth vector is sc-smooth on every sc-domain. For $e$ smooth and $U$ relatively open in $C$ with $U \subseteq G(e)$, the translate $U - e$ is relatively open in $C_e$ and $v \mapsto v - e$ is an sc-diffeomorphism $U \to U - e$. The restriction of a retraction $r$ to $U \cap r^{-1}(W) \cap W$ ($W$ open) is a retraction onto $r(U) \cap W$, tame if $r$ is. For a retract (tame retract) $(O, C, E)$ and a smooth $e \in E_0$, $((O \cap G(e)) - e, C_e, E)$ is a retract (tame retract), with the retraction $u \mapsto r(u + e) - e$. *Book:* not in the text. *Hidden hypotheses:* $e$ smooth — translation by $e$ preserves every level only then. *Deviation (not in the book).*

**V4.17 (charts centred at a smooth point;** `Polyfold.MChart.recentre`, `Polyfold.MChart.recentre_isChart`,
`Polyfold.MChart.recentre_apply_self`, `Polyfold.MChart.isTameRetract_recentre`,
`Polyfold.MChart.chartDegIndex_recentre`, `Polyfold.MPolyfold.exists_centredChart`,
`Polyfold.MPolyfold.IsTame.exists_centredChart`). For a chart $\varphi : V \to O \subseteq C$ of $X$, a presentation of $C$ and a smooth $e$, the chart $\varphi_e = \varphi - e$ on $V \cap \varphi^{-1}(G(e))$, modelled on $((O \cap G(e)) - e, C_e, E)$, is a chart of $X$, with $\varphi_e(x) = 0$ when $\varphi(x) = e$, the same degeneracy index as $\varphi$ on its domain, and a tame model when $\varphi$ has one. So every smooth point $x$ of the domain of a chart $\varphi$ lies in a chart $\chi$ of $X$ with domain inside that of $\varphi$ and $\chi(x) = 0$; in a tame M-polyfold every smooth point lies in a tame chart centred at it. *Book:* used without construction in Theorem 4.2 (ii), the proofs of Theorems 4.18 and 4.22, and Lemma 5.11. *Hidden hypotheses:* $x$ smooth; the model lies in $C_{\varphi(x)}$, not in $C$. *Deviation (not in the book).*

**V4.18 (strong bundle charts centred at a smooth point;** `Polyfold.StrongBundleChart.recentre`,
`Polyfold.StrongBundleChart.Compatible.recentre`, `Polyfold.StrongBundle.IsChart.recentre`,
`Polyfold.StrongBundle.IsChart.exists_centred`). For a strong bundle chart $c$ over $\varphi$, a presentation and a smooth $e$, the chart $c_e$ over $\varphi_e$ with bundle retraction $R_e(u, h) = (r(u + e) - e, \Gamma(u + e, h))$ and $\Phi_e(y) = (\Phi(y)_1 - e, \Phi(y)_2)$ is compatible with every chart $c$ is compatible with (the projection being continuous), and is a chart of the strong bundle. Every smooth point of the base in the domain of a strong bundle chart lies in a strong bundle chart centred at it, with smaller domains. *Book:* used in Lemma 5.11. *Hidden hypotheses:* $e$ smooth; $(e, 0)$ is then smooth in $E \oplus F$ and $E \oplus F^1$. *Deviation (not in the book).*

**V4.19 (the level data of an sc⁺-retraction and the mean value estimate;** `Polyfold.ScRetraction.lvlMap`,
`Polyfold.ScRetraction.lvlDeriv`, `Polyfold.ScRetraction.IsScPlus.hasFDerivWithinAt_lvlMap`,
`Polyfold.ScRetraction.IsScPlus.continuousOn_lvlDeriv`, `Polyfold.ScRetraction.IsScPlus.exists_mvt`,
`Polyfold.exists_norm_sub_sub_le_of_hasFDerivWithinAt`). Let $t$ be an sc⁺-retraction and $k \geq 0$. Then $t : U_{k+1} \to E_{k+1}$ is differentiable within $U_{k+1}$, with derivative $Dt(x) \in \mathcal{L}(E_{k+1})$ continuous in $x$ in the operator norm. For $u_0 \in U_{k+1}$ and $\theta > 0$ there is $r > 0$ with $C_{k+1} \cap B(u_0, r) \subseteq U_{k+1}$, $\|Dt(z) - Dt(u_0)\| \leq \theta$ there, and $|t(z') - t(z) - Dt(u_0)(z' - z)|_{k+1} \leq \theta |z' - z|_{k+1}$ for $z, z' \in C_{k+1} \cap B(u_0, r)$. *Book:* (76)–(77) and the estimates in the proofs of Lemmas

4.11 and 4.12, pp. 154–155. *Hidden hypotheses:* the mean value inequality is taken on the convex set $C_{k+1} \cap B(u_0, r)$, since $U_{k+1}$ need not be convex. *Deviation (restated):* the text's integrals are the mean value inequality.

**V4.20 (Proposition 4.9; `Polyfold.proposition_4_9`, `Polyfold.finiteDimensional_tangentSpace_zero`, `Polyfold.GraphData`).** Let $t : U \to U$ be an sc$^+$-retraction onto $O \ni 0$, $N = T_0O$ (finite-dimensional by Lemma 4.7), in good position to $C$, $Y$ a good complement of $N$, $Y$ **closed**, and $p : E_0 \to N$ continuous linear with $p = 1$ on $N$ and $p = 0$ on $Y$ (**the projection along** $Y$). There are an open $W \ni 0$ in $E_0$, a relatively open $V \ni 0$ in the partial quadrant $N \cap C$ of $N$ (constant structure) and $\Gamma : N \to E_0$, sc-smooth on $V$, with $p \circ \Gamma = 1$ on $V$, $\Gamma(a) - a \in Y$, $\Gamma(V) = O \cap W$, $\Gamma \circ p = 1$ on $O \cap W$, $\Gamma(0) = 0$, and $p$, $\Gamma$ sc-smooth between the retracts $(O \cap W, C, E)$ and $(V, N \cap C, N)$: $p : O \cap W \to V$ is an sc-diffeomorphism with inverse $\Gamma$. *Book:* Proposition 4.9, p. 150; proof pp. 150–156. *Hidden hypotheses:* $Y$ closed (Definition 3.21 asks for an sc-complement, which is closed; the formal good complement records only the splitting), and $p$ the projection along $Y$: the text writes $p = Dt(0)$, a projection along $\ker Dt(0)$, which nothing relates to $Y$, while Lemma 4.12 needs the projection along $Y$ (Findings, items 20, 21). *Deviation (hypotheses explicit):* the conclusion is stated on fixed sets, with $\Gamma$ sc-smooth on the fixed $V$. *Proof:* not the text's. The text applies Corollary 3.47 to the section $\mathbf{f}(u) = (1-p)(u - t(u))$ (Lemma 4.10), which gives a graph over a complement of its own choosing, sc-smooth only as a germ (Findings, item 22); the formal proof is direct (V4.21–V4.25).

**V4.21 (Lemma 4.11, uniqueness; `Polyfold.GraphData.qL_lvlDeriv_zero`, `Polyfold.GraphData.lvlDeriv_zer` `Polyfold.GraphData.exists_contraction_radius`, `Polyfold.GraphData.eq_of_p_eq`).** In the data of V4.20, on every level $k+1$: $(1-p)Dt(0) = 0$ and $Dt(0) = 1$ on $N$; for $\theta > 0$ there is $r > 0$ for which the estimates of V4.19 hold at $u_0 = 0$ and $|(1-p)(u'-u)|_{k+1} \le \|1-p\|\theta|u'-u|_{k+1}$ for $u, u' \in O_{k+1}$ with $|u|_{k+1}, |u'|_{k+1} < r$; and two points $u, u'$ with $p(u) = p(u')$ and $|(1-p)(u'-u)| \le c|u'-u|$, $c < 1$, coincide. *Book:* Lemma 4.11, p. 154: small solutions $a + y$ of $\mathbf{f} = 0$ have $y = \delta(a)$. *Hidden hypotheses:* none. *Deviation (restated):* uniqueness among points of $O$, without the $\delta$ of Corollary 3.47.

**V4.22 (Lemma 4.12, existence; `Polyfold.GraphData.eps'`, `Polyfold.GraphData.exists_graph_point`).** Let $\varepsilon$ be the constant of the good complement $Y$ and $\varepsilon' = \varepsilon/(\|\iota_1\|\|\mathrm{lift}_N\| + 1)$. If the estimates of V4.21 hold on $C_1 \cap B(0, r)$ with $\theta < 1$, $\|1-p\|\theta < 1$ and $\|1-p\|\theta(1+\varepsilon') \le \varepsilon'$, there is $\rho > 0$ such that every $a \in N \cap C$ with $|a|_1 < \rho$ has a $u \in O_1$ with $p(u) = a$, $u - a \in Y$, $|u-a|_1 \le \varepsilon'|a|_1$ and $|u|_1 < r$. *Book:* Lemma 4.12, pp. 154–155: $a + \delta(a) \in O$ for small $a$. *Hidden hypotheses:* none. *Deviation (restated):* the point is constructed as the fixed point of $y \mapsto (1-p)t(a+y)$ on $\{y \in Y_1 : |y|_1 \le \varepsilon'|a|_1\}$, where $a + y \in C$ by the good complement property; then the text's argument $a_1 = a$ shows $t(u) = u$.

**V4.23 ($p(O \cap W) \subseteq C$ and the neighbourhoods; `Polyfold.GraphData.p_mem_C`, `Polyfold.GraphData.Nbhd`, `Polyfold.GraphData.exists_nbhd`, `Polyfold.GraphData.Nbhd.image_V`, `Polyfold.GraphData.Nbhd.injOn_p`).** (a) If $u \in O_1$ satisfies $|(1-p)u|_1 \le \kappa|u|_1$ with $0 \le \kappa \le 1/2$ and $2\|\iota_1\|\kappa\|\mathrm{lift}_N\| \le \varepsilon$, then $p(u) \in C$. (b) For every $\theta_0 > 0$ there are $r > 0$, an open $W = B_0(0, \delta_0) \cap p^{-1}(V')$, an open $V' = \{a : |a|_1 < \rho'\}$ of $N$ and $\Gamma$ on $V = V' \cap C$ with $\Gamma(V) \subseteq O \cap W$, $p \circ \Gamma = 1$, $\Gamma(a) - a \in Y$, $|\Gamma(a)|_1 < r$, $p(O \cap W) \subseteq V$, $\Gamma \circ p = 1$ on $O \cap W$, $\|Dt(z) - Dt(0)\|_{\mathcal{L}(E_1)} \le \theta_0$ on $C_1 \cap B(0, r)$, and $|(1-p)(u'-u)|_1 \le \frac{1}{2}|u'-u|_1$ on $O_1 \cap B_1(0, r)$; then $\Gamma(V) = O \cap W$ and $p$ is injective on $O \cap W$. *Book:* the last paragraph of the proof of Proposition 4.9, p. 156 ("$V = \{a \in T_0O \cap C : |a|_0 < \varepsilon\}$"). *Hidden hypotheses:* $p(u) \in C$ for $u \in O$ near 0, which the text never checks (Findings, item 25). *Deviation (not in the book):* the neighbourhoods are open on level 0 and controlled on level 1

through the continuity of $t : U \to E_1$ at 0.

**V4.24 (the regularity of $\Gamma$: continuity and the frame formula; `Polyfold.GraphData.Nbhd.norm_Γ_lvl_one_sub_`, `Polyfold.GraphData.Nbhd.continuousOn_Γ_lvl`, `Polyfold.norm_sub_le_of_projection`, `Polyfold.hasFDerivWithinAt_of_projection`, `Polyfold.GraphData.A_injective`, `Polyfold.GraphData.eq_z`, `Polyfold.GraphData.scDeriv_sub_frame`, `Polyfold.GraphData.Nbhd.hasFDerivWithinAt_Γ_lvl`).** With the data of V4.23: $|\Gamma(a') - \Gamma(a)|_1 \le 2\|\mathrm{lift}_N\||a' - a|$, and $\Gamma$ is continuous into every level $E_m$ on $V$. If $\|Dt(u) - Dt(0)\|_{\mathcal{L}(E_1)} \le \theta_0$ with $\theta_0$ small (`FrameSmall`), then $A(u) = p \circ Dt(u)|_N$ is invertible, $T_u O \cap Y = 0$ and $(1 - B_u p)Dt(u) = 0$ for the frame $B_u = Dt(u)|_N \circ A(u)^{-1}$; and $\Gamma$ is differentiable into every level $E_{k+1}$ on $V$ with $D\Gamma(a) = Dt(\Gamma(a))|_N \circ (p \circ Dt(\Gamma(a))|_N)^{-1}$. The abstract step: if $s$ satisfies the mean value estimate at $u_0$ on a set $S$, $q \circ B = 1$, $(1 - Bq)Ds(u_0) = 0$ and $\theta\|1 - Bq\| \le 1/2$, then fixed points $u, u'$ of $s$ in $S$ satisfy $|u' - u| \le 2\|B\||q(u' - u)|$, and every continuous right inverse of $q$ with values in the fixed points has derivative $B$ at $q(u_0)$. *Book:* not in the text (Corollary 3.47 gives $\delta$ as an sc-smooth germ). *Hidden hypotheses:* none. *Deviation (not in the book).*

**V4.25 (the $C^k$-bootstrap and the sc-diffeomorphism; `Polyfold.GraphData.contDiffOn_lvlDeriv_apply`, `Polyfold.GraphData.Nbhd.contDiffOn_Γ_lvl`, `Polyfold.GraphData.Nbhd.isScSmooth_Γ`, `Polyfold.GraphData.Nbhd.V_isRelOpenInQuadrant`, `Polyfold.GraphData.Nbhd.p_isScSmoothRetractMap`, `Polyfold.GraphData.Nbhd.Γ_isScSmoothRetractMap`).** $(x, w) \mapsto Dt_{j+1}(x)w$ is $C^k$ on $U_{j+k+1} \times E_{j+k+1}$ (Proposition 1.18 for the tangent map of the lift); $\Gamma$ is $C^k$ into every $E_{j+1}$ on $V$ for every $k$, hence $C^\infty$ into every level and sc-smooth on $V$ (Proposition 1.19); $(V, N \cap C, N)$ is a local model; $p$ and $\Gamma$ are sc-smooth between the retracts $(O \cap W, C, E)$ and $(V, N \cap C, N)$. *Book:* the last sentence of the proof of Proposition 4.9, p. 156 ("having the sc-smooth map $a \mapsto a + \delta(a)$ as its inverse"). *Hidden hypotheses:* none. *Deviation (not in the book):* derivatives of $t$ are applied only to vectors of the smooth finite-dimensional $N$, so the level loss of $Tt$ does not accumulate.

**V4.26 (charts composed with an sc-diffeomorphism between retracts; `Polyfold.MChart.compRetract`, `Polyfold.MChart.compRetract_isChart`).** For a chart $\varphi$ onto $(O, C, E)$ and mutually inverse $f$, $g$ that are sc-smooth between $(O, C, E)$ and a retract $(O', C', E')$ (Definition 2.7), $f \circ \varphi$ is a chart onto $(O', C', E')$, and a chart of the M-polyfold whenever $\varphi$ is. *Book:* not stated; "we obtain a smoothly compatible atlas", p. 157. *Hidden hypotheses:* none. *Deviation (not in the book).*

**V4.27 (Theorem 4.2, (i) ⇒ (ii); `Polyfold.MChart.IsCornerChart.mem_smooth`, `Polyfold.MChart.IsCornerChart`, `Polyfold.MChart.IsCornerChart.exists_centred_tame`, `Polyfold.tangentSpace_relOpen`, `Polyfold.isInGoodPosition_top`).** Every point of a corner chart is smooth for its model. A point $x$ in the domain of a corner chart $\chi$ of $X$ lies in a chart $\psi$ of $X$ with $\psi(x) = 0$, modelled on a relatively open subset of the partial quadrant $C_{\chi(x)}$ in the same finite-dimensional space; this model is a tame retract (identity retraction), its tangent space at 0 is the whole space, and the whole space is in good position to any partial quadrant (complement 0). Together with V4.14 and V4.1, (i) implies (ii). *Book:* proof of Theorem 4.2, p. 156. *Hidden hypotheses:* the centred chart, the tangent space at its centre and its good position, which the text's proof does not address (Findings, item 53). *Deviation (hypotheses explicit).*

**V4.28 (Theorem 4.2, (ii) ⇒ (i); `Polyfold.exists_proj_goodComplement`, `Polyfold.CornerChartData`, `Polyfold.exists_cornerChartData`, `Polyfold.CornerChartData.chart`, `Polyfold.CornerChartData.chart_i`, `Polyfold.CornerChartData.chart_isChart`, `Polyfold.MPolyfold.theorem_4_2_chart`, `Polyfold.MPolyfold.hasCornerStructure_of_centred`).** If every point is smooth and $1_0^1$ is sc-smooth, then for every chart $\psi$ of $X$ with $\psi(x) = 0$ whose tangent space $N = T_0 O$ is finite-dimensional and in good position to $C$ there are a closed good complement $Y$ of $N$, the

projection $p$ along it, and $W$, $V$, $\Gamma$ of Proposition 4.9; the chart $p \circ \psi$ on $\psi^{-1}(W)$, modelled on $(V, N \cap C, N)$, is a corner chart and a chart of $X$, with $x$ in its domain and $(p \circ \psi)(x) = 0$. If every point has such a chart $\psi$ (tame or not), $X$ has a corner structure. *Book:* proof of Theorem 4.2, pp. 156–157. *Hidden hypotheses:* none; the closed good complement is chosen by the normal form of Proposition 3.22, and the compatibility of the charts $p \circ \psi$, not argued in the text, is V4.26 (Findings, items 54, 55). *Deviation (hypotheses explicit).*

**V4.29 (Definition 4.13, smooth finite-dimensional submanifolds;** `Polyfold.MPolyfold.IsSmoothSubmanifold`, `Polyfold.MPolyfold.SubmanifoldData`, `Polyfold.MPolyfold.isSmoothSubmanifold_iff`, `Polyfold.MPolyfold.SubmanifoldData.toSubData`, `Polyfold.MPolyfold.IsSmoothSubmanifold.isSubMPolyf`
$A \subseteq X$ is a *smooth finite-dimensional submanifold* if every $a \in A$ has an open $V \ni a$ and an sc$^+$-retraction $s$ on $V$ (V4.9) with $s(V) = A \cap V$. Such an $A$ is a sub-M-polyfold (Definition 2.20) and carries the induced structure of Proposition 2.21. *Book:* Definition 4.13, p. 157. *Hidden hypotheses:* none.

**V4.30 (the sc$^+$-retraction in a chart;** `Polyfold.MPolyfold.SubmanifoldData.isScSmoothRetractMap_lift`, `Polyfold.MPolyfold.SubmanifoldData.exists_chartRetraction`, `Polyfold.MPolyfold.SubmanifoldData.is` `Polyfold.MPolyfold.SubmanifoldData.apply_mem_smooth`**).** Let $(V, s)$ be data of Definition 4.13, $\varphi$ a chart of $X$ and $W = V \cap \operatorname{dom}\varphi \cap s^{-1}(V \cap \operatorname{dom}\varphi)$. The lift of $\varphi \circ s \circ \varphi^{-1}$ to $E^1$ is sc-smooth from $(\varphi(W), C, E)$ to $(O_1, C_1, E^1)$. For every retraction $r$ onto $O$ there is an open $G$ with $\varphi(W) = O \cap G$ and an sc$^+$-retraction $t$ on $U \cap r^{-1}(G)$, equal to $\varphi \circ s \circ \varphi^{-1} \circ r$ there, onto $\varphi(W \cap A)$. So $\varphi(W \cap A)$ is an sc$^+$-retract, and $\varphi(a)$ is smooth for $a \in A \cap V \cap \operatorname{dom}\varphi$. *Book:* proof of Theorem 4.18, pp. 160–161 ("taking $V$ and $U$ small, we may assume $V = \varphi^{-1}(O)$ … it is a sc$^+$-retraction since $s$ is"). *Hidden hypotheses:* the set $W$ of Proposition 2.21 in place of "$V = \varphi^{-1}(O)$", and the sc$^+$ condition read in the charts of $X$ and $X^1$ (Findings, item 60). *Deviation (hypotheses explicit).*

**V4.31 (Remark 4.14 in a chart;** `Polyfold.MPolyfold.SubmanifoldData.remark_4_14`**).** At $a \in A \cap V \cap \operatorname{dom}\varphi$ and the level-1 point $z$ over $\varphi(a)$: $z$ is smooth; the tangent map of $\rho = \varphi \circ s \circ \varphi^{-1}$ at $z$ maps $T_zO$ onto the tangent space $T_z(\varphi(W \cap A))$; this tangent space lies in $T_zO$; and it is a finite-dimensional linear subspace. *Book:* Remark 4.14, p. 157 ("$T_aA = Ts(a)(T_aX)$ … finite-dimensional"). *Hidden hypotheses:* $T_aX$ is read in a chart (gap G.7). *Deviation (restated):* the sc$^+$-operator is $Dt(z)$ for the chart retraction $t$ of V4.30, not $Ts(a)$.

**V4.32 (Remark 4.14 at points with $d_X = 0$;** `Polyfold.isInGoodPosition_univ`, `Polyfold.partialQuadrantAtWi` `Polyfold.MPolyfold.isGoodPositionAt_of_degIndex_eq_zero`, `Polyfold.MPolyfold.hasCornerStructure_o`
Every subspace is in good position to the whole space; at a point $e \in C$ with $d_C(e) = 0$, $C_e$ is the whole space. If $a \in A$ and $d_X(a) = 0$, $A$ is in good position at $a$ (Definition 4.16, V4.38). If $d_X = 0$ on all of $A$, the induced structure on $A$ has a corner structure; $X$ need not be tame. *Book:* Remark 4.14, p. 157 ("near points $a \in A$ satisfying $d_X(a) = 0$, necessarily smooth finite-dimensional manifolds in the classical sense"). *Hidden hypotheses:* none. *Deviation (narrower):* the local form, for the open piece $A \cap \{d_X = 0\}$ of an arbitrary $A$, is not formalized (gap G.33).

**V4.33 (Proposition 4.15 (1);** `Polyfold.MPolyfold.IsSmoothSubmanifold.subset_smoothSet`, `Polyfold.MPolyfold.IsSmoothSubmanifold.mem_levelSet_subPolyfold_iff`, `Polyfold.MPolyfold.IsSmooth`
$A \subseteq X_\infty$; the levels of the induced structure are the traces of the levels of $X$; every point of $A$ is smooth for the induced structure. *Book:* Proposition 4.15 (1), p. 158. *Hidden hypotheses:* none. *Deviation (hypotheses explicit):* "$A \subseteq X_\infty$" and "$A = A_\infty$ for the induced structure" are separated; the proof goes through the chart retraction of V4.30 and Lemma 4.7 (i).

**V4.34 (Proposition 4.15 (2);** `Polyfold.MPolyfold.IsSmoothSubmanifold.proposition_4_15_2`**).**

$A$ is locally closed in $X$: the intersection of an open and a closed set (equivalently, every $a \in A$ has an open $V$ with $V \cap \overline{A} \subseteq A$). *Book:* Proposition 4.15 (2), p. 158, with a garbled definition of "locally closed" (the letter $U$ used twice). *Hidden hypotheses:* none. *Deviation (restated):* Mathlib's notion; a filter replaces the text's sequence, so no metrizability is needed.

**V4.35 (Proposition 4.15 (3); `Polyfold.MPolyfold.IsSmoothSubmanifold.proposition_4_15_3`).** The boundary $\{d_A \geq 1\}$ of $A$ with its induced structure lies in $\partial X = \{d_X \geq 1\}$. *Book:* Proposition 4.15 (3), p. 158. *Hidden hypotheses:* none. *Proof:* $d_A \leq d_X$ (Lemma 2.26).

**V4.36 (Proposition 4.15 (4); `Polyfold.MPolyfold.IsSmoothSubmanifold.proposition_4_15_4`).** For the induced structure on $A$, every point is smooth and $1^1_0 : A \to A^1$ is sc-smooth (for any junk value). *Book:* Proposition 4.15 (4), p. 158. *Hidden hypotheses:* none. *Proof:* every point of $A$ lies in an induced chart whose model $\varphi(W \cap A)$ is an sc$^+$-retract (V4.30), and V4.14 applies; the text says only that the sc$^+$-retraction "is the identity on the retract $A$".

**V4.37 (tangent spaces of sub-retracts; `Polyfold.tangentSpace_inter_isOpen`, `Polyfold.tangentSpace_subset`, `Polyfold.scDeriv_mem_tangentSpace_of_subset`).** For a retract $Q$ and an open $K$, $T_z(Q \cap K) = T_zQ$ at points of $Q \cap K$. For retracts $Q \subseteq O = r(U)$, $T_zQ \subseteq T_zO$. If $F \circ r$ is sc-smooth and $F$ maps the retract $Q \subseteq O$ into a retract $Q'$, then $D(F \circ r)(z)$ maps $T_zQ$ into $T_{F(z)}Q'$. *Book:* not stated. *Hidden hypotheses:* none. *Deviation (not in the book).*

**V4.38 (Definition 4.16 in a chart; `Polyfold.IsGoodPositionRetractAt`, `Polyfold.MPolyfold.SubmanifoldData`, `Polyfold.MPolyfold.IsGoodPositionAt`, `Polyfold.MPolyfold.SubmanifoldData.isGoodPositionChart_iff`).** For retracts $Q \subseteq O$, a presentation $T$ and a level-1 point $z$: $T_zQ$ is in good position to $C_z$ inside $T_zO$ ($T_zQ \subseteq T_zO$, $T_zQ \cap C_z$ has interior points in $T_zQ$, and $T_zQ$ has a complement $P \subseteq T_zO$ in $T_zO$ with a constant $\varepsilon$ as in Definition 3.21). $A$ is *in good position at* $a \in A$ if this holds for $Q = \varphi(W \cap A)$ and the model $O$ of some chart $\varphi$ at $a$, some data of Definition 4.13 at $a$, some presentation and the level-1 point over $\varphi(a)$. For retractions $r$ onto $O$ and $t$ onto $\varphi(W \cap A)$ the condition says that $Dt(z)E$ is in good position to $C_z$ inside $Dr(z)E$. *Book:* Definition 4.16, p. 159. *Hidden hypotheses:* the chart form (gap G.7); the chart form needs no tameness. *Deviation (restated).*

**V4.39 (the chart independence of Definition 4.16; `Polyfold.IsGoodPositionRetractAt.transport`, `Polyfold.MPolyfold.chart_change_of_transport`, `Polyfold.MPolyfold.isGoodPositionRetractAt_chart_c`, `Polyfold.MPolyfold.IsGoodPositionAt.isGoodPositionChart`, `Polyfold.MPolyfold.isGoodPositionAt_iff`).** Good position of a sub-retract is carried by a germ of sc-diffeomorphisms between retracts that maps the sub-retracts into each other. Hence if $A$ is in good position at $a$, the chart condition holds for every data of Definition 4.13 at $a$, every chart at $a$, every presentation and the level-1 point over the image of $a$; and good position at $a$ is equivalent to the condition for all of them. *Book:* implicit in Definition 4.16. *Hidden hypotheses:* none. *Deviation (not in the book).*

**V4.40 (Definition 4.17; `Polyfold.MPolyfold.IsTameSubmanifold`).** $A$ is a *tame* submanifold if it is a smooth finite-dimensional submanifold whose induced M-polyfold structure is tame. *Book:* Definition 4.17, p. 159. *Hidden hypotheses:* none.

**V4.41 (Theorem 4.18; `Polyfold.MPolyfold.theorem_4_18`, `Polyfold.MPolyfold.hasCornerStructure_of_cent`, `Polyfold.MPolyfold.IsGoodPositionAt.isInGoodPosition_of_tame`).** Let $X$ be tame and $A$ a smooth finite-dimensional submanifold in good position at every point. Then the induced structure on $A$ has a corner structure (Definition 4.1) and $A$ is tame. More generally, the induced structure has a corner structure as soon as every $a \in A$ lies in a chart $\psi$ of $X$ with $\psi(a) = 0$ in which $T_0(\psi(W \cap A))$ is in good position to $C$ in $E$; and good position at $a$ gives this in every tame chart centred at $a$. *Book:* Theorem 4.18, p. 159; proof pp. 160–161. *Hidden hypotheses:* none.

*Proof:* the text's route through Theorem 4.2, with the steps it skips: the centred chart (V4.17), the chart retraction (V4.30), the closed complement (V4.28) and the compatibility of the charts $p \circ \psi$ (V4.26); tameness of $X$ enters only through Lemma 4.19.

**V4.42 (Lemma 4.19; `Polyfold.isGoodComplement_sup_range`, `Polyfold.lemma_4_19`, `Polyfold.IsInGoodPosit`**
Let $T$ present $C$, $\pi$ a bounded idempotent with $(1-\pi)E_0 \subseteq E_0^{\min}$, the minimal subspace at $0$ ($\{\lambda = 0\}$), and $N$ in good position to $C$ inside $\pi E_0$. Then $N$ is in good position to $C$ in $E_0$, and for every good complement $Y$ of $N$ inside $\pi E_0$ with constant $\varepsilon$, $Y \oplus (1-\pi)E_0$ is a good complement of $N$ in $E_0$, with constant $\varepsilon/(\|\pi\|+1)$. *Book:* Lemma 4.19, p. 160. *Hidden hypotheses:* $(1-\pi)E_0$ lies in the minimal subspace at the corner $0$ of the centred chart (Proposition 2.35), not merely in $C$. *Deviation (hypotheses explicit):* the text concludes $n+y \in C$ from $n+y+z \in C$ "since $z \in Z \subset C$", which is false ($C = [0,\infty)$, $z = 1$, $n+y = -\frac{1}{2}$); it holds because $Z \subseteq C \cap (-C)$. The text's norm $|y|_0 + |z|_0$ is replaced by the constant $\varepsilon/(\|\pi\|+1)$ (Findings, item 69).

**V4.43 (Remark 4.20; `Polyfold.tangentSpace_eq_range_graph`, `Polyfold.lemma_3_55_relOpen`, `Polyfold.IsInGoodPosition.isInGoodPositionIn_of_le`, `Polyfold.MPolyfold.remark_4_20`).**
If the retract $Q$ is the image of an sc-diffeomorphism $\Gamma$ from $(V, N \cap C, N)$ with inverse $p$, then at $v \in V$ and the level-1 point $y$ over $\Gamma(v)$, $T_y Q = D\Gamma(v)N$ and $p \circ D\Gamma(v) = 1$. For $X$ tame and $A$ in good position at $a$, there is an open $U \ni a$ such that $A$ is in good position at every point of $U \cap A$. *Book:* Remark 4.20, p. 161, without "tame". *Hidden hypotheses:* $X$ tame (Definition 4.16 is stated for tame $X$; Lemma 4.19 is used). *Deviation (hypotheses explicit):* "the tangent spaces move only slowly" is $D\sigma(0) = 0$ for $\Gamma = 1 + \sigma$, and Lemma 3.55 is applied with the closed complement of the normal form.

**V4.44 (Definition 4.21 in a chart; `Polyfold.IsGeneralPositionRetractAt`, `Polyfold.MPolyfold.SubmanifoldD`
`Polyfold.MPolyfold.IsGeneralPositionAt`).** For retracts $Q \subseteq O$, a presentation $T$ and a level-1 point $z$: there is an sc-subspace $B \subseteq T_z O \cap E_z$ with $(T_z O)_m = (T_z Q)_m \oplus B_m$ on every level $m$ ($T_z O \cap E_z = T_z^R O$). $A$ is *in general position at* $a \in A$ if this holds for $Q = \varphi(W \cap A)$ and the model of some chart at $a$, some data, presentation and lift. *Book:* Definition 4.21, p. 161. *Hidden hypotheses:* the sc-structure of $T_z O$ is the one induced from $E$ at the smooth point; the chart form (gap G.7). *Deviation (restated).*

**V4.45 (the chart independence of Definition 4.21; `Polyfold.IsGeneralPositionRetractAt.transport`, `Polyfold.MPolyfold.isGeneralPositionRetractAt_chart_change`, `Polyfold.MPolyfold.IsGeneralPositio`
`Polyfold.MPolyfold.isGeneralPositionAt_iff_forall`).** General position of a sub-retract is carried by a germ of sc-diffeomorphisms at a smooth point; hence general position at $a$ holds in every chart, for every data, presentation and lift, and is equivalent to the condition for all of them. *Book:* implicit in Definition 4.21. *Hidden hypotheses:* the point is smooth (true on $A$). *Deviation (not in the book).*

**V4.46 (linear algebra at the corner of a centred chart; `Polyfold.mem_minimalSubspaceWith_zero`, `Polyfold.minimalSubspaceWith_zero_le`, `Polyfold.add_mem_iff_of_mem_zero`, `Polyfold.degIndexWith_add`
`Polyfold.coordsOn_surjective`, `Polyfold.exists_isClosed_isCompl_le`, `Polyfold.exists_goodComplement`
`Polyfold.degIndex_slice`).** For $C = T^{-1}([0,\infty)^n \oplus W)$: the minimal subspace at $0$ is $E_0^{\min} = \{\lambda = 0\}$ and lies in every $E_x$; adding a vector of $E_0^{\min}$ changes neither membership in $C$ nor the degeneracy index. For a finite-dimensional $N$ with $N + E_0^{\min} = E_0$: $\lambda|_N$ is onto $\mathbb{R}^n$, $N$ is in good position to $C$ with a closed good complement $Y \subseteq E_0^{\min}$, and $d_{N\cap C}(v) = d_C(v)$ for $v \in N \cap C$. A finite-dimensional $N$ and a closed $K$ with $N + K$ the whole space give a closed complement of $N$ inside $K$. *Book:* proof of Theorem 4.22, pp. 162–163 ("$T_0 A$ has in $E$ an sc-complement $Y$ contained in $W$"). *Hidden hypotheses:* $Y$ closed. *Deviation (restated):* the text's

splitting $T_0A = N \oplus (T_0A \cap W)$ is replaced by the coordinates $\lambda|_{T_0A}$ (Findings, items 82, 83).

**V4.47 (Lemma 7.A of the exposition, and general position from $T_yQ + E_0^{\min} = E_0$; `Polyfold.exists_scComplement_le`, `Polyfold.isGeneralPositionRetractAt_of_sup_eq_top`).** (a) For $N \subseteq E_\infty$ finite-dimensional, an sc-subspace $R$ and $M = N + R$, there is an sc-subspace $B \subseteq R$ with $M_m = N_m \oplus B_m$ on every level. (b) For $r$ tame, a retract $Q \subseteq O$, a smooth $y \in Q_1$ with $T_yQ$ finite-dimensional, contained in $E_\infty$ and $T_yQ + E_0^{\min} = E_0$, the retract $Q$ is in general position at $y$ in $O$. *Book:* proof of Theorem 4.22 (3), p. 164 ("$T_aO = T_aA + T_a^RO$, implying that $T_aA$ has in $T_aO$ a sc-complement contained in $T_a^RO$"). *Hidden hypotheses:* $T_y^RO$ an sc-subspace (Proposition 2.36, for tame retracts). *Deviation (hypotheses explicit).*

**V4.48 (general position implies good position; `Polyfold.IsGeneralPositionRetractAt.isGoodPositionRetra`).** For $r$ tame and a smooth $z \in O_1$, if $Q$ is in general position at $z$ in $O$, it is in good position at $z$. *Book:* not stated; needed for the tameness in Theorem 4.22 (4) and for Theorem 4.23. *Hidden hypotheses:* none. *Deviation (not in the book).*

**V4.49 (Theorem 4.22 in a tame chart centred at the point; `Polyfold.sup_minimalSubspaceWith_eq_top`, `Polyfold.sup_minimalSubspaceWith_eq_top_graph`, `Polyfold.theorem_4_22_chart`).** Let $r$ be tame onto $O \ni 0$, $t$ an sc$^+$-retraction onto $Q \subseteq O$ with $0 \in Q$, and $N = T_0Q$, with $Q$ in general position at 0. Then $N + E_0^{\min} = E_0$, $N$ is in good position to $C$, $N$ has a closed good complement $Y \subseteq E_0^{\min}$ with a projection $p$ whose kernel is exactly $Y$, and the data $W$, $V$, $\Gamma$ of Proposition 4.9 exist for this $Y$, with (2) $d_{N\cap C}(v) = d_C(\Gamma(v))$ on $V$ and (3) general position of $Q$ at every level-1 point over $Q \cap W$. *Book:* proof of Theorem 4.22, pp. 162–164. *Hidden hypotheses:* none. *Deviation (restated):* $T_yQ + E_0^{\min} = E_0$ along the graph comes from $p \circ D\Gamma(v) = 1$ and $\ker p = Y$ (the text: "by means of the particular form of the map $\alpha$").

**V4.50 (Theorem 4.22, local structure in general position; `Polyfold.MPolyfold.theorem_4_22`).** Let $X$ be tame, $A$ a smooth finite-dimensional submanifold, in general position at $x \in A$. There are a chart $\psi$ of $X$ with $\psi(x) = 0$, a tame retraction $r$ onto its model for a presentation $T$ with $n$ half-line coordinates, and an open $U \ni x$ inside the domain of $\psi$ with: (1) $d_X(x) = n$, and $\psi(x)$ lies in exactly $d_X(x)$ faces of the M-polyfold $O$; (2) $d_X(a) = d_A(a)$ for $a \in A \cap U$ ($d_A$ for the induced structure); (3) $A$ is in general position at every point of $A \cap U$; (4) for every $\sigma \subseteq \{1, \dots, n\}$ there are faces $F_i \ni 0$ of the retract ($i \in \sigma$, $F_i$ in the coordinate hyperplane $\lambda_i = 0$) and an open $G \ni 0$ such that $F_\sigma = \psi^{-1}(G \cap \bigcap_{i\in\sigma} F_i)$ contains $x$, lies in $U$, and is a tame sub-M-polyfold with $d_{F_\sigma} + \#\sigma = d_X$ on it, and $A \cap F_\sigma$ is a smooth finite-dimensional submanifold of $F_\sigma$, in general position in $F_\sigma$ at each of its points, and tame. *Book:* Theorem 4.22, p. 162; proof pp. 162–164. *Hidden hypotheses:* the chart is re-centred at $x$; "smooth submanifold" is finite-dimensional. *Deviation (restated):* (1) is a statement about the chart model (local faces are not identified with global faces, gap G.35); $F_\sigma$ is a germ, taken inside $U$; (4), without proof in the text, is V4.51, V4.52. (1) and (4) differ from the text's in content, not only in form: the text's local faces are faces of the neighbourhood $U$ itself (the closures in $U$ of the components of $U \cap \{d_X = 1\}$, all through $x$), the formal ones faces of the model retract of the chart; (2) and (3) are the text's. Identifying the two needs a local face theorem for small neighbourhoods, believed true for tame $X$ and not formalized (Findings, Chapter 4, items 90, 130; round-trip item V4.50).

**V4.51 (Theorem 4.22 (4) in the chart; `Polyfold.killOn`, `Polyfold.exists_scPlus_retraction_corner`, `Polyfold.exists_cornerModel_retraction`, `Polyfold.theorem_4_22_four_chart`).** In the situation of V4.49, for $\sigma \subseteq \{1, \dots, n\}$ and an open $B \ni 0$: with $\kappa_\sigma(v) = v - s(\lambda(v)1_\sigma)$ ($s$ a linear section of $\lambda|_N$) there are an open $G \ni 0$, $G \subseteq W \cap B$, and an sc$^+$-retraction $t_\sigma = \Gamma \circ \kappa_\sigma \circ p \circ t$ onto $Q \cap G \cap f_\sigma$, $f_\sigma = \{\lambda_i = 0, i \in \sigma\}$, whose tangent spaces contain $T_yQ \cap f_\sigma$. A retraction

onto a subset of $f_\sigma$ has a corner model on $[0,\infty)^{n-\#\sigma}\oplus W$, sc$^+$ if the retraction is. For every open $B_0\ni 0$ there are faces $F_i\ni 0$ ($i\in\sigma$) and an open $G\ni 0$, $G\subseteq W\cap B_0$, such that $O\cap G\cap f_\sigma$ is the retract of a retraction whose corner model is tame with $d_{C'}(\pi_\sigma y)+\#\sigma=d_C(y)$, $Q\cap G\cap f_\sigma$ and its corner model are sc$^+$-retracts, and the corner model of $Q\cap G\cap f_\sigma$ is in general position at each of its level-1 points. *Book:* not in the text ("the previous arguments, but now applied to $A\cap F_\sigma$, conclude the proof"). *Hidden hypotheses:* none. *Deviation (not in the book).*

**V4.52 (sc$^+$-retractions in a chart, and Theorem 4.22 (4) for one $\sigma$; `Polyfold.MPolyfold.isScPlusRetractionOn_o` `Polyfold.MChart.cornerPreimage`, `Polyfold.MPolyfold.theorem_4_22_four_of_chart`).** For a chart $\chi$ of $X$ and an sc$^+$-retraction $\tau$ of its model with $\tau(U)\subseteq O$, $\chi^{-1}\circ\tau\circ\chi$ is an sc$^+$-retraction on the open set $\chi^{-1}(\tau.U)$ with image $\chi^{-1}(\tau(U))$. In a tame chart $\psi$ centred at $x$ with the data of V4.49 for the chart retraction of $A$, for every $\sigma$ there are faces $F_i\ni 0$ and an open $G\ni 0$ such that $\psi^{-1}(G\cap\bigcap_{i\in\sigma}F_i)$ contains $x$, lies in the open set $W$ of the submanifold chart, and has the properties (4) of V4.50. *Book:* not in the text. *Hidden hypotheses:* none. *Deviation (not in the book).*

**V4.53 (the family $\mathbb{R}^m\times X$ and stratified regular points; `Polyfold.prod_rn_isTame`, `Polyfold.fibreSet`, `Polyfold.rnPres`, `Polyfold.IsStratRegularPoint`, `Polyfold.IsStratRegularValue`).** For tame $X$, $\mathbb{R}^m\times X$ with the product charts is tame. The fibre of $A\subseteq\mathbb{R}^m\times X$ over $\lambda$ is $A_\lambda=\{x:(\lambda,x)\in A\}$. A point $z\in A$ is a *stratified regular point* if, for some data of Definition 4.13 at $z$, some $a\in\mathbb{R}^m$, some chart $\varphi$ of $X$ at $z_2$, some presentation $T$ of its partial quadrant and the level-1 point $y$ over the image of $z$ in the product chart $(r\mapsto r-a)\times\varphi$, every $\mu\in\mathbb{R}^m$ is the first component of a tangent vector $v$ of the chart model of $A$ at $y$ with $v_2\in E_{\varphi(z_2)}$ (the projection maps $T_zA\cap E_z$, the tangent space of the stratum through $z$, onto $\mathbb{R}^m$); $\lambda$ is a *stratified regular value* if every $z\in A$ over $\lambda$ is a stratified regular point. *Book:* §4.3, p. 164, and the proof of Theorem 4.23, p. 166 ("regular values of $p_\sigma$"). *Hidden hypotheses:* the chart form (gap G.7). *Deviation (restated):* the chart independence is not formalized (gap G.36).

**V4.54 (Theorem 4.23; `Polyfold.theorem_4_23`).** Let $X$ be tame and $A$ a smooth finite-dimensional submanifold of $\mathbb{R}^m\times X$ with tame induced structure and compact closure, $\varepsilon>0$, and suppose (1) for $|\lambda|\le\varepsilon$ ($|\lambda|^2=\sum\lambda_i^2$) the set $A\cap\{z_1=\lambda\}$ is compact and non-empty, (2) $A$ is in general position at every point. Then there is $\Sigma\subseteq\{|\lambda|<\varepsilon\}$ whose complement in $\{|\lambda|<\varepsilon\}$ is Lebesgue-null, consisting of stratified regular values, such that for $\lambda\in\Sigma$ the fibre $A_\lambda$ is compact, a smooth finite-dimensional submanifold of $X$, in general position at each of its points, and its induced structure has a corner structure and is tame. *Book:* Theorem 4.23, p. 165; proof pp. 165–166. *Hidden hypotheses:* hypothesis (2), misprinted in the text ("there exists a sc-complement $T_zA$ in $T_zZ$"), is Definition 4.21. *Deviation (stronger):* $\Sigma$ consists of stratified regular values, which are regular values; the tameness of $A$, the non-emptiness in (1) and $\varepsilon>0$ are not used, and $A_\lambda$ is moreover tame. With ordinary regular values the conclusion fails at the value: in $\mathbb{R}\times[0,\infty)$, $A=\{(x,x):0\le x<1\}\cup(-1,1)\times\{5\}$ satisfies the hypotheses, $Dp$ is onto everywhere, and $A_0=\{0,5\}$ meets $\partial X$ (Findings, item 102).

**V4.55 (Remark 4.24; `Polyfold.quotDim_le_finrank_of_eq_sup`, `Polyfold.remark_4_24_chart`, `Polyfold.MPolyfold.remark_4_24_codim`, `Polyfold.MPolyfold.remark_4_24`, `Polyfold.MPolyfold.remark_4`).** For tame $X$ and smooth $x$, in a tame chart at $x$ the reduced tangent space has codimension $d_X(x)$ in the tangent space. If $A$ is in general position at $a$, then $d_X(a)\le\dim T_aA$ (in any tame chart at $a$); in particular a submanifold in general position with $T_aA=0$ does not meet $\partial X$ at $a$. *Book:* Remark 4.24, p. 165. *Hidden hypotheses:* $X$ tame and $x$ smooth for the codimension. *Deviation (hypotheses explicit):* stated for any submanifold in general position, not only for the $A_\lambda$ of Theorem 4.23 (to which it applies by V4.54).

**V4.56 (Sard's theorem, vendored; `Polyfold.MilnorSard.Sard.critSet`, `Polyfold.MilnorSard.Sard.sardStat` `Polyfold.MilnorSard.sard`, `Polyfold.MilnorSard.dense_compl_image_critSet`).** Let $E$, $F$ be finite-dimensional real normed spaces (in the same universe), $F$ with its Borel σ-algebra and a Haar measure $\mu$, $U \subseteq E$ open and $f$ of class $C^\infty$ on $U$. Then $\mu(f(\mathrm{Crit})) = 0$, where Crit is the set of points of $U$ at which $Df$ is not onto, and the complement of $f(\mathrm{Crit})$ is dense. *Book:* "Sard's theorem", proof of Theorem 4.23, p. 166, without statement. *Hidden hypotheses:* Mathlib has only the cases $\dim E \le \dim F$; the general case is Milnor's induction, copied from the Milnor project (`math_faithful`) with only the namespace changed (Findings, item 100). *Deviation (not in the book).*

**V4.57 (the strata and Sard's theorem on them; `Polyfold.quadC`, `Polyfold.stratSub`, `Polyfold.zeroSetOf`, `Polyfold.finrank_stratSub`, `Polyfold.stratCritSet`, `Polyfold.measure_image_strat`** Let $V$ be finite-dimensional with linear coordinates $\ell : V \to \mathbb{R}^n$ and $C = \{\ell \ge 0\}$. The stratum subspace of $\sigma$ is $\{\ell_i = 0, i \in \sigma\}$, of dimension $\dim V - \#\sigma$ when $\ell$ is onto; $\sigma(v) = \{i : \ell_i(v) = 0\}$. For $f$ of class $C^\infty$ on $U = O \cap C$ ($O$ open) into a finite-dimensional $F$, the stratified critical points are the $v \in U$ at which $Df(v)$ restricted to the stratum subspace of $\sigma(v)$ is not onto, and their image is null for every Haar measure. *Book:* proof of Theorem 4.23, pp. 165–166 ("$A \cap F^\circ_\sigma$ is a smooth manifold … of dimension $\dim(A) - \#\sigma$"; "Using Sard's theorem we find a subset $\Sigma_\sigma$ … having full measure"). *Hidden hypotheses:* the dimension count, which Theorem 4.22 does not state (the text cites "Theorem 4.23" for it). *Deviation (restated):* in the model space $T_0 A$ of the chart.

**V4.58 (Theorem 3.78 in $C^\infty$ form; `Polyfold.ContractionData.exists_solution_smooth`, `Polyfold.theorem_3_78_smooth`).** Under the hypotheses of Theorem 3.78 (V3.63) with $f$ of class $C^\infty$, one solution $\tau$ is $C^\infty$ on $V'$, with the other conclusions of V3.63 (including the uniqueness clause). *Book:* Theorem 3.78, p. 141, states one $j$; the proof of Theorem 4.23 needs $C^\infty$. *Hidden hypotheses:* none. *Deviation (stronger):* the contraction radii of Lemmas 3.79, 3.80 and the fixed point do not depend on $j$.

**V4.59 (the kernel at a stratified regular point is in good position; `Polyfold.quadAt`, `Polyfold.isPartialQuadrant_quadAt`, `Polyfold.exists_goodComplement_stratSub`).** For $\ell$ onto, $\{\ell_i \ge 0, i \in \sigma\}$ is a partial quadrant. If $K + \{\ell_i = 0, i \in \sigma\} = V$, then $K$ is in good position to $\{\ell_i \ge 0, i \in \sigma\}$ with a closed good complement inside the stratum subspace. *Book:* not in the text here; the argument of Lemma 5.26. *Hidden hypotheses:* none. *Deviation (not in the book).*

**V4.60 (a retraction onto the fibre over a stratified regular point; `Polyfold.exists_fibre_retraction`).** For $\ell$ onto, $f$ of class $C^\infty$ on $O \cap C$ and $v_0 \in O \cap C$ not stratified-critical, there are an open $O' \ni v_0$, $O' \subseteq O$, and a map $\rho$, $C^\infty$ on $O' \cap C$, mapping $O' \cap C$ into itself and into $f^{-1}(f(v_0))$, equal to the identity on $O' \cap C \cap f^{-1}(f(v_0))$, with $D\rho(v_0)$ of range $\ker Df(v_0)$ and equal to the identity there. *Book:* "for $\lambda \in \Sigma$ … $A_\lambda$ is a smooth compact manifold with boundary with corners", p. 166, in the model. *Hidden hypotheses:* none. *Deviation (not in the book):* the implicit function theorem at corner points (V4.58 with V4.59).

**V4.61 (the sc⁺-retraction onto $A_\lambda$; `Polyfold.isScSmooth_slice`, `Polyfold.scDeriv_slice`, `Polyfold.exists_slice_retraction`).** In the model $\mathbb{R}^m \oplus E$ of a product chart, let $t$ be an sc⁺-retraction described near a point as the graph $\Gamma : V \to t(U) \cap W$ with inverse $p$ (Proposition 4.9), and $\rho$ a $C^\infty$ retraction of $O' \cap (\mathbb{R}^m \times C)$ onto the points $v$ with $\Gamma(v)_1 = c$. Then $\tau(e) = (\Gamma(\rho(p(t(c,e)))))_2$ is an sc⁺-retraction of $E$ (partial quadrant $C$) onto $\{e : (c,e) \in t(U) \cap W, p(c,e) \in O'\}$; if $D\rho(v_0) = 1$ on $\{h : (D\Gamma(v_0)h)_1 = 0\}$, the tangent space of its retract at the point over $\Gamma(v_0)_2$ contains $(D\Gamma(v_0)h)_2$ for all such $h$. The slice $e \mapsto (c,e)$ is sc-smooth with derivative $h \mapsto (0,h)$. *Book:* not constructed (Definition 4.13 requires it). *Hidden*

*hypotheses:* none. *Deviation (not in the book).*

**V4.62 (general position of $A_\lambda$; `Polyfold.isGeneralPositionRetractAt_slice`).** In a product chart with $r_X$ tame: if the chart model of $A$ is in general position at $z = (c, y)$, the projection maps the tangent space of the stratum $T_zA \cap E_z$ onto $\mathbb{R}^m$, and $\tau$ is an sc$^+$-retraction onto $Q \subseteq O$ whose tangent space at the smooth $y$ contains the second components of the tangent vectors of $A$ at $z$ with first component $0$, then $Q$ is in general position at $y$. *Book:* proof of Theorem 4.23, p. 166 ("in view of the assumption (2) … $T_{z'}(\{\lambda\} \times A_\lambda)$ has a sc-complement in $T^R_{z'}X$"). *Hidden hypotheses:* stratified regularity at $z$ itself. *Deviation (restated).*

**V4.63 (the local step of Theorem 4.23; `Polyfold.FibreLocal`, `Polyfold.exists_local_family`).** Let $X$ be tame and $A \subseteq \mathbb{R}^m \times X$ a smooth finite-dimensional submanifold in general position at every point. Every $z_0 \in A$ has an open neighbourhood $U$ and a Lebesgue-null CV $\subseteq \mathbb{R}^m$ such that at every $z \in A \cap U$ with $z_1 \notin$ CV: $z$ is a stratified regular point, $A_{z_1}$ is near $z_2$ the image of an sc$^+$-retraction on an open subset of $X$, and $A_{z_1}$ is in general position and in good position at $z_2$. *Book:* proof of Theorem 4.23, pp. 165–166, in one $U(z)$. *Hidden hypotheses:* none. *Deviation (restated):* in the product chart of a tame chart of $X$ centred at $z_{0,2}$, through V4.49, V4.57, V4.60–V4.62.

**V4.64 (the countable cover: Theorem 4.23 for any parameter set; `Polyfold.theorem_4_23_of_subset`).** For $X$ tame, $A \subseteq \mathbb{R}^m \times X$ a smooth finite-dimensional submanifold with compact closure, in general position at every point, and any $S \subseteq \mathbb{R}^m$, there is $\Sigma \subseteq S$ with $S \setminus \Sigma$ Lebesgue-null such that every $\lambda \in \Sigma$ is a stratified regular value and the conclusions of V4.63 hold at every point of $A_\lambda$. *Book:* proof of Theorem 4.23, p. 165 ("We cover the set $\{z \in A : |p(z)| \leq \varepsilon\}$ with the finitely many such neighborhoods"). *Hidden hypotheses:* none. *Deviation (stronger):* any $S$; the text's set need not be compact and need not have a finite subcover (in $\mathbb{R} \times \mathbb{R}$, $A = (-2\varepsilon, 2\varepsilon) \times \{2\} \cup \{(\varepsilon(1-x), x) : 0 < x < 1\}$); $A$ is second countable, its closure being compact in the metrizable $\mathbb{R}^m \times X$, and countably many neighbourhoods cover it (Findings, item 103).

**What is not verified in this chapter**

- p. 151, Lemma 4.10 (the section $u \mapsto (1-p)(u - t(u))$ of $U \lhd Y$ is sc-Fredholm): not formalized, because no longer used — Proposition 4.9 is proved directly (V4.20). It would need $U \lhd Y$ as a strong bundle, $Y$ and $Y \cap F$ as sc-subspaces, and at each smooth point the re-centred strong bundle chart, filler and conjugation (gap G.32).
- pp. 167–168, §4.4 in full: Definitions 4.25, 4.27, Proposition 4.26, the exterior derivative and the relative complex, Theorem 4.28. It needs $\oplus^k TX$ as an M-polyfold and a linear $T_xX$ (gap G.7), the M-polyfolds $X^i$ for $i \geq 2$ (gap G.6), and integration on manifolds with corners of varying dimension, which Mathlib lacks; Proposition 4.26 is quoted from another paper, $d^2 = 0$ is asserted, and Theorem 4.28 has no proof in the text (gaps G.37–G.40).
- p. 159, the open question after Proposition 4.15: a question, not a statement. Its last sentence, for a retract without boundary, is V4.32 with $A = O$; that the "T" is not a topological manifold is not formalized (gap G.34).
- p. 148, Remark 4.3: why an example of "$X^1 \neq X$" needs a retraction whose first derivative loses a level (one whose level maps are $C^1$ gives none, by a Baire category argument): not formalized; the example itself is verified (V4.6; Findings, item 141).
- The chart independence of stratified regularity (V4.53): true by the transport argument of V4.39, since the transition maps of product charts are id $\times\, \tau$ up to translation, but not formalized; Theorem 4.23 states it for some chart (gap G.36).

- p. 157, Remark 4.14, the local form "near points with $d_X(a) = 0$ $A$ is a classical manifold" for the open piece $A \cap \{d_X = 0\}$ of an arbitrary $A$: verified when $d_X = 0$ on all of $A$ (V4.32), not for the piece (gap G.33).
- p. 162, Theorem 4.22 (1) and (4) for the local faces of the neighbourhood $U$ itself: the identification of the faces of the chart model with (germs of) global faces of $X$, a local face theorem for small neighbourhoods; (1) and (4) are verified for the chart (V4.50) (gap G.35; Findings, Chapter 4, item 130).
- Definition 4.1: the classical manifold structure beyond the $C^\infty$ property of the transition maps, and the local constancy of the dimension (V4.1, V4.2).
- Definitions 4.16, 4.21 and Remark 4.14 in $T_aX$: verified in charts, with chart independence (V4.38, V4.39, V4.44, V4.45), since $T_aX$ carries no linear structure here (gap G.7 of Chapter 2).

---

## 6. Chapter 5 (HWZ §§5.1–5.4): verified except Theorems 5.22, 5.27 and the cited Theorems 5.32, 5.40, 5.43

Chapter 5 of Part I (preprint pp. 169–199) is the Fredholm package for M-polyfolds. §5.1 introduces auxiliary norms on the part $Y_{0,1}$ of biregularity $(0,1)$ of a strong bundle (Definition 5.1, Lemma 5.2, Proposition 5.3). §5.2 defines compact solution sets, weakly proper and proper sections (Definition 5.4) and proves that the three agree for sc-Fredholm sections (Theorem 5.5, proof in Appendix 5.4.1), that solution sets are locally compact on every level (Theorem 5.6) and compact in $X_\infty$ (Theorem 5.7), and that the shifted sections are sc-Fredholm (Corollary 5.8). §5.3 builds sc$^+$-sections from sc-smooth bump functions (Definition 5.9, Proposition 5.10, Lemma 5.11), introduces the allowable sections $\Gamma^{+,1}_U(P)$ (Remark 5.12), and proves the perturbation theorems: transversality when $\partial X = \emptyset$ (Theorem 5.13, Definition 5.15), general position (Definition 5.17, Theorem 5.18), good position relative to the boundary (Theorem 5.19), the Morse-type structure of homotopies (Definition 5.20, Theorem 5.22 with Lemmas 5.23, 5.24), perturbation into good position (Theorem 5.25, Lemma 5.26) and cobordisms (Theorem 5.27). Appendix 5.4.2 discusses sc-smooth bump functions and partitions of unity (Definitions 5.28–5.41, Theorems 5.32–5.43, Corollary 5.44).

What is verified. **Theorems 5.5, 5.6, 5.7, 5.13, 5.18, 5.19, 5.25, 5.34, 5.37 hold**, with the readings recorded below, and so do Lemmas 5.2, 5.11, 5.23, 5.24, 5.26, Propositions 5.3, 5.10, 5.30, 5.35, 5.38, 5.42, Remark 5.12, Corollary 5.8 (for each fixed $i$) and Corollary 5.44 (with Theorem 5.43 as an explicit hypothesis). Every definition of the chapter is formalized: Definitions 5.1 (V5.3), 5.4 (V5.7), 5.9 and 5.29 (V5.16), 5.15 (V5.23), 5.17 (V5.31), 5.20 (V5.41), 5.28 (V5.47), 5.31 (V5.50), 5.33 (V5.51), 5.36 (V5.54), 5.39 (V5.59), 5.41 (V5.60). Corollary 5.8 is verified on the shifted strong bundle $P^1 : Y^1 \to X^1$, which the text uses without defining it, and by iteration for every fixed $i$, with the filler condition on level 1 that its "it is a priori clear" hides (V5.13, V5.14). Theorems 5.5–5.7, Corollary 5.8 and Lemma 5.11 need no tameness (V5.10–V5.14, V5.20). Several places of the text needed more than a reading. (a) *Tameness.* The §3.6 implicit function theorems are proved for sections whose germ data lie in a tame strong bundle chart (Definitions 3.50, 3.51; Chapter 3, Findings 190–199); Theorems 5.13, 5.18, 5.19, 5.25 take $f$ with tame germ data, the reading of the tame strong bundle with which §5.3 opens, and nothing else (Findings 74–78). Theorem 5.18 also assumes $X$ tame, which its proof uses through Theorems 4.22 and 4.23 (V5.38). (b) *"Generic" and "dense".* The perturbations are $s_0 + \sum \lambda_i t_i$ for $\lambda$ outside a Lebesgue null set of a small ball (V5.29, V5.38), and density is the bound $\rho(s, s') = \sup_x N(s(x) - s'(x)) \leq \varepsilon$ in the metric space of $\rho$ (V5.21, V5.30, V5.39). (c) *Omitted steps proved.* The spanning sections (V5.24), the compactness

that keeps the zeros of small perturbations near $\{0\} \times S_{f+s_0}$ (V5.27), the parametrized solution set as a smooth submanifold in the sense of Definition 4.13 (V5.32), $T_yS = \ker f'(y)$ and with it the chart independence of Definition 5.17 (V5.33, V5.34), the openness of general position (V5.35), and the properties (i)–(ii) of the proof of Theorem 5.18 (V5.36). (d) *Repairs.* Definition 5.9 is read with $f(x) \neq 0$ (V5.16), which the weak reading gives at the smooth points of an sc-Banach space (V5.16); "$(\Gamma_U^{+,1}(P), \rho)$ is not complete" is false without a hypothesis and is proved in general, by an example (V5.21); the function $\sigma$ of Proposition 5.30 is misprinted (V5.17); the extension by 0 in Theorem 5.34 and both directions of Theorem 5.37 need smaller charts (V5.52, V5.55, V5.56); Proposition 5.38 is not "an immediate consequence of Theorem 5.37" (V5.58). (e) *Theorem 5.19* is proved without applying the implicit function theorem to the parametrized section at the boundary zeros, where it is in general not in good position (V5.40; Finding 88, on paper). (f) *Lemma 5.23* exposes a hidden hypothesis of Theorem 5.22, $dt(x) \neq 0$ (V5.42). Not verified: Theorems 5.22 and 5.27, Remarks 5.14, 5.16, 5.21, the cited Theorems 5.32, 5.40, 5.43, the Sobolev remark and a dense form of Theorem 5.19 (see the list at the end of this section). A fourth addendum (2026-11-02: the shifted strong bundle and Corollary 5.8, the metric $\rho$, weak against pointwise bump functions; Findings, items 140–164) updated V5.13, V5.14, V5.16, V5.21, V5.22 and V5.30 in place.

The setting. A section of $P : Y \to X$ is a map $s : X \to Y$ with $P \circ s = \mathrm{id}$; its support is the closure of $\{s \neq 0\}$. $X$ and $Y$ are paracompact Hausdorff spaces, as for every M-polyfold of the project. $Y_{0,1}$ is the M-polyfold $Y[1]$ of §2.5, an auxiliary norm is a function on it, and "$N(y) \leq r$" for a point $y$ of $Y$ includes $y \in Y_{0,1}$ (the text's convention $N(y) = \infty$ off $Y_{0,1}$). The linearization $f'(x) : T_xX \to Y_x$ is read in a strong bundle chart, as the operator $\mathrm{Lin}(z) : T_zO \to K_z$ of Chapter 3, because $T_xX$ carries no linear structure here (gap G.7 of Chapter 2); transversality is chart independent (Chapter 3), and general position is shown chart independent at the zeros near a zero in good position (V5.34). In the perturbation theorems $|\lambda|$ is the maximum norm on $\mathbb{R}^m$, and "the solution set is a manifold" is stated for a non-empty solution set (the empty set is trivially one). Findings referred to below are those of `Exposition/Findings.md`, "Chapter 5".

**V5.1 (the charts $\Phi[1]$ for every strong bundle chart; `Polyfold.StrongBundle.oneChartOf`, `Polyfold.StrongBundle.isChart_oneChartOf`, `Polyfold.StrongBundle.oneChartOf_compatible`, `Polyfold.StrongBundle.mem_oneSet_iff_of_isChart`).** For every strong bundle chart $\Phi$ of $P$ compatible with the atlas (not only those of the atlas), a point $y$ in its domain lies in $Y_{0,1}$ iff the fibre component of $\Phi(y)$ lies in $F_1$; the map $\Phi[1] : P^{-1}(V) \cap Y_{0,1} \to K[1]$ is a chart of the M-polyfold $Y[1]$, and the charts $\Phi[1]$ of a compatible chart and of a chart of the atlas are compatible. *Book:* §5.1, p. 169, and Lemma 5.2 (ii), which quantifies over strong bundle isomorphisms $\Phi : Y|V \to K$. *Hidden hypotheses:* none. *Deviation:* none; Chapter 2 built $\Phi[1]$ only for the charts of the atlas (Findings, item 2).

**V5.2 (the fibre $(Y_{0,1})_x$ and its chart norm $|\cdot|_1$; `Polyfold.StrongBundle.oneFiber`, `Polyfold.StrongBundle.toOn`
`Polyfold.StrongBundle.zeroOne`, `Polyfold.StrongBundle.liftOneFiber`, `Polyfold.StrongBundle.liftOneF`
`Polyfold.StrongBundle.mem_range_liftOneFiber`, `Polyfold.StrongBundle.isClosed_slice`,
`Polyfold.StrongBundle.complete_liftOneFiber`, `Polyfold.StrongBundle.exists_norm_liftOne_le_mul`).** The points of the fibre $P^{-1}(x)$ that lie in $Y_{0,1}$ form a linear subspace $(Y_{0,1})_x$; its zero is the point $0_x$ of $Y[1]$. In a strong bundle chart around $x$ the fibre component of $\Phi[1]$ is an injective linear map $(Y_{0,1})_x \to F_1$ onto the slice $\{h \in F_1 \mid (\varphi(x), h) \in K[1]\}$, which is closed in $F_1$, so $(Y_{0,1})_x$ is complete for $|h|_1$. The norms $|\cdot|_1$ of two charts around $x$ are comparable, with one constant, on the fibres over a neighbourhood of $x$. *Book:* §5.1, p. 169 ("the fibers $Y_x$ have the structure of Banach spaces"; $|h|_1$ in Lemma 5.2). *Hidden hypotheses:* none. *Deviation (restated):* the text takes the linear and Banach structure of the fibres of $Y_{0,1}$ for granted; it is constructed, and $|h|_1$ depends

on the chart up to equivalence (Findings, item 1).

**V5.3 (Definition 5.1, auxiliary norms; `Polyfold.StrongBundle.IsCompleteFiberNorm`, `Polyfold.StrongBundle.IsAuxNorm`, `Polyfold.StrongBundle.IsAuxNorm.nonneg`, `Polyfold.StrongBundle.Is`** An auxiliary norm is a continuous $N : Y[1] \to \mathbb{R}$ such that (1) on every fibre $(Y_{0,1})_x$, $N$ is a norm ($N(v) = 0 \Rightarrow v = 0$, $N(tv) = |t|N(v)$, the triangle inequality) for which every Cauchy sequence converges, and (2) if $P(w_k) \to x$ in $X$ and $N(w_k) \to 0$, then $w_k \to 0_x$ in $Y[1]$. It is non-negative and homogeneous. *Book:* Definition 5.1, p. 169. *Hidden hypotheses:* none. *Deviation (restated):* "complete norm" is read as a norm on $(Y_{0,1})_x$ with complete normed space; the topology of $Y_{0,1}$ is that of $Y[1]$ (Findings, item 1).

**V5.4 (Lemma 5.2; `Polyfold.lemma_5_2`, `Polyfold.StrongBundle.exists_le_mul_norm_liftOne`, `Polyfold.StrongBundle.IsAuxNorm.exists_norm_liftOne_le`, `Polyfold.StrongBundle.IsAuxNorm.exists_b`, `Polyfold.lemma_5_2_literal`).** Let $N : Y[1] \to \mathbb{R}$ be continuous and a complete norm on every fibre. Then $N$ is an auxiliary norm iff every $x \in X$ lies in the domain of a strong bundle chart $\Phi$ with an open neighbourhood $V \subseteq \operatorname{dom}\varphi$ of $x$ and constants $0 < a < A$ such that $a\,N(w) \le |h|_1 \le A\,N(w)$ for all $w \in Y_{0,1}$ with $P(w) \in V$, $\Phi(w) = (p, h)$. For an auxiliary norm such bounds hold in *every* strong bundle chart around $x$; in the text's literal form (`lemma_5_2_literal`) the bounds hold for all $w \in Y_{0,1}$ over the whole domain of the chart, obtained by restricting the chart to the neighbourhood (V2.57); the upper bound $g(w) \le C|h|_1$ holds for every $g$ continuous at $0_x$ and homogeneous on the fibres near $x$. *Book:* Lemma 5.2, p. 169; proof pp. 169–170. *Hidden hypotheses:* the proof gives the bounds near $x$ only; on all of $P^{-1}(V)$ for the domain $V$ of $\Phi$ they hold after restricting the chart, which the text does silently. *Deviation:* none for `lemma_5_2_literal`; the first form has $V$ a neighbourhood of $x$ on which the bounds hold; the contradiction argument needs no metrizability (Findings, items 2, 3; Chapter 2, items 42, 44). Formerly "restated", before the literal form was proved.

**V5.5 (Proposition 5.3, existence; `Polyfold.proposition_5_3_exists`, `Polyfold.StrongBundle.gluedNorm`, `Polyfold.StrongBundle.continuous_gluedNorm`, `Polyfold.StrongBundle.isCompleteFiberNorm_gluedNorm`, `Polyfold.StrongBundle.gluedNorm_bounds`).** Every strong bundle has an auxiliary norm: for the charts $\Phi_i$ of the atlas and a partition of unity $(\rho_i)$ subordinate to their domains, $N(w) = \sum_i \rho_i(P(w))\,|\Phi_i[1](w)_2|_1$ is continuous on $Y[1]$, a complete norm on every fibre, and comparable near each point with the norm $|\cdot|_1$ of any chart there. *Book:* Proposition 5.3, p. 171. *Hidden hypotheses:* none. *Deviation:* none; the text's proof is one sentence (Findings, item 4).

**V5.6 (Proposition 5.3, comparison; `Polyfold.proposition_5_3_compare`).** For auxiliary norms $N_0$, $N_1$ there is a continuous $f : X \to (0, \infty)$ with $f(x)N_0(w) \le N_1(w) \le f(x)^{-1}N_0(w)$ for every $w \in Y_{0,1}$ over $x$. *Book:* Proposition 5.3, p. 171. *Hidden hypotheses:* none. *Deviation:* none; the statement's $N_0(h)$ is $N_0(w)$ (Findings, item 5).

**V5.7 (Definition 5.4; `Polyfold.StrongBundle.AuxLe`, `Polyfold.StrongBundle.IsWeaklyProper`, `Polyfold.StrongBundle.IsProper`, `Polyfold.StrongBundle.IsProper.isWeaklyProper`, `Polyfold.StrongBundle.IsWeaklyProper.isCompact_solutionSet`).** A section $f$ has a compact solution set if $f^{-1}(0)$ is compact in $X$; it is weakly proper if moreover for every auxiliary norm $N$ there is an open $U \supseteq f^{-1}(0)$ such that $\{f = s\}$ is compact for every sc$^+$-section $s$ with support in $U$ and $N(s(y)) \le 1$ for all $y$; it is proper if moreover for every $N$ there is an open $U \supseteq f^{-1}(0)$ with $\mathrm{cl}_X\{x \in U \mid N(f(x)) \le 1\}$ compact, where $N(f(x)) \le 1$ includes $f(x) \in Y_{0,1}$. A continuous proper section is weakly proper, and a weakly proper section has a compact solution set. *Book:* Definition 5.4, p. 171. *Hidden hypotheses:* "proper implies weakly proper" needs $f$ continuous (so that $\{f = s\}$ is closed). *Deviation:* none (Findings, item 26).

**V5.8 (Theorem 5.5, the local form for basic germs; `Polyfold.IsSc0ContractionGerm.exists_isCompact_bdd`, `Polyfold.IsBasicGerm.exists_isCompact_bdd`).** Let $f$ be a basic germ on $\tilde{C} \oplus W$ with $P \circ f(a,w) = w - B(a,w)$, $\tilde{C}$ closed. There is an open $O \ni 0$, independent of $c$, such that for every $c$ the points $x$ of $U_0 \cap O$ with $f(x) \in \mathbb{R}^N \oplus W_1$ and $|f(x)|_1 \le c$ lie in a compact subset of $E_0$, namely the image of $\{(a,z) \mid a \in \tilde{C}, |a|_0 \le \rho, z \in K_c, |z|_0 \le r/8\}$ under $(a,z) \mapsto (a, \delta(a,z))$, with $K_c$ a compact set containing the level-1 ball of radius $c$ and $\delta$ the parametrized fixed point of $w \mapsto B(a,w) - z$. *Book:* Appendix 5.4.1, pp. 189–192, (82)–(90). *Hidden hypotheses:* $\tilde{C}$ closed, as in Theorem 3.40. *Deviation (restated):* only the level-0 contraction is used, and the neighbourhood does not depend on the bound (Findings, items 23, 25).

**V5.9 (Theorem 5.5, the local step; `Polyfold.IsStrongBundleMap.exists_fiber_bound_one`, `Polyfold.exists_nbhd_isCompact_closure_bdd`).** A strong bundle map $\Psi$ has a uniform $(0,1)$ fibre bound near each point, $|\Psi[1](u,k)_2|_1 \le M|k|_1$ for $|u-u_0|_0 < \delta$. If $(f, x_0)$ is an sc-Fredholm germ, then in the strong bundle chart $\Phi$ of the germ there is an open $V \ni x_0$ such that for every $C$ the closure in $X$ of $\{x \in V \mid \Phi(f(x))_2 \in F_1, |\Phi(f(x))_2|_1 \le C\}$ is compact. *Book:* Appendix 5.4.1, pp. 189–192 ("without loss of generality $f = h + t$", "it suffices to find in local coordinates"). *Hidden hypotheses:* the uniform fibre bound of the conjugating isomorphism. *Deviation (stronger):* $x_0$ need not be a zero; subtracting the sc$^+$-part before the conjugation replaces the text's $t$ and its estimates (83), (85), and the closure is pulled back to $X$ because the retract is the fixed point set of the retraction (Findings, items 23, 24).

**V5.10 (Theorem 5.5; `Polyfold.theorem_5_5`, `Polyfold.theorem_5_5_iff`).** Let $f$ be an sc-Fredholm section of a strong bundle over an M-polyfold $X$. If $f$ has a compact solution set, $f$ is proper; and for $f$ the properties proper, weakly proper and having a compact solution set are equivalent. *Book:* Theorem 5.5, p. 172; proof Appendix 5.4.1, pp. 189–192. *Hidden hypotheses:* none. *Deviation (stronger):* $X$ need not be tame (the appendix states the theorem without tameness); the finite cover of $f^{-1}(0)$ and the closure of a finite union are made explicit (Findings, items 24, 25).

**V5.11 (Theorem 5.6; `Polyfold.theorem_5_6`).** Let $f$ be an sc-Fredholm section and $f(x) = 0$. There are open neighbourhoods $x \in U(i+1) \subseteq U(i)$ of $x$ in $X$ such that for every $i$ the closure in $X$ of $\{y \in U(i) \mid f(y) = 0\}$ is the image of a compact subset of $X_i$ (with its level topology). *Book:* Theorem 5.6, p. 172. *Hidden hypotheses:* none. *Deviation:* none; it is Theorem 3.12 at a zero, which is a smooth point where $(f, x)$ is an sc-Fredholm germ.

**V5.12 (Theorem 5.7; `Polyfold.theorem_5_7`, `Polyfold.MPolyfold.tendsto_smoothLevel_iff`, `Polyfold.MPolyfold.smoothLevel_topology_eq`, `Polyfold.MPolyfold.metrizableSpace_smoothLevel`).** $X_\infty$ carries the initial topology of the inclusions $X_\infty \to X_m$; it is metrizable, and a sequence converges in it iff it converges on every level. If $f$ is an sc-Fredholm section whose solution set $S$ is compact in $X$, then $S \subseteq X_\infty$ and $S$ is a compact subset of $X_\infty$. *Book:* Theorem 5.7, p. 172. *Hidden hypotheses:* the topology of $X_\infty$, which the text does not define. *Deviation (stronger):* $X$ need not be tame. The convergence on every level of a sequence of zeros is Corollary 3.13 at its limit, not Theorem 5.6 as the text says (Findings, item 20).

**V5.13 (Corollary 5.8; `Polyfold.corollary_5_8`, `Polyfold.corollary_5_8_two`, `Polyfold.StrongBundle.solut` `Polyfold.isCompact_solutionSet_level`).** Let $f$ be an sc-Fredholm section of $P : Y \to X$ (Definition 3.14, in the text's germ reading of Definition 3.11) whose solution set is compact. Then $f^1 = f|_{X_1}$ is an sc-Fredholm section of the shifted strong bundle $P^1 : Y^1 \to X^1$ (V5.14) and its solution set is compact in $X^1$; $f^2 = (f^1)^1$ is an sc-Fredholm section of $P^2 = (P^1)^1$ with compact solution set; and for every $i$ the zeros of $f$ in $X_i$ form a compact subset of $X_i$. *Book:* Corollary

5.8, p. 173 (over a tame $X$, for every $i$). *Hidden hypotheses:* the bundle $P^i$, which the text does not define, is the iterated shift $P^i = (P^{i-1})^1$. *Deviation:* none; the first statement holds for every strong bundle, so every fixed $i$ follows by iterating it ($i = 2$ is stated); one statement for a family indexed by $i$ is not formalized, and the base $(X^1)^1$ of $P^2$ has the points of $X_2$, its topology not compared with that of $X_2$ (gap G.6). Tameness is not used; the conclusion is Definition 3.14 for $f^i$, not tame germ data, which would need tame charts of $P^1$ (Findings, items 140–146). Formerly "narrower": only the compactness half was verified (gap G.41, now closed for each fixed $i$).

**V5.14 (Corollary 5.8: the shifted strong bundle, and $f^1$ sc-Fredholm on it; `Polyfold.StrongBundle.shift`, `Polyfold.StrongBundleChart.shift`, `Polyfold.StrongBundleChart.Compatible.shift`, `Polyfold.StrongBundle.mem_shift_doubleLevel_iff`, `Polyfold.StrongBundle.sectionShift`, `Polyfold.StrongBundle.regularizing_sectionShift`, `Polyfold.ScNbhdGerm.shiftParam`, `Polyfold.IsSc0ContractionGerm.shift`, `Polyfold.IsBasicGerm.shift`, `Polyfold.IsConjugateToBasicGerm.`
`Polyfold.IsConjugateToBasicGerm.shift_germ`, `Polyfold.IsFillingGerm.mem_range_toBase_of_fillDefec`
`Polyfold.IsFillingGerm.germ_shift`, `Polyfold.isScFredholmGerm_sectionShift`, `Polyfold.corollary_5_8`
`Polyfold.corollary_5_8_chart`).** *The shifted strong bundle.* $P^1 : Y^1 \to X^1$ is a strong bundle over the M-polyfold $X^1$ (Definition 2.51): its total space is $Y_{1,1}$, the level 1 of the M-polyfold $Y[0]$ with its level topology; $P^1$ is the restriction of $P$; the fibre over $x$ is $P^{-1}(x) \cap Y_{1,1}$, a linear subspace, carrying the topology of $F_1$; and the atlas consists of the index raises $\Phi^1$ of the strong bundle charts $\Phi$ of $P$ (base chart the level chart $\varphi^1$, fibre model $F^1$, retraction $R$ restricted to $U^1 \lhd F^1$), compatible charts having compatible index raises. Its double levels are $(Y^1)_{m,k} = Y_{m+1,k+1}$. *$f^1$ is sc-Fredholm.* For every sc-Fredholm section $f$ (no compactness needed), $f^1 = f|_{X_1}$ is an sc-Fredholm section of $P^1$: it is sc-smooth, regularizing, and at every smooth point of $X^1$ it has the data of Definition 3.11 in the index raise $\Phi^1$ of the chart $\Phi$ of a datum $(g, s)$ of $f$ there. The principal part of $f^1$ is the level-1 map of that of $f$; $g^1$ is a filling germ of it, including condition (3) of Definition 3.4 on level 1, which comes from a regularity statement at the centre of every datum of Definition 3.11: $v \in \ker Dr(0)$ and $Dk(0)v \in F_m$ imply $v \in E_m$, $k$ the fill defect. *In a chart.* The level shift $f^1$ of a basic germ $f$ on $\tilde{C} \oplus W$, $\tilde{C}$ a partial quadrant, is a basic germ on $\tilde{C} \oplus W^1$ with $B^1$ in place of $B$; if a principal part $k$ is conjugated to a basic germ $h$ by $\Psi$, then $k^1$ is conjugated to $h^1$ by the index raise $\Psi^1$. Hence for an sc-Fredholm section with compact solution set: the zeros in $X_1$ form a compact subset of $X_1$, and at every smooth point $x$, in the chart $\Phi$ centred at $x$ of the germ $(f, x)$ with filling germ $g$ and sc$^+$-germ $s$ (the text's germ reading of Definition 3.11), the level-1 germ $(g - s)^1$ is conjugated to a basic germ (by the germ form of the index raise, which uses $g - s$ only near 0) and $s^1$ is an sc$^+$-germ on the shifted germ of $s$. *Book:* Corollary 5.8, p. 173 ("it is a priori clear that $f^i$ is sc-Fredholm", as a section of $P^i : Y^i \to X^i$). *Hidden hypotheses:* the bundle $Y^1 \to X^1$, which the text does not define ($Y[i]$ is defined for $i = 0, 1$ only): $Y^1 = Y_{1,1}$, the convention of $E^1$, forced by "$f^i$ is obtained by raising the index" (an sc-smooth section maps $X_1$ into $Y_{1,1}$, in general not into $Y_{1,2}$); $\tilde{C}$ a partial quadrant. *Deviation:* none; "it is a priori clear" hides condition (3) of the filling on level 1, which does not follow from level 0 alone (an sc-operator can be bijective on level 0 and not onto on level 1); it is proved for the data of Definition 3.11 by reducing to $g(0) = 0$ with Lemma 3.17 (subtracting the section $z \mapsto \Gamma(z, -g(0))$, which leaves the fill defect unchanged) and applying Proposition 3.52 in sc form. No hypothesis beyond the text's is added; tameness is not used, and the tame form (germ data in tame charts of $P^1$) is not stated (Findings, items 22, 140–146). Formerly "narrower": the bundle was not built and the filler on level 1 not verified (gap G.41, now closed for each fixed $i$). In the text's germ reading of Definition 3.11 (phase 2; Findings, Chapter 3, item 333) $s^1$ is an sc$^+$-germ on the shifted germ; in the whole-chart reading it was sc$^+$ on all of $U^1$.

**V5.15 ($\mathbb{R}$ as an M-polyfold; sc-smooth functions; `Polyfold.realPolyfold`, `Polyfold.realChart`, `Polyfold.isChart_realChart`, `Polyfold.MPolyfold.isScSmoothMap_of_local`, `Polyfold.MPolyfold.IsScSmo` `Polyfold.isScSmoothMap_real_of_local`, `Polyfold.isScSmoothMap_real_const`, `Polyfold.isScSmoothMap_r` `Polyfold.MPolyfold.IsScSmoothMap.comp_contDiff`).** $\mathbb{R}$ with the constant sc-structure and the single chart id is an M-polyfold, and an sc-smooth function on $X$ is an sc-smooth map $X \to \mathbb{R}$ (Definition 2.14). Sc-smoothness of a map between M-polyfolds can be checked in one pair of charts around each point (continuity included); an sc-smooth function read in a chart is sc-smooth on the retract; constants are sc-smooth, and $\sigma \circ f$ is sc-smooth for $\sigma : \mathbb{R} \to \mathbb{R}$ of class $C^\infty$. *Book:* Definitions 5.9, 5.29, 5.33 ("sc-smooth map $f : X \to \mathbb{R}$"). *Hidden hypotheses:* none. *Deviation (not in the book):* the M-polyfold $\mathbb{R}$ is presupposed by the text (Findings, item 40).

**V5.16 (Definitions 5.9 and 5.29; `Polyfold.MPolyfold.AdmitsBumpFunctions`, `Polyfold.MPolyfold.AdmitsBump` `Polyfold.MPolyfold.AdmitsBumpFunctions.weak`, `Polyfold.MPolyfold.HasBumpAt`, `Polyfold.MPolyfold.adm` `Polyfold.MPolyfold.AdmitsBumpFunctionsWeak.hasBumpAt_of_mem_smooth`, `Polyfold.MPolyfold.admitsBum`).** $X$ admits sc-smooth bump functions if for every $x$ and every open $U \ni x$ there is an sc-smooth $f : X \to \mathbb{R}$ with $f(x) \neq 0$ and $\operatorname{supp} f \subseteq U$. Definition 5.29 as written ("$f$ not identically zero") is a separate notion, implied by the first. A point $x$ is a *bump point* if every open $U \ni x$ carries an sc-smooth $f$ with $f(x) \neq 0$ and $\operatorname{supp} f \subseteq U$: the first notion says that every point is a bump point, and the second follows if the bump points are dense. On an sc-Banach space $E$ (as an M-polyfold with the identity chart) the second notion makes every smooth point a bump point; on M-polyfolds built on sc-Hilbert spaces the first notion holds. *Book:* Definition 5.9, p. 173; Definition 5.29, p. 192. *Hidden hypotheses:* none. *Deviation (restated):* "$f \neq 0$" is read as $f(x) \neq 0$, the reading Proposition 5.30's proof uses. Whether the weak reading implies it is open only at non-smooth points: at a smooth point of an sc-Banach space a bump that is non-zero near $x$ is translated to $x$ by a smooth vector, and translation by a non-smooth vector does not preserve the levels; that the weak reading is exactly the density of the bump points is argued, one direction formalized (Findings, items 41, 160–163).

**V5.17 (Propositions 5.10 and 5.30; `Polyfold.proposition_5_30`, `Polyfold.proposition_5_10`).** If $X$ admits sc-smooth bump functions, then for every $x$ and open $U \ni x$ there is an sc-smooth $f : X \to [0,1]$ with $f(x) = 1$, $\operatorname{supp} f \subseteq U$, and $f \equiv 1$ on a neighbourhood of $x$. *Book:* Proposition 5.10, p. 173; Proposition 5.30, p. 193. *Hidden hypotheses:* Definition 5.9 in the reading of V5.16. *Deviation:* none; the text's $\sigma$ ("0 for $s \leq 1$ and 1 for $s \geq 1$") is misprinted and $g(x) = 1$ is not given; the proof uses $f = \tau(3g/g(x) - 1)$ with a smooth transition $\tau$ (Findings, item 42).

**V5.18 (the sc-calculus of sections: $\Gamma^+(P)$, supports, $f \cdot s$ and sums; `Polyfold.sectionSupport`, `Polyfold.sectionSmul`, `Polyfold.sectionSum`, `Polyfold.StrongBundle.scPlusSections`, `Polyfold.StrongBundle.IsScPlusSection.mem_oneSet`, `Polyfold.StrongBundle.isScPlusSection_fiberZer` `Polyfold.StrongBundle.IsScPlusSection.smul_left`, `Polyfold.StrongBundle.IsScPlusSection.add`, `Polyfold.StrongBundle.isScPlusSection_sectionSum`).** $\Gamma^+(P)$ is the set of sc$^+$-sections (Definition 2.52); their values lie in $Y_{0,1}$. The zero section is an sc$^+$-section; for an sc-smooth function $f$ and an sc$^+$-section $s$, $f \cdot s$ (fibrewise) is an sc$^+$-section; sums and finite sums of sc$^+$-sections are sc$^+$-sections. *Book:* §5.3, p. 173 ($\Gamma^+(P)$); used in the proof of Lemma 5.11 ("$s(y) = \beta(y)t(y)$ is a sc$^+$-section") and in the finite combinations $\sum \lambda_i s_i$ of Theorems 5.13, 5.18. *Hidden hypotheses:* none. *Deviation (not in the book):* the text uses these rules without stating them; in a chart $(f \cdot s)^\wedge = (f \circ \varphi^{-1}) \cdot \hat{s}$ and the product rule of the sc-calculus applies (Findings, item 45).

**V5.19 (local sc$^+$-sections cut off and extended by $0$; `Polyfold.StrongBundle.cutoffSection`, `Polyfold.StrongBundle.isScPlusSection_cutoffSection`).** Let $\Phi$ be a strong bundle chart of the atlas, $\tau : O \to F_1$ such that $z \mapsto (z, \tau(z))$ is sc-smooth $O \to K[1]$, and $\beta : X \to \mathbb{R}$ sc-smooth

with supp $\beta \subseteq \operatorname{dom}\varphi$. Then $y \mapsto \Phi^{-1}(\varphi(y), \beta(y)\tau(\varphi(y)))$ on $\operatorname{dom}\varphi$, $0_y$ outside, is an sc$^+$-section. *Book:* proof of Lemma 5.11, p. 174 ("if we construct $s$ with support in $Q'$ we can extend it by 0"). *Hidden hypotheses:* none. *Deviation (not in the book):* the extension by 0 is proved (Findings, item 46 (c)).

**V5.20 (Lemma 5.11; `Polyfold.lemma_5_11`).** Let $X$ admit sc-smooth bump functions and $N$ be an auxiliary norm. For a smooth point $x$, a smooth point $e$ of $Y$ with $P(e) = x$, $\varepsilon > 0$ and an open $U \ni x$, there is an sc$^+$-section $s$ with $s(x) = e$, $\operatorname{supp} s \subseteq U$ and $N(s(y)) < N(e) + \varepsilon$ **for all** $y$. *Book:* Lemma 5.11, p. 173; proof p. 174. *Hidden hypotheses:* Definition 5.9 in the reading of V5.16. *Deviation (stronger):* the bound holds at every point (the text's bound at $x$ is trivial; the proof gives the uniform one, which §5.3 needs); tameness of $P$ and $X$, normality of $X$ and a chart centred at $x$ are not needed (Findings, item 46).

**V5.21 ($\Gamma^{+,1}_U(P)$ and the metric $\rho$; `Polyfold.StrongBundle.allowableSections`, `Polyfold.StrongBundle.secti` `Polyfold.StrongBundle.allowableDist`, `Polyfold.StrongBundle.allowableMetric`, `Polyfold.StrongBundle` `Polyfold.TrivialBundle.not_completeSpace_allowableSections`, `Polyfold.StrongBundle.completeSpace_` `Polyfold.TrivialBundle.completeSpace_allowableSections_point`).** For an auxiliary norm $N$ and an open $U$, $\Gamma^{+,1}_U(P)$ is the set of sc$^+$-sections $s$ with $\operatorname{supp} s \subseteq U$ and $N(s(x)) \le 1$ for all $x$ (the allowable sections). For two sc$^+$-sections $N(s(x) - s'(x))$ is the norm of the difference in the fibre $(Y_{0,1})_x$. The uniform distance $\rho(s, s') = \sup_x N(s(x) - s'(x))$ is finite ($\rho \le 2$) and, for an auxiliary norm, a metric on $\Gamma^{+,1}_U(P)$; $\rho(s, s') \le \varepsilon$ iff $N(s(x) - s'(x)) \le \varepsilon$ for all $x$. For the trivial bundle $\mathbb{R} \times \mathbb{R} \to \mathbb{R}$ with $N(x, v) = |v|$ and $U = \mathbb{R}$ the metric space is not complete; for $U = \emptyset$ (any bundle) and over a point with fibre $\mathbb{R}^k$ (any $U$) it is complete. *Book:* §5.3, p. 174 ("a metric space … not complete"). *Hidden hypotheses:* none for the metric. *Deviation (restated):* the text's "not complete" is false without a hypothesis (the cases $U = \emptyset$ and a point base) and is read as "not complete in general", proved by the example: $s_j(x) = (x, \sqrt{x^2 + 1/(j+1)^2}/(2\sqrt{1+x^2}))$ is a $\rho$-Cauchy sequence of allowable sections whose pointwise limit has the fibre part $|x|/(2\sqrt{1+x^2})$, not differentiable at 0, while the fibre part of an sc$^+$-section of the trivial bundle is $C^1$. Non-completeness in the text's setting ($U$ containing a non-isolated point over which $Y_{0,1} \neq 0$) is expected by the same mechanism and not formalized (Findings, items 47, 150–153).

**V5.22 (Remark 5.12; `Polyfold.remark_5_12`, `Polyfold.StrongBundle.IsAuxNorm.auxLe_of_sectionAdd`, `Polyfold.StrongBundle.IsInAllowableCompletion`, `Polyfold.remark_5_12_completion`, `Polyfold.remark_5_12_of_isScFredholmSection`).** Let $f$ have its solution set inside the open $U$ with $\operatorname{cl}\{x \in U \mid N(f(x)) \le 1\}$ compact. If a section $s$ vanishes outside $U$ and has $N(s(x)) \le 1$ everywhere — as the uniform limits of allowable sections do — then the solutions of $f + s = 0$ lie in $\{x \in U \mid N(f(x)) \le 1\}$, and their closure is compact. As stated (`remark_5_12_completion`): with the points of the completion represented by sections $s$ with values in $Y_{0,1}$ that are uniform $N$-limits of allowable sections (`IsInAllowableCompletion`), for $f$ continuous the solution set of $f + s = 0$ is closed, hence compact; for $f$ sc-Fredholm with compact solution set, with the $U$ of Theorem 5.5 (`remark_5_12_of_isScFredholmSection`). *Book:* Remark 5.12, p. 174. *Hidden hypotheses:* none. *Deviation (restated):* the completion of $\Gamma^{+,1}_U$ is represented by sections, uniform $N$-limits of allowable sections; the metric space itself is V5.21, and that every $\rho$-Cauchy sequence has such a limit section is not formalized (Findings, items 131, 153). The closedness of the solution set, which the remark does not discuss, is proved by a chart argument: Lemma 5.2's bound turns $N(s - s') \le \varepsilon$ into a level-0 bound in a chart, and $f + s'$ is continuous for allowable $s'$; "adapted to $N$" is read as the properness of V5.7 (Findings, items 27, 130, 131).

**V5.23 (Definition 5.15; `Polyfold.StrongBundle.IsTransversal`, `Polyfold.StrongBundle.IsLinSurjAt.linAt`**

A section $g$ is transversal to the zero section if at every zero $x$ the linearization is onto, in a strong bundle chart around $x$; it is then onto in every such chart. *Book:* Definition 5.15, p. 175. *Hidden hypotheses:* none. *Deviation (restated):* $g'(x) : T_xX \to Y_x$ is read in charts (gap G.7 of Chapter 2) (Findings, item 70).

**V5.24 (spanning sections at a zero; `Polyfold.exists_fin_mem_of_dense_of_closedComplemented`, `Polyfold.exists_spanning_sections`).** If $R$ is a closed, complemented subspace of finite codimension of a Banach space $K$ and $D$ a dense subspace, finitely many vectors of $D$ span a complement of $R$. Hence, if $X$ admits sc-smooth bump functions, $N$ is an auxiliary norm, $g$ is sc-Fredholm and $g(y) = 0$, then for every open $U \ni y$ there are finitely many allowable sc$^+$-sections $s_1, \dots, s_k$ supported in $U$ and a strong bundle chart centred at $y$ in which $\mathrm{Lin}(0)(T_0O) + \mathrm{span}\{\hat{s}_i(0)\} = K_0$. *Book:* proof of Theorem 5.13, p. 175 ("we find finitely many allowable sc$^+$-sections such that the range of $(f + s_0)'(x)$ together with the sections span $Y_x$"). *Hidden hypotheses:* none. *Deviation (not in the book):* the text does not say how; the density of the smooth vectors in $K_0$ is the point (Findings, item 66).

**V5.25 (the parametrized section and its slices; `Polyfold.paramSlice`, `Polyfold.isZeroAt_paramSection_iff`, `Polyfold.isScFredholmSection_paramSlice`, `Polyfold.isScPlusSection_paramSlice`, `Polyfold.sectionSupport_paramSlice_subset`, `Polyfold.auxLe_paramSlice`).** For a section $g$ and sections $t_1, \dots, t_m$, the slice $g + \sum \lambda_i t_i$ of $F(\lambda, y) = g(y) + \sum \lambda_i t_i(y)$ (Theorem 3.18) is a section of $P$, and $(\lambda, y)$ is a zero of $F$ iff $y$ is a zero of the slice. For $g$ sc-Fredholm and $t_i$ sc$^+$ the slices are sc-Fredholm; slices of sc$^+$-sections are sc$^+$ with support in the union of the supports; and if $N(s_0) \le 1 - \varepsilon$, $N(t_i) \le 1$ and $m|\lambda| \le \varepsilon$, then $N(s_0 + \sum \lambda_i t_i) \le 1$. *Book:* proof of Theorem 5.13, p. 176 ("there is a number $\delta_0 > 0$ such that $|\lambda| < \delta_0$ implies $N(\sum \lambda_j t_j(y)) < \varepsilon/2$"). *Hidden hypotheses:* none. *Deviation (not in the book):* bookkeeping the text leaves implicit.

**V5.26 ("regular value $\Leftrightarrow$ transversal" and transversality of $F$ at $(0, y)$; `Polyfold.surjective_of_forall_exists`, `Polyfold.forall_exists_ker_fst_of_surjective`, `Polyfold.isLinSurjAt_paramSlice_of_ker_fst`, `Polyfold.isLinSurjAt_paramSection_zero_of_spanning`, `Polyfold.isLinSurjAt_paramSection_zero_of_co`).** For $M(a, u) = Lu + \sum a_i e_i$ onto, the projection $\ker M \to \mathbb{R}^m$ is onto iff $L$ is onto. Consequently, if the linearization of $F$ at a zero $(\lambda, y)$ is onto and the kernel of $\mathrm{Lin}_F(0)$ in the product chart $(r \mapsto r - \lambda) \times \Phi$ projects onto $\mathbb{R}^m$, the slice $g + \sum \lambda_i t_i$ is transversal at $y$. Conversely, if the range of $\mathrm{Lin}_g(0)$ and the $\hat{t}_i(0)$ span $K_0$ in a chart centred at the zero $y$ of $g$, $F$ is transversal at $(0, y)$, and adding sections keeps this. *Book:* proof of Theorem 5.13, pp. 175–176 ("the linearization at $(0, x)$ is surjective"; "it is easily verified that $f + s_0 + s(\lambda_0, \cdot)$ is transversal"). *Hidden hypotheses:* none. *Deviation (not in the book):* only "regular value $\Rightarrow$ transversal" is needed and proved for slices; the text's "$\Leftrightarrow$" for $\tilde{S}$ is not formalized (Findings, items 64, 67).

**V5.27 (the openness steps and the omitted compactness step; `Polyfold.isWithoutBoundary_prod_rn`, `Polyfold.isClosed_solutionSet`, `Polyfold.exists_isOpen_isLinSurjAt_zero_near`, `Polyfold.exists_isOpen_goodPosition_of_isLinSurjAt`, `Polyfold.exists_pos_forall_mem_of_isCompact`).** If $\partial X = \emptyset$ then $\partial(\mathbb{R}^m \times X) = \emptyset$; the solution set of an sc-smooth section is closed. For $\partial X = \emptyset$ and $g$ with tame germ data: if $F$ is transversal at $(0, x)$, it is transversal at $(0, y)$ for the zeros $y$ of $g$ near $x$; if $F$ is transversal at every $(0, y)$, $y \in S_g$, there is an open $V \supseteq \{0\} \times S_g$ all of whose zeros of $F$ are in good position in a tame chart. If $Z \subseteq \mathbb{R}^m \times X$ is compact and the open $V$ contains $Z \cap \{\lambda = 0\}$, then $V$ contains $Z \cap \{|\lambda| < \delta\}$ for some $\delta > 0$. *Book:* proof of Theorem 5.13, pp. 175–176 ("in view of the interior case of the implicit function theorem, Theorem 3.19"; "$U \subset \{|\lambda| < \delta_0\} \oplus X$"). *Hidden hypotheses:* tame germ data, the reading of the tame strong bundle of §5.3 (Findings, item 75). *Deviation (not in the book):* the passage to $|\lambda| < \delta_0$ is the omitted step,

proved with Remark 5.12 (Findings, items 63, 68).

**V5.28 (the Sard step; `Polyfold.exists_null_isLinSurjAt_paramSlice_near`).** Let $g$ have tame germ data, $t_i$ be sc$^+$-sections, and $z_0$ a zero of $F(\lambda, y) = g(y) + \sum \lambda_i t_i(y)$ in good position in a tame chart. There are an open $W \ni z_0$ and a Lebesgue null set $CV \subseteq \mathbb{R}^m$ such that for every zero $(\lambda, y) \in W$ of $F$ with $\lambda \notin CV$ the slice $g + \sum \lambda_i t_i$ is transversal at $y$. *Book:* proof of Theorem 5.13, p. 176 ("taking a regular value $\lambda_0$ for the projection $\tilde{S} \to \mathbb{R}^m$"). *Hidden hypotheses:* none. *Deviation (not in the book):* $\tilde{S}$ is not built as a manifold; Sard's theorem is applied in its stratified form on the model of the regular tangent germ chart of Theorem 3.54 (2), which avoids showing that the points over the zeros are interior (Findings, items 64, 65).

**V5.29 (Theorem 5.13, generic form; `Polyfold.theorem_5_13_generic`, `Polyfold.StrongBundle.SolutionSetIs`).** Let $\partial X = \emptyset$, $X$ admit sc-smooth bump functions, $f$ be an sc-Fredholm section with tame germ data and compact solution set, and $N$ an auxiliary norm. There is an open $U \supseteq S_f$ such that (1) $S_{f+s}$ is compact for every $s \in \Gamma^{+,1}_U(P)$, and (2) for every sc$^+$-section $s_0$ with $\operatorname{supp} s_0 \subseteq U$ and $N(s_0(x)) \leq 1 - \varepsilon$ ($\varepsilon > 0$) there are allowable $t_1, \dots, t_m \in \Gamma^{+,1}_U(P)$, $\delta > 0$ and a Lebesgue null set $CV \subseteq \mathbb{R}^m$ such that for every $\lambda$ with $|\lambda| < \delta$, $\lambda \notin CV$: $s_\lambda = s_0 + \sum \lambda_i t_i$ is allowable, $f + s_\lambda$ is transversal, $S_{f+s_\lambda}$ is compact and, if non-empty, a sub-M-polyfold whose induced structure is tame and that of a manifold without boundary of locally constant dimension, with regular tangent germ data at every zero of dimension the Fredholm index of $(f + s_\lambda)'(y)$. *Book:* Theorem 5.13, pp. 174–175; proof pp. 175–176. *Hidden hypotheses:* $f$ with tame germ data (Definition 3.51), the reading of §5.3's tame strong bundle; the statement's "$N(f(x)) \leq 1$" is $N(s(x)) \leq 1$ and "$(f, x)$ in good position" is $(f + s, x)$. *Deviation (restated):* the dense set of the text is described as the parameters off a null set of a ball (Findings, items 60, 62, 74, 75).

**V5.30 (Theorem 5.13, the text's dense form; `Polyfold.theorem_5_13`, `Polyfold.StrongBundle.allowableDist_`).** Under the hypotheses of V5.29 there is an open $U \supseteq S_f$ such that $S_{f+s}$ is compact for every $s \in \Gamma^{+,1}_U(P)$, and for every allowable $s'$ and $\varepsilon > 0$ there is an allowable $s$ with $N(s(x) - s'(x)) \leq \varepsilon$ for all $x$ such that $f + s$ is transversal and $S_{f+s}$ is compact and, if non-empty, a manifold without boundary as in V5.29. *Book:* Theorem 5.13, pp. 174–175. *Hidden hypotheses:* as in V5.29. *Deviation (restated):* density in $(\Gamma^{+,1}_U(P), \rho)$ (V5.21) is the bound $\rho(s, s') \leq \varepsilon$, stated pointwise and equivalent to it; the text's "$N(s_0(y) + s(\lambda_0, y)) \leq 1 - \varepsilon/2$" is $\leq 1$ (Findings, items 60, 62).

**V5.31 (Definition 5.17, in a tame chart; `Polyfold.StrongBundle.IsGeneralPositionGerm`, `Polyfold.StrongBundleChart.linKer`, `Polyfold.StrongBundle.IsGeneralPositionGerm.isLinSurjAt`).** $(f, y)$ with $f(y) = 0$ is in general position if in some tame strong bundle chart around $y$, at the level-1 point $z$ over $\varphi(y)$, $\operatorname{Lin}(z) : T_z O \to K_z$ is onto, and for every presentation $T$ of the partial quadrant the kernel of $\operatorname{Lin}(z)$ has an sc-complement in $T_z O$ contained in $T^R_z O = T_z O \cap E_z$: an sc-subspace $B$ with $(T_z O)_m = (\ker \operatorname{Lin}(z))_m \oplus B_m$ on every level. General position includes transversality at $y$. *Book:* Definition 5.17, p. 177. *Hidden hypotheses:* the chart is tame (the reduced tangent space of Definition 2.38 is that of a tame M-polyfold). *Deviation (restated):* stated in a chart (gap G.7 of Chapter 2); chart independence at the zeros near a zero in good position is V5.34 (Findings, items 71, 76, 82).

**V5.32 (the bridge lemma: solution sets as smooth submanifolds; `Polyfold.isScPlus_of_isScSmooth_const`, `Polyfold.MPolyfold.TangentGermChart.exists_scPlusRetraction`, `Polyfold.MPolyfold.TangentGermChart`, `Polyfold.MPolyfold.isSmoothSubmanifold_of_isRegular`, `Polyfold.StrongBundle.isSmoothSubmanifold_s`).** An sc-smooth map on a finite-dimensional space with the constant structure is sc$^+$. For regular tangent germ data of Definition 3.25 (2), the retraction $\tau = \Gamma \circ \pi \circ r$ of the proof of Theorem

3.27 is an sc⁺-retraction, so a set with regular tangent germ data at every point is a smooth finite-dimensional submanifold (Definition 4.13). In particular, if every zero of a section in the open $W$ is in good position in a tame chart, $S_f \cap W$ is a smooth finite-dimensional submanifold. *Book:* proof of Theorem 5.18, p. 179 ("hence we can apply Theorem 4.23"). *Hidden hypotheses:* none. *Deviation (not in the book):* the text never verifies Definition 4.13 for solution sets (Findings, item 81).

**V5.33 ($T_yS = \ker f'(y)$; `Polyfold.tangentSpace_subset_linKer`, `Polyfold.MPolyfold.TangentGermChart.ex` `Polyfold.HasIndepTangent.transport`, `Polyfold.exists_tangentSpace_eq_linKer`).** In a strong bundle chart the tangent space of a retract on which the principal part vanishes lies in $\ker \mathrm{Lin}(z)$. For $f$ with tame germ data and $(f, x_0)$ in good position in a tame chart there is an open $V \ni x_0$ whose zeros are all in good position such that at every zero $y \in V$, in every strong bundle chart around $y$ and for every open $G \ni y$ in its domain with $\varphi(G \cap S_f \cap V)$ a retract, the tangent space of that retract at the point over $\varphi(y)$ equals $\ker \mathrm{Lin}(z)$. *Book:* used in the proofs of Theorems 5.18 and 5.25 ("$T_zS_\lambda$ has a sc-complement contained in $T^R_xX$"); Lemma 5.24 ("$T_xS = \ker f'(x)$"). *Hidden hypotheses:* none. *Deviation (not in the book):* "$\supseteq$" is a dimension count, $d$ independent tangent vectors of the bridge retraction carried to every chart (Findings, item 82).

**V5.34 (Definition 5.17 is Definition 4.21 for the solution set; `Polyfold.exists_isGeneralPositionGerm_iff`, `Polyfold.isGoodPositionGermTame_of_isGeneralPositionAt`).** For $f$ with tame germ data and $(f, x_0)$ a zero in good position in a tame chart there is an open $V \ni x_0$ such that every zero in $V$ is in good position, $S_f \cap V$ is a smooth finite-dimensional submanifold, and at every zero $y \in V$, $(f, y)$ is in general position (Definition 5.17) iff $S_f \cap V$ is in general position at $y$ (Definition 4.21); both are then independent of the chart. If $f'(y)$ is onto and a smooth submanifold $A \subseteq S_f$ is in general position at $y$, then $(f, y)$ is in good position in a tame chart. *Book:* Definition 5.17, p. 177, and the proofs of Theorems 5.18, 5.25. *Hidden hypotheses:* none. *Deviation (not in the book):* the chart independence of Definition 5.17 holds near zeros in good position (Findings, items 82, 85).

**V5.35 (openness of general position; `Polyfold.exists_isOpen_isGeneralPositionGerm`, `Polyfold.MPolyfold.IsGeneralPositionAt.congr_nhds`).** Definition 4.21 depends only on the germ of the submanifold. For $X$ tame, $f$ with tame germ data and $(f, x_0)$ in general position at the zero $x_0$, there is an open $W \ni x_0$ such that $S_f \cap W$ is a smooth finite-dimensional submanifold and at every zero $y \in W$, $(f, y)$ is in good and in general position and $S_f \cap W$ is in general position at $y$. *Book:* proof of Theorem 5.18, p. 178 ("they will also hold at $(0, y)$ for $y \in S_f$ close to $x$"; "the properties (i)–(ii) hold for all $(\lambda, y) \in \tilde{S}$"). *Hidden hypotheses:* $X$ tame. *Deviation (not in the book):* used without proof in the text; proved through V5.34 and Theorem 4.22 (3) (Findings, item 83).

**V5.36 (properties (i)–(ii) of the proof of Theorem 5.18; `Polyfold.StrongBundleChart.IsParamGP`, `Polyfold.StrongBundleChart.IsParamGP.comp`, `Polyfold.isGeneralPositionGerm_paramSection_of_isPara` `Polyfold.isParamGP_of_isGeneralPositionGerm`, `Polyfold.exists_sections_isParamGP`).** In a chart centred at $y$, (i) is $\mathrm{Lin}_g(0)(T_0O) + \mathrm{span}\{\hat{t}_i(0)\} = K_0$ and (ii) is: every $w \in T_0O$ is $u + e$ with $e \in E_0$ and $\mathrm{Lin}_g(0)u + \sum a_i\hat{t}_i(0) = 0$ for some $a$. Adding sections keeps (i)–(ii). For $g$ with tame germ data and a tame chart centred at the zero $y$, (i)–(ii) hold iff $F$ is in general position at $(0, y)$ (Definition 5.17). If $X$ admits sc-smooth bump functions and $N$ is an auxiliary norm, finitely many allowable sc⁺-sections supported in any given open $U \ni y$ achieve (i)–(ii). *Book:* proof of Theorem 5.18, p. 178. *Hidden hypotheses:* none. *Deviation (restated):* the complement $L$ of $T^R_xX$ is spanned by smooth vectors, so that Lemma 5.11 applies; (i)–(ii) give good position of $F$ first, and then, with $\ker F' = T\tilde{S}$, general position (Findings, item 84).

**V5.37 (Theorem 5.18 from a family; `Polyfold.theorem_5_18_of_family`, `Polyfold.StrongBundle.SolutionSet`).** Let $X$ be tame, $f$ an sc-Fredholm section with tame germ data and compact solution set, $N$ an auxiliary norm. There is an open $U \supseteq S_f$ such that $S_{f+s}$ is compact for every $s \in \Gamma_U^{+,1}(P)$, and for every sc$^+$-section $s_0$ with $\operatorname{supp} s_0 \subseteq U$, $N(s_0) \leq 1-\varepsilon$, and every family of allowable $t_1, \dots, t_m$ for which $F(\lambda, y) = f(y) + s_0(y) + \sum \lambda_i t_i(y)$ is in general position at every $(0, x)$, $x \in S_{f+s_0}$, there are $\delta > 0$ and a null set $CV$ such that for $|\lambda| < \delta$, $\lambda \notin CV$, $s_\lambda = s_0 + \sum \lambda_i t_i$ is allowable and $f + s_\lambda$ is transversal, in general position at every zero, and $S_{f+s_\lambda}$ is compact, a smooth finite-dimensional submanifold in general position at each point, whose induced structure has a corner structure and is tame, with $d_{S_{f+s_\lambda}} = d_X$ on it. *Book:* Theorem 5.18, p. 177; proof pp. 178–179. *Hidden hypotheses:* as in V5.38. *Deviation (not in the book):* the theorem with the family as a hypothesis; Theorem 4.23 is used in its countable-cover form (Findings, item 85).

**V5.38 (Theorem 5.18, generic form; `Polyfold.theorem_5_18`).** Let $X$ be tame and admit sc-smooth bump functions, $f$ an sc-Fredholm section with tame germ data and compact solution set, $N$ an auxiliary norm. There is an open $U \supseteq S_f$ such that (1) $S_{f+s}$ is compact for every $s \in \Gamma_U^{+,1}(P)$, and (2) for every sc$^+$-section $s_0$ with $\operatorname{supp} s_0 \subseteq U$ and $N(s_0(x)) \leq 1-\varepsilon$ there are allowable $t_1, \dots, t_m$, $\delta > 0$ and a Lebesgue null set $CV$ such that for every $\lambda$ with $|\lambda| < \delta$, $\lambda \notin CV$, $s_\lambda = s_0 + \sum \lambda_i t_i$ is allowable and $f + s_\lambda$ has all the conclusions of V5.37. *Book:* Theorem 5.18, p. 177; proof pp. 177–179. *Hidden hypotheses:* $X$ tame (the proof uses Theorems 4.22 and 4.23; the opening of §5.3 places the discussion over a tame strong bundle); $f$ with tame germ data. *Deviation (restated):* the dense set is described as the parameters off a null set, as in V5.29; "replacing $f + s_0$ by $f$" is not needed (Findings, items 80, 85).

**V5.39 (Theorem 5.18, the text's dense form; `Polyfold.theorem_5_18_dense`).** Under the hypotheses of V5.38 there is an open $U \supseteq S_f$ such that $S_{f+s}$ is compact for every $s \in \Gamma_U^{+,1}(P)$, and for every allowable $s'$ and $\varepsilon > 0$ an allowable $s$ with $N(s(x) - s'(x)) \leq \varepsilon$ for all $x$ such that $f + s$ is transversal, $(f + s, x)$ is in general position at every $x \in S_{f+s}$, and $S_{f+s}$ is a compact smooth submanifold in general position with a tame corner structure and $d_{S_{f+s}} = d_X$. *Book:* Theorem 5.18, p. 177. *Hidden hypotheses:* as in V5.38. *Deviation (restated):* density as in V5.30 (Findings, item 80).

**V5.40 (Theorem 5.19, generic form; `Polyfold.theorem_5_19`, `Polyfold.StrongBundle.IsGoodPositionGermTam`).** Good position depends only on the germ of the section. Let $X$ admit sc-smooth bump functions, $f$ be an sc-Fredholm section with tame germ data and compact solution set, $N$ an auxiliary norm, and $(f, x)$ in good position in a tame chart at every zero $x \in \partial X$. There is an open $U \supseteq S_f$ with $S_{f+s}$ compact for every $s \in \Gamma_U^{+,1}(P)$, and there are allowable $t_1, \dots, t_m$ with supports disjoint from $\partial X$, $\delta > 0$ and a null set $CV$ such that for $|\lambda| < \delta$, $\lambda \notin CV$, $s_\lambda = \sum \lambda_i t_i$ is allowable, vanishes on the neighbourhood $X \setminus \operatorname{supp} s_\lambda$ of $\partial X$, $f + s_\lambda$ is transversal and in good position at every zero, and $S_{f+s_\lambda}$ is compact and, if non-empty, a sub-M-polyfold whose induced structure is tame and a manifold with boundary with corners. *Book:* Theorem 5.19, p. 180. *Hidden hypotheses:* none. *Deviation (stronger):* $X$ need not be tame; "arbitrarily small" is $\lambda \to 0$; "$(f, x)$ in good position" is $(f + s, x)$. The text's "as before" would apply the implicit function theorem to $F$ at the boundary zeros, where $F$ is in general not in good position (gap G.46, on paper); the proof uses it only at interior points and keeps $f + s = f$ near the boundary zeros. No dense form (gap G.45) (Findings, items 87, 88).

**V5.41 (Definition 5.20; `Polyfold.restrictSection`, `Polyfold.StrongBundle.IsGenCompactHomotopy`, `Polyfold.StrongBundle.IsGenCompactHomotopy.isCompact_solutionSet_restrict`).** $(f, t)$ is a generalized compact homotopy if $X$ is tame, $f$ is an sc-Fredholm section with tame germ data

and compact solution set, $t : X \to \mathbb{R}$ is sc-smooth with $t(X) = [0,1]$, and for every $\tau \in [0,1]$ the fibre $X_\tau = t^{-1}(\tau)$ is a sub-M-polyfold with tame induced structure on which the restriction $f|X_\tau$ — the section $a \mapsto (a, f(a))$ of the pull-back of $P$ along the inclusion — is an sc-Fredholm section with tame germ data. The restrictions then have compact solution sets. *Book:* Definition 5.20, pp. 180–181. *Hidden hypotheses:* none. *Deviation (restated):* $Y|X_\tau$ is the pull-back bundle (Proposition 2.54); the end problems are the restrictions to $t^{-1}(0)$, $t^{-1}(1)$, and the identification up to isomorphism (Remark 5.21) is not formalized (gap G.47; Findings, item 100).

**V5.42 (Lemma 5.23; `Polyfold.lemma_5_23`, `Polyfold.ne_zero_of_isCompl_ker`, `Polyfold.exists_isCompl_ke`** Let $\ell : T \to \mathbb{R}$ be linear and non-zero and $Z$ a complement of $\ker \ell$. For every linear $\tau : S \to \mathbb{R}$ there is a unique linear $b : S \to Z$ with $\ell \circ b = \tau$, and $\ell \circ (b + a) = \tau$ whenever $\ell \circ b = \tau$ and $a$ takes values in $\ker \ell$. A non-zero complement of $\ker \ell$ exists only if $\ell \neq 0$, and then it can be spanned by a vector of any spanning set. *Book:* Lemma 5.23, p. 182, and the sentence before it ("since $x$ is a smooth point, we find a one-dimensional smooth linear sc-subspace $Z$ with $T_xX = Z \oplus \ker dt(x)$"). *Hidden hypotheses:* $dt(x) \neq 0$, which Theorem 5.22 does not assume. *Deviation (explicit):* linear algebra; the hidden hypothesis is exposed (Findings, item 101).

**V5.43 (the product rule at a level-1 point; `Polyfold.scDeriv_smul_apply`, `Polyfold.scDeriv_clm_sub_apply`, `Polyfold.chartLinCutoff`, `Polyfold.isScSmoothMap_chartLinCutoff`).** For sc-smooth $a : U \to \mathbb{R}$ and $w : U \to F$, $D(a \cdot w)(z)v = Da(z)v \cdot w(z) + a(z) \cdot Dw(z)v$; $D(\ell(\rho(\cdot) - p))(z) = \ell \circ D\rho(z)$; and $g = \beta \cdot \ell(\varphi(\cdot) - \varphi(x))$ on $\operatorname{dom}\varphi$, 0 outside, is sc-smooth when $\beta$ is and $\operatorname{supp}\beta \subseteq \operatorname{dom}\varphi$. *Book:* proof of Lemma 5.24, pp. 182–183. *Hidden hypotheses:* none. *Deviation (not in the book):* tools for V5.44.

**V5.44 (Lemma 5.24; `Polyfold.exists_scPlusSection_linAt`, `Polyfold.lemma_5_24`).** Let $X$ admit sc-smooth bump functions, $\Phi$ a strong bundle chart around the smooth point $x$, $z$ the level-1 point over $\varphi(x)$, and $e$ a smooth point of the fibre over $x$. For every continuous linear $\ell$ on the model space and open $U \ni x$ there is an sc$^+$-section $s$ with $s(x) = 0$, $\operatorname{supp} s \subseteq U$ and $\operatorname{Lin}_s(z)v = \ell(v)\Phi(e)_2$ on $T_zO$. Hence, for a section $f$, a finite-dimensional $K \subseteq T_zO$ and a linear $b : K \to T_zO$ with $\operatorname{Lin}_f(z)(bv) \in \mathbb{R}\,\Phi(e)_2$ for all $v$, there is an sc$^+$-section $s$ with $s(x) = 0$, $\operatorname{supp} s \subseteq U$ and $\operatorname{Lin}_f(z)(bv) + \operatorname{Lin}_s(z)v = 0$ for $v \in K$. *Book:* Lemma 5.24, pp. 182–183. *Hidden hypotheses:* that $f'(x) \circ b_\tau$ has its image in the line of a smooth point $e$ is a hypothesis (the text's justification, "the image of any vector in $T_xS$ belongs to $Y_\infty$", is not formalized). *Deviation (restated):* $s'(x)$ is read in a chart, $T_xS$ as a finite-dimensional $K$; the graph of $S$ over $T_xS$ and the case distinction are not needed (Findings, item 102).

**V5.45 (Theorem 5.25; `Polyfold.theorem_5_25`).** Let $X$ be tame and admit sc-smooth bump functions, $f$ a proper sc-Fredholm section with tame germ data, $N$ an auxiliary norm. There is an open $U \supseteq S_f$ with $S_{f+s}$ compact for every $s \in \Gamma^{+,1}_U(P)$, and an $s \in \Gamma^{+,1}_U(P)$ such that $(f + s, x)$ is in good position in a tame chart at every $x \in S_{f+s}$, and $f + s$ has the conclusions of V5.37; in particular $S_{f+s}$ has a corner structure. *Book:* Theorem 5.25, pp. 186–187; proof pp. 187–188. *Hidden hypotheses:* $f$ with tame germ data. *Deviation (restated):* $U$ is the neighbourhood the proof produces (that of Theorem 5.18), not an arbitrary one with the compactness property; the text's "in view of Theorem 3.54 … in general position" is V5.35, and "$x \in S_f$" is $x \in S_{f+s_0}$ (Findings, item 86).

**V5.46 (Lemma 5.26; `Polyfold.isInGoodPositionIn_of_scComplement_le_reduced`, `Polyfold.StrongBundle.I` `Polyfold.StrongBundle.IsGeneralPositionGerm.isGoodPositionGermTame`).** In a tame retract, a subspace $N$ of $T_zO$ with an algebraic complement $B \subseteq T^R_zO$ is in good position to the partial cone, with good complement $B$ and $N \cap C_z$ of non-empty interior in $N$. Hence for an sc-Fredholm section $f$, a zero $y$ with $(f, y)$ in general position (Definition 5.17) is in good position (Definition

3.53), and in the tame chart of the germ data when $f$ has tame germ data. *Book:* Lemma 5.26, p. 187; "clearly general position implies good position", p. 179. *Hidden hypotheses:* none beyond the tame chart of Definition 5.17. *Deviation (stronger):* $X$ need not be tame; the interior condition of Definition 3.21, which the text omits, is proved (Findings, items 72, 76).

**V5.47 (Definition 5.28; `Polyfold.MPolyfold.IsScSmoothPartitionOfUnity`, `Polyfold.MPolyfold.AdmitsScPa`.** $X$ admits sc-smooth partitions of unity if for every open cover $(U_i)_{i\in\iota}$ (with $\iota$ in the universe of $X$) there is a partition of unity $(\rho_i)$ — continuous, $\rho_i \geq 0$, locally finite supports, $\sum \rho_i = 1$ — with $\operatorname{supp}\rho_i \subseteq U_i$ and every $\rho_i$ sc-smooth. *Book:* Definition 5.28, p. 192. *Hidden hypotheses:* none. *Deviation (restated):* the index type is restricted to the universe of $X$, which loses nothing (every cover can be indexed by subsets of $X$) (Findings, item 110).

**V5.48 (sc-smooth functions: algebra, local finiteness, extension by $0$; `Polyfold.MPolyfold.IsScSmoothMap.rea`, `Polyfold.MPolyfold.IsScSmoothMap.real_sub`, `Polyfold.MPolyfold.IsScSmoothMap.real_mul`, `Polyfold.isScSmoothMap_real_finsetSum`, `Polyfold.isScSmoothMap_real_finsetProd`, `Polyfold.isScSmoothMap_real_finsum`, `Polyfold.isScSmoothMap_real_of_eventuallyEq`, `Polyfold.isScSmoothMap_real_of_chart`).** Sums, differences, products, finite sums and finite products of sc-smooth functions are sc-smooth; so is a locally finite sum; a function that near every point agrees with an sc-smooth function is sc-smooth; a function supported in the domain of a chart whose representative is sc-smooth on the retract is sc-smooth on $X$. *Book:* proof of Theorem 5.37, pp. 195–197. *Hidden hypotheses:* none. *Deviation (not in the book):* used and not stated (Findings, item 112).

**V5.49 (partitions of unity give bump functions; `Polyfold.MPolyfold.AdmitsScPartitionsOfUnity.admitsBum`.** If $X$ admits sc-smooth partitions of unity, it admits sc-smooth bump functions (Definition 5.9 in the reading of V5.16). *Book:* §5.4.2, p. 192 (the open problem whether the converse holds). *Hidden hypotheses:* none. *Deviation (not in the book):* the easy direction, which the text implies; the converse is not addressed (Findings, item 119).

**V5.50 (Definition 5.31; `Polyfold.HasContDiffBump`, `Polyfold.AdmitsContDiffPartitionsOfUnity`).** A Banach space admits a $C^k$-bump function ($k \in \mathbb{N}\cup\{\infty\}$) if some $C^k$ function $V \to \mathbb{R}$, not identically zero, has bounded support; it admits $C^k$ partitions of unity if every open cover has a subordinate partition of unity of $C^k$ functions. *Book:* Definition 5.31, p. 193; partitions of unity of Banach spaces, pp. 192, 197–198. *Hidden hypotheses:* none. *Deviation:* none (Findings, item 114).

**V5.51 (Definition 5.33; `Polyfold.LocalModel.HasBumpProperty`, `Polyfold.closure_inter_ball_subset`).** A local model $(O, C, E)$ has the sc-smooth bump function property if for every $x \in O$ and open $W \ni x$ of $E_0$ with $\mathrm{cl}_E(O\cap W) \subseteq O$ there is an sc-smooth $f : O \to \mathbb{R}$ with $f(x) \neq 0$ and support (in $O$) inside $W$. A retract is closed near each of its points: $O \cap \mathrm{cl}\, B(z, r/2) \subseteq O$ when $B(z, r)$ lies in the open set of the domain of a retraction onto $O$. *Book:* Definition 5.33, p. 194. *Hidden hypotheses:* none. *Deviation (restated):* "$f \neq 0$" is read as $f(x) \neq 0$, as in V5.16 (Findings, items 41, 43).

**V5.52 (Theorem 5.34; `Polyfold.theorem_5_34`).** $X$ admits sc-smooth bump functions iff some family of charts of $X$ covering $X$ has local models with the sc-smooth bump function property. *Book:* Theorem 5.34, p. 194 ("clearly"). *Hidden hypotheses:* none. *Deviation (explicit):* the extension by 0 of a local bump can fail to be continuous (a chart of the unit ball onto the whole space); the proof takes the local bump in a small ball whose preimage has closure inside the chart domain; "atlas" is read as a covering family of charts (Findings, item 43).

**V5.53 (Proposition 5.35 and M-polyfolds on sc-Hilbert spaces; `Polyfold.LocalModel.hasBumpProperty_of_qua`.**

`Polyfold.proposition_5_35, Polyfold.MPolyfold.admitsBumpFunctions_of_hilbert`**).** If $\kappa\|v\|^2 \le b(v,v)$ for a continuous bilinear $b$ on $E_0$ and some $\kappa > 0$ — in particular if $E_0$ is a Hilbert space, $\|v\|^2 = b(v,v)$ — the local model $(O, C, E)$ has the sc-smooth bump function property. An M-polyfold covered by charts on sc-Hilbert spaces admits sc-smooth bump functions. *Book:* Proposition 5.35, p. 194; §5.3, p. 173 ("for example, the case if $X$ is built on sc-Hilbert spaces"). *Hidden hypotheses:* none. *Deviation (restated):* "Hilbert space" is a norm given by a continuous bilinear form, not a second normed structure; translation by a non-smooth point is legitimate because the bump is $C^\infty$ on $E_0$ (Findings, item 44).

**V5.54 (Definition 5.36;** `Polyfold.LocalModel.HasApproxProperty`**).** A local model $(O, C, E)$ has the sc-smooth approximation property if for every open $W$ of $E_0$ with $\mathrm{cl}(O \cap W) \subseteq O$, every continuous $f : O \to [0,1]$ with support (in $O$) inside $W$ and every $\varepsilon > 0$ there is an sc-smooth $g : O \to [0,1]$ with support inside $W$ and $|f - g| \le \varepsilon$ on $O$. *Book:* Definition 5.36, p. 195. *Hidden hypotheses:* none. *Deviation (restated):* $V = O \cap W$ and the closure is taken in $E_0$ ($C$ is closed) (Findings, item 110).

**V5.55 (Theorem 5.37 "(i) ⇒ (ii)";** `Polyfold.hasApproxProperty_restrict, Polyfold.MChart.isClosed_inter_`**).** If $\mathrm{cl}_X(s)$ lies in the domain of a chart $\varphi$, then $\varphi^{-1}(A)$ is closed in $X$ for closed $A \subseteq \varphi(s)$. If $X$ admits sc-smooth partitions of unity, every chart restricted to an open $s$ with $\mathrm{cl}_X(s) \subseteq \mathrm{dom}\,\varphi$ has a local model with the sc-smooth approximation property. *Book:* proof of Theorem 5.37, pp. 195–196. *Hidden hypotheses:* none. *Deviation (narrower):* the text claims the property for the charts of any atlas; its step "$\mathrm{cl}_X(\varphi^{-1}(V)) \subseteq U$" fails (the chart of the unit ball onto the whole space), so the restricted charts are used; whether every chart has the property is not settled (Findings, item 111).

**V5.56 (Theorem 5.37 "(ii) ⇒ (i)";** `Polyfold.MChart.exists_small_nhds, Polyfold.LocalModel.HasApproxProp`
`Polyfold.exists_isScSmoothMap_urysohn, Polyfold.admitsScPartitionsOfUnity_of_approx`**).** Every point of a chart domain has arbitrarily small open neighbourhoods $N$ with $\mathrm{cl}_X(N) \subseteq \mathrm{dom}\,\varphi$ and $\mathrm{cl}(\varphi(N)) \subseteq O$; the approximation property transports to continuous functions supported in such $N$. If a family of charts covering $X$ has local models with the approximation property, then disjoint closed sets of $X$ are separated by sc-smooth Urysohn functions $X \to [0,1]$, and $X$ admits sc-smooth partitions of unity. *Book:* proof of Theorem 5.37, pp. 196–197. *Hidden hypotheses:* none. *Deviation (restated):* the refinement "with the same index set" does not exist in general and the closure condition is not arranged in the text; the proof goes through Urysohn functions and Mathlib's bump coverings; metrizability is not used (Findings, item 112).

**V5.57 (Theorem 5.37;** `Polyfold.theorem_5_37`**).** $X$ admits sc-smooth partitions of unity subordinate to any open cover iff some family of charts of $X$ covering $X$ has local models with the sc-smooth approximation property. *Book:* Theorem 5.37, p. 195. *Hidden hypotheses:* none. *Deviation (restated):* "atlas" is a covering family of charts; for "(i) ⇒ (ii)" it is the family of V5.55 (Findings, items 111, 112).

**V5.58 (Proposition 5.38;** `Polyfold.proposition_5_38`**).** If $E_0$ admits $C^\infty$ partitions of unity, every local model $(O, C, E)$ has the sc-smooth approximation property. *Book:* Proposition 5.38, p. 197. *Hidden hypotheses:* "the sc-Banach space admits smooth partitions of unity" is read for $E_0$. *Deviation (explicit):* not "an immediate consequence of Theorem 5.37" (which yields some atlas, and would need partitions of unity on an open subset of $C$); the argument of "(i) ⇒ (ii)" is repeated in $E_0$ (Findings, item 113).

**V5.59 (Definition 5.39 and the statement of Theorem 5.40;** `Polyfold.cZero, Polyfold.mem_cZero_iff,`
`Polyfold.norm_cZero, Polyfold.IsCoordinatewiseContDiffEmbedding, Polyfold.ToruńczykStatement`**).**

$c_0(\Gamma)$ is the closed subspace of $\ell^\infty(\Gamma)$ of the functions tending to 0 along the cofinite filter, equivalently those with $\{\gamma \mid |f(\gamma)| > \varepsilon\}$ finite for every $\varepsilon > 0$, with the supremum norm; a homeomorphic embedding into $c_0(\Gamma)$ is coordinate-wise $C^k$ if every coordinate is $C^k$. Theorem 5.40 is recorded as a statement, not proved and not used. *Book:* Definition 5.39, Theorem 5.40, p. 198. *Hidden hypotheses:* none. *Deviation (restated):* the text's "max" norm is the supremum (attained for $f \neq 0$) (gap G.52; Findings, items 114, 117).

**V5.60 (Definition 5.41; `Polyfold.IsWCG`, `Polyfold.IsReflexive`).** A Banach space is weakly compactly generated if some set compact in the weak topology has dense linear span; it is reflexive if the canonical map into the bidual is onto. *Book:* Definition 5.41, p. 198 (reflexivity is used in Proposition 5.42, not defined). *Hidden hypotheses:* none. *Deviation:* none (Findings, item 114).

**V5.61 (Proposition 5.42; `Polyfold.proposition_5_42_reflexive`, `Polyfold.proposition_5_42_separable`, `Polyfold.span_closedBall_eq_top`).** (i) A reflexive Banach space is WCG; (ii) a separable Banach space is WCG. *Book:* Proposition 5.42, p. 198. *Hidden hypotheses:* none. *Deviation:* none; Kakutani's theorem is Mathlib's Banach–Alaoglu argument, and in (ii) a dense sequence of the space, suitably scaled, replaces the text's $\{0\} \cup \{x_n/n\}$ (Findings, item 115).

**V5.62 (the statement of Theorem 5.43, and Hilbert spaces; `Polyfold.GodefroyTroyanskiWhitfieldZizlerState` `Polyfold.hasContDiffBump_of_hilbert`, `Polyfold.isReflexive_of_hilbert`).** Theorem 5.43 is recorded as a statement: a WCG space with a $C^k$-bump function admits $C^k$ partitions of unity. A Banach space whose norm comes from a continuous bilinear form has a $C^\infty$-bump function and, if complete, is reflexive. *Book:* Theorem 5.43, p. 198; proof of Corollary 5.44, p. 199 ("a Hilbert space is reflexive"). *Hidden hypotheses:* none. *Deviation:* Theorem 5.43 is not proved (gap G.53) (Findings, items 116, 117).

**V5.63 (Corollary 5.44; `Polyfold.corollary_5_44`, `Polyfold.MPolyfold.admitsScPartitionsOfUnity_of_hilb`** If $E_0$ is a Hilbert space ($\|v\|^2 = b(v,v)$) and Theorem 5.43 holds for $E_0$ and $k = \infty$, the local model $(O, C, E)$ has the sc-smooth approximation property. An M-polyfold covered by charts on such spaces admits sc-smooth partitions of unity. *Book:* Corollary 5.44, p. 199. *Hidden hypotheses:* Theorem 5.43 for $E_0$ (cited, not proved). *Deviation (narrower):* Theorem 5.43 is an explicit hypothesis; the proof's "in view of Theorem 5.37" is Proposition 5.38, and the bump Theorem 5.43 needs is the classical $C^\infty$ one (Findings, item 116).

**What is not verified in this chapter**

- p. 173, Corollary 5.8 as one statement for all $i$: it is verified for each fixed $i$ by iterating the shift ($i = 1, 2$ stated; V5.13, V5.14); a family $P^i$ indexed by $i$, the identification of the iterated base $(X^1)^1$ with $X_2$ as a topological space (the sets agree; gap G.6), and the tame form of the conclusion (the tameness of the index raises of tame charts) are not formalized (Findings, items 145, 146).
- p. 174, the non-completeness of $(\Gamma^{+,1}_U(P), \rho)$ in the text's setting: it is proved for the trivial bundle over $\mathbb{R}$, and it fails for $U = \emptyset$ and over a point (V5.21); a general sufficient condition is not formalized, and neither is the identification of the completion of Remark 5.12 with the uniform $N$-limits of allowable sections (V5.22) (Findings, items 152, 153).
- p. 173 and p. 192, Definitions 5.9 and 5.29: whether the weak reading gives a bump function non-zero at a non-smooth point (V5.16; Findings, item 163).
- p. 175, Remark 5.14, and p. 176, Remark 5.16: comments on [21] and a sketch of the homotopy argument "once we have finished our general discussion of the boundary case"; not formalized (gaps G.43, G.44).

- p. 180, a dense form of Theorem 5.19: the generic form gives perturbations $\sum \lambda_i t_i$ with $\lambda$ arbitrarily small (V5.40), not density around a given allowable $s'$ (gap G.45).
- p. 180, Finding 88: at a zero $(0, x)$, $x \in \partial X$, where all $t_i$ vanish, the parametrized section is in good position only if $\ker f'(x) + E_x \supseteq T_x X$, a general position condition that good position does not imply; checked on paper only. The proof of Theorem 5.19 avoids the point (V5.40) (gap G.46).
- p. 181, Remark 5.21 (end problems up to isomorphism): not formalized; Definition 5.20 takes the restrictions as the end problems (V5.41) (gap G.47).
- p. 181, Theorem 5.22 (Morse-type structure): needs finite-dimensional Morse theory on manifolds with boundary, the cotangent bundle of the solution set, a parametric Sard theorem and a trivialization $\Phi_\lambda : S \to S_\lambda$ of the family, and the hidden hypothesis $dt(x) \neq 0$ of V5.42; Lemmas 5.23 and 5.24 are verified (V5.42–V5.44) (gap G.48).
- p. 188, Theorem 5.27 (cobordism between good position perturbations): its proof is left to the reader (gap G.49).
- p. 193, Theorem 5.32 (Bonic–Frampton, cited from [9]): not stated (gap G.50).
- pp. 193–194, the $L^p$ and Sobolev remark: the norms of $W^{k,p}(\Omega)$, the bump function of $L^p$, $W^{1,4}(\Omega)$ against $W^{1,3/2}(\Omega)$ and the question it raises; it needs Sobolev spaces and their sc-structures (gap G.1) and Theorem 5.32 (gap G.51).
- p. 198, Theorems 5.40 (Toruńczyk) and 5.43 (Godefroy–Troyanski–Whitfield–Zizler), cited from [9], [12]: recorded as statements (V5.59, V5.62), not proved; Theorem 5.43 is an explicit hypothesis of Corollary 5.44 (V5.63) (gaps G.52, G.53).
- Definitions 5.15 and 5.17 in $T_x X$: verified in charts (V5.23, V5.31), with chart independence for Definition 5.17 at the zeros near a zero in good position (V5.34), since $T_x X$ carries no linear structure here (gap G.7 of Chapter 2).

---

## 7. Sign-off

The statements below are the ones the reader is asked to confirm say what HWZ mean. Each is a summary of the V-item it names; the standing hypotheses of that item are not repeated.

1. (V1.1) An sc-Banach space is a sequence of Banach spaces $E_0 \supset E_1 \supset \cdots$ with compact inclusions and $E_\infty$ dense in every $E_m$, taken as an indexed family with injective compact inclusions; finite-dimensional spaces carry the constant structure, sums have levels $E_m \oplus F_m$, and $E^k$ has levels $E_{k+m}$.
2. (V1.2) An sc-operator is a linear map continuous $E_m \to F_m$ on every level, given by its level maps; an sc-isomorphism is one that is a continuous linear equivalence on every level.
3. (V1.3) An sc$^+$-operator is an sc-operator $E \to F^1$; it is compact $E_m \to F_m$ on every level.
4. (V1.4) An sc-subspace is a closed subspace whose smooth points are dense in each $F \cap E_m$; an sc-splitting $E = F \oplus G$ consists of two sc-subspaces with $F_m$ and $G_m$ complementary on every level.
5. (V1.5) A finite-dimensional subspace is an sc-subspace iff it lies in $E_\infty$, and then has an sc-complement.
6. (V1.6) For a closed subspace $A$ of an sc-Banach space $E$, the quotients $E_m/A_m$ form an sc-Banach space $E/A$ with the quotient map an sc-operator whose level kernels are the $A_m$.
7. (V1.7) An sc-Fredholm operator is an sc-operator with sc-splittings $E = \ker T \oplus X$, $F = C \oplus \operatorname{range} T$, $\ker T$ and $C$ finite-dimensional, and $T$ bijective $X_m \to \operatorname{range}(T)_m$ on every level; its index $\dim \ker T - \dim C$ is $\dim \ker T_0 - \dim \operatorname{coker} T_0$, and each $T_m$ is Fredholm.

8. (V1.8) An sc-Fredholm operator is regularizing: $Te \in F_m$ implies $e \in E_m$; conversely, an sc-operator that is Fredholm on level 0 and regularizing is sc-Fredholm.
9. (V1.9) The sum of an sc-Fredholm operator and an sc$^+$-operator is sc-Fredholm.
10. (V1.10) A partial quadrant is a set $C = T^{-1}([0,\infty)^n \oplus W)$ for an sc-isomorphism $T : E \to \mathbb{R}^n \oplus W$; a tuple $(U, C, E)$ has $U = O \cap C$ with $O$ open.
11. (V1.11) A map is sc$^0$ on $U$ if it maps $U_m$ into $F_m$ continuously for every $m$.
12. (V1.12) The tangent $TE$ is $E^1 \oplus E$ and $TU = U^1 \oplus E$, with levels $U_{m+1} \times E_m$.
13. (V1.13) A map is sc$^1$ on $U$ if it is sc$^0$, differentiable from $U_1$ into $F_0$ at every point of $U_1$ with a derivative $Df(x)$ bounded on $E_0$, and $(x,h) \mapsto (f(x), Df(x)h)$ is sc$^0$ on $TU$; $Df(x)$ is unique on relatively open subsets of partial quadrants; $x \mapsto Df(x)$ need not be norm continuous (an example on a weighted $c_0$ scale).
14. (V1.14) $f$ is sc^{k+1} if it is sc$^1$ with sc^k tangent map, and sc-smooth if sc^k for all $k$.
15. (V1.15) An sc$^0$ map on an sc-domain is sc$^1$ iff on every level $f : U_{m+1} \to F_m$ is differentiable with a derivative extending to a bounded $Df(x) : E_m \to F_m$ that depends jointly continuously on $(x,h) \in U_{m+1} \times E_m$; then $f : U_{m+1} \to F_m$ is $C^1$, and $Df(x)$ is an sc-operator at smooth points.
16. (V1.16) If $f$ is sc^k on $U$ then $f : U^1 \to F^1$ is sc^k.
17. (V1.17) If $f$ is sc^k then $f : U_{m+k} \to F_m$ is $C^k$, and $f : U_{m+l} \to F_m$ is $C^l$ for $l \le k$.
18. (V1.18) If $f : U_{m+l} \to F_m$ is $C^{l+1}$ for all $m \ge 0$ and $0 \le l \le k$, then $f$ is sc^{k+1}, as in the text; if this holds for every $l$, $f$ is sc-smooth.
19. (V1.19) A map into $\mathbb{R}^N$ that is $C^{l+1}$ on $U_l$ for $0 \le l \le k$ is sc^{k+1}, as in the text; one that is $C^{l+1}$ on every $U_l$, or $C^\infty$ on $U$, is sc-smooth.
20. (V1.20) The composition of sc$^1$ maps between relatively open subsets of partial quadrants is sc$^1$ with $T(g \circ f) = Tg \circ Tf$; sc^k and sc-smooth maps compose likewise.
21. (V1.21) The degeneracy index $d_C(x)$ is the number of vanishing half-line coordinates of $T(x)$.
22. (V1.22) $d_C$ does not depend on the presentation $T$, and neither does the number $n$ of half-line factors.
23. (V1.23) An sc$^1$ map with sc$^1$ inverse between relatively open subsets of partial quadrants preserves the degeneracy index at every point.
24. (V1.24) For a compact operator $c$ on a Banach space, $1 + c$ is Fredholm (Riesz–Schauder); a continuous linear map with finite-dimensional kernel and closed range of finite codimension is Fredholm.
25. (V1.25) A compact perturbation of a Fredholm operator between Banach spaces is Fredholm.
26. (V1.26) If $(x,y) \mapsto A(x)y$ is $C^k$ on $s \times Y$ and $\iota : Y' \to Y$ is compact, then $x \mapsto A(x) \circ \iota$ is $C^k$ into $\mathcal{L}(Y', Z)$; for an sc$^1$ map the level operators $Df(x) : E_m \to F_m$ exist for $x \in U_{m+1}$ and are the derivatives of $f : U_{m+1} \to F_m$.
27. (V1.27) An sc-domain has level sets that are open-in-convex with unique differentiability; relatively open subsets of partial quadrants are sc-domains, and $T$ and the level shift preserve the property.
28. (V1.28) The index $\dim\ker - \dim\operatorname{coker}$ is additive under composition, and the Fredholm index between Banach spaces is invariant under small and under compact perturbations.
29. (V1.29) Every level map $T_m$ of an sc-Fredholm operator is Fredholm of index $\operatorname{ind}(T) = \dim K - \dim C$.
30. (V1.30) For $T$ sc-Fredholm and $S$ sc$^+$, $\operatorname{ind}(T+S) = \operatorname{ind}(T)$, on every level.
31. (V1.31) An sc-structure on a finite-dimensional space is sc-isomorphic, through its inclusions, to the constant one; an onto inclusion $E_{m+1} \to E_m$ forces $E_m$ finite-dimensional.
32. (V1.32) $E = A \oplus B$ is an sc-splitting iff some idempotent sc-operator has range $A$ and kernel

$B$.

33. (V1.33) For an sc-subspace $A$, a class $x + A$ meeting every $E_j$ contains a smooth point: the smooth points of $E/A$ are the classes of smooth points.
34. (V1.34) $(x, h) \mapsto A(x)h$ is jointly continuous iff $x \mapsto A(x)$ is continuous for uniform convergence on compact sets; so $Df$ is compact-open continuous on $U_1$, and not in general norm continuous (V1.13).
35. (V1.35) Between finite-dimensional spaces, on relatively open subsets of quadrants, sc$^1$ is $C^1$ within $U$ (one-sided derivatives at the boundary) and sc-smooth is $C^\infty$.
36. (V1.36) $C_1$ is a partial quadrant in $E^1$ and $TC$ one in $TE$; $U_1$ and $TU$ are relatively open in them.
37. (V1.37) If $f : U_{m+l} \to F_m$ is $C^{l+1}$ for all $m$ and $l \leq k$, the derivatives $d^i f(x)$ on $E_{m+l}$ into $F_m$ ($i \leq l + 1$, $i \leq k + 1$) are restrictions of one base-level map, continuous in $x$, and differentiable in $x$ with derivative $d^{i+1} f(x)$ for $i \leq l$, $i \leq k$.
38. (V1.38) An elementary map $z \mapsto d^i f(\pi_0 z)(\pi_1 z, \dots, \pi_i z)$ with $\pi_0 : X \to E^p$, $\pi_a : X \to E^{n_a}$, into $F^m$, with $m + i \leq p$, $m + i \leq n_a + 1$, $i \leq k$ and an sc-domain as domain, is sc^s for $i + s \leq k + 1$; its tangent map is a pair of elementary maps and sums of them.
39. (V1.39) Sums of sc^k maps on an sc-domain are sc^k; the evaluation of continuous multilinear maps has the product-rule derivative.
40. (V2.1) An sc-smooth retraction is an sc-smooth $r : U \to U$, $U$ relatively open in a partial quadrant, with $r \circ r = r$; its retract $O = r(U)$ is its fixed-point set, and $Dr(x)$ is a projection for $x \in O_1$.
41. (V2.2) $(O, C, E)$ is a retract if $O = r(U)$ for such an $r$.
42. (V2.3) Two retractions onto the same $O$ have the same tangent image $Tr(TU) = Tr'(TU')$.
43. (V2.4) $TO = Tr(TU) = \{(r(x), Dr(x)h)\}$, and $T_x O$ is the range of the projection $Dr(x)$.
44. (V2.5) Sc-smoothness of $f \circ r$ and $T(f \circ r)$ on $TO$ do not depend on the retraction $r$ onto $O$, and $T(f \circ r)$ maps $TO$ into $TO'$.
45. (V2.6) $f : O \to O'$ is sc-smooth if $f(O) \subseteq O'$ and $f \circ r$ is sc-smooth for a (every) retraction onto $O$; $Tf = T(f \circ r)|_{TO}$.
46. (V2.7) Sc-smooth maps between retracts compose, with $T(g \circ f) = Tg \circ Tf$.
47. (V2.8) $(TO, TC, TE)$ is a retract with retraction $Tr$, and $Tf$ is sc-smooth between tangents.
48. (V2.9) Open subsets of retracts, and images of idempotent sc-smooth maps on retracts, are retracts.
49. (V2.10) A chart is a homeomorphism of an open set onto a retract; charts are compatible when both transition maps are sc-smooth between the overlap retracts.
50. (V2.11) An atlas is a set of pairwise compatible charts covering $X$; atlases are equivalent when their union is an atlas, and this is an equivalence relation.
51. (V2.12) An M-polyfold is a Hausdorff paracompact space with an atlas, its charts being all charts compatible with it.
52. (V2.13) $f$ is sc-smooth if it is continuous and all its coordinate representations are sc-smooth between retracts; one pair of charts per point suffices.
53. (V2.14) $X_m$ consists of the points that lie on level $m$ in one (every) chart, with the topology generated by the chart preimages of open subsets of $O_m$.
54. (V2.15) The inclusions $X_{m+1} \to X_m$ are continuous.
55. (V2.16) The level charts $(V_1, \varphi, (O_1, C_1, E^1))$ make $X_1$ an M-polyfold.
56. (V2.17) For every M-polyfold, $X$ and every $X_m$ are metrizable, $X_m$ is paracompact, and $X_\infty$ (initial topology of the inclusions) is metrizable; no hypothesis beyond the text's.
57. (V2.18) Tangent tuples $(x, \varphi, h)$ are equivalent when one is the transport of the other by $T$ of the transition map; this is an equivalence relation.

58. (V2.19) $TX$ is the set of classes, with $p : TX \to X_1$ and the bijections $T\varphi : TV \to TO$.
59. (V2.20) $TX$ is Hausdorff, metrizable and paracompact, and $p$ is continuous (openness of $p$ not verified).
60. (V2.21) The charts $(TV, T\varphi, (TO, TC, TE))$ make $TX$ an M-polyfold, with transition maps $T(\varphi' \circ \varphi^{-1})$, and $p : TX \to X^1$ is sc-smooth.
61. (V2.22) An sc-smooth $f : X \to Y$ has a well-defined sc-smooth tangent map $Tf : TX \to TY$ covering $f$, with $T(g \circ f) = Tg \circ Tf$.
62. (V2.23) A sub-M-polyfold is locally the image of an sc-smooth retraction of an open set.
63. (V2.24) A sub-M-polyfold, with the subspace topology, is an M-polyfold for which the inclusion and the maps $i^{-1} \circ r$ are sc-smooth.
64. (V2.25) In a chart, at a smooth $a$, $T_aO = Dt(a)E \oplus (Dr(a) - Dt(a))E$ is an sc-splitting with $Dt(a)E = T_aA$, and $Ts$ maps $T_aO$ onto $T_aA$.
65. (V2.26) $d_X(x)$ is the least $d_C(\varphi(x))$ over the charts $\varphi$ around $x$.
66. (V2.27) $d_X$ is upper semicontinuous.
67. (V2.28) Sc-diffeomorphisms between open sets preserve $d_X$.
68. (V2.29) $\partial X = \{d_X \geq 1\}$ is closed; $X$ is without boundary iff $d_X \equiv 0$.
69. (V2.30) On a sub-M-polyfold $A$, $d_A \leq d_X$.
70. (V2.31) $E_x$ is the closed subspace cut out by the coordinates vanishing at $x$, of codimension $d_C(x)$; $C_x$ the cone on which they are $\geq 0$.
71. (V2.32) $T^R_xO = T_xO \cap E_x$ and $C_xO = T_xO \cap C_x$, defined at every $x \in O_1$.
72. (V2.33) $T^R_xO \subseteq C_xO \subseteq T_xO$, and $C_xO$ is a partial quadrant in $T_xO$ at every smooth point of every sc-smooth retract: the remark's claim that in general it is not is false.
73. (V2.34) For an sc¹ map between quadrants, $Df(x)(E_x) \subseteq E'_{f(x)}$ and $Df(x)(C_x) \subseteq C'_{f(x)}$ at every level-1 point.
74. (V2.35) A germ of sc¹-diffeomorphisms between retracts maps $T^R_xO$ onto $T^R_yO'$ and $C_xO$ onto $C_yO'$, at every level-1 point.
75. (V2.36) A nonempty retract, and a nonempty relatively open subset of a partial quadrant, is an M-polyfold with the chart id.
76. (V2.37) $\dim(T_xO/T^R_xO) \leq d_X(x)$ in every chart of every M-polyfold at points on level 1; the second statement holds without its hypothesis (V2.33).
77. (V2.38) On a relatively open subset of a partial quadrant, the degeneracy index of the M-polyfold equals $d_C$.
78. (V2.39) $C_e$ and $E_e$ are the cone and the subspace cut out by the coordinates vanishing at $e$, in a presentation of $C$.
79. (V2.40) A retraction is tame if it preserves $d_C$ and, at smooth points of $O$, $T_xO$ has an sc-complement inside $E_x$ (for every presentation of $C$).
80. (V2.41) For a tame $r$ at a smooth $x \in O$, $(1 - Dr(x))E \subseteq E_x$ is an sc-complement of $T_xO$; only the second tameness condition is used.
81. (V2.42) For a tame retract, $\dim(T_xO/T^R_xO) = d_O(x)$ at smooth points, where $T^R_xO$ is an sc-subspace and $C_xO$ a partial quadrant in $T_xO$, and $d_O = d_C$ everywhere; a tame chart of any M-polyfold realizes $d_X$.
82. (V2.43) A tame M-polyfold has an equivalent atlas of charts with tame models.
83. (V2.44) A face is the closure of a component of $\{d_X = 1\}$; face-structured means every $x$ lies in exactly $d_X(x)$ faces.
84. (V2.45) Near a point of a tame retract, the retraction maps points of a coordinate hyperplane $f_i$ into the closure of a single component of $\{d_C = 1\} \cap O \cap f_i$.
85. (V2.46) The faces of a tame retract are the components of the sets $f_i \cap O$, distinct for distinct

$i$ through a common point.

86. (V2.47) A tame retract is face-structured.
87. (V2.48) In a tame M-polyfold each point has a tame chart realizing $d_X$ whose model is face-structured, and lies in at most $d_X(x)$ global faces.
88. (V2.49) A face of a tame retract is the retract of $r \circ P_i$ on a neighbourhood of the face, hence a sub-M-polyfold, and with its induced structure it is tame with $d_F = d_O - 1$.
89. (V2.50) Every point of a tame M-polyfold lies in exactly $d_X(x)$ local faces, each tame with $d_F(x) = d_X(x) - 1$; the intersection of $\#\sigma$ of them is, near $x$, a tame sub-M-polyfold $A$ with $d_A = d_X - \#\sigma$.
90. (V2.51) In a tame M-polyfold the interior $F \cap \{d_X = 1\}$ of a face is a component $G$ and a sub-M-polyfold; if $X$ is face-structured, every face is a tame sub-M-polyfold with $d_X = d_F + 1$.
91. (V2.52) $X \times Y$ is an M-polyfold with the product charts, $d_{X\times Y} \leq d_X + d_Y$, with equality and $X \times Y$ tame when $X$ and $Y$ are tame.
92. (V2.53) For a local sc-diffeomorphism $f$ and an sc-smooth $g$ the fibred product is a sub-M-polyfold of $X \times Y$ with $d = d_Y$, tame if $Y$ is; if both are local sc-diffeomorphisms, $d = d_X = d_Y$, and $2d = d_{X\times Y}$ for tame $X, Y$.
93. (V2.54) A strong bundle map preserves the double filtration, is linear in the fibre and sc-smooth on $[0]$ and $[1]$; a strong bundle retraction is an idempotent one, covering an sc-smooth retraction.
94. (V2.55) A local strong bundle retract $K = R(U \lhd F)$ and its part $K[1]$ are sc-smooth retracts, with sc-smooth projections to $O$.
95. (V2.56) Sections of $K$ are sc-smooth into $K[0]$, sc$^+$ into $K[1]$; sc$^+$ implies sc-smooth.
96. (V2.57) A strong bundle chart is a fibrewise linear homeomorphism $P^{-1}(V) \to K$ over a chart of $X$; compatibility means sc-smooth transitions on $[0]$ and $[1]$; equivalence of strong bundle atlases is an equivalence relation, and charts restrict to open subsets of the base.
97. (V2.58) A strong bundle is $P : Y \to X$ with a strong bundle atlas; $Y[0]$ and $Y[1]$ are M-polyfolds, $P[0]$ and $P[1]$ are sc-smooth, and one chart decides whether $y \in Y_{m,k}$; the charts compatible with the atlas depend only on its class.
98. (V2.59) Sections of a strong bundle are sc-smooth into $Y[0]$, sc$^+$ in the charts $\Phi[1]$, which is the same as sc-smooth into $Y[1]$.
99. (V2.60) The pull-back $f^*Y$ is the fibre product of $P$ and $f$.
100. (V2.61) The pull-back of a strong bundle along an sc-smooth map is a strong bundle, with the pulled-back charts and retractions $R_g$.
101. (V2.62) A paracompact Hausdorff locally metrizable space is metrizable (Smirnov), proved in a pull-back form.
102. (V2.63) Sc^k-smoothness restricts to smaller domains and is a local property.
103. (V2.64) $Y[1]$ is Hausdorff and paracompact (metrizable), and the inclusion $Y[1] \to Y[0]$ is sc-smooth.
104. (V3.1) A germ of neighbourhoods is a nested sequence of relatively open neighbourhoods $U_m$ of 0 in $C_m$, with tangent germ $(U_{m+1} \times E_m)_m$; a germ is a fixed sequence of sets.
105. (V3.2) sc$^0$-, sc$^1$-, sc^k- and sc-smooth germs are the notions of Chapter 1 with $U \cap E_m$ replaced by $U_m$.
106. (V3.3) Germs restrict, compose (when the inner germ maps $U_m$ into $V_m$ level by level), pair, and sc-operators are sc-smooth germs.
107. (V3.4) Propositions 1.16–1.18 hold for germs.
108. (V3.5) An sc-smooth section germ is a section whose principal part composed with $r$ is an sc-smooth germ, on some germ of neighbourhoods inside $U$, in some centred strong bundle chart compatible with the atlas.

109. (V3.6) Sums of sections have sums of principal parts; sc$^+$-sections have sc-smooth representatives $O \to K[1]$ in every compatible chart.
110. (V3.7) A filling is an sc-smooth germ satisfying (1)–(3) of Definition 3.4 near $0$, the map in (3) an sc$^1$-germ (the text's germ reading).
111. (V3.8) For a filler with $g(0) = 0$, $Tf(0) = Dg(0)|_{Dr(0)E}$ is onto iff $Dg(0)$ is, Fredholm iff $Dg(0)$ is, with the same index; if $Dg(0)$ is sc-Fredholm, so is $Tf(0)$, and condition (3) holds on every level.
112. (V3.9) A filled version is a filling of the principal part in the chart of Definition 3.11, an sc-smooth germ whose levels may shrink with $m$.
113. (V3.10) A basic germ is an sc-smooth germ with $f(0) = 0$ whose $W$-component is $w - B(a, w)$ with $B$ an sc$^0$-contraction germ.
114. (V3.11) For sc$^1$ sc$^0$-contraction germs, Lemma 3.9 as stated: $|D_2B(x)\zeta|_m \le \varepsilon|\zeta|_m$ for $x \in U_{m+1}$ small on level $m$, for every $\varepsilon$ and $m$; and $D_2B(0,0) = 0$.
115. (V3.12) $D(h+s)(0)$ is sc-Fredholm of index $n-N$ for a basic germ $h$ and an sc$^+$-germ $s$, and on every level $m$, $D(h+s)(x)$ is Fredholm of index $n-N$ for $x \in U_{m+1}$ small on level $m$.
116. (V3.13) An sc-Fredholm germ has a filling germ $g$ and an sc$^+$-germ $s$, $s(0) = g(0)$, with $g - s$ conjugated to a basic germ near $0$ (the text's germ reading).
117. (V3.14) Near a zero of an sc-Fredholm germ, the closures of the zeros in $O(i)$ are compact subsets of $X_i$.
118. (V3.15) Zeros converging to the base point (or, for sc-Fredholm sections, to any point) converge on every level.
119. (V3.16) An sc-Fredholm section is sc-smooth, regularizing and an sc-Fredholm germ at every smooth point.
120. (V3.17) An sc-Fredholm section plus an sc$^+$-section is sc-Fredholm, for any strong bundle.
121. (V3.18) $s \circ r$ agrees with $s$ on $O$ and $R$ fixes $(r(u), s(r(u)))$.
122. (V3.19) $g + s \circ r$ is a filling germ of $f + s$ when $g$ is one of $f$, for $s$ an sc-smooth section of $K$ (a germ suffices).
123. (V3.20) Good position: $N \cap C$ has interior in $N$ and an algebraic complement satisfies the $\varepsilon$-condition; the text's sc-complement form is also formalized and holds for smooth $N$ on a partial quadrant.
124. (V3.21) If $N$ is in good position to $C$, then $N \cap C$ is a partial quadrant in $N$.
125. (V3.22) The tangent germ property, with $Y$ closed and "sc^∞-germ" read with $k$-dependent neighbourhoods.
126. (V3.23) The tangent germ property is invariant under sc-diffeomorphisms.
127. (V3.24) Under the regularity conditions (a) $\delta$ sc-smooth on $V$, (b) $\pi(O) \subseteq C$, a set with the tangent germ property is a sub-M-polyfold, tame, with corner charts $\pi\varphi$; without (a), (b) the text's proof fails in examples.
128. (V3.25) An sc$^0$-contraction germ has the contraction estimate on small balls on every level, and a working form with explicit regions.
129. (V3.26) The solution germ exists, is sc$^0$, and is unique in the contraction region.
130. (V3.27) If $B$ is sc$^1$, the solution germ is sc$^1$ on the germ shifted by one level.
131. (V3.28) $f^{(1)}$ is a contraction germ of class sc^{min(k, j)} with solution germ $T\delta$; it satisfies Definition 3.28.
132. (V3.29) If $B$ is sc^k, the solution germ is sc^j on some shifted germ for every $j \le k$.
133. (V3.30) If $B$ is sc^k, the solution germ is sc^k on the germ shifted by $2^k - 1$ levels; the text's hypotheses suffice.
134. (V3.31) If $B$ is sc-smooth, $\delta$ is $C^k$ into $W_m$ on a neighbourhood $V_{m,k}$ of $0$; $\delta$ is an sc-smooth

germ on one germ.

135. (V3.32) For a quadrant-like $\widetilde{C}$, any representative of the solution germ of an sc^k (sc-smooth) contraction germ is sc^k (an sc-smooth germ) on a smaller germ.
136. (V3.33) The basic class is Definition 3.7.
137. (V3.34) A linear coordinate change makes the index-raised push-forward of a basic germ plus an sc$^+$-germ a basic germ, with $n - N$ unchanged.
138. (V3.35) The solutions near 0 of a contraction germ plus an sc$^+$-germ have compact closures on every level, for closed $\widetilde{C}$.
139. (V3.36) Zeros of $h + s$ near 0 lie on level 1, where the linearization is surjective Fredholm of index $n - N$.
140. (V3.37) Zeros of the push-forward near 0 have their $X$-component small on level 1.
141. (V3.38) Under Lemma 3.42's regularity, the zeros of $g$ near 0 form the graph of a $C^1$ map $\tau$ over $\ker Dg(0) \cap C'$ into a good complement, $\tau(0) = 0$, $D\tau(0) = 0$, and $\tau$ is $C^k$ into level $k$ near 0 for every $k$.
142. (V3.39) $\ker Dg(0)$ is the graph of $D\delta(0)$ over $\ker DH(0)$.
143. (V3.40) Linear isomorphisms carrying $C$ onto $C'$ carry good complements and good position.
144. (V3.41) $\ker DH(0)$ has a good complement $Z$ with $Z \times W$ a good complement of $K'$.
145. (V3.42) The zeros of $f + s$ near 0 form the graph of a $C^1$ map $\sigma$ over $\ker D(f + s)(0) \cap C$, $\sigma(0) = 0$, $D\sigma(0) = 0$, $C^k$ into level $k$ near 0 for every $k$.
146. (V3.43) Definition 3.48 is Definition 3.4.
147. (V3.44) After shrinking, (1) and (2) of the filler hold on all of $O \cap W$, $U \cap W$.
148. (V3.45) Definitions 3.50, 3.51 are Definitions 3.11, 3.14 for a tame strong bundle; the theorems of §3.6 use them with the chart of the germ data tame (V3.103).
149. (V3.46) In a centred chart, $Dg(0)$ and $Tf(x_0)$ are sc-Fredholm with index $n - N$ on every level, the dimensions of any basic germ to which $g - s$ is conjugated; at a zero of an sc-Fredholm section the linearization is sc-Fredholm in every chart.
150. (V3.47) Good position of the germ: $Tf(0)$ onto and $\ker Dg(0)$ in good position to $T_0O \cap C$ in $T_0O$, in a centred chart; also with the text's sc-complement, the same at zeros of sc-Fredholm sections with tame germ data.
151. (V3.48) Near a zero in good position in a tame chart, the zero set is carried injectively by $\psi\varphi$ onto the graph of a $C^1$ map $\sigma$ over $\ker D(h + t)(0) \cap C'$, with $Y$ closed, the kernel smooth, and $C^k$-lifts of $\sigma$ into level $k$.
152. (V3.49) Along the graph of $\tau$ near 0, $Y$ (closed) stays a good complement of $N_v$ with respect to $C_{v+\tau(v)}$; also for $\tau$ on a relatively open $V$ and any presentation of $C$.
153. (V3.50) Without boundary, a zero set with the regular tangent germ property is a manifold without boundary.
154. (V3.51) A zero set with the regular tangent germ property is a tame sub-M-polyfold and a manifold with boundary with corners.
155. (V3.52) Injectivity of $L(0)$ and the compactness property make $L(b)$ injective for $b$ near 0; in Theorem 3.58 both hypotheses follow from its condition (2).
156. (V3.53) On a compact parameter set with injective $L(b)_m$ there is a uniform lower bound $c_m$.
157. (V3.54) $(v, k) \mapsto L(v)^{-1}k$ is an sc-smooth germ under uniform lower bounds $c_m$.
158. (V3.55) A good complement has vanishing coordinates in $\Sigma$.
159. (V3.56) $d_{N\cap C} = d_C$ off $N \cap W$, and $d_{N\cap C} + \dim(N \cap W) = \dim N$ on it.
160. (V3.57) $d_{N\cap C}(v) = d_C(v + \tau(v))$ for $v \notin N \cap W$ near 0.
161. (V3.58) The quadrant inverse function theorem holds for $C^j$ maps.
162. (V3.59) A $C^j$ map with $\|Df - 1\| \leq \frac{1}{2}$ preserving interior points is a $C^j$-diffeomorphism

onto a relatively open set.

163. (V3.60) Such a map is bi-Lipschitz, injective, and has relatively open image.
164. (V3.61) Such a map is a homeomorphism onto its image.
165. (V3.62) Its inverse has derivative $[Df(g(y))]^{-1}$.
166. (V3.63) The implicit function theorem in partial quadrants, with $f(k+\tau(k)) = 0$ and $k+y \in U$.
167. (V3.64) There is a region on which $B(k,\cdot)$ is a $\frac{1}{2}$-contraction and fixed points are unique.
168. (V3.65) $B(k,\cdot)$ maps $\{|y| \leq \varepsilon|k|\}$ into itself.
169. (V3.66) $\tau$ is continuous with $|\tau(k)| \leq \varepsilon|k|/2$, Lipschitz with constant $\varepsilon/(1+\varepsilon) \leq \varepsilon$ on all of $V_0$.
170. (V3.67) $\tau$ is $C^j$ on the whole $V_0$, with the derivative formula.
171. (V3.68) Under the filler condition at the nearby zeros (a consequence of good position, V3.102), near a zero in good position in a tame chart of an sc-Fredholm section with tame germ data over an M-polyfold without boundary the zeros are smooth points in good position in tame charts and form a manifold without boundary; the forms with the text's hypotheses, the index and the kernel are V3.95, V3.97, V3.99, V3.101.
172. (V3.69) Near a zero in good position in a tame chart whose nearby zeros are in good position in tame charts (or, for tame germ data, under the filler condition of V3.82), the local solution set consists of smooth points and is a tame sub-M-polyfold equivalent to a $d$-dimensional manifold with boundary with corners.
173. (V3.70) With good position in a tame chart at every zero ($S \neq \emptyset$), the solution set is a sub-M-polyfold with tame induced structure, a manifold with boundary with corners of locally constant dimension; no hypothesis is added.
174. (V3.71) The image of a tame retract under an sc-diffeomorphism, and an open subset of a tame retract, are tame retracts.
175. (V3.72) A nonempty set with regular data of Definition 3.25 of some dimension at every point is a sub-M-polyfold, tame, with an atlas of finite-dimensional corner charts $\pi\varphi$.
176. (V3.73) Two graph descriptions of one set, the first $C^1$ and the second centred at a point of it with $C^k$-lifts at $0$, have the same dimension, and the first is $C^k$ into every level near that point.
177. (V3.74) Data of Definition 3.25 with $\pi(O) \subseteq C$, at a point whose neighbours all have data, can be shrunk to regular data, and the dimension is locally constant.
178. (V3.75) A finite-dimensional subspace in good position has a closed good complement along which the projection maps the partial quadrant into itself.
179. (V3.76) If $g$ is conjugated to a basic germ and $s$ is an sc$^+$-map with $s(0) = 0$, then $(g+s)^1$ is conjugated to a basic germ, by $\Lambda \circ \Phi^1$.
180. (V3.77) Theorems 3.43, 3.47 and the graph description of Theorem 3.54 hold with a closed good complement whose projection maps the partial quadrant into itself.
181. (V3.78) At a zero, the linearizations in two strong bundle charts are conjugate by isomorphisms respecting the partial quadrants: surjectivity, Fredholm index and good position of the kernel do not depend on the chart.
182. (V3.79) Surjectivity onto a dense part of $K_z$ and good position of $\ker Dg(z) \cap T_z O$, in a chart not centred at the zero $y$ ($z$ in the germ of $g$), give good position of $(f,y)$ in a tame chart, for tame germ data.
183. (V3.80) In the chart at $x_0$, at the zeros near $x_0$, $D(h+t)$ is surjective Fredholm of index $n-N$ and the tangent space $N_v$ of the graph is in good position with complement $Y$.
184. (V3.81) In the chart at $x_0$, at the zeros near $x_0$, $\ker Dg(z_0) = D\psi^{-1}(N_v)$ lies in $T_{z_0}O$ and is

in good position there; no hypothesis is added.

185. (V3.82) For tame germ data, if the filler in the tame chart at $x_0$ also satisfies condition (3) of Definition 3.4 at its zeros near 0, every zero near $x_0$ is in good position in a tame chart; the text's argument needs that condition, which holds whenever $(f, x_0)$ is in good position (V3.102).
186. (V3.83) A good complement of $\mathbb{R} \times N$ projects into the lineality space; the tilted kernel of the parameter extension is in good position.
187. (V3.84) At a zero $x_0$ in good position in a tame chart, the solution set near $x_0$ has data of Definition 3.25 at $x_0$ in the chart $\psi\varphi$, with $\pi(O) \subseteq C$.
188. (V3.85) If $x_0$ and the zeros near it are in good position in tame charts, the solution set has regular data at $x_0$ of a dimension $d$ that every nearby datum shares; Theorem 3.54 (1)–(2) under the filler condition of V3.82.
189. (V3.86) Over an M-polyfold without boundary, with good position in a tame chart at every zero, the solution set is a manifold without boundary of locally constant dimension.
190. (V3.87) With good position in a tame chart at every zero (the text's hypotheses), the solution set is a sub-M-polyfold with tame induced structure, a manifold with boundary with corners of locally constant dimension, the disjoint union of its open and closed parts $S_n$ of dimension $n$.
191. (V3.88) Given sc-isomorphisms $L(b) = P_Y Df(b,0)|_X$ with uniform lower bounds and condition (3), $\Phi_*(f - f(0))$ is a basic germ with contraction $x - L(b)^{-1}P_Y(f - f(0))$.
192. (V3.89) Under conditions (1)–(3) of Theorem 3.58, in the coordinates of its proof, $f - f(0)$ is conjugated near 0 to a basic germ.
193. (V3.90) A $C^j$ map with $\|Df - 1\| \leq \frac{1}{2}$ on a convex set, whose image has unique differentiability, has a $C^j$ inverse with $Dg = [Df \circ g]^{-1}$.
194. (V3.91) For a strong bundle (tameness not needed), an sc-Fredholm section $f$ and sc$^+$-sections $s_i$, $(r, x) \mapsto f(x) + \sum r_i s_i(x)$ is an sc-Fredholm section of the pull-back bundle over $\mathbb{R}^n \times X$.
195. (V3.92) In the product chart at $(0, x)$ the linearization of the perturbed section is $(\delta r, \xi) \mapsto f'(x)\xi + \sum \delta r_i s_i(x)$ (chart form); at $(r_0, x)$ the first term is that of $f + \sum r_{0,i} s_i$.
196. (V3.93) In the chart at a zero $x_0$ in good position, at every zero $z_0$ of the filler near 0 and for every smooth $k$, $\rho(z_0)k = Dg(z_0)x$ for some $x \in T_{z_0}O$, by a curve of zeros of $g + \mu\rho(r(\cdot))k$ through $z_0$ from an implicit function theorem at $z_0$.
197. (V3.94) With the text's hypotheses (tame germ data, good position in a tame chart), every zero near a zero in good position is in good position in a tame chart, and the solution set has there regular data of a dimension that every nearby datum shares; also with good position in the text's sense (sc-complement) in the conclusion.
198. (V3.95) With the text's hypotheses (tame germ data, good position in a tame chart), the local solution set near a zero in good position consists of smooth points in good position in tame charts and is a tame sub-M-polyfold, a $d$-dimensional manifold with boundary with corners, without boundary if $d_X \equiv 0$.
199. (V3.96) At a zero, the linearization in every chart is sc-Fredholm of the index $n - N$ of any conjugating basic germ on every level, and onto on every level with $\dim\ker = n - N$ if the germ is in good position.
200. (V3.97) The dimension of the solution set near a zero in good position in a tame chart is the Fredholm index, $\dim\ker f'(y) = \operatorname{ind} f'(y)$, at every zero near it and in every chart; also in Theorems 3.23, 3.19, 3.57.
201. (V3.98) For tame germ data, near a zero in good position in a tame chart, every zero $y$ has $f'(y)$ onto with $\operatorname{ind} f'(y) = \operatorname{ind} f'(x_0)$: the index is locally constant along the solution set;

for tame germ data, without bump functions, the index is locally constant along the zero set without good position, the index of Definition 6.1 with a locally defined sc$^+$-section is well defined and exists at every smooth point, and it is constant along every sc-smooth path $\mathbb{R} \to X$ (Proposition 6.2); under bump functions the local and the global reading of Definition 6.1 agree.

202. (V3.99) At the zeros near $x_0$, $D\psi$ maps $\ker f'(y)$, read in the chart at $x_0$, onto the tangent space of the solution graph.
203. (V3.100) For an sc-Fredholm section with tame germ data and a zero $y$ with $d_X(y) = 0$, $(f, y)$ is in good position in a tame chart if and only if $f'(y)$ is onto.
204. (V3.101) Over an M-polyfold without boundary, Theorems 3.19 and 3.56 hold for tame germ data with the hypothesis "$f'(x)$ surjective", with tame induced structure and the dimension the index.
205. (V3.102) At a point of the retract with $\ker Dg \subseteq T_{z_0}O$, $\rho(z_0)F \subseteq Dg(T_{z_0}O)$ implies condition (3) of the filler there (and conversely when $Dg$ is onto); so the filler condition at the nearby zeros holds whenever $f$ has tame germ data and $(f, x_0)$ is in good position in a tame chart.
206. (V3.103) The theorems of §3.6 assume, as Definitions 3.50, 3.51 and 3.53 do, germ data and good position in a strong bundle chart whose base retraction is tame; compatible charts whose base retraction is the identity of $U$ are tame, so the hypotheses can be met.
207. (V3.104) The product of the constant retraction onto $(1, 0)$ of $[0, \infty) \oplus W$ with any sc-smooth retraction of a nonempty set is not tame: the degeneracy index drops by one under it.
208. (V3.105) If $f$ is an sc-Fredholm section with tame germ data and $s$ an sc$^+$-section, $f + s$ has tame germ data, in the same charts.
209. (V3.106) If $f$ has tame germ data, $(r, x) \mapsto f(x) + \sum r_i s_i(x)$ is an sc-Fredholm section of $\pi^*P$ with tame germ data: the product chart of a tame chart is tame.
210. (V3.107) The good position of $(f, y)$ obtained in one chart holds in every chart carrying germ data at $y$, in particular in the tame one.
211. (V4.1) A corner chart is a chart onto a relatively open subset of a partial quadrant in a finite-dimensional space with the constant structure; a corner structure is an equivalent atlas of corner charts, of possibly varying dimension, and makes $X$ tame.
212. (V4.2) The transition maps between compatible corner charts are $C^\infty$ on their relatively open overlaps.
213. (V4.3) An sc-Banach space with finite-dimensional $E_0$ has all levels equal and is sc-isomorphic to $E_0$ with the constant structure; all its points are smooth.
214. (V4.4) Corner structures are the atlases of charts onto relatively open subsets of partial quadrants in sc-Banach spaces with finite-dimensional $E_0$; those of fixed dimension $n$ are the manifolds with corners of Theorem 3.27.
215. (V4.5) An M-polyfold, tame or not, has a corner structure iff all its points are smooth, $1_0^1$ is sc-smooth, and every point is the centre of a tame chart whose $T_0O$ is finite-dimensional and in good position.
216. (V4.6) $1_1^0 : X^1 \to X$ is sc-smooth; if $X = X_1$ and $1_0^1$ is sc-smooth it is an sc-diffeomorphism; some M-polyfold with $X_0 = X_\infty$ has $1_0^1$ not sc-smooth; a finite-dimensional space with the constant structure has $V^1 = V$, and all its retracts are sc$^+$-retracts.
217. (V4.7) An sc$^+$-map is a map with sc-smooth lift into $F^1$; it raises levels, is sc-smooth, and is preserved by composition with sc-smooth maps on either side.
218. (V4.8) At a smooth point the derivative of an sc$^+$-map is an sc$^+$-operator, compact on every level; a compact projection has finite-dimensional range.

219. (V4.9) An sc$^+$-retraction is an sc-smooth retraction that is an sc$^+$-map (on an open subset of an M-polyfold: with values in $X_1$ and $1_0^1 \circ s$ sc-smooth).
220. (V4.10) A retraction with the same retract as an sc$^+$-retraction is an sc$^+$-retraction.
221. (V4.11) An sc$^+$-retract is the retract of some, equivalently of every, retraction, which is sc$^+$.
222. (V4.12) The retract of an sc$^+$-retraction consists of smooth points and its tangent spaces are finite-dimensional.
223. (V4.13) A retract is an sc$^+$-retract iff it consists of smooth points and the lift $O \to O^1$ is sc-smooth between retracts.
224. (V4.14) A chart's model is an sc$^+$-retract iff its domain is smooth and $1_0^1$ is sc-smooth on it; "$X_0 = X_\infty$ and $1_0^1$ sc-smooth" iff every chart's model is an sc$^+$-retract.
225. (V4.15) $C_e$ is a partial quadrant with $C_e - e = C_e$; near $e$ it agrees with $C$, and re-centring preserves degeneracy indices, minimal subspaces and the cones $C_v$.
226. (V4.16) Translation by a smooth $e$ is an sc-diffeomorphism from relatively open subsets of $C$ near $e$ onto relatively open subsets of $C_e$, and carries retracts (tame retracts) to retracts (tame retracts).
227. (V4.17) Every smooth point of a chart's domain is the centre of a re-centred chart of $X$, tame if the original is.
228. (V4.18) Re-centring a strong bundle chart in the base gives a strong bundle chart; every smooth point of the base is the centre of one.
229. (V4.19) An sc$^+$-retraction is $C^1$ on every level $k+1$ with operator-norm-continuous derivative, and satisfies the mean value estimate with any constant $\theta$ on small balls of $C_{k+1}$.
230. (V4.20) For an sc$^+$-retraction onto $O \ni 0$ with $T_0O$ in good position, a closed good complement $Y$ and the projection $p$ along $Y$, $p$ is an sc-diffeomorphism from $O \cap W$ onto a relatively open $V \ni 0$ of $T_0O \cap C$, with inverse $\Gamma$, $\Gamma(a) - a \in Y$.
231. (V4.21) Near 0, $(1-p)$ contracts differences of points of $O$ on level $k+1$, so two such points with the same projection coincide.
232. (V4.22) For small $a \in N \cap C$ there is a point $u \in O$ with $p(u) = a$, $u - a \in Y$ small on level 1.
233. (V4.23) Projections of points of $O$ near 0 lie in $C$, and the neighbourhoods $W$, $V$ and the inverse $\Gamma$ exist with the required level-1 control.
234. (V4.24) $\Gamma$ is Lipschitz on level 1, continuous into every level, and differentiable into every level with the frame formula $D\Gamma(a) = Dt(\Gamma(a))|_N (p\, Dt(\Gamma(a))|_N)^{-1}$.
235. (V4.25) $\Gamma$ is $C^\infty$ into every level, hence sc-smooth on the fixed $V$; $p$ and $\Gamma$ are sc-smooth between the retracts.
236. (V4.26) A chart composed with an sc-diffeomorphism between retracts is a chart of the M-polyfold.
237. (V4.27) Corner charts re-centre to tame centred charts whose $T_0O$ is the whole finite-dimensional space, in good position; so (i) implies (ii).
238. (V4.28) Under (ii), $p \circ \psi$ (with a closed good complement) is a corner chart of $X$ at every point; so (ii) implies (i).
239. (V4.29) A smooth finite-dimensional submanifold is locally the image of an sc$^+$-retraction on an open set; it is a sub-M-polyfold.
240. (V4.30) In every chart, $\varphi \circ s \circ \varphi^{-1} \circ r$ is an sc$^+$-retraction onto the chart image of $A$, whose points are smooth.
241. (V4.31) In a chart, $T_aA$ is the image of $T_aX$ under the tangent map of $s$, lies in $T_aX$, and is finite-dimensional.
242. (V4.32) At points with $d_X = 0$ a submanifold is in good position; if $d_X = 0$ on all of $A$, $A$ has a corner structure, without tameness of $X$.

243. (V4.33) $A \subseteq X_\infty$, and all points of $A$ are smooth for its induced structure.
244. (V4.34) $A$ is locally closed in $X$.
245. (V4.35) $\partial A \subseteq \partial X$.
246. (V4.36) $1^1_0 : A \to A^1$ is sc-smooth for the induced structure.
247. (V4.37) Tangent spaces of retracts are germs, are monotone under inclusion of retracts, and are mapped into each other by derivatives of maps between retracts.
248. (V4.38) Good position of $A$ at $a$ means, in a chart, that $T_z A$ is in good position to $C_z$ inside $T_z O$.
249. (V4.39) Good position of $A$ at $a$ does not depend on the chart, the data of Definition 4.13 or the presentation.
250. (V4.40) A tame submanifold is one whose induced M-polyfold structure is tame.
251. (V4.41) A smooth finite-dimensional submanifold of a tame M-polyfold, in good position everywhere, has a corner structure and is tame.
252. (V4.42) In a tame centred chart, good position inside $T_0 O$ gives good position in $E$, with good complement $Y \oplus (1-\pi)E$ and constant $\varepsilon/(\|\pi\|+1)$.
253. (V4.43) For tame $X$, good position at a point holds at all nearby points of $A$.
254. (V4.44) General position of $A$ at $a$ means, in a chart, an sc-complement of $T_z A$ in $T_z O$ inside $T^R_z O$, level by level.
255. (V4.45) General position does not depend on the chart, the data or the presentation.
256. (V4.46) At the corner of a centred chart, $N + E^{\min}_0 = E_0$ gives onto coordinates, good position with a closed complement in $E^{\min}_0$, and $d_{N\cap C} = d_C$ on $N \cap C$.
257. (V4.47) A finite-dimensional smooth $N$ has in $N + R$ an sc-complement inside the sc-subspace $R$; $T_y Q + E^{\min}_0 = E_0$ gives general position at $y$.
258. (V4.48) In a tame retract, general position at a smooth point implies good position.
259. (V4.49) In a tame centred chart, general position at the centre gives the graph description with $Y \subseteq E^{\min}_0$, equal degeneracy indices along the graph, and general position along it.
260. (V4.50) Theorem 4.22 (1)–(4) holds in a tame chart centred at a point of general position, with $F_\sigma$ a germ inside $U$; (1) and (4) for the faces of the chart model, not of $U$ itself.
261. (V4.51) $\Gamma \circ \kappa_\sigma \circ p \circ t$ is an sc$^+$-retraction onto $A \cap F_\sigma$ in the chart; its corner model is sc$^+$ and in general position in the face model.
262. (V4.52) An sc$^+$-retraction in a chart is an sc$^+$-retraction on an open subset of $X$; $A \cap F_\sigma$ is a tame submanifold of the tame $F_\sigma$, in general position in it.
263. (V4.53) A stratified regular point is one where the projection maps the tangent space of the stratum through it onto $\mathbb{R}^m$, in some product chart.
264. (V4.54) Theorem 4.23: almost every $\lambda$ with $|\lambda| < \varepsilon$ is a stratified regular value whose fibre is a compact submanifold in general position with a tame corner structure.
265. (V4.55) In a tame chart $\operatorname{codim} T^R_x X = d_X(x)$; general position at $a$ gives $d_X(a) \le \dim T_a A$, and a zero-dimensional submanifold in general position avoids $\partial X$.
266. (V4.56) Sard: the critical values of a $C^\infty$ map between finite-dimensional spaces form a Haar-null set.
267. (V4.57) The strata of $\{\ell \ge 0\}$ have dimension $\dim V - \#\sigma$, and the stratified critical values of a $C^\infty$ map form a null set.
268. (V4.58) Theorem 3.78 for $C^\infty$ data has a $C^\infty$ solution.
269. (V4.59) At a stratified regular point the kernel of $Df$ is in good position to the local partial quadrant, with a closed complement in the stratum.
270. (V4.60) Near a stratified regular point the fibre of $f$ is the image of a $C^\infty$ retraction whose derivative at the point is the identity on $\ker Df$.
271. (V4.61) $e \mapsto (\Gamma(\rho(p(t(c,e)))))_2$ is an sc$^+$-retraction onto the chart image of $A_c$, whose tangent

space contains the vertical tangent vectors of $A$.

272. (V4.62) General position of $A$ and stratified regularity at $z$ give general position of the fibre at $z_2$.
273. (V4.63) Every point of $A$ has a neighbourhood and a null set of parameters off which the conclusions of Theorem 4.23 hold locally.
274. (V4.64) Countably many such neighbourhoods cover $A$, so for any parameter set the good parameters have null complement.
275. (V5.1) Every strong bundle chart compatible with the atlas reads $Y_{0,1}$ by the fibre component in $F_1$, and its $\Phi[1]$ is a chart of $Y[1]$.
276. (V5.2) The points of $P^{-1}(x)$ in $Y_{0,1}$ form a linear subspace, complete for the chart norm $|\cdot|_1$; the norms of two charts are locally uniformly comparable.
277. (V5.3) An auxiliary norm is a continuous function on $Y[1]$, a complete norm on every fibre, with $P(w_k) \to x$ and $N(w_k) \to 0$ forcing $w_k \to 0_x$.
278. (V5.4) A continuous fibrewise complete norm is auxiliary iff near every point it is two-sided comparable with $|h|_1$ in some strong bundle chart, also literally on the whole domain of the chart; auxiliary norms are so in every chart.
279. (V5.5) Every strong bundle has an auxiliary norm, glued from the chart norms by a partition of unity.
280. (V5.6) Two auxiliary norms are comparable by a continuous positive function $f$: $fN_0 \le N_1 \le f^{-1}N_0$.
281. (V5.7) Compact solution set, weakly proper and proper are as in the text, with $N = \infty$ off $Y_{0,1}$; for continuous $f$ proper implies weakly proper implies compact solution set.
282. (V5.8) Near 0, the points where a basic germ (on a closed $\tilde{C}$) has level-1 norm $\le c$ lie in a compact set, on a neighbourhood independent of $c$.
283. (V5.9) Strong bundle maps have locally uniform $(0,1)$ fibre bounds; at an sc-Fredholm germ the points with $|\Phi(f(x))_2|_1 \le C$ near $x_0$ have compact closure, for every $C$.
284. (V5.10) An sc-Fredholm section with compact solution set is proper, without tameness; proper, weakly proper and compact solution set are equivalent for sc-Fredholm sections.
285. (V5.11) At a zero of an sc-Fredholm section there are nested neighbourhoods $U(i)$ with $\mathrm{cl}\{y \in U(i) \mid f(y) = 0\}$ compact in $X_i$.
286. (V5.12) With the initial topology of the inclusions, $X_\infty$ is metrizable, and a compact solution set of an sc-Fredholm section is compact in $X_\infty$, without tameness.
287. (V5.13) For an sc-Fredholm section with compact solution set, $f^1$ is an sc-Fredholm section of $P^1$ with compact solution set, and so is $f^2$ of $P^2$; the solution set is compact in every $X_i$.
288. (V5.14) $P^1 : Y_{1,1} \to X^1$ with the index raises of the strong bundle charts is a strong bundle, and for every sc-Fredholm section $f$, $f^1$ is an sc-Fredholm section of it, the filler condition on level 1 included.
289. (V5.15) $\mathbb{R}$ with the constant structure is an M-polyfold; sc-smoothness of maps is checked in local charts; $C^\infty$ functions of sc-smooth functions are sc-smooth.
290. (V5.16) Admitting sc-smooth bump functions means a bump with $f(x) \neq 0$ inside every neighbourhood of $x$; Definition 5.29 as written is weaker, follows from the density of such points, and gives them at the smooth points of an sc-Banach space.
291. (V5.17) With bump functions, every $x$ has an sc-smooth $f : X \to [0,1]$ supported in a given neighbourhood, equal to 1 near $x$.
292. (V5.18) The zero section, products of sc-smooth functions and sc$^+$-sections, and finite sums of sc$^+$-sections are sc$^+$-sections; sc$^+$-sections take values in $Y_{0,1}$.
293. (V5.19) A local sc$^+$-section of a chart of the atlas, cut off by an sc-smooth function supported

in the chart domain and extended by $0$, is an sc$^+$-section.

294. (V5.20) With bump functions and an auxiliary norm, every smooth $e$ over a smooth $x$ is the value at $x$ of an sc$^+$-section supported in $U$ with $N(s(y)) < N(e) + \varepsilon$ everywhere.
295. (V5.21) $\Gamma^{+,1}_U(P)$ is the set of sc$^+$-sections supported in $U$ with $N \le 1$; $\rho = \sup_x N(s(x) - s'(x))$ is a metric on it, not complete for the trivial bundle over $\mathbb{R}$, complete for $U = \emptyset$.
296. (V5.22) A section vanishing off an adapted $U$ with $N(s) \le 1$ has the zeros of $f + s$ in $\{x \in U \mid N(f(x)) \le 1\}$, with compact closure; for $s$ in the completion (uniform $N$-limits of allowable sections) the solution set of $f + s$ is closed, hence compact.
297. (V5.23) Transversality to the zero section is surjectivity of the linearization at every zero, in some and then every chart.
298. (V5.24) At a zero of an sc-Fredholm section, finitely many allowable sc$^+$-sections supported in a given $U$ span a complement of the range of the linearization.
299. (V5.25) The slices $g + \sum \lambda_i t_i$ carry the zeros of $F$, are sc-Fredholm, and stay allowable for $m|\lambda| \le \varepsilon$ when $N(s_0) \le 1 - \varepsilon$.
300. (V5.26) A slice over a parameter where $\ker \mathrm{Lin}_F$ projects onto $\mathbb{R}^m$ is transversal; spanning sections make $F$ transversal at $(0, y)$, also after adding sections.
301. (V5.27) $\mathbb{R}^m \times X$ has no boundary if $X$ has none; transversality of $F$ spreads to nearby zeros and gives good position on an open $V \supseteq \{0\} \times S_g$, which contains the zeros over small $\lambda$.
302. (V5.28) Near a zero of $F$ in good position, the slices are transversal at their zeros over all $\lambda$ off a null set.
303. (V5.29) Theorem 5.13: for $\partial X = \emptyset$, $f$ with tame germ data, there is $U$ such that allowable perturbations have compact solution sets and $s_0 + \sum \lambda_i t_i$ is transversal with a manifold solution set for small $\lambda$ off a null set.
304. (V5.30) Theorem 5.13, dense form: such good allowable perturbations exist within $\rho$-distance $\varepsilon$ of every allowable $s'$.
305. (V5.31) General position of $(f, y)$: in a tame chart, $\mathrm{Lin}(z)$ onto and its kernel with an sc-complement inside $T^R_z O$, for every presentation.
306. (V5.32) Regular tangent germ data give sc$^+$-retractions, so solution sets in good position are smooth finite-dimensional submanifolds (Definition 4.13).
307. (V5.33) Near a zero in good position, the tangent space of the solution set equals $\ker \mathrm{Lin}(z)$ in every chart.
308. (V5.34) Near a zero in good position, general position of $(f, y)$ is general position of the solution set at $y$, and is chart independent.
309. (V5.35) For $X$ tame, general position of $(f, x_0)$ persists at the nearby zeros, with the solution set a submanifold in general position there.
310. (V5.36) Properties (i)–(ii) are general position of $F$ at $(0, y)$, survive added sections, and are achieved by finitely many allowable sections.
311. (V5.37) For $X$ tame, a family with $F$ in general position along $\{0\} \times S_{f+s_0}$ gives, for small $\lambda$ off a null set, $f + s_\lambda$ with all the conclusions of Theorem 5.18.
312. (V5.38) Theorem 5.18, generic form, for $X$ tame with bump functions and $f$ with tame germ data.
313. (V5.39) Theorem 5.18, dense form: perturbations in general position exist within $\rho$-distance $\varepsilon$ of every allowable $s'$.
314. (V5.40) Theorem 5.19: with good position at the boundary zeros, small $\sum \lambda_i t_i$ vanishing near $\partial X$ give transversality and good position everywhere, a compact manifold with corners.
315. (V5.41) A generalized compact homotopy is a tame $X$, $f$ with tame germ data and compact solution set, and $t$ onto $[0, 1]$ with tame fibres on which $f$ restricts to sc-Fredholm sections of

the pull-backs.

316. (V5.42) Lemma 5.23 is linear algebra, and the one-dimensional complement $Z$ of $\ker dt(x)$ exists iff $dt(x) \neq 0$.
317. (V5.43) The product rule and the affine chain rule hold at level-1 points; $\beta \cdot \ell(\varphi(\cdot) - \varphi(x))$ is sc-smooth.
318. (V5.44) Lemma 5.24: for $\mathrm{Lin}_f(z) \circ b$ with values in the line of a smooth $e$, an sc$^+$-section vanishing at $x$ with small support cancels it on $K$.
319. (V5.45) Theorem 5.25: for $X$ tame and $f$ proper with tame germ data, some allowable $s$ puts $f + s$ in good position at every zero.
320. (V5.46) Lemma 5.26: general position implies good position, with the interior condition, without tameness of $X$.
321. (V5.47) Admitting sc-smooth partitions of unity: every open cover has a subordinate partition of unity of sc-smooth functions.
322. (V5.48) Sums, products, locally finite sums and chart-supported extensions by 0 of sc-smooth functions are sc-smooth.
323. (V5.49) Sc-smooth partitions of unity give sc-smooth bump functions.
324. (V5.50) $C^k$-bump functions and $C^k$ partitions of unity of a Banach space are as in the text.
325. (V5.51) The sc-smooth bump function property of a local model, with $f(x) \neq 0$; a retract is closed near each of its points.
326. (V5.52) Theorem 5.34: bump functions on $X$ iff a covering family of charts with the local bump property.
327. (V5.53) A local model on a Hilbert $E_0$ (or a norm dominated by a continuous quadratic form) has the bump property; M-polyfolds on sc-Hilbert spaces admit bump functions.
328. (V5.54) The sc-smooth approximation property of a local model, with $V = O \cap W$ and closures in $E_0$.
329. (V5.55) With sc-smooth partitions of unity, charts restricted to sets with closure in the chart domain have the approximation property.
330. (V5.56) A covering family of charts with the approximation property gives sc-smooth Urysohn functions and sc-smooth partitions of unity.
331. (V5.57) Theorem 5.37: sc-smooth partitions of unity iff a covering family of charts with the approximation property.
332. (V5.58) If $E_0$ admits $C^\infty$ partitions of unity, every local model on $E$ has the approximation property.
333. (V5.59) $c_0(\Gamma)$ and coordinate-wise $C^k$ embeddings are as in the text; Theorem 5.40 is only stated.
334. (V5.60) WCG: a weakly compact set with dense span; reflexive: the bidual map is onto.
335. (V5.61) Reflexive and separable Banach spaces are WCG.
336. (V5.62) Theorem 5.43 is only stated; Hilbert spaces have $C^\infty$-bump functions and are reflexive.
337. (V5.63) Corollary 5.44 holds with Theorem 5.43 for $E_0$ as an explicit hypothesis.