

2062

B.E. (Mechanical Engineering)

Fourth Semester

MEC-406: Numerical Analysis

Time allowed: 3 Hours

Max. Marks: 50

**NOTE:** Attempt five questions in all, including Question No. 1 which is compulsory and selecting two questions from each Section. Use of simple calculator is allowed. All questions carry equal marks.

X-X-X

1. (a) What is total numerical error? How do you avoid numerical error?
- (b) Define simple and multiple roots of a nonlinear equation mathematically.  
Which method do you find useful in multiple root and how?
- (c) Explain the difference between curve fitting and interpolation. What are the methods used in curve fitting and interpolation?
- (d) What are the disadvantages of the Simpson's  $3/8^{\text{th}}$  rule compared with the Simpson's  $1/3^{\text{rd}}$ ? When does the Simpson's  $1/3^{\text{rd}}$  rule gives exact rule?
- (e) Write the second order difference approximation for first and second order derivatives based on central differences.

### SECTION-A

2. (a) Given that  $x = 10.00 \pm 0.05$ ;  $y = 0.0356 \pm 0.0002$ ;  $z = 15300 \pm 100$ ;  $t = 62000 \pm 500$ . Find the maximum value of the absolute error in  
(i)  $x + y + z + t$ ; (ii)  $z^3$ .
- (b) Find the root of multiplicity 2 near 0.5 for the equation:  $x^3 - x^2 - x + 1 = 0$ .
3. (a) State the condition of convergence of Gauss-Seidel iterative method. Apply this method, to solve the system:  
$$x + 3y + 10z = 24, 2x + 17y + 4z = 25, 28x + 4y - z = 32.$$
- (b) Define norm of a matrix. List the different types of norms of a matrix. What is condition number of a matrix? Explain how the condition number is useful in determining whether the matrix is ill-conditioned. Compute it for  $A = \begin{bmatrix} 9 & 8 \\ 1 & 1 \end{bmatrix}$ .
4. (a) Fit a second degree parabola  $y = a + bx + cx^2$  to the data  $(x_i, y_i)$ :  
(1, 0.63), (3, 2.05), (4, 4.08), (6, 10.78).
- (b) Lagrange's formula can be used to express a rational function as a sum of partial fractions. Express  $f(x) = \frac{x^2 + x - 3}{x^3 - 2x^2 - x + 2}$  as a sum of partial fractions.

(2)

**SECTION-B**

5. (a) Use trapezoidal rule to evaluate  $\int_1^2 \int_1^2 \frac{dx dy}{x+y}$  taking four sub-intervals.

(b) Given the following values of  $f(x) = \ln x$ , find the approximate value of  $f^1(2.0)$  and  $f^{11}(2.0)$ :

x	2.0	2.2	2.6
f(x)	0.69315	0.78846	0.95551

6. (a) State the difference between single step and multistep methods in solving ODEs. Use Adams-Basforth formula to find  $y(0.4)$  for the equation  $\frac{dy}{dx} = \frac{1}{2}xy$  using the data:

x	0	0.1	0.2	0.3
y	1	1.0025	1.0101	1.00228

(b) Solve the BVP:  $y^{11} - 64y + 10 = 0$ ,  $y(0) = 0$ ,  $y(1) = 0$  by finite difference method.

7. Using Crank-Nicolson implicit scheme to solve:  $16 \frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$ ,  $0 < x < 1$ ,  $t > 0$ , given that  $u(x, 0) = 0$ ,  $u(0, t) = 0$ ,  $u(1, t) = 100t$ , compute  $u$  for one time step.

x-x-x