

THE FULL ADJOINT-GENERATED CASE OF GINIBRE'S CONDITION Q_3 FORMAL PROOF AND COMPUTATIONAL SUPPLEMENT FOR PART III

LESLIE P. POLZER

ABSTRACT. Let G be a compact connected Lie group with simple Lie algebra and let $\mathcal{Q}_G$ be the closed multiplicative convex cone generated by its real adjoint character. We reduce Ginibre's full condition Q_3 on $\mathcal{Q}_G$ to one explicit two-parameter hierarchy and prove that hierarchy for every compact connected G with simple Lie algebra. The proof treats every finite tuple length, every sign assignment, and every even number of minus factors; Ginibre's Proposition 1 then promotes the adjoint generator to $\mathcal{Q}_G$. The last finite classical boxes are closed by bounded Littlewood determinant identities and an exact prime-field/GMP certificate covering 17,862 residual inequalities. Here "full" refers to Ginibre's finite-tuple and sign-pattern quantifiers on the explicitly stated adjoint-generated cone; it does not mean the cone of all positive-definite functions. We also give an exact $\mathrm{SO}(3)$ counterexample showing that Ginibre's abelian cone of all real positive-definite functions cannot be carried over literally to nonabelian compact groups.

Remark 0.1 (Submission architecture and audit status). The unified manuscript `submission.pdf` contains its own abstract and proof map, followed by `paper.pdf` (Parts I–II) and the present formal proof and computational supplement for Part III. The separately rebuilt `full_q3_main.pdf` is a navigation aid, not a third proof component. The present file is a load-bearing theorem dependency included in the unified manuscript, while the optional `paper_full.pdf` derivation archive is not. In particular, the finite B/C correction-prefix contract and the proved half-stable bridge occur in `paper.pdf`; the exact higher-minus tails, finite ledgers, and replay semantics occur here. The concise dependency spine is recorded in `PUBLICATION_PROOF_SPINE.md`. Diagnostic and superseded routes are not printed in the formal supplement and are not inputs to that spine; the source-only development archive can be typeset separately by defining `\GinibreDevelopmentArchive`. Files under `referee_audits/` and the historical report in the source archive are author-controlled self-audits. They are reproducibility aids, not independent peer review. The theorem depends only on the numbered results and authenticated computations mapped in the code and data availability section.

7. THE EXACT ADJOINT-GENERATED TARGET

The cone studied here is the smallest uniformly closed cone containing the adjoint character and closed under multiplication and nonnegative linear combination. It is natural for two independent reasons. Representation theoretically, its polynomial part consists exactly of nonnegative linear combinations of tensor-power characters $1, \chi_{\mathrm{ad}}, \chi_{\mathrm{ad}}^2, \dots$; hence its Haar moments are invariant-tensor multiplicities in tensor powers of the adjoint module. From the viewpoint of Ginibre's Proposition 1, it is precisely the generated cone to which the single-generator proof promotes. A statement on any larger cone requires additional inequalities not supplied by that promotion. The theorem therefore isolates the full signed-product consequences of the semiring cone generated by the single adjoint character. This cone is smaller than the character cone of the tensor category generated by the adjoint module: an irreducible summand of an adjoint tensor power need not have character equal to a nonnegative polynomial in χ_{ad} . It does not assert that a general central

positive-definite function is a uniform limit of these polynomials, and it supplies no conclusion for a model whose interaction or observables do not belong to this cone.

Let G be as in the abstract, let μ_G be normalized Haar measure, and write $X = \chi_{\text{ad}}(g)$ and $Y = \chi_{\text{ad}}(h)$ for independent Haar variables. Since the adjoint representation is real, X and Y are real-valued.

Definition 7.1 (Adjoint-generated cone). The adjoint-generated multiplicative convex cone is

$$\mathcal{Q}_G = \overline{\left\{ \sum_{j=0}^d c_j \chi_{\text{ad}}^j : d \geq 0, c_j \geq 0 \right\}}^{\|\cdot\|_\infty} \subset C(G).$$

For integers $a, n \geq 0$, define

$$I_{a,n}^G = \int_{G \times G} (\chi_{\text{ad}}(g) - \chi_{\text{ad}}(h))^{2a} (\chi_{\text{ad}}(g) + \chi_{\text{ad}}(h))^n d\mu_G(g) d\mu_G(h).$$

Lemma 7.2 (Concrete nonpolynomial interactions in the cone). *Let $R = \max_G |\chi_{\text{ad}}|$. If*

$$A(z) = \sum_{j \geq 0} c_j z^j, \quad c_j \geq 0, \quad \sum_{j \geq 0} c_j R^j < \infty,$$

then $A(\chi_{\text{ad}}) \in \mathcal{Q}_G$. In particular, $e^{\beta \chi_{\text{ad}}} \in \mathcal{Q}_G$ for every $\beta \geq 0$. Thus the theorem applies not only to finite tensor-power characters but also to the standard exponential adjoint interaction and to finite products and nonnegative mixtures of such interactions.

Proof. The partial sums of $A(\chi_{\text{ad}})$ are nonnegative linear combinations of the powers of χ_{ad} . The displayed scalar bound and the Weierstrass test give uniform convergence on G , so the limit belongs to $\mathcal{Q}_G$ by definition. Taking $c_j = \beta^j/j!$ proves the exponential assertion. Closure under products and nonnegative linear combinations is built into the cone. $\square$

Lemma 7.3 (The adjoint-generated cone is positive definite). *Every element of $\mathcal{Q}_G$ is a real continuous positive-definite function on G . In particular, $\mathcal{Q}_G$ is a subcone of the cone $\mathcal{P}_G$ used in Proposition 7.6 below.*

Proof. Choose an orthonormal basis (e_j) of the real adjoint module. Then

$$\chi_{\text{ad}}(g) = \sum_j \langle e_j, \text{Ad}(g)e_j \rangle$$

is positive definite: for arbitrary $g_1, \dots, g_r \in G$ and $z_1, \dots, z_r \in \mathbb{C}$ its defining quadratic form is

$$\sum_j \left\| \sum_{k=1}^r z_k \text{Ad}(g_k) e_j \right\|^2 \geq 0.$$

Products of positive-definite functions are positive definite by tensoring their unitary representations, and nonnegative sums are positive definite by direct sums and scaling. Finally, a uniform limit preserves every finite positive-semidefinite Gram matrix entrywise. The character and all its powers are real and continuous, proving the assertion from the definition of $\mathcal{Q}_G$. $\square$

Theorem 7.4 (Full Q_3 reduction on the adjoint-generated cone). *Ginibre's condition Q_3 holds for $\mathcal{Q}_G$ if and only if*

$$I_{a,n}^G \geq 0 \quad (a, n \geq 0).$$

The only nonautomatic members of this hierarchy are

$$a \geq 1, \quad n \text{ odd}.$$

Theorem 7.5 below is exactly the row $a = 1$.

Proof. For the singleton generating set $S_G = \{\chi_{\text{ad}}\}$, a signed product with r minus factors and n plus factors is

$$\int_{G \times G} (X - Y)^r (X + Y)^n d\mu_G d\mu_G.$$

It vanishes when r is odd, by interchanging X and Y , and for $r = 2a$ it is $I_{a,n}^G$. Thus the displayed hierarchy is precisely condition Q_3 for the generating set S_G .

For completeness, the passage from the generator to the cone uses the factorization

$$\begin{aligned} (fg)(x) \pm (fg)(y) &= \frac{1}{2} [(f(x) + f(y))(g(x) \pm g(y)) \\ &\quad + (f(x) - f(y))(g(x) \mp g(y))]. \end{aligned}$$

Iterating this identity expands every signed factor belonging to a product of generators as a sum, with nonnegative coefficients, of products of signed generator factors. Convex combinations preserve the inequality, and uniform limits preserve it because Haar measure is finite. This is the argument of Ginibre [3, Proposition 1 and (1.5)]; it proves that S_G satisfies Q_3 if and only if $\mathcal{Q}_G$ does.

It remains to identify the automatic sectors. If n is even, the integrand defining $I_{a,n}^G$ is pointwise nonnegative. When $a = 0$, expand

$$\mathbf{E}(X + Y)^n = \sum_{j=0}^n \binom{n}{j} m_j^G m_{n-j}^G, \quad m_j^G = \dim((\mathfrak{g}_{\mathbb{C}}^{\otimes j})^G) \geq 0.$$

Hence the all-plus sector is nonnegative as well. $\square$

Theorem 7.5 (Two-minus adjoint-character theorem). *For every compact connected Lie group G with simple Lie algebra and every integer $n \geq 0$,*

$$I_{1,n}^G \geq 0.$$

Proof. This is the main theorem of the formal Parts I–II manuscript, [1, Theorem 2.2]. In the notation of that theorem its functional $Q_3^G(n)$ is exactly $I_{1,n}^G$ above. That manuscript is generated as `paper.pdf` from `paper.tex` and is submitted with the present Part III. Its correction-prefix certificate contract and local half-stable bridge are stated and proved in that manuscript; no result is imported from the optional derivation archive. $\square$

Proposition 7.6 (The all-positive-definite analogue is false). *Let $G = \text{SO}(3)$ with normalized Haar measure, and let $\mathcal{P}_G$ denote the cone of all real continuous positive-definite functions on G . Then $\mathcal{P}_G$ does not satisfy Ginibre’s condition Q_3 . More precisely, four functions in $\mathcal{P}_G$ and the sign pattern $--++$ give the value*

$$-\frac{8}{279}.$$

Consequently the adjoint-generated restriction in Theorem 7.4 cannot be removed, even for a compact connected group with simple Lie algebra.

Proof. For a unit vector $v \in \mathbb{R}^3$, put

$$f_v(g) = \langle v, gv \rangle, \quad g \in \text{SO}(3).$$

This function is real and continuous. It is positive definite because, for arbitrary $g_1, \dots, g_r \in G$ and $z_1, \dots, z_r \in \mathbb{C}$,

$$\sum_{i,j=1}^r \bar{z}_i z_j f_v(g_i^{-1} g_j) = \left\| \sum_{j=1}^r z_j g_j v \right\|^2 \geq 0.$$

We first record the Haar integrals needed below. For unit vectors $v_1, \dots, v_4$, define

$$c_1 = \langle v_1, v_2 \rangle \langle v_3, v_4 \rangle, \quad c_2 = \langle v_1, v_3 \rangle \langle v_2, v_4 \rangle, \quad c_3 = \langle v_1, v_4 \rangle \langle v_2, v_3 \rangle.$$

Haar projection on the standard representation gives

$$\int_G f_{v_i} = 0, \quad \int_G f_{v_i} f_{v_j} = \frac{\langle v_i, v_j \rangle^2}{3}.$$

For the fourth integral, the three pairing tensors in $(\mathbb{R}^3)^{\otimes 4}$ have Gram matrix with diagonal entries 9 and off-diagonal entries 3. Its inverse has diagonal entries $2/15$ and off-diagonal entries $-1/30$. Since Haar averaging is the orthogonal projection onto the invariant tensors, contraction with $v_1 \otimes \cdots \otimes v_4$ on both sides gives

$$\int_G \prod_{i=1}^4 f_{v_i} = \frac{2}{15} \sum_{i=1}^3 c_i^2 - \frac{1}{15} \sum_{1 \leq i < j \leq 3} c_i c_j. \quad (1)$$

Take

$$v_1 = \left(\sqrt{\frac{19}{31}}, \sqrt{\frac{12}{31}}, 0 \right), \quad v_2 = \left(\sqrt{\frac{19}{31}}, -\sqrt{\frac{12}{31}}, 0 \right), \quad v_3 = v_4 = (1, 0, 0).$$

Then $(c_1, c_2, c_3) = (7, 19, 19)/31$, and (1) equals $61/961$. Expanding the signed product and using the vanishing one-point integrals yields, with $M_{ij} = \int_G f_{v_i} f_{v_j} d\mu_G$,

$$\begin{aligned} \mathcal{J} &= \int_{G \times G} (f_{v_1}(g) - f_{v_1}(h)) \\ &\quad \cdot (f_{v_2}(g) - f_{v_2}(h)) \\ &\quad \cdot (f_{v_3}(g) + f_{v_3}(h)) \\ &\quad \cdot (f_{v_4}(g) + f_{v_4}(h)) \\ &= 2 \left[\int_G \prod_{i=1}^4 f_{v_i} d\mu_G + M_{12}M_{34} - M_{13}M_{24} - M_{14}M_{23} \right] \\ &= 2 \left[\frac{61}{961} + \frac{49 - 361 - 361}{9 \cdot 961} \right] = -\frac{8}{279}. \end{aligned}$$

This is a strict violation of Q_3 inside $\mathcal{P}_G$. □

Remark 7.7 (Comparison with Ginibre's abelian Example 4). Ginibre [3, Example 4] proves Q_3 for the cone of all real positive-definite functions when the compact group is a finite product of circles and finite cyclic groups. At the end of that paper he proposes the same cone on a noncommutative compact group only as a candidate. Proposition 7.6 shows that this proposed extension is false. The result proved here matches Ginibre's complete Q_3 quantifiers and his Proposition 1 cone-promotion mechanism, but on the explicitly smaller multiplicative cone generated by the adjoint character. The matrix coefficients in Proposition 7.6 are not central. Consequently neither that counterexample nor the affirmative theorem decides Q_3 for the intermediate cone of all real continuous central positive-definite functions on a nonabelian compact group.

Remark 7.8 (Relation to earlier nonabelian correlation inequalities). Sylvester [4, Secs. 1, 3–4] had already disproved the general Ginibre inequality for higher-dimensional $O(N)$ spin systems by constructing graph-cycle counterexamples. His variables are spins on a product of spheres and his signed factors are indexed by graph edges. The present Proposition 7.6 supplies a different, exact witness directly in the compact-group formulation proposed at the end of Ginibre's 1970 paper: four explicit real positive-definite functions on $SO(3)$, integrated over two normalized Haar variables, give $-8/279$. We therefore make no claim that this is the first nonabelian failure of a Ginibre inequality.

Abdessalam [5, Sec. 1 and Thms. 2.1–2.3] subsequently formulated general and padded Ginibre inequalities for invariant observables of $O(N)$ and $\mathbb{CP}^{N-1}$ models. He proves large-power asymptotics in terms of Kirchhoff polynomials and proves all padded Ginibre inequalities for inverse half-integer powers of stable determinantal polynomials. Those results concern an asymptotic regime

and a determinantal-polynomial substitute; they do not establish the finite-rank compact-group Haar hierarchy proved here. Conversely, the present theorem is restricted to the adjoint-generated cone and does not settle the invariant-observable GKS2 problem for finite $O(N)$ spin systems, which Abdesselam records as open for $N \geq 3$.

Remark 7.9 (Scope of potential quantum applications). Theorem 14.23 is an algebraic Haar-positivity theorem. Through Ginibre's general mechanism it can serve as an input for group-valued lattice models whose interactions genuinely lie in $\mathcal{Q}_G$. It does not by itself prove correlation inequalities for quantum Heisenberg or general $O(N)$ vector-spin systems, nonlinear sigma models, lattice gauge theories, or continuum quantum field theories. In particular, the occurrence of an adjoint representation in a gauge model neither places its plaquette interactions and observables in $\mathcal{Q}_G$ nor removes the shared-link constraints. This distinction is visible in the recent work of Chevyrev and Garban [6, Thm. 1.5, Cor. 1.6, and Rem. 1.7]: their Villain-action limit applies to nonabelian lattice gauge theories, while the Ginibre monotonicity consequence established there is abelian.

Such applications require additional model-specific results: Gibbs-measure promotion on the relevant observable cone, a thermodynamic-limit argument, and, for a quantum reconstruction, a reflection-positive transfer-matrix or equivalent construction. A direct route toward the conventional finite- N $O(N)$ problem would instead require the two-minus or padded inequality for defining-representation matrix coefficients on products of spheres. That is a separate theorem and is not asserted here.

Lemma 7.10 (Moment and generating-function forms). *For $a, n \geq 0$,*

$$I_{a,n}^G = \sum_{j=0}^{2a} \sum_{k=0}^n (-1)^j \binom{2a}{j} \binom{n}{k} m_{2a-j+k}^G m_{j+n-k}^G.$$

If

$$M_G(t) = \mathbf{E} e^{tX} = \sum_{r \geq 0} m_r^G \frac{t^r}{r!},$$

then, as an identity of convergent power series,

$$F_a^G(t) := \mathbf{E}[(X - Y)^{2a} e^{t(X+Y)}] = \sum_{j=0}^{2a} (-1)^j \binom{2a}{j} M_G^{(2a-j)}(t) M_G^{(j)}(t) = \sum_{n \geq 0} I_{a,n}^G \frac{t^n}{n!}.$$

Thus the full hierarchy is equivalent to coefficientwise nonnegativity of every F_a^G .

Proof. Expand $(X - Y)^{2a}(X + Y)^n$ twice by the binomial theorem and use the independence of X and Y . Expanding only $(X - Y)^{2a}$ after inserting $e^{tX} e^{tY}$ gives the derivative formula. Finally, expansion of $e^{t(X+Y)}$ gives the last equality. The adjoint character is bounded on the compact group, so all series converge absolutely and termwise integration is justified. $\square$

Lemma 7.11 (Central-cover invariance). *If $\pi : \tilde{G} \rightarrow G$ is a finite central covering of compact connected Lie groups, then*

$$I_{a,n}^{\tilde{G}} = I_{a,n}^G \quad (a, n \geq 0).$$

Thus every result below depends only on the simple Lie algebra and applies to all of its compact connected global forms.

Proof. The adjoint characters satisfy $\chi_{\text{ad}}^{\tilde{G}} = \chi_{\text{ad}}^G \circ \pi$, and the push-forward of normalized Haar measure under π is normalized Haar measure. Apply these facts in both variables of the defining integral. $\square$

Lemma 7.12 (Nonnegative cumulants give nonnegative moments). *Let*

$$K(t) = \sum_{r \geq 1} \kappa_r \frac{t^r}{r!} \quad \text{with} \quad \kappa_r \geq 0,$$

and write

$$e^{K(t)} = \sum_{n \geq 0} q_n \frac{t^n}{n!}.$$

Then $q_n \geq 0$ for every n .

Proof. The exponential formula gives

$$q_n = \sum_{\substack{k_1, \dots, k_n \geq 0 \\ k_1 + 2k_2 + \dots + nk_n = n}} n! \prod_{r=1}^n \frac{1}{k_r!} \left(\frac{\kappa_r}{r!} \right)^{k_r}.$$

Every summand is nonnegative. □

Lemma 7.13 (Adjoint-character normalization). *For G compact and connected with simple Lie algebra,*

$$\int_G \chi_{\text{ad}}(g) d\mu_G(g) = 0, \quad \int_G \chi_{\text{ad}}(g)^2 d\mu_G(g) = 1, \quad \max_G \chi_{\text{ad}} = \dim \mathfrak{g}.$$

Proof. The complexified adjoint representation is irreducible: a G -invariant complex subspace differentiates, by connectedness, to an $\text{ad}(\mathfrak{g}_{\mathbb{C}})$ -invariant subspace and hence is an ideal of the simple Lie algebra $\mathfrak{g}_{\mathbb{C}}$. It is nontrivial, so Schur orthogonality gives zero for its character mean. The compact real form has an invariant positive-definite inner product, making the adjoint representation self-dual and its character real. Schur orthogonality therefore also gives $\int_G \chi_{\text{ad}}^2 = \int_G |\chi_{\text{ad}}|^2 = 1$. Finally, an orthogonal operator on the d_G -dimensional real adjoint module has trace at most d_G , with equality at the identity. □

Lemma 7.14 (Even-moment lower bound for a positive cap). *Let X be real-valued with $-c \leq X \leq d$, let $0 < \varepsilon < d - c$, and put $t = d - \varepsilon$. If $t \geq c$, then for every integer $s \geq 1$,*

$$\mathbf{P}\{X > t\} \geq \frac{\mathbf{E}X^{2s} - t^{2s}}{d^{2s} - t^{2s}}.$$

If moreover $\mathbf{E}X^2 = 1$ and $\varepsilon > 1$, then

$$\mathbf{P}\{|X| < \varepsilon\} \geq 1 - \varepsilon^{-2}.$$

Proof. On the complement of $\{X > t\}$ one has $|X| \leq t$, while everywhere $|X| \leq d$. Thus, with $p = \mathbf{P}\{X > t\}$,

$$\mathbf{E}X^{2s} \leq pd^{2s} + (1-p)t^{2s}.$$

Rearrangement proves the first inequality. The second is Markov's inequality applied to X^2 : $\mathbf{P}\{|X| \geq \varepsilon\} \leq \varepsilon^{-2} \mathbf{E}X^2$. □

Theorem 7.15 (Fixed-group total-degree tail). *Let G be a compact connected Lie group with simple Lie algebra, put*

$$d_G = \dim \mathfrak{g}, \quad c_G = -\min_{g \in G} \chi_{\text{ad}}(g),$$

and choose

$$0 < \varepsilon < \frac{d_G - 2c_G}{2}.$$

Define the positive Haar masses

$$p_{G,\varepsilon} = \mu_G\{g : \chi_{\text{ad}}(g) > d_G - \varepsilon\} \mu_G\{g : |\chi_{\text{ad}}(g)| < \varepsilon\}$$

and put $q_{G,\varepsilon} = d_G - 2\varepsilon$. If $a \geq 1$, n is odd, and

$$2a + n > \frac{\log(1/p_{G,\varepsilon})}{\log(q_{G,\varepsilon}/(2c_G))},$$

then $I_{a,n}^G > 0$.

Proof. First we verify that all constants in the statement exist. Lemma 7.13 shows that the adjoint character has Haar mean zero and second moment one, so it takes both positive and negative values. Since G is connected, its continuous image $\chi_{\text{ad}}(G)$ is an interval; hence that interval contains zero and $c_G > 0$. Garibaldi–Guralnick–Rains [9, Theorem 2.1] prove $\chi_{\text{ad}}(g) \geq -\text{rank}(G)$. If $r = \text{rank}(G)$, then

$$d_G = r + |R| \geq 3r > 2r \geq 2c_G,$$

because a rank- r root system contains the positive and negative simple roots. Thus the interval prescribed for ε is nonempty and $q_{G,\varepsilon} > 2c_G$. Both sets defining $p_{G,\varepsilon}$ are nonempty open sets: the first contains the identity and the second contains a point at which χ_{ad} vanishes. Haar measure has full support on a compact group: finitely many left translates of any nonempty open set cover G , so such a set cannot have Haar measure zero. Hence $p_{G,\varepsilon} > 0$.

Write $S = X + Y$ and $D = X - Y$. On the negative half-plane $S < 0$, the bounds $X, Y \geq -c_G$ imply

$$|S| \leq 2c_G, \quad |D| \leq 2c_G.$$

Indeed, if $X \geq Y$, then $S < 0$ gives $X < -Y \leq c_G$, whence $0 \leq X - Y < 2c_G$; the other ordering is symmetric. Therefore the absolute value of the entire negative contribution is at most

$$(2c_G)^{2a+n}.$$

On the positive rectangle

$$X > d_G - \varepsilon, \quad |Y| < \varepsilon,$$

one has $D > q_{G,\varepsilon}$ and $S > q_{G,\varepsilon}$. Its contribution is strictly greater than

$$p_{G,\varepsilon} q_{G,\varepsilon}^{2a+n}.$$

The remainder of the half-plane $S \geq 0$ is nonnegative because n is odd and $2a$ is even. Consequently

$$I_{a,n}^G > (2c_G)^{2a+n} \left[p_{G,\varepsilon} \left(\frac{q_{G,\varepsilon}}{2c_G} \right)^{2a+n} - 1 \right],$$

and the assumed logarithmic inequality makes the bracket positive. $\square$

Corollary 7.16 (Each fixed group has only a finite residual box). *For a fixed compact connected G with simple Lie algebra, only finitely many pairs (a, n) with $a \geq 2$ and odd n remain after combining the two-minus Theorem 7.5 with Theorem 7.15.*

Proof. Theorem 7.5 supplies $a = 1$. The preceding theorem supplies every pair with total degree $2a + n$ above one fixed finite threshold. There are only finitely many nonnegative integer pairs below that threshold. $\square$

8. COMPLETE CLOSURE IN TYPE A_1

Lemma 8.1 (An effective $SU(2)$ Haar rectangle). *For the adjoint character of $SU(2)$ one may take*

$$c_G = 1, \quad d_G = 3, \quad \varepsilon = \frac{1}{8}, \quad q_{G,\varepsilon} = \frac{11}{4},$$

and the Haar mass in Theorem 7.15 satisfies

$$p_{G,1/8} > \frac{3789107}{38106288000} > \left(\frac{8}{11} \right)^{29}.$$

Consequently $I_{a,n}^{SU(2)} > 0$ whenever $a \geq 1$, n is odd, and $2a + n \geq 29$.

Proof. Parametrize conjugacy classes by $0 \leq \alpha \leq \pi$. The normalized class density and adjoint character are

$$d\mu(\alpha) = \frac{2}{\pi} \sin^2 \alpha d\alpha, \quad \chi_{\text{ad}}(\alpha) = 1 + 2 \cos(2\alpha) = 3 - 4 \sin^2 \alpha.$$

Thus $c_G = 1$ and $d_G = 3$. Put

$$A_0 = (0, 1/6) \cup (\pi - 1/6, \pi).$$

Since $\sin \alpha < \alpha$ on $(0, 1/6)$, every point of A_0 satisfies $\chi_{\text{ad}} > 3 - 1/8$. The elementary identity

$$\frac{22}{7} - \pi = \int_0^1 \frac{x^4(1-x)^4}{1+x^2} dx > 0$$

gives $1/\pi > 7/22$. Taylor's theorem gives $\sin x \geq x - x^3/6$ for $0 \leq x \leq 1/6$, and hence

$$\begin{aligned} \mu(A_0) &= \frac{4}{\pi} \int_0^{1/6} \sin^2 x dx \\ &> \frac{14}{11} \int_0^{1/6} (x - x^3/6)^2 dx = \frac{541301}{277136640}. \end{aligned}$$

Next put

$$B_0 = \{\alpha : |\alpha - \pi/3| < 1/32\} \cup \{\alpha : |\alpha - 2\pi/3| < 1/32\}.$$

The character vanishes at the two interval centers and $|\chi'_{\text{ad}}(\alpha)| \leq 4$, so $|\chi_{\text{ad}}| < 1/8$ on B_0 . For $0 \leq u \leq 1/32$,

$$\sin(\pi/3 - u) = \frac{\sqrt{3}}{2} \cos u - \frac{1}{2} \sin u > \frac{5}{6} \left(1 - \frac{1}{2048}\right) - \frac{1}{64} = \frac{10043}{12288} > \frac{4}{5}.$$

Here we used $\sqrt{3} > 5/3$, $\cos u \geq 1 - u^2/2$, and $\sin u \leq u$. Symmetry gives the same bound on the second interval. Their total length is $1/8$, so

$$\mu(B_0) > \frac{2}{\pi} \frac{1}{8} \frac{16}{25} > \frac{14}{275}.$$

It follows that

$$p_{G,1/8} \geq \mu(A_0)\mu(B_0) > \frac{3789107}{38106288000}.$$

The second displayed comparison in the statement follows from the exact integer calculation

$$3789107 \cdot 11^{29} - 38106288000 \cdot 8^{29} = 114033204110077056951613191119358337 > 0.$$

It is independently checked afresh by `character_ring_iter/verify_full_q3_su2_gmp.cpp`. Since $q_{G,1/8}/(2c_G) = 11/8$, the fixed-group tail theorem applies at every total degree at least 29. $\square$

Proposition 8.2 (Exact $SU(2)$ residual certificate). *For every $a \geq 2$ and every odd $n \geq 1$ with*

$$2a + n \leq 27,$$

one has $I_{a,n}^{\text{SU}(2)} > 0$. There are exactly 78 such pairs; their smallest value is

$$I_{2,1}^{\text{SU}(2)} = 16.$$

Proof. Let $c_k(j)$ be the multiplicity of the spin- j irreducible in the k th tensor power of the spin-one representation. The $SU(2)$ Clebsch–Gordan rule gives the exact integer recurrence

$$c_0(j) = [j = 0], \quad c_{k+1}(0) = c_k(1), \quad c_{k+1}(j) = c_k(j-1) + c_k(j) + c_k(j+1) \quad (j \geq 1).$$

The adjoint moment is $m_k = c_k(0)$. Substitution in Lemma 7.10 reduces every claimed inequality to a finite integer sum.

The proof companion is

$$\text{character_ring_iter/verify_full_q3_su2_gmp.cpp}.$$

It implements precisely this recurrence and the double sum in Lemma 7.10. It uses GMP integers throughout, checks every generated moment against the independent Catalan binomial transform

$$m_k = \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} \frac{1}{j+1} \binom{2j}{j},$$

which follows from $\chi_{\text{ad}} = |\text{tr } U|^2 - 1$, and checks the independently known prefix

$$(m_0, \dots, m_9) = (1, 0, 1, 1, 3, 6, 15, 36, 91, 232),$$

enumerates the complete quantified domain, asserts that its cardinality is 78, rejects any nonpositive value, and asserts the stated exact minimum. It also cross-multiplies the rational tail comparison in Lemma 8.1. The command `make full-q3-su2-audit` compiles the source with warnings promoted to errors and reruns every check. Its terminal success marker is `SU2_FULL_Q3 VERIFICATION: ALL PASS`. $\square$

Theorem 8.3 (Full adjoint-generated Q_3 in type A_1). *Let G be any compact connected group with Lie algebra of type A_1 . Then Ginibre's full condition Q_3 holds on $\mathcal{Q}_G$.*

Proof. By Lemma 7.11, it suffices to use $G = \text{SU}(2)$. The automatic sectors are covered by Theorem 7.4. The row $a = 1$ is Theorem 7.5. For $a \geq 2$ and odd n , the total degree $2a + n$ is odd. Proposition 8.2 covers $2a + n \leq 27$, while Lemma 8.1 covers $2a + n \geq 29$. Hence the complete generator hierarchy is nonnegative, and Theorem 7.4 promotes it to $\mathcal{Q}_G$. $\square$

9. COMPLETE CLOSURE IN TYPE A_2

Proposition 9.1 (Exact $\text{SU}(3)$ cap and residual certificate). *For the adjoint character of $\text{SU}(3)$,*

$$m_{54} = 1073719774632588649045995816896752254587680.$$

This moment gives

$$\mathbf{P}\{X > 6\} \geq \frac{m_{54} - 6^{54}}{8^{54} - 6^{54}}, \quad \mathbf{P}\{|X| < 2\} \geq \frac{3}{4},$$

and the exact comparison

$$\frac{3}{4} \frac{m_{54} - 6^{54}}{8^{54} - 6^{54}} \cdot 2^{29} > 1$$

holds. Moreover,

$$I_{a,n}^{\text{SU}(3)} > 0 \quad \text{for all } a \geq 2, \text{ } n \text{ odd, } 2a + n \leq 27.$$

There are exactly 78 inequalities in the last display, and their smallest value is $I_{2,1}^{\text{SU}(3)} = 72$.

Proof. For Haar $U \in \text{SU}(3)$, put $Z = |\text{tr } U|^2$. Scalar invariance identifies its moments with the corresponding $\text{U}(3)$ moments, and Schur–Weyl duality gives

$$E_j := \mathbf{E} Z^j = \sum_{\substack{\lambda \vdash j \\ \ell(\lambda) \leq 3}} (f^\lambda)^2, \quad m_k = \mathbf{E} (Z - 1)^k = \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} E_j,$$

where f^λ is computed by the hook-length formula.

The proof companion is

`character_ring_iter/verify_full_q3_su3_gmp.cpp`.

It enumerates every partition in these formulas and performs all arithmetic with GMP integers. It reconstructs $m_0, \dots, m_{54}$, checks the prefix

$$(m_0, \dots, m_9) = (1, 0, 1, 2, 8, 32, 145, 702, 3598, 19280),$$

and independently verifies every generated moment against the proved recurrence

$$(k+3)^2 m_{k+1} = k(7k+11)m_k + 8k(k+1)m_{k-1} \quad (k \geq 1).$$

It asserts the displayed value of m_{54} and cross-multiplies the tail comparison exactly.

For the residual box it substitutes the reconstructed moments into Lemma 7.10, enumerates the complete 78-pair domain, rejects any nonpositive result, and asserts the exact minimum. The command `make full-q3-su3-audit` rebuilds and reruns the verifier. Its terminal marker is `SU3_FULL_Q3 VERIFICATION: ALL PASS`.

It remains only to justify that the probability bounds consumed by the integer comparison are valid. The identity $\chi_{\text{ad}}(U) = |\text{tr } U|^2 - 1$ gives $-1 \leq X \leq 8$, and equality at -1 occurs at $\text{diag}(1, \omega, \omega^2)$ for a primitive cube root ω . Apply Lemma 7.14 with $(c, d, \varepsilon, s) = (1, 8, 2, 27)$. Its two conclusions are precisely the two displayed probability bounds. $\square$

Theorem 9.2 (Full adjoint-generated Q_3 in type A_2). *Let G be any compact connected group with Lie algebra of type A_2 . Then Ginibre's full condition Q_3 holds on $\mathcal{Q}_G$.*

Proof. Central-cover invariance reduces the proof to $G = \text{SU}(3)$. The automatic sectors and cone promotion are given by Theorem 7.4, and the $a = 1$ row is Theorem 7.5. Let $a \geq 2$ and let n be odd. The character bounds are $c_G = 1, d_G = 8$. Taking $\varepsilon = 2$ in Theorem 7.15 gives $q_{G,\varepsilon}/(2c_G) = 2$. The exact probability comparison in Proposition 9.1 therefore proves $I_{a,n} > 0$ for total degree $2a + n \geq 29$. The same proposition supplies every odd total degree at most 27. This exhausts the hierarchy and proves the theorem. $\square$

10. COMPLETE CLOSURE IN TYPES A_3 AND A_4

Proposition 10.1 (Exact $\text{SU}(4)$ and $\text{SU}(5)$ cap and residual certificates). *Put*

$$M_4 = m_{94}^{\text{SU}(4)}, \quad M_5 = m_{96}^{\text{SU}(5)}.$$

The following exact rational inequalities hold:

$$\frac{15}{16} \frac{M_4 - 11^{94}}{15^{94} - 11^{94}} \left(\frac{7}{2}\right)^{27} > 1,$$

and

$$\frac{285}{289} \frac{2^{96} M_5 - 31^{96}}{48^{96} - 31^{96}} \left(\frac{7}{2}\right)^{37} > 1.$$

In addition,

G	residual domain	number of pairs	minimum
$\text{SU}(4)$	$a \geq 2, n \text{ odd}, 2a + n \leq 25$	66	$I_{2,1} = 94,$
$\text{SU}(5)$	$a \geq 2, n \text{ odd}, 2a + n \leq 35$	136	$I_{2,1} = 96.$

Every residual value in the table is strictly positive.

Proof. For $N = 4, 5$, Schur–Weyl duality and the hook-length formula give the exact source

$$E_j^{(N)} = \sum_{\substack{\lambda \vdash j \\ \ell(\lambda) \leq N}} (f^\lambda)^2, \quad m_k^{(N)} = \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} E_j^{(N)}.$$

The proof companion is

`character_ring_iter/verify_full_q3_su45_gmp.cpp`.

It enumerates these partitions, evaluates every hook product and binomial transform with GMP integers, and reconstructs the moments through degrees 94 and 96. As independent checks, it verifies the prefixes

$$(m_0^{(4)}, \dots, m_9^{(4)}) = (1, 0, 1, 2, 9, 43, 245, 1557, 10829, 80958),$$

$$(m_0^{(5)}, \dots, m_9^{(5)}) = (1, 0, 1, 2, 9, 44, 264, 1824, 14210, 122224),$$

and every instance in range of the two proved low-rank recurrences from Parts I–II. It then substitutes the moments in Lemma 7.10, checks the complete residual domains and their exact minima, and cross-multiplies both displayed tail inequalities. No floating-point comparison is used. The command `make full-q3-su45-audit` rebuilds the verifier; its terminal marker is `SU45_FULL_Q3 VERIFICATION: ALL PASS`.

To connect the exact comparisons to Haar probabilities, use $\chi_{\text{ad}} = |\text{tr } U|^2 - 1$, so $c_G = 1$ and $d_G = N^2 - 1$. Trace-zero elements exist in both groups. For $SU(4)$ apply Lemma 7.14 with

$$(d, \varepsilon, s) = (15, 4, 47).$$

It gives cap mass at least $(M_4 - 11^{94})/(15^{94} - 11^{94})$ and central mass at least $15/16$; the tail ratio is $(d - 2\varepsilon)/(2c_G) = 7/2$. For $SU(5)$ use

$$(d, \varepsilon, s) = (24, 17/2, 48).$$

After multiplying numerator and denominator by 2^{96} , its cap bound is $(2^{96}M_5 - 31^{96})/(48^{96} - 31^{96})$; the central mass is at least $285/289$, and the tail ratio is again $7/2$. These are exactly the quantities checked by the verifier. $\square$

Theorem 10.2 (Full adjoint-generated Q_3 in types A_3 and A_4). *Let G be a compact connected group whose simple Lie algebra has type A_3 or A_4 . Then Ginibre’s full condition Q_3 holds on $\mathcal{Q}_G$.*

Proof. Central-cover invariance reduces the two cases to $SU(4)$ and $SU(5)$. The automatic sectors and cone promotion follow from Theorem 7.4; the $a = 1$ row is supplied by Theorem 7.5. For $SU(4)$, the first exact comparison of Proposition 10.1 and Theorem 7.15 supply every odd total degree at least 27, while the residual certificate supplies every odd total degree at most 25. The corresponding split for $SU(5)$ is $2a + n \geq 37$ and $2a + n \leq 35$. Thus no generator case remains in either group, and Theorem 7.4 completes the proof. $\square$

11. STABLE TYPE A

Theorem 11.1 (Full stable-rank type- A hierarchy). *Let $G = SU(N)$ and let $a \geq 1, n \geq 0$. If*

$$N \geq 2a + n,$$

then $I_{a,n}^G > 0$. In fact,

$$F_a^G(t) = (2a)! \frac{e^{-2t}}{(1-t)^{2a+2}} \quad \text{through degree } n.$$

Proof. For every $r \leq N$, Schur–Weyl duality gives

$$\mathbf{E}_{SU(N)} |\text{tr } U|^{2r} = r!.$$

Since $\chi_{\text{ad}}(U) = |\text{tr } U|^2 - 1$, all adjoint moments through degree N agree with those of $A = E - 1$, where E is exponential with mean one. The integral $I_{a,n}$ uses no moment above degree $2a + n$, so the rank hypothesis permits replacement of X, Y by independent copies of A .

Let E_1, E_2 be independent mean-one exponentials, put $R = E_1 + E_2$ and $T = E_1/R$. Then R and T are independent, R has the gamma density $re^{-r} dr$, and T is uniform on $[0, 1]$. Moreover,

$$X - Y = R(2T - 1), \quad X + Y = R - 2.$$

Consequently

$$\begin{aligned} F_a^G(t) &= \mathbf{E}(2T - 1)^{2a} \mathbf{E}[R^{2a} e^{t(R-2)}] \\ &= \frac{1}{2a+1} e^{-2t} \frac{(2a+1)!}{(1-t)^{2a+2}} = (2a)! \frac{e^{-2t}}{(1-t)^{2a+2}}. \end{aligned}$$

The logarithm of the last factor after removal of the positive constant is

$$-2t - (2a+2) \log(1-t).$$

Its first cumulant is $2a$, and its r th cumulant for $r \geq 2$ is $(2a+2)(r-1)!$. They are all positive, so Lemma 7.12 proves strict positivity of every coefficient. $\square$

Lemma 11.2 (Paley–Zygmund cap bound). *If $W \geq 0$ has finite second moment and $0 \leq t < \mathbf{E}W$, then*

$$\mathbf{P}\{W > t\} \geq \frac{(\mathbf{E}W - t)^2}{\mathbf{E}W^2}.$$

Proof. Write $A = \{W > t\}$. Then

$$\mathbf{E}W \leq t + \mathbf{E}(W \mathbf{1}_A) \leq t + (\mathbf{E}W^2)^{1/2} \mathbf{P}(A)^{1/2}$$

by Cauchy–Schwarz. Rearrangement and squaring prove the claim. $\square$

Lemma 11.3 (Polynomial one-sided tail bound). *Let X be real-valued, let $T \in \mathbb{R}$, and let P be a real polynomial for which*

$$A = \mathbf{E}[(X - T)P(X)^2] > 0, \quad B = \mathbf{E}[(X - T)^2 P(X)^4] > 0.$$

Then

$$\mathbf{P}\{X > T\} \geq \frac{A^2}{B}.$$

Proof. Put

$$R(x) = 1 - \frac{A}{B}(x - T)P(x)^2.$$

If $x \leq T$, then $R(x) \geq 1$, so $\mathbf{1}_{\{x \leq T\}} \leq R(x)^2$. Therefore

$$\begin{aligned} \mathbf{P}\{X \leq T\} &\leq \mathbf{E}R(X)^2 \\ &= 1 - 2\frac{A}{B}A + \frac{A^2}{B^2}B = 1 - \frac{A^2}{B}. \end{aligned}$$

Taking complements proves the assertion. $\square$

Lemma 11.4 (One-sided cap propagation for every even-minus sector). *Let X, Y be independent copies of a real random variable satisfying $X \geq -C$ and $\mathbf{E}X^2 = 1$, and let T, δ, p_0 satisfy*

$$\delta > 1, \quad \mathbf{P}\{X > T\} \geq p_0 > 0, \quad q := T - \delta > 2C.$$

If K is odd and

$$p_0(1 - \delta^{-2})q^K > (2C)^K, \tag{2}$$

then

$$\mathbf{E}[(X - Y)^{2a}(X + Y)^n] > 0$$

for every $a \geq 1$, every odd n , and every $2a + n \geq K$.

Proof. Chebyshev's inequality and $\mathbf{E}Y^2 = 1$ give $\mathbf{P}\{|Y| < \delta\} \geq 1 - \delta^{-2}$. On the product of this event with $\{X > T\}$, both $X - Y$ and $X + Y$ exceed q . On the negative half-plane $X + Y < 0$, the lower support bound gives $|X - Y|, |X + Y| \leq 2C$, as in the proof of Theorem 7.15. The rest of the positive half-plane contributes nonnegatively. Hence, for $L = 2a + n$,

$$\mathbf{E}[(X - Y)^{2a}(X + Y)^n] > p_0(1 - \delta^{-2})q^L - (2C)^L.$$

The right side is positive at $L = K$ by (2); after division by $(2C)^L$ it is strictly increasing in L because $q/(2C) > 1$. $\square$

Proposition 11.5 (Uniform higher-minus tail for $SU(N)$, $N \geq 6$). *Let $a \geq 1$ and let n be odd. Put $L = 2a + n$ and define*

$$K_N = \begin{cases} 27, & N = 6, \\ 29, & N = 7, 8, \\ 31, & N \geq 9. \end{cases}$$

If $N \geq 6$ and $L \geq K_N$, then

$$I_{a,n}^{\text{SU}(N)} > 0.$$

Proof. Put $Z = |\text{tr } U|^2$ and $X = Z - 1$. For

$$E_r^{(N)} = \mathbf{E} Z^r = \sum_{\substack{\lambda \vdash r \\ \ell(\lambda) \leq N}} (f^\lambda)^2,$$

apply Lemma 11.2 to $W = Z^{16}$ and $t = (33/5)^{16}$. The exact inequalities verified below include $E_{16}^{(N)} > t$ and give

$$\mathbf{P}\{X > 28/5\} = \mathbf{P}\{Z > 33/5\} \geq \frac{(E_{16}^{(N)} - (33/5)^{16})^2}{E_{32}^{(N)}}.$$

Since $\mathbf{E} X^2 = 1$, Lemma 7.14 also gives

$$\mathbf{P}\{|X| < 11/10\} \geq \frac{21}{121}.$$

On the product of these two events, both $X - Y$ and $X + Y$ exceed

$$\frac{28}{5} - \frac{11}{10} = \frac{9}{2}.$$

On the negative half-plane $X + Y < 0$, the identity $X = Z - 1 \geq -1$ gives $|X - Y|, |X + Y| \leq 2$. Exactly as in Theorem 7.15, it therefore suffices that

$$\frac{21}{121} \frac{(E_{16}^{(N)} - (33/5)^{16})^2}{E_{32}^{(N)}} \left(\frac{9}{4}\right)^L > 1.$$

The exact GMP verifier

`character_ring_iter/verify_full_q3_sun_ge6_gmp.cpp`

clears denominators and checks this inequality at $L = K_N$ for every $6 \leq N \leq 31$. For $N \geq 32$, Schur–Weyl stability gives $E_{16}^{(N)} = 16!$ and $E_{32}^{(N)} = 32!$, so the single checked stable-row comparison at $N = 32$ applies to every larger N . The left side increases with L , because $9/4 > 1$, completing the tail proof. $\square$

Proposition 11.6 (Exact residual type- A certificate for $N \geq 6$). *For every $6 \leq N \leq 28$ and every pair $a \geq 2$, n odd, satisfying*

$$N < 2a + n < K_N,$$

one has $I_{a,n}^{\text{SU}(N)} > 0$. The quantified domain contains 1315 pairs. Its smallest value is

$$I_{2,3}^{\text{SU}(6)} = 3550.$$

Proof. The proof companion is

`character_ring_iter/verify_full_q3_sun_ge6_gmp.cpp`.

For each $6 \leq N \leq 28$, it reconstructs $E_0^{(N)}, \dots, E_{29}^{(N)}$ by enumerating all partitions of bounded length and evaluating the hook-length formula, then forms

$$m_k^{(N)} = \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} E_j^{(N)}.$$

It checks $E_j^{(N)} = j!$ independently whenever $j \leq N$, substitutes the moments in Lemma 7.10, enforces the exact stable/tail complement defining the residual domain, checks the per-rank and total scope counts, rejects every nonpositive value, and asserts the displayed minimum.

The same source reconstructs $E_{16}^{(N)}$ and $E_{32}^{(N)}$ for $6 \leq N \leq 31$, uses $16!, 32!$ at the stable endpoint, and checks the tail comparison of Proposition 11.5 after the exact integer clearing

$$21(5^{16}E_{16}^{(N)} - 33^{16})^2 9^{K_N} > 121 5^{32} E_{32}^{(N)} 4^{K_N}.$$

The command `make full-q3-sun-ge6-audit` rebuilds and reruns the complete calculation. Its terminal marker is `SUN_GE6_FULL_Q3 VERIFICATION: ALL PASS.` $\square$

Theorem 11.7 (Full adjoint-generated Q_3 in every type A). *Let G be a compact connected group whose simple Lie algebra has type A_r for some $r \geq 1$. Then Ginibre's full condition Q_3 holds on $\mathcal{Q}_G$.*

Proof. Types A_1 – A_4 are Theorems 8.3, 9.2, and 10.2. Let $G = SU(N)$ with $N \geq 6$; central-cover invariance handles every other global form. The automatic sectors follow from Theorem 7.4, and Theorem 7.5 supplies $a = 1$.

It remains to take $a \geq 2$, n odd, and $L = 2a + n$. If $L \leq N$, Theorem 11.1 applies. If $L \geq K_N$, Proposition 11.5 applies. A pair in neither range can occur only for $6 \leq N \leq 28$, and every such pair is supplied by Proposition 11.6. Indeed, for $N = 29$ there is no odd integer strictly between 29 and 31, and for $N \geq 30$ the stable and tail ranges overlap. Thus the generator hierarchy is complete for every N . Theorem 7.4 promotes it to $\mathcal{Q}_G$. $\square$

12. COMPLETE CLOSURE FOR THE EXCEPTIONAL GROUPS

For an exceptional group G , write $m_j = m_j^G$ and set

$$A_G = m_{2k+1} - T m_{2k}, \quad B_G = m_{4k+2} - 2T m_{4k+1} + T^2 m_{4k}.$$

The following table fixes all parameters used below. The column M is the largest moment consumed by the tail calculation, and K is an odd total degree.

G	c	T	δ	k	M	K
G_2	2	91/10	6/5	9	38	25
F_4	4	211/10	13/10	12	50	37
E_6	3	98/5	7/5	10	42	29
E_7	7	333/10	13/10	17	70	63
E_8	8	461/10	13/10	23	94	71

TABLE 1. Parameters for the exceptional higher-minus tail certificates.

Proposition 12.1 (Exact exceptional higher-minus tails). *Take the row of Table 1 corresponding to G . If $a \geq 1$, n is odd, and $2a + n \geq K$, then*

$$I_{a,n}^G > 0.$$

Proof. The bounds $\chi_{\text{ad}} \geq -c$ used in the five rows are rigorous: for G_2, F_4, E_7, E_8 they follow from the rank bound of Garibaldi–Guralnick–Rains [9, Theorem 2.1], and for E_6 the sharper value $\min \chi_{\text{ad}} = -3$ is [9, Theorem 2.3 and Table 1].

Apply Lemma 11.3 with $P(x) = x^k$. It gives

$$\mathbf{P}\{X > T\} \geq \frac{A_G^2}{B_G}.$$

By Lemma 7.13 and Markov’s inequality,

$$\mathbf{P}\{|Y| < \delta\} \geq 1 - \delta^{-2}.$$

On the product of these two events,

$$X - Y > T - \delta, \quad X + Y > T - \delta.$$

On $X + Y < 0$, the common lower bound $X, Y \geq -c$ instead gives $|X - Y|, |X + Y| \leq 2c$, exactly as in Theorem 7.15. Thus, with $L = 2a + n$, it suffices that

$$\rho_G R_G^L > 1, \quad \rho_G = \frac{A_G^2}{B_G}(1 - \delta^{-2}), \quad R_G = \frac{T - \delta}{2c}. \quad (3)$$

The exact verifier

`character_ring_iter/verify_full_q3_exceptional_gmp.cpp`

checks $A_G, B_G > 0$, $R_G > 1$, and $\rho_G R_G^K > 1$ by GMP rational arithmetic in every row. Hence (3) holds for every $L \geq K$. $\square$

Proposition 12.2 (Exact exceptional residual certificate). *For each row of Table 1, one has*

$$I_{a,n}^G > 0 \quad \text{whenever} \quad a \geq 2, \quad n \text{ is odd}, \quad 2a + n < K.$$

The exact scopes and minima are

G	largest residual degree	number of pairs	$\min I_{a,n}^G$
G_2	23	55	36
F_4	35	136	36
E_6	27	78	38
E_7	61	435	36
E_8	69	561	36

and every displayed minimum occurs at $(a, n) = (2, 1)$.

Proof. The proof companion is

`character_ring_iter/verify_full_q3_exceptional_gmp.cpp`.

It reads the exact invariant-tensor moments audited in Parts I–II. For every exceptional type, the complete theorem-consumed range is regenerated from the Cartan datum by exact Racah–Speiser iteration and the character-pairing identities: through degrees 38, 65, 80, 70, 100 for G_2, F_4, E_6, E_7, E_8 , respectively. Only after that independent regeneration, a separate control compares every row termwise with the corresponding ancillary tensor-rank block of Bourjaily–Plesser–Vergu [10, ancillary file `adjoint_tensor_ranks.tex`]. For E_8 , the control also compares the maintained transcription of degrees 71–100 directly with the primary ancillary file. Thus no ancillary entry is a premise of the residual certificate.

For every pair in the stated boxes, the verifier evaluates the integer formula of Lemma 7.10, rejects a nonpositive value, checks the five scope counts, and asserts the five minima and their coordinates. Independently, it forms A_G, B_G, ρ_G, R_G as reduced GMP rationals, checks the tail inequality at K , and checks that the same chosen certificate does not yet pass at $K - 2$. The command `make`

full-q3-exceptional-audit rebuilds and reruns the arithmetic audit; its terminal marker is EXCEPTIONAL_FULL_Q3 VERIFICATION: ALL PASS. $\square$

Theorem 12.3 (Full adjoint-generated Q_3 for every exceptional type). *Let G be a compact connected group whose simple Lie algebra has type G_2, F_4, E_6, E_7 , or E_8 . Then Ginibre's full condition Q_3 holds on $\mathcal{Q}_G$.*

Proof. By central-cover invariance, it suffices to use the compact form employed in the moment computation. The automatic sectors follow from Theorem 7.4, and Theorem 7.5 supplies $a = 1$. For $a \geq 2$ and odd n , Proposition 12.1 applies when $2a + n \geq K$, while Proposition 12.2 applies below K . Thus the entire generator hierarchy holds, and Theorem 7.4 promotes it to $\mathcal{Q}_G$. $\square$

13. STABLE CLASSICAL TYPES

Theorem 13.1 (Full stable classical hierarchy). *Let $a \geq 1$ and $n \geq 0$, and put $L = 2a + n$. The inequality $I_{a,n}^G \geq 0$ holds in each of the following ranges:*

G	stable-rank condition
$B_\ell = \mathrm{SO}(2\ell + 1)$	$L \leq 2\ell + 1,$
$C_\ell = \mathrm{Sp}(2\ell)$	$L \leq 2\ell + 1,$
$D_\ell = \mathrm{SO}(2\ell)$	$L \leq \ell - 1.$

More precisely, throughout these ranges

$$F_a^G(t) = e^{-t+t^2/2} \sum_{j=0}^a \frac{\gamma_{a,j}}{(1-t)^{2j+1}} \quad \text{through degree } n,$$

where

$$\gamma_{a,j} = \binom{2a}{2j} (2a - 2j - 1)!! \frac{\binom{2j}{j}}{4^j} (2j)! > 0.$$

Here $(-1)!! = 1$.

Proof. The exact stable joint trace laws of Diaconis–Shahshahani [2, Theorems 4 and 6] identify the stable orthogonal and symplectic adjoint variables with the common law

$$X = \frac{Z_1^2 - 1 + \sqrt{2}Z_2}{2},$$

where Z_1, Z_2 are independent standard normal variables. The sign of Z_2 is immaterial.

We first check that the stated ranks supply every moment used by $I_{a,n}$. For type B , the orthogonal Schur integral criterion [11, Section 3, paragraph preceding Lemma 3.1] applied to $s_{(1,1)}^s$ is stable when $s \leq 2\ell + 1$. For type C , the symplectic criterion in the same paragraph is stable when $\ell \geq \lfloor s/2 \rfloor$, equivalently $s \leq 2\ell + 1$. In type D , the trace degree of χ_{ad}^s is $2s$; the exact orthogonal trace law and the equality of the $\mathrm{SO}(2\ell)$ and $\mathrm{O}(2\ell)$ integrals apply when $2s < 2\ell$, namely $s \leq \ell - 1$. Since Lemma 7.10 uses only moments of total degree at most L , the displayed hypotheses give the stable law throughout the required calculation.

Take independent copies

$$X = \frac{Z_1^2 - 1 + \sqrt{2}Z_2}{2}, \quad Y = \frac{W_1^2 - 1 + \sqrt{2}W_2}{2}.$$

In polar coordinates for (Z_1, W_1) , put

$$R = \frac{Z_1^2 + W_1^2}{2}, \quad C = \frac{Z_1^2 - W_1^2}{Z_1^2 + W_1^2}.$$

Then R is mean-one exponential, $C = \cos(2\Theta)$ with Θ uniform, and R, C are independent. Also

$$A = \frac{Z_2 + W_2}{\sqrt{2}}, \quad B = \frac{Z_2 - W_2}{\sqrt{2}}$$

are independent standard normals, independent of (R, C) , and

$$X + Y = R - 1 + A, \quad X - Y = RC + B.$$

Only even powers survive in the conditional expansion of $(RC + B)^{2a}$. Using

$$\mathbf{E}B^{2q} = (2q - 1)!!, \quad \mathbf{E}C^{2j} = \frac{\binom{2j}{j}}{4^j}, \quad \mathbf{E}[R^{2j}e^{tR}] = \frac{(2j)!}{(1 - t)^{2j+1}},$$

and $\mathbf{E}e^{tA} = e^{t^2/2}$ gives the asserted formula for F_a^G .

It remains to prove coefficientwise positivity. For each $j \geq 0$, put

$$H_j(t) = \frac{e^{-t+t^2/2}}{(1 - t)^{2j+1}}.$$

Its logarithm has cumulants

$$\kappa_1 = 2j, \quad \kappa_2 = 2j + 2, \quad \kappa_r = (2j + 1)(r - 1)! \quad (r \geq 3).$$

They are nonnegative, so every coefficient of H_j is nonnegative by Lemma 7.12. The coefficients $\gamma_{a,j}$ are strictly positive; hence their finite linear combination has nonnegative coefficients as claimed. $\square$

14. A SIMULTANEOUS NONSTABLE CLASSICAL TAIL

For the rest of the classical argument let $X = \chi_{\text{ad}}(U)$ and $Y = \chi_{\text{ad}}(V)$ for independent Haar elements, and put

$$S = X + Y, \quad D = X - Y.$$

For $G = B_b, C_b, D_b$, use the defining trace T_1 and the signed square trace

$$\tau_G = \begin{cases} \text{Tr}(U^2), & G = B_b, D_b, \\ -\text{Tr}(U^2), & G = C_b. \end{cases}$$

Thus in all three families

$$X = \frac{T_1^2 - \tau_G}{2}. \tag{4}$$

Lemma 14.1 (Exact classical negative endpoints). *For the standard representatives,*

$$C_{B_b} = b, \quad C_{C_b} = b, \quad C_{D_b} = \begin{cases} b, & b \text{ even}, \\ b - 2, & b \text{ odd}. \end{cases}$$

Proof. Write the defining eigenvalue pairs as z_j, z_j^{-1} , put $x_j = (z_j + z_j^{-1})/2 \in [-1, 1]$, and set $R = \sum_j x_j$ and $Q = \sum_j x_j^2$. Direct substitution in (4) gives

$$\chi_{\text{ad}}^{B_b} = b + 2R + 2(R^2 - Q), \quad \chi_{\text{ad}}^{D_b} = b + 2(R^2 - Q), \quad \chi_{\text{ad}}^{C_b} = 2R^2 + 2Q - b.$$

The first two expressions are affine in each variable separately, so their minima occur at vertices of the cube. At a vertex $Q = b$ and $R = b - 2k$. The type- B value is $2R^2 + 2R - b$, whose minimum is $-b$; the type- D value is $2R^2 - b$, whose minimum is $-b$ for even b and $2 - b$ for odd b . The type- C expression is at least $-b$, with equality at $x_1 = \dots = x_b = 0$. $\square$

Lemma 14.2 (Root-normalized classical Weyl cap). *Let $G = B_b$ or D_b , put $d = \dim G$, and use the positive roots*

G	κ in $Q(\theta) = \sum_{\alpha>0} \alpha(\theta)^2 = \kappa \theta ^2$	$ W $	$(d_1, \dots, d_b)$
B_b	$2b - 1$	$2^b b!$	$(2, 4, \dots, 2b)$
D_b	$2b - 2$	$2^{b-1} b!$	$(2, 4, \dots, 2b - 2, b)$

Set $z_{B_b} = 2^{-b}$ and $z_{D_b} = 1$. Suppose that $Q(\theta) \leq R$ lies in the standard eigenangle chart and that throughout this ball

$$\prod_{\alpha>0} \left(\frac{\sin(\alpha(\theta)/2)}{\alpha(\theta)/2} \right)^2 \geq \eta > 0.$$

Then, for the adjoint character X ,

$$\mathbf{P}\{X > d - R\} \geq \frac{z_G \eta}{|W|} \frac{\prod_{i=1}^b d_i!}{(2\pi)^{b/2} \Gamma(d/2 + 1)} \left(\frac{R}{2\kappa} \right)^{d/2}. \quad (5)$$

If $0 < R < 24$, one may take

$$\eta = \left(1 - \frac{R}{24} \right)^2. \quad (6)$$

For D_b , one may instead take, whenever $R < 12\kappa$,

$$\eta = \left(1 - \frac{R}{12\kappa} \right)^\kappa. \quad (7)$$

Proof. The character estimate $2(1 - \cos u) \leq u^2$ gives $d - X \leq Q(\theta)$. If $Q(\theta) = R > 0$, at least one root coordinate is nonzero and $2(1 - \cos u) < u^2$ for that coordinate; if $Q(\theta) < R$, the conclusion is immediate. Hence $X > d - R$ throughout the cap. Weyl integration and the stated sine-product bound therefore reduce the cap mass to

$$\frac{z_G \eta}{|W|(2\pi)^b} \int_{Q(\theta) \leq R} \prod_{\alpha>0} \alpha(\theta)^2 d\theta.$$

The factor $z_{B_b} = 2^{-b}$ occurs because the squared-length-two Macdonald–Mehta normalization replaces each short root e_i by $\sqrt{2}e_i$; all type- D roots already have squared length two. The Macdonald–Mehta identity [8, Theorem 3.1(i)], followed by radial integration of the homogeneous discriminant, gives

$$\int_{Q(\theta) \leq R} \prod_{\alpha>0} \alpha(\theta)^2 d\theta = (2\pi)^{b/2} \frac{\prod_i d_i!}{\Gamma(d/2 + 1)} \left(\frac{R}{2\kappa} \right)^{d/2},$$

which proves (5).

For $u = \alpha(\theta)$, the elementary bound $\sin(u/2)/(u/2) \geq 1 - u^2/24$ applies. If $R < 24$, all factors on the right are positive, and

$$\prod_{\alpha>0} (1 - u_\alpha)^2 \geq \left(1 - \sum_{\alpha>0} u_\alpha \right)^2, \quad u_\alpha = \frac{\alpha(\theta)^2}{24},$$

proves (6). In type D , every root has squared length two, so $u_\alpha \leq R/(12\kappa)$. Concavity of $\log(1 - u)$ on the interval from zero to this upper bound, together with $\sum_\alpha u_\alpha = Q(\theta)/24 \leq R/24$, gives (7). $\square$

Lemma 14.3 (All-even-minus trace-tail reduction). *Let $C_G = -\min_G \chi_{\text{ad}}$, and choose $c > 2C_G$, $d > 1$, and $q \in \mathbb{R}$. Define*

$$A = \{|T_1| \geq \sqrt{2c + 2d + q}, \tau_G \leq q\}, \quad B = \{|X| < d\}, \quad p = \mathbf{P}(A)\mathbf{P}(B).$$

If $a \geq 1$, n is odd, and $L = 2a + n$, then

$$I_{a,n}^G \geq 2pc^L - 2\mathbf{E}(\tau_G)_+^L. \quad (8)$$

More generally, if $L_0 \leq L$, then

$$I_{a,n}^G \geq 2pc^L - 2(2C_G)^{L-L_0} \mathbf{E}(\tau_G)_+^{L_0}. \quad (9)$$

Consequently, if L_0 is odd and

$$pc^{L_0} > \mathbf{E}(\tau_G)_+^{L_0}, \quad (10)$$

then $I_{a,n}^G > 0$ for every $a \geq 1$, odd n , and $2a + n \geq L_0$.

Proof. On A , equation (4) gives $X \geq c + d$. Hence on the rectangle $A \times B$ both S and D are at least c . On the transposed rectangle $B \times A$, one has $S \geq c$ and $D \leq -c$. These rectangles are disjoint because $X \geq c + d > d$ on A , and each has probability p . Their combined contribution is therefore at least $2pc^L$.

The rest of the half-plane $S \geq 0$ contributes nonnegatively. On $S < 0$, split into $X \leq Y$ and $Y < X$. The contributions have equal absolute value. On the first piece $X < 0$ and

$$|D| = Y - X \leq -2X, \quad |S| = -X - Y \leq -2X.$$

Equation (4) also gives $-2X \leq \tau_G$, so $|D|, |S| \leq (\tau_G)_+$. This proves (8) after doubling. Independently, $X, Y \geq -C_G$ implies $|D|, |S| \leq 2C_G$ on $S < 0$. Retaining L_0 of the L factors under the first bound and the other $L - L_0$ factors under the second proves (9).

Finally, if (10) holds, then $c > 2C_G$ makes the right side of (9) strictly positive for every $L \geq L_0$. Every relevant total degree is odd, which proves the last assertion. $\square$

Lemma 14.4 (Exponential bound for the square-trace moment). *For every integer $L \geq 1$ and every $v > 0$,*

$$\mathbf{E}(\tau_G)_+^L \leq \left(\frac{L}{ev}\right)^L \mathbf{E}e^{v\tau_G}.$$

For the classical square traces and $C_G \geq 296$, one has in particular

$$\mathbf{E}(\tau_G)_+^L \leq 4e^{20} \left(\frac{L}{4e}\right)^L. \quad (11)$$

Proof. For $z \geq 0$, the maximum of $z^L e^{-vz}$ occurs at $z = L/v$ and equals $(L/(ev))^L$. Multiplication by e^{vz} and expectation prove the first display.

For B_b , the power-two identity of Rains [12, Theorem 1.5, equation (23)] gives

$$\tau_G \stackrel{d}{=} 1 + 2 \operatorname{Re} \operatorname{Tr} W, \quad W \sim \operatorname{Haar}(\operatorname{U}(b)).$$

The strong Szegő upper bound of Courteaut–Johansson–Lambert [13, Lemma 2.12] therefore gives $\mathbf{E}e^{v\tau_G} \leq e^{v+v^2}$.

For C_b, D_b , the corresponding power-two decomposition [12, Theorem 1.5, equation (22), and equation (36)] writes $\tau_G - 1$ as a sum, up to harmless sign changes, of two independent centered full orthogonal traces. The Fredholm estimate obtained from [13, Proposition 3.3 and equations (3.4)–(3.6)] is

$$\mathbf{E}e^{v\tau_G} \leq \frac{e^{v+v^2}}{(1 - \mathbf{b}_{d_0}(v))(1 - \mathbf{b}_{d_1}(v))}, \quad \mathbf{b}_d(v) = \frac{e^{v^2/(d+1)} v^d}{1 - v^2/(d+1)^2 d!},$$

provided the companion trace-norm envelope is below one. When $C_G \geq 296$, both component dimensions are at least 11. At $v = 4$, the directed verifier cited below checks

$$\mathbf{a}_d(v) = \frac{e^{v^2/(d+1)} v^d}{d!(1 - v/(d+1))\sqrt{1 - v^2/(d+1)^2}}, \quad \mathbf{a}_{11}(4) < 1, \quad \mathbf{b}_{11}(4) < \frac{1}{2}.$$

Both quantities decrease with $d \geq 11$: after replacing d by $d+1$, the factorial contributes $4/(d+1)$, while the exponential and each remaining rational factor also decrease. Thus each denominator is at least $1/2$. In every family $\mathbf{E}e^{4\tau_G} \leq 4e^{20}$, and substitution in the first display proves (11). $\square$

Lemma 14.5 (Source-audited high-rank trace rectangle). *Put*

$$r = \frac{2000001}{1000000}, \quad \eta = \frac{359}{2000}, \quad A_0 = \frac{4571}{2000}.$$

For $G = B_b, C_b, D_b$ with $C = C_G \geq 296$, set

$$c = rC, \quad d = q = \eta C.$$

Then

$$\mathbf{P}\{|T_1| \geq \sqrt{2c + 3d}, \tau_G \leq d\} \geq \frac{1}{2}e^{-A_0C}, \quad \mathbf{P}\{|X| < d\} \geq \frac{3}{4}.$$

Proof. The second inequality is Chebyshev's inequality, since $\mathbf{E}X^2 = 1$ and $d > 2$.

The first is the Rains-square trace certificate of Parts I–II, whose inputs are recalled here. The one-trace density estimate [13, Theorem 1.4], the Mills lower bound, and the checked orthogonal parity shifts give

$$\mathbf{P}\{|T_1| \geq \sqrt{(2r + 3\eta)C}\} \geq e^{-A_0C}.$$

Rains' power-two decomposition and the same density estimate give

$$\mathbf{P}\{\tau_G \geq \eta C\} \leq \exp\left(-\frac{(\eta C - 1)^2}{4}\right) + \frac{10(\log(2\lfloor C/2 \rfloor))^{1/4}}{\sqrt{\Gamma(2\lfloor C/2 \rfloor + 1)}} \leq \frac{1}{2}e^{-A_0C}.$$

The last two displayed comparisons, including the unitary projection in type B , are checked with directed rounding for every $296 \leq C \leq 10000$ and with explicit derivative signs thereafter by `character_ring_iter/trace_square_cutoff_mpfr.cpp`. Subtracting the square-trace exceptional event proves the result. $\square$

Proposition 14.6 (Full higher-minus tail for large classical rank). *Let $G = B_b, C_b, D_b$ and suppose $C_G \geq 296$. Then*

$$I_{a,n}^G > 0 \quad (a \geq 1, n \text{ odd})$$

outside the stable range of Theorem 13.1. Hence the complete generator hierarchy holds in every such row.

Proof. Let L_0 be the first odd total degree after the stable range. The exact negative endpoints give

$$L_0 = \begin{cases} 2b + 3 = 2C_G + 3, & G = B_b, C_b, \\ b + 1 = C_G + 1, & G = D_b, b \text{ even}, \\ b = C_G + 2, & G = D_b, b \text{ odd}. \end{cases}$$

In particular $C_G + 1 \leq L_0 \leq 2C_G + 3$.

Use the parameters of Lemma 14.5. In the notation of Lemma 14.3, its two probability estimates give

$$p \geq \frac{3}{8}e^{-A_0C}, \quad C = C_G.$$

Lemma 14.4 shows that the ratio of the two sides in (10) is at least

$$\frac{3}{32}e^{-A_0C-20} \left(\frac{4e r C}{L_0}\right)^{L_0}.$$

Since the base is greater than one and $C + 1 \leq L_0 \leq 2C + 3$, the logarithm of this ratio is bounded below by

$$H(C) = \log \frac{3}{32} - A_0C - 20 + (C + 1) \log \left(\frac{4e r C}{2C + 3}\right). \quad (12)$$

The directed-rounding companion

$$\text{character_ring_iter/verify_full_q3_bcd_high_mpfr.cpp}$$

checks $H(296) > 8$, $H'(296) > 1/10$, and, more importantly for the monotonicity argument,

$$\log \left(\frac{4er \cdot 296}{2 \cdot 296 + 3} \right) - A_0 > \frac{9}{100}.$$

The factor $4erC/(2C + 3)$ is increasing, and the remaining term in

$$H'(C) = -A_0 + \log \left(\frac{4erC}{2C + 3} \right) + (C + 1) \left(\frac{1}{C} - \frac{2}{2C + 3} \right)$$

is positive. Consequently the first two terms in $H'(C)$ have sum greater than $9/100$ for every $C \geq 296$, while the last term is positive. Hence $H'(C) > 0$ throughout that range. Thus (10) holds at L_0 , and Lemma 14.3 proves every later odd total degree. Theorem 13.1 supplies the prefix.

The same verifier checks the two Fredholm values consumed in Lemma 14.4. The command `make full-q3-bcd-high-audit` rebuilds both directed-rounding programs and requires their terminal zero-failure markers. The source-matched machine-C transcript `certificates/full_q3/fullq3bcdanalytic0003_machine_c.log` authenticates both sources and records zero compile and run statuses; its adjacent SHA-256 file authenticates the transcript. $\square$

Lemma 14.7 (Finite-rank Fredholm MGF bounds). *Let*

$$d_T = \begin{cases} 2b + 1, & G = B_b, \\ 2b, & G = C_b, D_b, \end{cases} \quad \mathbf{b}_d(v) = \frac{e^{v^2/(d+1)} v^d}{1 - v^2/(d+1)^2 d!},$$

and put

$$\mathbf{a}_d(v) = \frac{e^{v^2/(d+1)} v^d}{d!(1 - v/(d+1))\sqrt{1 - v^2/(d+1)^2}}, \quad D_d(v) = 1 - \mathbf{b}_d(v).$$

If $0 < v < d_T + 1$ and $\mathbf{a}_{d_T}(v) < 1$, then

$$e^{v^2/2} D_{d_T}(v) \leq \mathbf{E} e^{\pm v T_1} \leq \frac{e^{v^2/2}}{D_{d_T}(v)}. \quad (13)$$

For C_b put

$$d_0 = 2 \left\lceil \frac{b+1}{2} \right\rceil, \quad d_1 = 2 \left\lceil \frac{b}{2} \right\rceil + 1,$$

and for D_b put

$$d_0 = 2 \left\lceil \frac{b}{2} \right\rceil, \quad d_1 = 2 \left\lceil \frac{b-1}{2} \right\rceil + 1.$$

Whenever the corresponding two trace-norm conditions hold,

$$\mathbf{E} e^{v\tau_G} \leq \begin{cases} e^{v+v^2}, & G = B_b, \\ \frac{e^{v+v^2}}{D_{d_0}(v)D_{d_1}(v)}, & G = C_b, D_b. \end{cases} \quad (14)$$

Proof. We spell out the Fredholm step because every finite-middle trace estimate uses it. Write (a_J, b_J) for the two Jacobi parameters in Courteaut–Johansson–Lambert, to distinguish them from the rank b used here. Their Proposition 3.3 gives, for every complex ξ ,

$$F_n^{a_J b_J}(\xi) = e^{i(b_J - a_J)\xi - \xi^2/2} \det(I + Q_n K_{i\xi}^{a_J b_J} Q_n). \quad (15)$$

The group-to-kernel conversion, including all deterministic eigenvalues, is as follows. The column D is the first Bessel order after applying Q_n .

G	(a_J, b_J)	n	K	D	fixed trace
$B_b = \text{SO}(2b+1)$	$(1/2, -1/2)$	b	K^{+-}	$2b+1$	$+1$
$C_b = \text{Sp}(2b)$	$(1/2, 1/2)$	b	K^{++}	$2b+2$	0
$D_b = \text{SO}(2b)$	$(-1/2, -1/2)$	b	K^{--}	$2b$	0

Indeed, Proposition 3.3 of the cited source defines

$$K_z^{+-} = (-J_{r+s+1}(-2z))_{r,s \geq 0}, \quad K_z^{++} = (-J_{r+s+2}(-2z))_{r,s \geq 0}, \quad K_z^{--} = (J_{r+s}(-2z))_{r,s \geq 0}.$$

Deleting the first n rows and columns changes the orders to $2n+1+r+s$, $2n+2+r+s$, and $2n+r+s$, respectively, proving the table. Substitution of $\xi = \mp iv$ in (15) writes the random-trace MGF as a Gaussian exponential times $\det(I + QK_{\pm v}Q)$.

For reference, the fourth Jacobi kernel, needed for the orthogonal power-map components below, has the same conversion. After applying the projection, the Bessel order in entry (r, s) , with $r, s \geq 0$, is

kernel	d	Bessel order
K^{--}	$2n$	$2n+r+s$
K^{-+}, K^{+-}	$2n+1$	$2n+1+r+s$
K^{++}	$2n+2$	$2n+2+r+s$.

Here d is the initial Bessel order. It agrees with the full orthogonal component dimension; for the K^{++} representation of $\mathrm{Sp}(2n)$, it is two larger than the defining dimension. Entry signs do not affect the following absolute-value estimates.

Apply the Bessel estimate used in the proof of [13, Lemma 3.6, equations (2.16) and (3.4)], followed by

$$(d+r+s)! \geq d!(d+1)^{r+s}, \quad e^{v^2/(d+r+s+1)} \leq e^{v^2/(d+1)}.$$

Writing $x = v/(d+1) < 1$, the squared row sum is a geometric series in x^{2s} , and the sum of its square roots is a geometric series in x^r . Hence, in each of the four cases,

$$\|QK_{\pm v}Q\|_1 \leq \frac{e^{v^2/(d+1)}v^d}{d!(1-x)\sqrt{1-x^2}} = \mathfrak{a}_d(v).$$

This is the explicit estimate obtained before, and independently of, the source's simpler sufficient restriction $|z| < 2e^{-5/4}n$. The use of $d_T = 2b$, rather than the exact order $2b+2$, in the symplectic row is conservative. For every integer $d \geq 0$ and $0 < v < d+1$, direct division gives

$$\frac{\mathfrak{a}_{d+1}(v)}{\mathfrak{a}_d(v)} = \frac{v}{d+1} e^{-v^2/((d+1)(d+2))} \frac{1-v/(d+1)}{1-v/(d+2)} \sqrt{\frac{1-v^2/(d+1)^2}{1-v^2/(d+2)^2}} < 1$$

and

$$\frac{\mathfrak{b}_{d+1}(v)}{\mathfrak{b}_d(v)} = \frac{v}{d+1} e^{-v^2/((d+1)(d+2))} \frac{1-v^2/(d+1)^2}{1-v^2/(d+2)^2} < 1.$$

Every factor is positive, and each displayed rational ratio is less than one. Hence both envelopes decrease with d , so the order- $2b+2$ symplectic kernel is bounded by the displayed order- $2b$ envelopes. Thus the hypothesis $\mathfrak{a}_{d_T}(v) < 1$ justifies the absolutely convergent Plemelj series

$$\log \det(I + QK_{\pm v}Q) = \sum_{j \geq 1} \frac{(-1)^{j+1}}{j} \mathrm{tr}(QK_{\pm v}Q)^j.$$

For a trace of the j -th power, the cyclic index sum uses the same factorial and Bessel bounds. Each cyclic index occurs twice, so the two geometric sums contribute $(1-x^2)^{-1}$. The calculation of [13, Lemma 3.7, equation (3.6)] therefore gives, for all four rows,

$$\left| \mathrm{tr}(QK_{\pm v}Q)^j \right| \leq \mathfrak{b}_d(v)^j, \quad \sum_{j \geq 1} \frac{1}{j} \left| \mathrm{tr}(QK_{\pm v}Q)^j \right| \leq -\log(1 - \mathfrak{b}_d(v)).$$

Notice that $0 \leq \mathfrak{b}_d(v) \leq \mathfrak{a}_d(v) < 1$: after writing $x = v/(d+1)$, the ratio $\mathfrak{a}_d/\mathfrak{b}_d$ is $\sqrt{(1+x)/(1-x)}$. Exponentiating the last absolute bound therefore gives

$$D_{d_T}(v) \leq \det(I + QK_{\pm v}Q) \leq D_{d_T}(v)^{-1}.$$

The determinant is positive because the Gaussian prefactor is positive and the left side of the Fredholm identity is an MGF. This proves the determinant sandwich directly from the cited estimates.

It remains to account for the deterministic eigenvalues. The source formula is written for the random eigenvalues only. In type B , the table has $b_J - a_J = -1$, so the random-trace factors at $\xi = \mp iv$ are $e^{\mp v}$. Restoring the fixed $+1$ eigenvalue multiplies the two MGFs by $e^{\pm v}$, cancelling those factors exactly. In types C, D one has $b_J - a_J = 0$ and no deterministic trace. Thus the full defining character is centered in every family, in agreement with character orthogonality, and the Gaussian prefactor is $e^{v^2/2}$. This proves (13) for both signs.

For completeness, we also derive the deterministic shifts in the square trace. Specializing Rains' Theorem 1.5, equations (22)–(23), to $p = 2$, and using his equation (36), gives independent Haar blocks with

$$\begin{aligned}\tau_{B_b} &\stackrel{d}{=} 1 + 2 \operatorname{Re} \operatorname{Tr} W, & W &\sim \operatorname{Haar}(\operatorname{U}(b)), \\ \tau_{C_b} - 1 &\stackrel{d}{=} -(\operatorname{Tr} O_{d_0}^- + \operatorname{Tr} O_{d_1}^+), \\ \tau_{D_b} - 1 &\stackrel{d}{=} \operatorname{Tr} O_{d_0}^+ + \operatorname{Tr} O_{d_1}^-.\end{aligned}$$

Here the superscript specifies the determinant component, and the dimensions d_0, d_1 are exactly those in the statement. The constants can also be read directly from the fixed eigenvalues. The $O^-(2b+2)$ model for $\operatorname{Sp}(2b)$ has fixed eigenvalues $+1, -1$; after squaring they contribute 2, whereas the odd $O^+(d_1)$ block contributes one fixed $+1$. Since $\tau_{C_b} = -\operatorname{Tr}(U^2)$, this yields the second line. In type D , the odd $O^-(d_1)$ block contributes a fixed -1 , yielding the third line. These are precisely the shifts encoded by Rains' convention of ignoring fixed ± 1 eigenvalues.

Apply the preceding upper determinant bound to each orthogonal component and multiply, using independence; the leading constant 1 contributes e^v . This gives the C, D clause of (14). The type- B clause is the unitary strong Szegő bound already proved in Lemma 14.4. $\square$

Lemma 14.8 (Exact defining-trace determinant formulas). *For $s \geq 0$, write $I_k = I_k(2s)$ for the modified Bessel function. If T_1 is the full defining trace, then*

$$\mathbf{E}_{B_b} e^{sT_1} = e^s \det[I_{|j-k|} - I_{j+k+1}]_{j,k=0}^{b-1}, \quad (16)$$

$$\mathbf{E}_{B_b} e^{-sT_1} = e^{-s} \det[I_{|j-k|} + I_{j+k+1}]_{j,k=0}^{b-1}, \quad (17)$$

$$\mathbf{E}_{C_b} e^{sT_1} = \det[I_{|j-k|} - I_{j+k+2}]_{j,k=0}^{b-1}, \quad (18)$$

$$\mathbf{E}_{D_b} e^{sT_1} = \frac{1}{2} \det[I_{|j-k|} + I_{j+k}]_{j,k=0}^{b-1}. \quad (19)$$

The type- C and type- D trace laws are symmetric.

Proof. Courteaut–Johansson–Lambert [13, Lemma 3.1] give the four Toeplitz–Hankel determinants for the Jacobi parameters in their equation (3.1). Substitution of $\psi(x) = e^{2sx}$ gives Fourier coefficients $I_k(2s)$. Their E_n^{+-} case is $\operatorname{SO}(2b+1)$ after restoring the fixed $+1$ eigenvalue, giving (16); replacing s by $-s$ gives (17). Their E_n^{++} case simultaneously describes $\operatorname{Sp}(2b)$ and the random part of $O^-(2b+2)$, giving (18). Their E_n^{--} case is $\operatorname{SO}(2b)$; its determinant equals 2 at $s = 0$, which accounts for the factor $1/2$ in (19). Finally, $-I$ belongs to the type- C and even-orthogonal groups, proving symmetry. $\square$

Lemma 14.9 (Finite layered recovery from an exact MGF). *Let Z be bounded, let $M(s) = \mathbf{E}e^{sZ}$, and fix $s > 0$ and a lower endpoint t . Choose $\lambda_- > 0$, increasing upper endpoints $u_0 < \dots < u_r$, and positive $\lambda_0, \dots, \lambda_r$. Put*

$$L_- = e^{\lambda_- t} \frac{M(s - \lambda_-)}{M(s)}, \quad U_i = e^{-\lambda_i u_i} \frac{M(s + \lambda_i)}{M(s)}, \quad F_i = 1 - L_- - U_i, \quad w_i = e^{-s u_i}.$$

If all F_i are positive, then

$$\mathbf{P}\{Z > t\} \geq M(s) \left[w_r F_r + \sum_{i=0}^{r-1} (w_i - w_{i+1}) F_i \right]. \quad (20)$$

An endpoint at or above the upper support of Z may be used with $U_i = 0$.

Proof. Under the tilted probability $d\mathbf{P}_s = e^{sZ} M(s)^{-1} d\mathbf{P}$, exponential Markov gives

$$\mathbf{P}_s\{Z \leq t\} \leq L_-, \quad \mathbf{P}_s\{Z \geq u_i\} \leq U_i.$$

Thus $\mathbf{P}_s\{t < Z < u_i\} \geq F_i$. On the other hand, summation by parts over the layers

$$(t, u_0), [u_0, u_1), \dots, [u_{r-1}, u_r)$$

gives

$$\mathbf{E}_s[e^{-sZ} \mathbf{1}_{\{t < Z < u_r\}}] \geq w_r \mathbf{P}_s\{t < Z < u_r\} + \sum_{i=0}^{r-1} (w_i - w_{i+1}) \mathbf{P}_s\{t < Z < u_i\}.$$

Insert the preceding lower bounds and multiply by $M(s)$. □

Lemma 14.10 (One-step tilted defining-trace lower bound). *Let $t, h > 0$ and put*

$$s = t + h, \quad \rho(t, h) = 1 - \frac{e^{-h^2/2}}{D(s)} \left(\frac{1}{D(s-h)} + \frac{1}{D(s+h)} \right),$$

where $D(v) = D_{d_T}(v)$. Assume $s - h > 0$, the sandwich (13) at $s - h, s, s + h$, and $\rho(t, h) > 0$. Then

$$\mathbf{P}\{|T_1| \geq t\} \geq 2D(s)\rho(t, h) \exp \left\{ \frac{s^2}{2} - s(t + 2h) \right\}. \quad (21)$$

Proof. For $\sigma \in \{+1, -1\}$, tilt probability by $M_\sigma(s)^{-1} e^{\sigma s T_1}$. Exponential Markov at parameter h and the two sides of (13) show that the tilted probability of leaving the interval $t < \sigma T_1 < t + 2h$ is at most $1 - \rho(t, h)$. The choice $s = t + h$ centers the Gaussian comparison at the midpoint of this interval; both exit losses are exactly $e^{-h^2/2}$. On the interval the inverse tilt is at least $e^{-s(t+2h)}$. The lower MGF bound at s therefore gives

$$\mathbf{P}\{\sigma T_1 \geq t\} \geq D(s)\rho(t, h) e^{s^2/2 - s(t+2h)}.$$

The positive and negative tails are disjoint. Summing the two bounds proves the display; no symmetry of the trace law is being assumed. □

Proposition 14.11 (Full higher-minus tail for the finite middle ranks). *The complete generator hierarchy holds in the following classical rows:*

$$B_b \ (22 \leq b \leq 295), \quad C_b \ (29 \leq b \leq 295), \quad D_b \ (71 \leq b \leq 297).$$

Proof. Let L_0 be the first odd degree after the stable range, as in the proof of Proposition 14.6. In every row put

$$r = \frac{2000001}{1000000}, \quad c = rC_G, \quad d = \frac{3}{2}, \quad t = \sqrt{2c + 2d + q},$$

and use the family-dependent parameters

G	rank range	q	h
B_b	$22 \leq b \leq 295$	$\frac{21}{5}\sqrt{b}$	$\frac{13}{10}$
C_b	$29 \leq b \leq 295$	$4\sqrt{b}$	$\frac{13}{10}$
D_b	$71 \leq b \leq 297$	$\frac{18}{5}\sqrt{b}$	$\frac{6}{5}$.

Let V_G denote the right side of (21). At

$$v_q = \frac{q-1}{2},$$

exponential Markov and (14) give an upper bound U_G for $\mathbf{P}\{\tau_G \geq q\}$. Hence the event A in Lemma 14.3 has probability at least $V_G - U_G$. Chebyshev gives $\mathbf{P}(B) \geq 5/9$. Finally, at

$$v_m = \frac{\sqrt{1+8L_0}-1}{4},$$

Lemmas 14.4 and 14.7 bound $\mathbf{E}(\tau_G)_+^{L_0}$.

The directed-rounding companion

`character_ring_iter/verify_full_q3_bcd_mid_mpfr.cpp`

evaluates precisely these formulas. It checks every trace-norm domain, the positive ρ value for both one-sided tilts, the strict difference $V_G - U_G$, and

$$\frac{5}{9}(V_G - U_G)c^{L_0} > \mathbf{E}(\tau_G)_+^{L_0}$$

in all 768 rows. Its minimum directed logarithmic margin is

$$1.0047268114794421860\dots > 0 \quad \text{at } D_{71};$$

the minimum log separation between V_G and U_G is $0.16429898256055109\dots$ at C_{29} , and the largest trace-norm envelope is $0.39958077942628398\dots < 1$, also at C_{29} . Thus (10) holds at L_0 in every row. Lemma 14.3 supplies the entire tail and Theorem 13.1 supplies the prefix. $\square$

Proposition 14.12 (Additional finite classical tails). *For the rows $B_{18}, \dots, B_{21}$, let the odd tail onsets be*

$$(K_{18}, K_{19}, K_{20}, K_{21}) = (47, 45, 45, 45).$$

For $D_{31}, \dots, D_{70}$, in increasing rank order, let the onsets be

$$(49, 51, 51, 51, 51, 53, 53, 55, 53, 55, 55, 57, 55, 57, 57, 59, 59, 61, 61, 61, \\ 61, 63, 61, 63, 63, 65, 63, 65, 65, 67, 67, 69, 67, 69, 69, 71, 71, 71, 71, 73).$$

Then $I_{a,n}^G > 0$ whenever $a \geq 1$, n is odd, and $2a + n \geq K_b$ in the corresponding row.

Proof. Use $c = (2000001/1000000)C_G$ in every row. For B_{18} take

$$d = \frac{13}{10}, \quad q = \frac{6}{5}b, \quad h = \frac{5}{4};$$

for B_{19}, B_{20}, B_{21} take

$$d = \frac{3}{2}, \quad q = b, \quad h = \frac{6}{5}.$$

For the type- D rows take $d = 3/2$, $h = 6/5$, and

$$q = \begin{cases} 7b/10, & 31 \leq b \leq 38, \\ 3b/5, & 39 \leq b \leq 52, \\ b/2, & 53 \leq b \leq 70. \end{cases}$$

The probability and moment bounds are exactly those in the proof of Proposition 14.11, now evaluated at the displayed onset rather than at the first post-stable degree. The same directed-rounding verifier checks all 44 rows. Its least logarithmic tail margin is

$$0.07529217823349327949\dots > 0 \quad \text{at } D_{43}, K_{43} = 55;$$

its least tilted mass parameter is $0.02269453416254314484\dots > 0$ at B_{19} . Its least logarithmic separation between the defining-trace lower bound and square-trace upper bound, attained at D_{40} , is

$$0.83270005200917997982\dots > 0.$$

Consequently (10) holds at every displayed onset. Lemma 14.3 propagates each inequality to all later odd total degrees. $\square$

Lemma 14.13 (Exact finite classical adjoint moments). *Write*

$$e_2^j = \sum_{\lambda \vdash 2j} c_j(\lambda) s_\lambda, \quad c_j(\lambda) \in \mathbb{Z}_{\geq 0},$$

and let s_j be the stable classical adjoint moment. Then

$$m_j^{B_b} = s_j - \sum_{\substack{\nu \vdash j \\ \ell(\nu) > 2b+1}} c_j(2\nu).$$

For type C one has

$$m_j^{C_b} = s_j - \sum_{\substack{\nu \vdash j \\ \ell(\nu) > b}} c_j((\nu_1, \nu_1, \nu_2, \nu_2, \dots)').$$

For type D one has, without a restriction on j ,

$$m_j^{D_b} = s_j - \sum_{\substack{\nu \vdash j \\ \ell(\nu) > 2b}} c_j(2\nu) + \sum_{\substack{\nu \vdash j-b \\ \ell(\nu) \leq 2b}} c_j(\lambda(b, \nu)),$$

where, after padding ν by zeros to length $2b$,

$$\lambda(b, \nu) = (1 + 2\nu_1, \dots, 1 + 2\nu_{2b}).$$

An empty partition sum is zero.

Proof. For both orthogonal families the adjoint representation is $\bigwedge^2 V$, so the displayed Schur expansion is the character decomposition of $(\bigwedge^2 V)^{\otimes j}$; iterated vertical-two-strip Pieri proves $c_j(\lambda) \in \mathbb{Z}_{\geq 0}$. The orthogonal Schur integral criterion in Rains [11, Section 3, paragraph preceding Lemma 3.1] says that the $O(N)$ integral of s_λ is one precisely for even λ of length at most N , and is zero otherwise. For odd $N = 2b + 1$, the central element $-I$ identifies the $O(N)$ and $SO(N)$ integrals of every power of $\bigwedge^2 V$. Subtracting the even partitions excluded by the length wall gives the type- B formula.

For type C , the adjoint representation is $\text{Sym}^2 V$. The standard involution ω of the symmetric-function ring sends h_2 to e_2 and s_λ to $s_{\lambda'}$. Thus the coefficient of s_λ in h_2^j is $c_j(\lambda')$. Rains' symplectic criterion in the same paragraph says that the integral of s_λ over $\text{Sp}(2b)$ is one precisely when $\ell(\lambda) \leq 2b$ and every row length of λ occurs an even number of times, and is zero otherwise. Such a partition is uniquely $\lambda = (\nu_1, \nu_1, \nu_2, \nu_2, \dots)$; its length constraint is $\ell(\nu) \leq b$. Subtraction from the unrestricted stable sum proves the type- C formula.

For $SO(2b)$, averaging a polynomial character over the identity component equals its $O(2b)$ average plus its determinant-twisted $O(2b)$ average. The first term admits the even partitions of length at most $2b$. The second admits precisely the partitions of length $2b$ whose rows are all odd: subtracting one full column leaves an even partition, to which Rains' orthogonal criterion applies. Every such determinant-class partition is uniquely $\lambda(b, \nu)$, and its size equation is $2b + 2|\nu| = 2j$. Hence $|\nu| = j - b$, proving the type- D formula. $\square$

Lemma 14.14 (Bounded Littlewood determinants for the finite classical moments). *Work in the completed ring of symmetric functions over $\mathbb{Q}$, put $e_r = h_r = 0$ for $r < 0$, and define*

$$f_r^e = \sum_{k \in \mathbb{Z}} e_k e_{k+r}, \quad f_r^h = \sum_{k \in \mathbb{Z}} h_k h_{k+r}, \quad e = \sum_{k \geq 0} e_k, \quad \bar{e} = \sum_{k \geq 0} (-1)^k e_k.$$

For $b \geq 1$, let

$$\begin{aligned} \mathcal{B}_b &= \frac{1}{2} \left\{ e \det_{1 \leq i, j \leq b} (f_{i-j}^e - f_{i+j-1}^e) + \bar{e} \det_{1 \leq i, j \leq b} (f_{i-j}^e + f_{i+j-1}^e) \right\}, \\ \mathcal{C}_b &= \det_{1 \leq i, j \leq b} (f_{i-j}^h - f_{i+j}^h), \\ \mathcal{D}_b &= \frac{1}{2} \det_{1 \leq i, j \leq b} (f_{i-j}^e + f_{i+j-2}^e). \end{aligned}$$

Then, for every $j \geq 0$,

$$m_j^{B_b} = [m_{(2^j)}] \mathcal{B}_b, \quad m_j^{C_b} = [m_{(2^j)}] \mathcal{C}_b, \quad m_j^{D_b} = [m_{(2^j)}] \mathcal{D}_b.$$

Here $[m_{(2^j)}]$ denotes the coefficient of the monomial symmetric function $m_{(2^j)}$, and every displayed completed series is interpreted degree by degree.

Proof. Let $c(\lambda)$ and $r(\lambda)$ denote the numbers of odd columns and odd rows of λ . Hall duality gives

$$\langle h_2^j, F \rangle = [m_{(2^j)}] F.$$

Transpose the admitted partitions in the type- B formula of Lemma 14.13. Since the involution ω interchanges e_2 and h_2 , this gives

$$m_j^{B_b} = \left\langle h_2^j, \sum_{\substack{\lambda_1 \leq 2b+1 \\ c(\lambda)=0}} s_\lambda \right\rangle.$$

Huh–Kim–Krattenthaler–Okada [7, Theorem 1.3, equations (1.9)–(1.10)], specialized at $u = 0$, give respectively the sum of the $c(\lambda) = 0$ and $c(\lambda) = 2b + 1$ sectors and their difference. Taking their half-sum gives exactly $\mathcal{B}_b$.

For type C , transposition gives

$$m_j^{C_b} = \left\langle e_2^j, \sum_{\substack{\lambda_1 \leq 2b \\ r(\lambda)=0}} s_\lambda \right\rangle.$$

Equation (1.6) of the same source identifies the Schur sum with $\det(f_{i-j}^e - f_{i+j}^e)$. Apply the Hall-isometric involution ω ; it sends e_2 to h_2 and f_r^e to f_r^h . Hall duality then gives the formula for $\mathcal{C}_b$.

Finally, the two components in the type- D formula transpose to the sectors

$$\lambda_1 \leq 2b, \quad c(\lambda) = 0 \text{ or } c(\lambda) = 2b.$$

The second sector is precisely the determinant-twisted component: its $2b$ columns all have odd length. Equation (1.7) of the cited theorem at $u = 0$ is the sum of these two sectors and equals $\mathcal{D}_b$. A final use of Hall duality proves the type- D identity. $\square$

Lemma 14.15 (Two-power coefficient functional). *Let F be homogeneous of symmetric-function degree $2j$. Define the specialization π_2 on power sums by*

$$\pi_2(p_1) = p, \quad \pi_2(p_2) = q, \quad \pi_2(p_r) = 0 \quad (r \geq 3).$$

Then

$$[m_{(2^j)}] F = j! \sum_{a=0}^j (2(j-a)-1)!! [p^{2(j-a)} q^a] \pi_2(F). \quad (22)$$

Under this specialization,

$$\sum_{r \geq 0} \pi_2(e_r)u^r = \exp(pu - qu^2/2), \quad \sum_{r \geq 0} \pi_2(h_r)u^r = \exp(pu + qu^2/2),$$

and hence, with $e_{-1} = h_{-1} = 0$,

$$re_r = pe_{r-1} - qe_{r-2}, \quad rh_r = ph_{r-1} + qh_{r-2}.$$

In the last display e_r and h_r abbreviate their specialized images $\pi_2(e_r)$ and $\pi_2(h_r)$.

Proof. Expand F in the power-sum basis. A power-sum monomial containing p_r with $r \geq 3$ cannot contain the ordinary monomial $x_1^2 \cdots x_j^2$, so it contributes nothing to $[m_{(2j)}]F$. The remaining power-sum monomials are $p_1^{2(j-a)}p_2^a$. In their expansion, the a ordered p_2 factors must choose distinct variables, while the remaining $2(j-a)$ ordered p_1 factors must hit each of the other variables twice. Thus the coefficient of $x_1^2 \cdots x_j^2$ is

$$\frac{j!}{(j-a)!} \frac{(2(j-a))!}{2^{j-a}} = j!(2(j-a)-1)!!,$$

which proves (22). The two generating series are the standard power-sum formulas for elementary and complete symmetric functions after applying π_2 . Differentiating them with respect to u and comparing coefficients gives the recurrences. $\square$

Proposition 14.16 (Exact polynomial and Weyl-cap tails in the residual rows). *For each row in the following table, $I_{a,n}^G > 0$ whenever $a \geq 1$, n is odd, and $2a + n \geq K$:*

G	T	δ	k	K	method	G	R	δ	q	K
B_3	227/20	23/20	8	37	polynomial	B_2, C_2	—	6/5	39/5	47
B_4	331/20	5/4	10	41	polynomial	B_6	1053/50	3/2	1386/25	51
B_5	75/4	6/5	10	55	polynomial	D_6	371/20	7/5	921/20	49
C_3	57/5	6/5	8	37	polynomial	D_7	18	8/5	357/5	53
D_4	57/4	23/20	9	47	polynomial	D_9	763/10	8/5	751/10	61
D_5	131/10	6/5	7	33	polynomial					

In the right-hand table, $d = \dim G$ and $q = d - R - \delta$ except in the B_2, C_2 row, where the cap threshold is $T = 9$ and $q = T - \delta$.

Proof. For a polynomial row, let

$$A = m_{2k+1} - Tm_{2k}, \quad B = m_{4k+2} - 2Tm_{4k+1} + T^2m_{4k}.$$

Lemma 11.3, with $P(x) = x^k$, gives $\mathbf{P}\{X > T\} \geq A^2/B$ whenever $A, B > 0$. The exact verifier cited below regenerates all these moments from Lemmas 14.14 and 14.15 and checks, in reduced GMP rational arithmetic,

$$A > 0, \quad B > 0, \quad \frac{A^2}{B}(1 - \delta^{-2})(T - \delta)^K > (2C_G)^K.$$

Lemma 11.4 proves the six polynomial rows.

For B_2 and C_2 , use the type- B_2 positive roots $\theta_1, \theta_2, \theta_1 + \theta_2, \theta_1 - \theta_2$. On $[0, 1/4]^2$, the inequality $\cos u \geq 1 - u^2/2$ gives $X > 9$. Moreover, $4\sin^2(u/2) \geq u^2/4$ and $\pi < 4$ give

$$\begin{aligned} \mathbf{P}\{X > 9\} &\geq \frac{1}{512} \left(\frac{1}{4}\right)^4 \int_0^{1/4} \int_0^{1/4} \theta_1^2 \theta_2^2 (\theta_1 + \theta_2)^2 (\theta_1 - \theta_2)^2 d\theta_1 d\theta_2 \\ &= \frac{1}{512} \left(\frac{1}{4}\right)^{14} \left(\frac{1}{21} - \frac{2}{25} + \frac{1}{21}\right) = \frac{1}{9019431321600}. \end{aligned}$$

The two rows have the same adjoint-character law because their Lie algebras are isomorphic and Lemma 7.11 removes the global-form choice.

For the remaining four cap rows, apply Lemma 14.2. Its exact rational lower bound $\hat{p}_G$ is obtained as follows:

$$\begin{aligned}\hat{p}_{B_6} &= \frac{(1 - R/24)^2 \prod_{i=1}^6 (2i)!}{2^6 6! 2^6 (44/7)^3 39!} \left(\frac{R}{22}\right)^{39}, \\ \hat{p}_{D_6} &= \frac{(1 - R/24)^2 (2!4!6!8!10!6!)}{2^5 6! (44/7)^3 33!} \left(\frac{R}{20}\right)^{33}, \\ \hat{p}_{D_7} &= \frac{(1 - R/24)^2 (2!4!6!8!10!12!7!)}{2^6 7!} \frac{4^{46} 46!}{12(22/7)^{49} 2!} \left(\frac{R}{24}\right)^{45} \frac{433}{500}, \\ \hat{p}_{D_9} &= \frac{(1 - R/192)^{16} (2!4!6!8!10!12!14!16!9!)}{2^8 9!} \frac{(7/5)^{477} 77!}{(44/7)^5 154!} \left(\frac{R}{32}\right)^{76} \frac{193}{125}.\end{aligned}$$

Here R has the row-specific value in the table. These are strict lower bounds in (5): besides $\pi < 22/7$, the odd-dimensional rows use

$$\sqrt{2} < \frac{3}{2}, \quad \sqrt{\frac{3}{4}} > \frac{433}{500} \quad (D_7),$$

and

$$\sqrt{2} < \frac{10}{7}, \quad \sqrt{\frac{763}{320}} > \frac{193}{125} \quad (D_9),$$

all verified by squaring positive rationals. The sine factors are (6) in the first three rows and (7) in D_9 . The same exact ledger checks $R < 9\kappa$ in all four rows; since $\pi > 3$, this places the Euclidean ball strictly inside the standard eigenangle chart.

Finally, the same GMP verifier checks for each cap row

$$\hat{p}_G(1 - \delta^{-2})q^K > (2C_G)^K,$$

using $C_{B_2} = C_{C_2} = 2$, $C_{B_6} = C_{D_6} = 6$, $C_{D_7} = 5$, and $C_{D_9} = 7$. Lemma 11.4 completes the proof. Every rational comparison in this proof is implemented in

`character_ring_iter/verify_full_q3_bcd_bounded_littlewood_gmp.cpp`.

The accepted transcript `certificates/full_q3/fullq3bcd0010_machine_c.log` asserts the exact scope of six polynomial rows and six rational-cap rows, prints a positive reduced rational margin for each, and ends with exit status zero and the terminal marker

`FULL_Q3_BCD_BOUNDED_LITTLEWOODVERIFICATION:ALLPASS.`

□

Proposition 14.17 (Directed exact-MGF tails in the residual rows). *For the ranks in the following displays, let K_b be the corresponding entry:*

b	7	8	9	10	11	12	13	14	15	16	17
$K_b(B)$	53	51	53	57	57	61	61	45	45	45	45

b	4	5	6	7	8	9	10	11	12	13	14	15	16
$K_b(C)$	61	55	39	39	49	49	53	55	53	57	59	55	55

b	17	18	19	20	21	22	23	24	25	26	27	28
$K_b(C)$	57	45	45	45	47	47	49	51	53	55	57	59

and

b	8	10	11	12	13	14	15	16	17	18	19	20
$K_b(D)$	57	51	57	45	45	45	45	43	43	43	45	45

b	21	22	23	24	25	26	27	28	29	30
$K_b(D)$	45	45	45	47	45	47	47	49	47	49.

Then, in every displayed row,

$$I_{a,n}^G > 0 \quad (a \geq 1, n \text{ odd}, 2a + n \geq K_b).$$

Proof. For each row, the certificate chooses exact rational numbers $r, q, \delta > 0$, sets $c = rb$ and

$$t = \sqrt{2c + 2\delta + q},$$

and applies Lemma 14.9 with eight layers to the exact determinants in Lemma 14.8. In type B the positive and negative defining-trace tails are evaluated separately; in types C and D symmetry doubles the positive tail. Let V be the resulting lower bound for $\mathbf{P}\{|T_1| \geq t\}$.

Rains' power-two decompositions [12, Theorem 1.5, equations (22), (23), and (36)] give an upper bound U for $\mathbf{P}\{\tau_G > q\}$ by exponential Markov. In type B the bound is

$$U \leq \exp\{-\lambda(q - 1) + \lambda^2\}.$$

In types C, D , the MGF of $\tau_G - 1$ is a product of two exact orthogonal-component MGFs from Lemma 14.8, with the component sizes and signs specified by the cited decomposition. Thus the event A in Lemma 14.3 has probability at least $V - U$. Every row certificate checks $V - U > 0$.

Chebyshev gives $\mathbf{P}\{|X| < \delta\} \geq 1 - \delta^{-2}$. At the displayed onset $K = K_b$, the certificate bounds

$$\mathbf{E}(\tau_G)_+^K \leq \min \left\{ d_T^K, \left(\frac{K}{ev} \right)^K \mathbf{E}e^{v\tau_G} \right\}, \quad v = \frac{\sqrt{1 + 8K} - 1}{4},$$

where $d_T = 2b + 1$ in type B and $d_T = 2b$ in types C, D ; the MGF is the same source-audited product just described. It then checks

$$(1 - \delta^{-2})(V - U)c^K > \mathbf{E}(\tau_G)_+^K, \quad c > 2C_G. \quad (23)$$

Lemma 14.3 propagates each row from K to every later odd total degree.

The proof companion is

`character_ring_iter/verify_full_q3_bcd_low_tail_mpfr.cpp`.

It evaluates each Bessel series

$$I_k(2s) = \sum_{m \geq 0} \frac{s^{2m+k}}{m!(m+k)!}$$

through 180 recurrence updates and encloses the remaining positive tail by a geometric series. It evaluates every determinant by interval Cholesky; the matrices are Gram matrices for positive Jacobi weights, and the program rejects a nonpositive pivot. Every rational operation, square root, exponential, logarithm, Bessel bound, and Cholesky operation is rounded outward at 384-bit MPFR precision. The raw certificate prints the full rational schedule for each row before its result and cross-checks all onsets against the independent header `character_ring_iter/full_q3_bcd_remaining_data.hpp`. It verifies all 58 rows; its least directed logarithmic margin in (23) is

$$0.0971194051797054477804174 \dots > 0 \quad \text{at } C_{15}.$$

Its terminal marker is

`FULL_Q3_BCD_LOW_TAIL VERIFICATION: ALL PASS.`

The complete transcript is

`certificates/full_q3/fullq3bcdlowtail0001_optimus.log`,

authenticated by its adjacent SHA-256 file. □

Proposition 14.18 (Exact bounded-Littlewood residual certificate). *For a row $G = B_b, C_b, D_b$, put*

$$s_G = \begin{cases} 2b + 1, & G = B_b, C_b, \\ b - 1, & G = D_b. \end{cases}$$

In the rows

$$B_2, \dots, B_{21}, \quad C_2, \dots, C_{28}, \quad D_4, \dots, D_{70},$$

let K_G be the tail onset assigned to that row in Proposition 14.16 or Proposition 14.17. Then

$$I_{a,n}^G > 0 \quad \text{whenever} \quad a \geq 2, \quad n \text{ is odd}, \quad s_G < 2a + n < K_G.$$

This domain contains exactly 17,862 pairs. Its smallest value is

$$I_{2,1}^{D_5} = 50.$$

Proof. We describe the exact finite calculation and why it evaluates the formulas in the preceding lemmas. After applying π_2 , substitute

$$p = t, \quad q = zt^2.$$

For the homogeneous degree- $2j$ part of any of B_b, C_b, D_b , let $P_{G,j}(z)$ be the coefficient of t^{2j} . It has degree at most j , and Lemma 14.15 becomes

$$m_j^G = j! \sum_{a=0}^j (2(j-a)-1)!! [z^a] P_{G,j}(z). \quad (24)$$

Thus the values at any $j+1$ distinct rational points determine the moment exactly.

The verifier

`character_ring_iter/verify_full_q3_bcd_bounded_littlewood_gmp.cpp`

works simultaneously through $j = 71$. For each of the integer nodes $z = 0, 1, \dots, 71$, it uses the recurrences in Lemma 14.15 to form the specialized e_r, h_r, f_r^e, f_r^h as truncated power series through degree 142, then evaluates every leading determinant in Lemma 14.14. Lagrange interpolation applies the functional in (24). All of this is done in exact prime fields; the primes exceed 2^{30} , and therefore every integer divided by the recurrences and every nonzero node difference is invertible.

The 31 prime-field evaluations are independent OpenMP tasks. Their residues are merged in one fixed descending-prime order by exact GMP CRT, giving a modulus of 961 bits. If $d_G = \dim G$, invariant multiplicity and $|\chi_{\text{ad}}| \leq d_G$ give the elementary reconstruction bound

$$0 \leq m_j^G \leq d_G^j.$$

The program chooses primes until their product is strictly larger than the largest such bound it consumes, and rejects any reconstructed moment outside its own bound. Hence the CRT representative is the unique integer moment, not a probable reconstruction.

The program next regenerates the stable recurrence, checks every stable prefix, checks the $B_2 = C_2$ Lie-isomorphism row, and substitutes the moments into the exact GMP sum of Lemma 7.10. The row ledger is `character_ring_iter/full_q3_bcd_remaining_data.hpp`; it asserts the stable endpoint, tail onset, and moment endpoint for every one of the 114 rows in the statement. The verifier rejects a nonpositive hierarchy value, an incorrect row ledger, or a scope other than 17,862.

The current-source transcript

`certificates/full_q3/fullq3bcdboundedfinal0001_current_source.log`

authenticates the unified determinant run from the exact verifier and row-data hashes printed in its header. It reports all 17,862 values positive with the stated minimum and ends with the terminal marker

`FULL_Q3_BCD_BOUNDED_LITTLEWOODVERIFICATION:ALLPASS.`

The wrapper records exit status zero, and the adjacent SHA-256 sidecar binds the raw transcript. The historical machine-C transcripts are

```
certificates/full_q3/fullq3bcd0010_machine_c.log,
certificates/full_q3/fullq3bcd0002_machine_c.log.
```

They separately cover the first 12,993 determinant cases and the remaining 4,869 B/D cases by reverse Pieri. They remain independent historical overlaps, not substitutes for the unified current-source transcript. As an implementation-independent control for the original low-row subledger, the earlier reverse-Pieri verifier agrees exactly on all 2,845 moments and all 7,484 residual pairs through degree 40. Only 538 required moments remain above that prefix. The independent modular checker described in the code-and-data section evaluates their determinant formulas using ordinary modular arithmetic and Newton forward differences, rather than the promotion verifier's Montgomery arithmetic, Lagrange weights, and GMP reconstruction. It requires agreement modulo 21 distinct primes whose product has 628 bits. This exceeds the independently checked 607-bit maximum of the bounds d_G^j . Both the candidate and the invariant multiplicity lie in $[0, d_G^j]$, so their difference has absolute value smaller than the modulus; the residue agreements therefore prove equality as integers. The checker then recomputes all remaining 5,509 hierarchy values with GMP integers and requires strict positivity. Thus no degree-41–59 reverse-Pieri frontier is needed; the degree-40 prefix remains the structurally independent overlap test. Its current-source transcript is

```
certificates/full_q3/fullq3bcdmodularfinal0001_current_source.log.
```

The header binds the checker, row data, and two input ledgers; the terminal marker and wrapper status are both zero, and the adjacent SHA-256 sidecar binds the transcript. $\square$

Proposition 14.19 (Exact B_{18} – B_{21} and D_{31} – D_{70} residual certificate). *For every B_b with $18 \leq b \leq 21$ and every D_b with $31 \leq b \leq 70$,*

$$I_{a,n}^G > 0 \quad (a \geq 2, n \text{ odd}).$$

Thus the complete generator hierarchy holds in all 44 rows.

Proof. Theorem 13.1 supplies the stable prefix, and Proposition 14.12 supplies the tail. The intervening $a \geq 2$, odd- n box is the B_{18} – B_{21} and D_{31} – D_{70} subledger of Proposition 14.18. The case $a = 1$ is Theorem 7.5. These ranges are exhaustive, proving the proposition.

The older reverse-Pieri source `character_ring_iter/verify_full_q3_bd_residual_gmp.cpp` and its authenticated machine-C transcript remain an algorithmically independent control: they check exactly 137 type- B and 4732 type- D residual pairs, 4869 in total, and report minimum

$$I_{8,15}^{D_{31}} = 1272988116891284109857142182380802 > 0.$$

The current aggregate rebuilds and runs this source as the mandatory secondary checker for the complementary 4,869-case partition. It remains a cross-check rather than an additional premise of the bounded-Littlewood proposition. $\square$

Corollary 14.20 (Completed middle orthogonal rows). *The complete generator hierarchy holds for*

$$B_b \quad (b \geq 18), \quad D_b \quad (b \geq 31).$$

Proof. Combine Propositions 14.6, 14.11, 14.12, and 14.19 with Theorem 13.1. $\square$

Theorem 14.21 (Full hierarchy in types B , C , and D). *Let G be a compact connected group with simple Lie algebra of type B_b ($b \geq 2$), C_b ($b \geq 2$), or D_b ($b \geq 4$). Then*

$$I_{a,n}^G \geq 0 \quad (a, n \geq 0).$$

Consequently Ginibre's full condition Q_3 holds on $\mathcal{Q}_G$.

Proof. First consider the low rows $B_2, \dots, B_{17}$, $C_2, \dots, C_{28}$, and $D_4, \dots, D_{30}$. The all-plus and even- n sectors are automatic by Theorem 7.4, while Theorem 7.5 supplies $a = 1$. Let $a \geq 2$, let n be odd, and put $L = 2a + n$. Theorem 13.1 applies when $L \leq s_G$. Proposition 14.18 applies when $s_G < L < K_G$. At $L \geq K_G$, the row is covered by either Proposition 14.16 or Proposition 14.17. These three ranges are exhaustive.

For B_b with $b \geq 18$ and D_b with $b \geq 31$, use Corollary 14.20. For C_b with $29 \leq b \leq 295$, use Proposition 14.11; the same proposition covers the still-needed middle ranges $22 \leq b \leq 295$ in type B and $71 \leq b \leq 297$ in type D . All remaining higher ranks satisfy the hypothesis of Proposition 14.6. Thus the generator hierarchy holds in every row. Central-cover invariance gives every compact connected global form, and Theorem 7.4 promotes the generator to Q_G . $\square$

Proposition 14.22 (Auditable computational interface). *Every machine-decided sign used in the classification closure is the value of one of the following fully specified mathematical predicates.*

- (1) *The target is the exact integer double sum in Lemma 7.10; no program-specific rearrangement changes the quantity being decided.*
- (2) *Type-A moments are supplied by the Schur–Weyl and hook-length formulas in Theorem 11.1 and Propositions 11.5–11.6. Exceptional moments are supplied by the Cartan-datum Racah–Speiser and character-pairing reconstruction stated in Proposition 12.2. Finite classical moments are supplied by Lemmas 14.13–14.15.*
- (3) *Every analytic tail is reduced to a positive lower bound by Theorem 7.15, Lemma 11.4, or Lemma 14.3. Their hypotheses are discharged by the exact cap formulas or by the Fredholm and exact-MGF inequalities in Lemmas 14.2, 14.4, 14.7, and 14.8. Directed arithmetic decides only the final explicitly stated interval inequalities.*
- (4) *The disjoint stable, finite, and tail regions are recorded row by row in the committed file*

`full_q3_classification_ledger.csv`.

Its generator

`generate_full_q3_classification_ledger.py`

reconstructs the classical rows from the same published cutoff header consumed by the exact verifier, derives every pair count from $L = 2a + n$, and fails unless the totals are exactly 1,315 general type-A, 1,265 exceptional, and 17,862 finite classical pairs. It also prints the four fixed low type-A rows and the parametric high-rank ranges, so no coverage claim is available only in C++ source.

- (5) *The mandatory secondary finite-classical replay is partitioned into the 12,993 low rows checked by the modular/Newton implementation and the 4,869 B_{18} – B_{21} , D_{31} – D_{70} rows checked by reverse Pieri. The independent classification audit reconstructs both domains from the theorem cutoffs, rejects an overlap, and proves that their disjoint union is the complete 17,862-case primary ledger.*

Thus hashes and transcripts authenticate source and execution, but the mathematical interfaces above determine what is proved. A replay is accepted only after evaluating these predicates over every ledger row with exact integer/rational arithmetic or with an outward-rounded positive interval.

Proof. Items (1)–(3) are precisely the backward mathematical dependencies of the family closures already proved above. For items (4)–(5), the ledger generator parses all 114 finite classical rows, recomputes the stable endpoint and the number of pairs at every intervening odd total degree, asserts the three displayed totals, and separately checks the $12,993 + 4,869$ secondary partition before comparing its byte-for-byte output with the committed ledger. The publication preflight and Part III aggregate both run this check. The exact and directed verifier semantics are stated in the proofs of the cited residual and tail propositions and summarized in the code-and-data section below. Consequently every accepted computational conclusion has both a mathematical predicate and an exhaustive ledger entry. $\square$

Theorem 14.23 (Full adjoint-generated Ginibre Q_3). *Let G be any compact connected Lie group with simple Lie algebra. For every integer $r \geq 0$, every tuple $f_1, \dots, f_r \in \mathcal{Q}_G$, and every choice of signs $\varepsilon_i \in \{+1, -1\}$ containing an even number of minus signs,*

$$\int_{G \times G} \prod_{i=1}^r (f_i(g) + \varepsilon_i f_i(h)) d\mu_G(g) d\mu_G(h) \geq 0.$$

If the number of minus signs is odd, the integral is zero. Hence $\mathcal{Q}_G$ satisfies Ginibre’s condition Q_3 with all of its original finite-tuple and sign-pattern quantifiers. In particular, this includes every finite tuple of exponential adjoint interactions $e^{\beta_i \chi_{\text{ad}}}$ with $\beta_i \geq 0$.

Proof. Classification leaves type A , types B, C, D , and the five exceptional types. Theorem 11.7, Theorem 14.21, and Theorem 12.3, respectively, prove the full generator hierarchy in those cases. Lemma 7.11 removes the choice of compact connected global form. Proposition 14.22 fixes the exhaustive row ledger and the exact mathematical predicates behind every machine-decided sign. Finally, Theorem 7.4 gives the asserted cone promotion and the vanishing of every odd-minus sector. The final exponential specialization is admissible by Lemma 7.2. $\square$

Remark 14.24 (Exact sense in which the result matches Ginibre). Theorem 14.23 matches the full quantifier structure of Ginibre’s 1970 condition Q_3 and uses his Proposition 1 multiplicative-cone promotion. It does not replace $\mathcal{Q}_G$ by the cone of all real continuous positive-definite functions. Such a replacement is impossible: Proposition 7.6 gives the exact $\text{SO}(3)$ value $-8/279$.

15. CODE AND DATA AVAILABILITY

Certificate semantics and trusted base. The computer-assisted results have three separate layers. First, each numbered proposition states the mathematical input formula, the finite quantified domain, and the sign or interval comparison to be decided. Second, the named verifier reconstructs the inputs and evaluates that comparison; a stored output value is not accepted as an input merely because it appears in a transcript. Third, the transcript records the source identities, scope counts, arithmetic outputs, terminal marker, and process status of one execution. A transcript authenticates a completed run but is not, by itself, the proof of the formula implemented by the verifier.

The trusted computational base is consequently explicit: a conforming C++ implementation, GMP integer and rational arithmetic, MPFR directed rounding, and the short orchestration and manifest parsers named below. No claim is made that these programs are formally verified. The exact stages avoid probabilistic reconstruction: when prime fields and CRT are used, the proved modulus bound makes the reconstructed integer unique. The interval stages accept a sign only when the outward-rounded lower endpoint is positive. Independent recurrence checks, alternative low-degree implementations, sanitizer runs, and mutation tests are redundancy controls; their stated cutoffs are not represented as full second proofs of computations beyond those cutoffs.

The historical execution snapshot for Parts I–III is rooted at the immutable commit

https://github.com/skypher/ginibre-q3-adjoint/tree/91363c3cebb37a44349f613ab0a5aa6dcd412af3/ginibre_q3.

Its full identifier is

91363c3cebb37a44349f613ab0a5aa6dcd412af3.

That commit is retained only as a historical execution snapshot. It is not the source authority for the present theorem; the historical tag `v1.0.0` identifies that snapshot. For the present submission, the canonical source bytes are the files in the submitted archive whose hashes are listed in

`certificates/full_q3/full_q3_source_manifest.sha256`.

The generated formal PDFs are separately bound by `publication_artifacts.sha256`. A public release of the final submission must preserve the submitted archive and reproduce both manifests exactly.

The formal submission is one unified file, `submission.pdf`. Its source wrapper `submission.tex` includes `paper.pdf` and the present `full_q3_extension.pdf` after a unified abstract and proof map. The standalone `full_q3_main.pdf` remains a manifested navigation aid. The optional `paper_full.pdf` preserves longer combinatorial derivations and ledgers but is not a theorem dependency. The archived computation-bearing sources include `paper.tex` and `full_q3_extension.tex`. The historical commit authenticates only its arithmetic snapshot. The current publication sources are bound directly by the source manifest displayed above and are the inputs rebuilt by the commands below. The final publication artifacts are rebuilt before their hashes are checked against `publication_artifacts.sha256`; mere existence of a pre-existing PDF is not an acceptance condition. The historical aggregate transcript predates publication-only prose and metadata revisions; the fail-closed `verify_full_q3_related_work_rebind.py` records that boundary and rejects any change to an arithmetic-bearing source. That rebind is a provenance test, not a step required to construct the submitted source: the final files are bound directly by the final-source manifest. A final release archive must preserve both the immutable execution snapshot and those exact final-source bytes. Part III is not offered as a standalone proof: its Theorem 7.5 imports Theorem 2.2 of Parts I–II, which is included in the submission and archive.

The optional derivation archive `paper_full.pdf` is rebuilt from the same source for provenance. The direct determinant/formula certificate contract, accepted-certificate boundary, half-stable bridge, and family closures all appear in `paper.pdf`.

The exact replay contract, software requirements, host/thread guidance, and terminal success markers are documented in `REPLAY.md` and `FULL_Q3_DISTRIBUTED_REPLAY.md`. The principal requirements are a C++20 compiler with OpenMP, GMP, MPFR, Python 3, GNU Make, and a $\text{\LaTeX}$ installation. From the repository root, the three aggregate commands are

```
python3 ginibre_q3/run_publication_short_audits.py
make -C ginibre_q3 clean-room-replay
make -C ginibre_q3 full-q3-extension.
```

The exact validated Ubuntu, compiler, GMP/MPFR, OpenMP, Python, and $\text{\TeX}$ versions used for the July 2026 publication rebuild are recorded in `ENVIRONMENT.md`; the Makefiles remain authoritative for compilation and link flags. These commands implement three reproducibility tiers. The first command runs the document, dependency, interface, scope, moment-formula, cone-promotion, $SO(3)$, and frontier-formula checks as one fail-closed short suite. The Part III command regenerates its exact and directed-rounding arithmetic with parallel suppliers. The separate Parts I–II clean-room command is the full certificate-regeneration tier. Both workflows enforce a ten-minute ceiling for each underlying computation on the validated host; an overrun must be profiled and optimized or replaced by a proved analytic reduction before release. The resource envelope is documented in `REPLAY.md`. Part III does not silently substitute its faster audit for that imported computation.

For the final public archive, the clean-worktree driver `run_final_publication_replay.sh` runs the preflight and both arithmetic aggregates, records the exact commit and final-source-manifest identity in one persistent log, and rejects any replay that changes the source tree. The deterministic packager `build_final_release_archive.py` then includes every tracked publication file, that successful replay log, and the exact commit metadata in one checksum-addressed archive. The commands and terminal markers are given in `REPLAY.md`. The archive deposited for publication must be the output of this final-source workflow; the historical execution snapshot is retained as audit history, not substituted for the final replay.

The independent check of the 12,993-case low-row subledger is supplied without constructing the large degree-41–59 reverse-Pieri frontier. Its source is

`character_ring_iter/verify_full_q3_bcd_modular_moment_checker.cpp`.

It checks the archived degree-40 reverse-Pieri overlap, verifies the 538 higher candidates by the uniquely bounded 21-prime calculation described above, and checks the 5,509 higher hierarchy values exactly. It rejects any incorrect row, degree, count, residue, character bound, modulus bound, or nonpositive hierarchy value before emitting `FULL_Q3_MODULAR_CHECKER VERIFICATION: ALL PASS`. The aggregate target `full-q3-extension` also rebuilds and runs the reverse-Pieri supplier for the complementary 4,869-case B/D subledger. The two secondary programs have disjoint domains of 12,993 and 4,869 cases, and the independent classification audit proves that their union is exactly the complete 17,862-case box. They are mandatory release cross-checks; the primary proof predicate remains the unified bounded-Littlewood calculation. The alias

`full-q3-extension-independent-controls`

runs the same secondary pair; `run_full_q3_bcd_independent_audit.sh` is the persistent-log wrapper.

SHA-256 manifests authenticate every load-bearing source, accepted transcript, generated ledger, and cited source artifact. Compiler and resource checks and author self-audits are integrity or redundancy tests; they are not independent peer review and do not by themselves decide a sign. The mathematical certificate claims are the exact integer/rational or directed-rounding comparisons described in the numbered results above.

The executable proof obligations are the following. This list fixes the input/output contract independently of program names.

- (1) An *exact moment supplier* receives a root datum or one of the bounded Littlewood determinants in Lemmas 14.13–14.15; it must output every requested integer moment together with its rank and degree. A CRT output is accepted only after the product modulus exceeds the displayed character bound d_G^j , making the representative unique.
- (2) An *exact hierarchy evaluator* receives only those moments and a finite list of triples (G, a, n) ; it evaluates the double sum of Lemma 7.10, rejects a missing or duplicate triple, and accepts only a strictly positive integer for every listed residual case. The row ledger must prove that the listed triples are exactly the complement of the stable and analytic-tail ranges.
- (3) A *directed analytic supplier* receives rational tail parameters and the exact determinant or Fredholm formulas proved above. Every elementary operation, Bessel remainder, determinant pivot, exponential, and logarithm is enclosed with outward MPFR rounding. It accepts only when the final lower endpoint is positive and when all trace-norm and support hypotheses are also certified by intervals.
- (4) An *independent control* may consume candidate moments but not a candidate sign. It must reconstruct the same mathematical object by a different recurrence, interpolation, or modular path and then reevaluate the hierarchy formula. Shared formulas and shared input ledgers are part of the trusted base and are not described as independent.

Table 2 gives the compact trusted-computation map. Here “exact” means integer or rational arithmetic, including prime-field evaluation followed by uniquely bounded GMP reconstruction; “directed” means outward-rounded 384-bit MPFR interval arithmetic.

The identifiers in the table resolve to the following authenticated transcripts:

- `certificates/full_q3/distributed0001/fullq3distributed0001.log`;
- `certificates/full_q3/fullq3complete0002_optimus.log`;
- `certificates/full_q3/fullq3bcdanalytic0003_machine_c.log`;
- `certificates/full_q3/fullq3bcdlowtail0001_optimus.log`;
- `certificates/full_q3/fullq3bcdboundedfinal0001_current_source.log`;
- `certificates/full_q3/fullq3bcdmodularfinal0001_current_source.log`;
- `certificates/full_q3/fullq3bcd0010_machine_c.log`; and
- `certificates/full_q3/fullq3bcd0002_machine_c.log`.

Result block	Program/transcript identifier	Arithmetic	Verified scope
Imported two-minus theorem	distributed0001	exact/directed	200 replay stages; all compact connected simple types; every $n \geq 0$
Low and stable type A	su2, su3, su45, sun-ge6	exact GMP	all residual boxes and exact tail comparisons; 1,673 residual pairs
Exceptional types	exceptional	exact GMP	five tails and 1,265 residual pairs
High and middle classical tails	bcdanalytic0003	directed MPFR	all $C_G \geq 296$ rows, 768 middle rows, and 44 supplemental rows
Remaining classical tails	bcdlowtail0001	directed MPFR	58 finite-rank tail rows
Finite classical residual boxes (primary)	current bounded replay	exact prime/GMP	114 rows and 17,862 residual inequalities
Finite classical controls	current modular/Newton and reverse-Pieri replays	exact prime/GMP	disjoint scopes $12,993 + 4,869 = 17,862$; historical transcripts retained

TABLE 2. Map from the computer-assisted proof blocks to their accepted program/transcript families and arithmetic boundaries.

REFERENCES

- [1] L. P. Polzer, The two-minus adjoint-character case of Ginibre’s Q_3 condition, Parts I–II: classification and exact certificates, formal journal manuscript, 2026; Parts I–II component of the present unified manuscript. The exact submitted source bytes are bound by `certificates/full_q3/full_q3_source_manifest.sha256` in the inseparable submission archive; the historical execution snapshot is https://github.com/skypher/ginibre-q3-adjoint/tree/91363c3cebb37a44349f613ab0a5aa6dcd412af3/ginibre_q3.
- [2] P. Diaconis and M. Shahshahani, On the eigenvalues of random matrices, *J. Appl. Probab.* **31A** (1994), 49–62.
- [3] J. Ginibre, General formulation of Griffiths’ inequalities, *Comm. Math. Phys.* **16** (1970), 310–328.
- [4] G. S. Sylvester, The Ginibre inequality, *Comm. Math. Phys.* **73** (1980), no. 2, 105–114.
- [5] A. Abdesselam, Non-Abelian correlation inequalities and stable determinantal polynomials, arXiv:2207.07603, 2022.
- [6] I. Chevyrev and C. Garban, Villain action in lattice gauge theory, *J. Stat. Phys.* **192** (2025), Paper 38.
- [7] J. Huh, J. S. Kim, C. Krattenthaler, and S. Okada, Bounded Littlewood identities with fixed number of odd rows or odd columns, *Electron. J. Combin.* **33** (2026), no. 3, Paper P3.5, doi:10.37236/14854.
- [8] P. Etingof, A uniform proof of the Macdonald–Mehta–Opdam identity for finite Coxeter groups, arXiv:0903.5084, 2009.
- [9] S. Garibaldi, R. M. Guralnick, and E. M. Rains, Invariant derivations and trace bounds, *J. Éc. polytech. Math.* **12** (2025), 1229–1287.
- [10] J. L. Bourjaily, M. R. Plesser, and C. Vergu, The many colours of amplitudes, arXiv:2412.21189.
- [11] E. M. Rains, Increasing subsequences and the classical groups, *Electron. J. Combin.* **5** (1998), Research Paper 12.
- [12] E. M. Rains, Images of eigenvalue distributions under power maps, *Probab. Theory Related Fields* **125** (2003), 522–538.
- [13] C. Courteaut, K. Johansson, and G. Lambert, From Berry–Esseen to super-exponential, *Electron. J. Probab.* **29** (2024), no. 11, 1–48, doi:10.1214/23-EJP1068.

Email address: `polzer@fastmail.com`