

Algebraic tunnelling for the restricted fractional Laplacian on distant domains

LESLIE P. POLZER

Independent Researcher

polzer@fastmail.com

August 27, 2026

Abstract. Let $D \subset \mathbb{R}^d$ be a bounded centrally symmetric Lipschitz domain, and let Ω_L be the union of two reflected translates of D whose centers are separated by L . For the restricted fractional Laplacian $(-\Delta)^s$, $0 < s < 1$, we determine the leading splitting of the two eigenvalues converging to the one-well ground-state energy. If μ_1 is the one-well ground-state eigenvalue, ϕ_1 is its positive L^2 -normalized eigenfunction, and $m_1 = \int_D \phi_1$, then

$$\lambda_1(\Omega_L) = \mu_1 - c_{d,s} m_1^2 L^{-d-2s} + O(L^{-d-2s-2} + L^{-2d-4s}),$$

$$\lambda_2(\Omega_L) = \mu_1 + c_{d,s} m_1^2 L^{-d-2s} + O(L^{-d-2s-2} + L^{-2d-4s}).$$

The proof uses an exact reflection reduction to the two operators $A_D \pm B_L$, an elementary isolated-eigenvalue estimate, and a second-order Taylor expansion of the interaction kernel. For two equal balls, the formula for λ_2 gives the leading rate in the nonlocal Hong–Krahn–Szegő minimizing sequence. More generally, central symmetry is unnecessary for the finite-well result: for any bounded Lipschitz reference domain and any fixed collection of distinct translation sites $a_1, \dots, a_N$, the first N eigenvalues are

$$\lambda_k = \mu_1 + L^{-d-2s} \theta_k + O(L^{-d-2s-2} + L^{-2d-4s}), \quad 1 \leq k \leq N,$$

where $\theta_1 \leq \dots \leq \theta_N$ are the eigenvalues of the explicit matrix with zero diagonal and off-diagonal entries $-c_{d,s} m_1^2 |a_i - a_j|^{-d-2s}$. Exact cell-integral Galerkin computations on two and three intervals illustrate the corresponding fixed-mesh limits, while a mesh study records stabilization of the one-well interaction coefficient. In the displayed spectral asymptotics, $L \rightarrow \infty$. The implicit constants in the two-well formulas depend only on d, s, D, R , and g ; in the multi-well formula they may also depend on the fixed N and configuration $(a_1, \dots, a_N)$.

Keywords. restricted fractional Laplacian; eigenvalue splitting; nonlocal double wells; spectral perturbation; shape optimization.

MSC2020 Classification. 35P15; 35R11.

1. Introduction

The restricted fractional Laplacian couples disjoint components through its integral kernel. This differs from the local Dirichlet Laplacian, whose spectrum on a disconnected union is

exactly the multiset union of the component spectra. Two distant congruent components therefore form a nonlocal analogue of a double well: their limiting degeneracies split at an algebraic, rather than exponential, scale.

Brasco and Parini proved that the fractional Hong–Kahn–Szegő bound for the second eigenvalue is strict and that two equal balls form a minimizing sequence when their mutual distance tends to infinity [BP16, Theorem 6.2]. Biagi, Dipierro, Valdinoci, and Vecchi proved the mixed local/nonlocal analogue [BDVV23, Theorem 1.1]. Parini and Salort placed such distant components in a general concentration–compactness dichotomy [PS20, Theorems 1.1 and 1.2]. Together these works establish the qualitative limiting and dichotomy picture but do not specify the leading interaction coefficient for the standard restricted fractional Laplacian. The purpose of this article is to extract that coefficient and the associated effective finite-dimensional interaction.

The influence of mutual position on the restricted fractional spectrum is also emphasized by Abatangelo, Felli, and Noris [AFN20, Introduction, p. 3], in a study of small fractional-capacity perturbations. In the shape-optimization setting, Zahl isolated the exact cross-energy under translations and proposed a signed point-mass interaction with kernel $|x - y|^{-d-2s}$ [Zah26, Section 10, equations (10.3) and (10.7)]. Wu subsequently settled Zahl’s no-local-minimum conjecture for that discrete signed-mass energy [Wu26, abstract]. In another recent direction, Dipierro, Proietti Lippi, Sportelli, and Valdinoci studied disconnected domains for generalized eigenproblems having a superposition of lower-order operators on the right-hand side [DPLSV26, Theorems 1.4–1.7]. None of these works derives a large-separation spectral expansion for the standard restricted fractional Laplacian. Zahl’s discrete energy anticipates the matrix in (4.9); the result here derives it from the full restricted fractional operator, bounds the off-diagonal block in operator norm, controls its coupling to the complementary one-well modes through the spectral gap, and identifies every ordered eigenvalue in the cluster. We are not aware of an earlier result giving either the two-component coefficient with a quantified remainder or this finite-well spectral reduction.

The first result, Theorem 3.3, treats the double ground-state cluster. Theorem 4.6 identifies the entire ground-state cluster for finitely many translated components with the spectrum of an explicit weighted interaction matrix; unlike the reflection reduction, this result does not require central symmetry of the reference component. Both proofs are self-contained after the standard one-well spectral facts. The interval provides an explicit benchmark: one-well eigenvalue and eigenfunction estimates are available in [Kwa12, Theorem 1 and Propositions 1–2]; sharper eigenvalue asymptotics and an eigenfunction bound uniform in both the index and the fractional order are proved in [Zha26, Theorems 1 and 2].

2. The one-well operator and exact two-well reduction

Fix $d \geq 1$ and $0 < s < 1$, and put

$$\kappa = d + 2s.$$

For an open set $G \subset \mathbb{R}^d$, let

$$\tilde{H}^s(G) = \{u \in H^s(\mathbb{R}^d) : u = 0 \text{ a.e. on } \mathbb{R}^d \setminus G\}.$$

For bounded Lipschitz G , this space agrees with the completion of $C_c^\infty(G)$ in the global $H^s(\mathbb{R}^d)$ norm [DFW21, Section 2, equation (2.1)]. The zero-exterior quadratic form is

$$\Pi_G[u] = \frac{c_{d,s}}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^d} \frac{|u(x) - u(y)|^2}{|x - y|^\kappa} dx dy, \quad u \in \tilde{H}^s(G), \quad (2.1)$$

where $c_{d,s} > 0$ is the normalization in the singular-integral definition of $(-\Delta)^s$. The form norm $(\Pi_G[u] + \|u\|_{L^2(G)}^2)^{1/2}$ is equivalent to the norm inherited from $H^s(\mathbb{R}^d)$ on $\tilde{H}^s(G)$. Thus (2.1) is a densely defined closed form, and we denote its associated self-adjoint operator on $L^2(G)$ by A_G . We write $\lambda_1(G) \leq \lambda_2(G) \leq \dots$ for the eigenvalues of A_G , repeated according to multiplicity. For an abstract self-adjoint operator T with compact resolvent, $\lambda_k(T)$ has the analogous meaning.

Let $D \subset B_R(0)$ be a bounded Lipschitz domain satisfying $D = -D$. The Lipschitz extension property and compactness of the form embedding follow from [DNPV12, Theorems 5.4 and 7.1]. Hence A_D has compact resolvent. Write its eigenvalues with multiplicity and denote the first two distinct levels by

$$\mu_1 < \mu_2, \quad g = \mu_2 - \mu_1 > 0.$$

The first eigenvalue is simple and has a strictly positive normalized eigenfunction by [BP16, Theorem 2.8], specialized to the linear case. Fix that eigenfunction and set

$$A_D \phi_1 = \mu_1 \phi_1, \quad \|\phi_1\|_{L^2(D)} = 1, \quad m_1 = \int_D \phi_1(x) dx > 0. \quad (2.2)$$

Simplicity and reflection invariance imply $\phi_1(-x) = \phi_1(x)$.

Fix a unit vector $e \in \mathbb{R}^d$. For $L > 2R$, define

$$D_L^- = D - \frac{L}{2}e, \quad D_L^+ = D + \frac{L}{2}e, \quad \Omega_L = D_L^- \cup D_L^+.$$

On $L^2(D)$ define the bounded integral operator

$$(B_L v)(x) = -c_{d,s} \int_D \frac{v(y)}{|Le - x - y|^\kappa} dy. \quad (2.3)$$

Its kernel is real and symmetric, so B_L is self-adjoint.

Lemma 2.1 (*Exact reflection reduction*). *The operator A_{Ω_L} is unitarily equivalent to*

$$\begin{pmatrix} A_D & B_L \\ B_L & A_D \end{pmatrix} \quad \text{on } L^2(D) \oplus L^2(D),$$

and hence to

$$(A_D + B_L) \oplus (A_D - B_L). \quad (2.4)$$

Proof. For $f \in \tilde{H}^s(\Omega_L)$ set

$$u(x) = f\left(x - \frac{L}{2}e\right), \quad v(x) = f\left(-x + \frac{L}{2}e\right), \quad x \in D.$$

This defines a unitary map from $L^2(\Omega_L)$ to $L^2(D) \oplus L^2(D)$. Translational and reflection invariance identify the two diagonal parts of the form with $\Pi_D[u]$ and $\Pi_D[v]$. Positive separation permits smooth compactly supported cutoffs isolating the two components. Multiplication by these cutoffs is bounded on $H^s(\mathbb{R}^d)$, while conversely the sum of separated translated and reflected one-well functions lies in $H^s(\mathbb{R}^d)$. Hence the unitary identifies $\tilde{H}^s(\Omega_L)$ with $\tilde{H}^s(D) \oplus \tilde{H}^s(D)$. The cross-kernel is bounded by positive separation. Each separate zero-exterior form already contains the square interaction with the location occupied by the other component. Thus, after the above changes of variables, expanding the square in (2.1) gives exactly

$$\begin{aligned} & \Pi_{\Omega_L}[f] - \Pi_D[u] - \Pi_D[v] \\ &= c_{d,s} \iint_{D \times D} \frac{|u(x) - v(y)|^2 - |u(x)|^2 - |v(y)|^2}{|Le - x - y|^\kappa} dx dy \\ &= -2c_{d,s} \operatorname{Re} \iint_{D \times D} \frac{u(x)\overline{v(y)}}{|Le - x - y|^\kappa} dx dy = 2 \operatorname{Re} \langle u, B_L v \rangle. \end{aligned}$$

The square terms therefore cancel, leaving no additional diagonal contribution. This proves the block representation in the sense of closed forms. Since B_L is bounded, it also gives the stated operator representation. The unitary change of variables $(u, v) \mapsto 2^{-1/2}(u + v, u - v)$ yields (2.4). $\blacksquare$

3. The doublet splitting

We first record the elementary perturbation estimate used for the two scalar branches.

Lemma 3.1 (*Ground-state perturbation bound*). *Let A be self-adjoint with compact resolvent, simple lowest eigenvalue μ , normalized ground state ϕ , and spectral gap $g > 0$. Let W be bounded and self-adjoint, put $b = \|W\|$ and $w = \langle \phi, W\phi \rangle$, and suppose $b \leq g/4$. Then*

$$-\frac{2b^2}{g} \leq \lambda_1(A + W) - (\mu + w) \leq 0. \quad (3.1)$$

Proof. Let $\mathfrak{a}$ be the closed quadratic form associated with A . The upper bound follows by testing the Rayleigh quotient with ϕ . For a normalized vector $\psi = a\phi + \eta$ in the form domain of $\mathfrak{a}$, where $\eta \perp \phi$ and $t = \|\eta\|$, the spectral gap gives

$$\mathfrak{a}[\psi] \geq \mu + gt^2.$$

Since $|a|^2 = 1 - t^2$, self-adjointness gives

$$\langle \psi, W\psi \rangle - w = -t^2w + 2 \operatorname{Re} \langle a\phi, W\eta \rangle + \langle \eta, W\eta \rangle.$$

Using $|w| \leq b$, $|a| \leq 1$, and $\|W\| = b$,

$$\begin{aligned} \mathfrak{a}[\psi] + \langle \psi, W\psi \rangle - (\mu + w) &\geq gt^2 - 2bt^2 - 2bt \\ &= (g - 2b)t^2 - 2bt \\ &= (g - 2b) \left(t - \frac{b}{g - 2b} \right)^2 - \frac{b^2}{g - 2b} \\ &\geq -\frac{b^2}{g - 2b} \geq -\frac{2b^2}{g}. \end{aligned}$$

Taking the infimum over normalized ψ proves the lower bound. ■

The next lemma extracts the interaction coefficient and records why central symmetry improves the kernel error by one power of L .

Lemma 3.2 (*Interaction-kernel expansion*). *Let*

$$I_L = \iint_{D \times D} \frac{\phi_1(x)\phi_1(y)}{|Le - x - y|^\kappa} dx dy.$$

For $L \geq 4R$,

$$|I_L - m_1^2 L^{-\kappa}| \leq C_{\kappa,R} m_1^2 L^{-\kappa-2}, \quad C_{\kappa,R} = 2^{\kappa+3} \kappa(\kappa+3) R^2. \quad (3.2)$$

Moreover,

$$\|B_L\| \leq \beta_L, \quad \beta_L = c_{d,s} 2^\kappa |D| L^{-\kappa}. \quad (3.3)$$

Proof. Set $F_L(z) = |Le - z|^{-\kappa}$. For $|z| \leq 2R$ and $L \geq 4R$, the segment from 0 to z remains at distance at least $L/2$ from Le . Direct differentiation gives

$$\|D^2 F_L(z)\| \leq \kappa(\kappa+3) |Le - z|^{-\kappa-2}.$$

Taylor's theorem therefore yields

$$|F_L(z) - L^{-\kappa} - \kappa L^{-\kappa-1} e \cdot z| \leq C_{\kappa,R} L^{-\kappa-2}$$

for $|z| \leq 2R$, with the displayed value of $C_{\kappa,R}$. Because ϕ_1 is even,

$$\int_D x \phi_1(x) dx = 0.$$

Consequently the integral of the linear Taylor term with $z = x + y$ vanishes. Since ϕ_1 is positive, integration of the remainder gives (3.2).

Finally, $|Le - x - y| \geq L - 2R \geq L/2$. The Schur test applied to (2.3) gives

$$\|B_L\| \leq c_{d,s} |D| (L - 2R)^{-\kappa} \leq c_{d,s} 2^\kappa |D| L^{-\kappa},$$

which is (3.3). ■

Theorem 3.3 (*Two-well algebraic splitting*). *Let $D \subset B_R(0)$ be a bounded centrally symmetric Lipschitz domain, and let Ω_L be defined above. There is $L_0 < \infty$, determined by d, s, D, R and the one-well gap g , such that for $L \geq L_0$ the first two eigenvalues of A_{Ω_L} satisfy*

$$|\lambda_1(\Omega_L) - (\mu_1 - c_{d,s}m_1^2L^{-\kappa})| \leq c_{d,s}C_{\kappa,R}m_1^2L^{-\kappa-2} + \frac{2\beta_L^2}{g}, \quad (3.4)$$

$$|\lambda_2(\Omega_L) - (\mu_1 + c_{d,s}m_1^2L^{-\kappa})| \leq c_{d,s}C_{\kappa,R}m_1^2L^{-\kappa-2} + \frac{2\beta_L^2}{g}. \quad (3.5)$$

In particular, as $L \rightarrow \infty$,

$$L^{d+2s}(\lambda_2(\Omega_L) - \lambda_1(\Omega_L)) \longrightarrow 2c_{d,s}m_1^2. \quad (3.6)$$

Proof. By Lemma 2.1, the spectrum is the union of the spectra of $A_D + B_L$ and $A_D - B_L$. Put

$$t_L = \langle \phi_1, B_L \phi_1 \rangle = -c_{d,s}I_L < 0.$$

Choose $L_0 \geq 4R$ so large that, for $L \geq L_0$,

$$\beta_L \leq \frac{g}{4}, \quad 2c_{d,s}m_1^2(L + 2R)^{-\kappa} > \frac{4\beta_L^2}{g}. \quad (3.7)$$

Such a choice exists because the left-hand side of the second inequality is of order $L^{-\kappa}$ and its right-hand side is of order $L^{-2\kappa}$.

Apply Lemma 3.1 first with $W = B_L$ and then with $W = -B_L$. The two branch ground-state energies have the form

$$E_+(L) = \mu_1 + t_L + \rho_+(L), \quad E_-(L) = \mu_1 - t_L + \rho_-(L), \quad (3.8)$$

where

$$|\rho_{\pm}(L)| \leq \frac{2\beta_L^2}{g}.$$

The second eigenvalue of either branch is at least $\mu_2 - \beta_L \geq \mu_1 + 3g/4$, whereas both branch ground-state energies are at most $\mu_1 + \beta_L \leq \mu_1 + g/4$. Thus the first two eigenvalues of the full operator are the two numbers in (3.8).

Since $|Le - x - y| \leq L + 2R$ and $\phi_1 > 0$,

$$I_L \geq m_1^2(L + 2R)^{-\kappa}.$$

The second condition in (3.7) and (3.8) imply $E_+(L) < E_-(L)$. Hence $\lambda_1(\Omega_L) = E_+(L)$ and $\lambda_2(\Omega_L) = E_-(L)$. Equations (3.4) and (3.5) now follow from Lemma 3.2. Subtracting the two expansions and multiplying by L^κ proves (3.6), since $\kappa > 0$. ■

Corollary 3.4 (*Rate for the distant-ball minimizing sequence*). *Let $D = B_R(0)$ and let Ω_L be the union of two radius- R balls whose centers are separated by L , so that $|\Omega_L| = 2|B_R|$. Then, as $L \rightarrow \infty$,*

$$L^{d+2s}(\lambda_2(\Omega_L) - \lambda_1(B_R)) \longrightarrow c_{d,s} \left(\int_{B_R} \phi_1(x) dx \right)^2.$$

Thus the minimizing sequence in [BP16, Theorem 6.2] approaches its limiting value at the explicit leading order L^{-d-2s} .

Proof. This is (3.5) with $D = B_R(0)$ and $\mu_1 = \lambda_1(B_R)$. ■

4. Finitely many wells

The two-well reflection decomposition is special to a pair. For finitely many components we instead keep the full block operator and compare its lowest spectral cluster with its compression to the translated one-well ground states.

Throughout this section, let $D \subset B_R(0)$ be any bounded Lipschitz domain; central symmetry is not assumed. The compactness and ground-state results [DNPV12, Theorems 5.4 and 7.1] and [BP16, Theorem 2.8] apply unchanged. Write $\mu_1 < \mu_2$, $g = \mu_2 - \mu_1 > 0$, choose the positive normalized ground state $A_D \phi_1 = \mu_1 \phi_1$, and set $m_1 = \int_D \phi_1(x) dx > 0$.

Definition 4.1 (Multi-well geometry). Fix an integer $N \geq 2$ and distinct points $\mathbf{a} = (a_1, \dots, a_N)$ in $\mathbb{R}^d$. Set

$$\delta_{\mathbf{a}} = \min_{i \neq j} |a_i - a_j|, \quad \Sigma_p(\mathbf{a}) = \max_{1 \leq i \leq N} \sum_{j \neq i} |a_i - a_j|^{-p} \quad (p > 0). \quad (4.1)$$

For $L\delta_{\mathbf{a}} > 2R$, define

$$D_{j,L} = D + La_j, \quad \Omega_{\mathbf{a},L} = \bigcup_{j=1}^N D_{j,L}. \quad (4.2)$$

The sets in this union are pairwise disjoint.

On $\mathcal{H}_N = \bigoplus_{j=1}^N L^2(D)$, let $\mathcal{A}_0 = \bigoplus_{j=1}^N A_D$. For $i \neq j$ define

$$(B_{ij,L}v)(x) = -c_{d,s} \int_D \frac{v(y)}{|L(a_i - a_j) + x - y|^\kappa} dy, \quad B_{ii,L} = 0, \quad (4.3)$$

and let $\mathcal{V}_L = (B_{ij,L})_{i,j=1}^N$.

Lemma 4.2 (Exact multi-well reduction and norm bound). If $L\delta_{\mathbf{a}} \geq 4R$, then $A_{\Omega_{\mathbf{a},L}}$ is unitarily equivalent to $\mathcal{A}_0 + \mathcal{V}_L$. Moreover,

$$\|\mathcal{V}_L\| \leq \gamma_L, \quad \gamma_L = c_{d,s} 2^\kappa |D| \Sigma_\kappa(\mathbf{a}) L^{-\kappa}. \quad (4.4)$$

Proof. For $f \in L^2(\Omega_{\mathbf{a},L})$, put $u_j(x) = f(x + La_j)$ for $x \in D$. This is a unitary identification with $\mathcal{H}_N$. Translation invariance gives the diagonal operator $\mathcal{A}_0$. The cutoff argument in the proof of Lemma 2.1 identifies the form domain with $\bigoplus_{j=1}^N \tilde{H}^s(D)$, and positive separation makes every cross-kernel bounded. For $i < j$, direct expansion of the form (2.1) gives the cross term

$$-2c_{d,s} \operatorname{Re} \iint_{D \times D} \frac{u_i(x) \overline{u_j(y)}}{|L(a_i - a_j) + x - y|^\kappa} dx dy.$$

Since $B_{ji,L} = B_{ij,L}^*$, summing these terms proves the exact block representation in the sense of closed forms and hence as operators.

For $i \neq j$ and $x, y \in D$,

$$|L(a_i - a_j) + x - y| \geq L|a_i - a_j| - 2R \geq \frac{L}{2}|a_i - a_j|.$$

The Schur test therefore gives

$$\|B_{ij,L}\| \leq c_{d,s} 2^\kappa |D| L^{-\kappa} |a_i - a_j|^{-\kappa}. \quad (4.5)$$

For $u = (u_1, \dots, u_N)$, set $r_j = \|u_j\|_{L^2(D)}$ and let G be the symmetric matrix with entries $G_{ij} = \|B_{ij,L}\|$. Then

$$\|\mathcal{V}_L u\|_{\mathcal{H}_N} \leq \|Gr\|_{\ell^2} \leq \left(\max_i \sum_j G_{ij} \right) \|r\|_{\ell^2}.$$

Combining this inequality with (4.5) proves (4.4). ■

We next record the finite-multiplicity perturbation estimate that replaces Lemma 3.1.

Lemma 4.3 (*Isolated spectral-cluster bound*). *Let A be self-adjoint with compact resolvent. Suppose its lowest eigenvalue μ has multiplicity N , let P be the corresponding spectral projection, and suppose*

$$A|_{P^\perp} \geq \mu + g$$

in the quadratic-form sense for some $g > 0$. Let W be bounded and self-adjoint, set $b = \|W\|$, and assume $b \leq g/4$. If $\eta_1 \leq \dots \leq \eta_N$ are the eigenvalues of PWP on $\text{ran } P$, then, for $1 \leq k \leq N$,

$$-\frac{2b^2}{g} \leq \lambda_k(A + W) - (\mu + \eta_k) \leq 0. \quad (4.6)$$

In addition,

$$\lambda_{N+1}(A + W) \geq \mu + g - b \geq \mu + \frac{3g}{4}. \quad (4.7)$$

Proof. The min-max principle applied to the span of the first k eigenvectors of PWP gives the upper bound in (4.6). For the lower bound, write $\psi = p + q$ with $p = P\psi$ and $q = (I - P)\psi$. Then

$$\begin{aligned} \langle \psi, (A + W)\psi \rangle &\geq \mu \|\psi\|^2 + \langle p, PWPp \rangle + (g - b)\|q\|^2 - 2b\|p\|\|q\| \\ &\geq \mu \|\psi\|^2 + \langle p, PWPp \rangle - \frac{2b^2}{g}\|p\|^2 + \left(\frac{g}{2} - b\right)\|q\|^2. \end{aligned}$$

The second line uses $2b\|p\|\|q\| \leq 2b^2g^{-1}\|p\|^2 + (g/2)\|q\|^2$. Thus, in the form sense, $A + W$ dominates the direct sum of

$$\mu I + PWP - \frac{2b^2}{g}I \quad \text{on } \text{ran } P$$

and $(\mu + g/2 - b)I$ on $P^\perp$. Since $\eta_N \leq b \leq g/4$, we have

$$\eta_N - \frac{2b^2}{g} \leq b - \frac{2b^2}{g} < \frac{g}{2} - b.$$

The strict inequality follows because the difference between the last two quantities is $(g - 2b)^2/(2g) > 0$. Hence the first N eigenvalues of this comparison operator are the N eigenvalues of its P -block. Another application of the min-max principle gives the lower bound in (4.6). Finally, $A + W \geq A - bI$ and the min-max principle give (4.7). $\blacksquare$

Let P_1 be the orthogonal projection in $\mathcal{H}_N$ onto the span of the N vectors whose j th component is ϕ_1 and whose other components vanish. In this basis the compression $P_1 \mathcal{V}_L P_1$ is represented by the real symmetric matrix $T_L = (t_{ij}(L))$, where

$$t_{ii}(L) = 0, \quad t_{ij}(L) = -c_{d,s} \iint_{D \times D} \frac{\phi_1(x)\phi_1(y)}{|L(a_i - a_j) + x - y|^\kappa} dx dy \quad (i \neq j). \quad (4.8)$$

Definition 4.4 (Effective interaction matrix). Define the real symmetric $N \times N$ matrix $M_{\mathbf{a}}$ by

$$(M_{\mathbf{a}})_{ii} = 0, \quad (M_{\mathbf{a}})_{ij} = -c_{d,s} m_1^2 |a_i - a_j|^{-\kappa} \quad (i \neq j). \quad (4.9)$$

Write its ordered eigenvalues as $\theta_1 \leq \dots \leq \theta_N$.

Lemma 4.5 (Compression error). If $L\delta_{\mathbf{a}} \geq 4R$, then

$$\|T_L - L^{-\kappa} M_{\mathbf{a}}\|_{\ell^2 \rightarrow \ell^2} \leq \varepsilon_L, \quad \varepsilon_L = c_{d,s} C_{\kappa,R} m_1^2 \Sigma_{\kappa+2}(\mathbf{a}) L^{-\kappa-2}. \quad (4.10)$$

Proof. Fix $i \neq j$, put $a = a_i - a_j$, and set $F(z) = |La + z|^{-\kappa}$. For $|z| \leq 2R$, the segment from 0 to z stays at distance at least $L|a|/2$ from $-La$. As in the proof of Lemma 3.2, Taylor's theorem gives

$$|F(z) - F(0) - DF(0)z| \leq C_{\kappa,R} L^{-\kappa-2} |a|^{-\kappa-2}. \quad (4.11)$$

The linear term vanishes after integration because

$$\iint_{D \times D} \phi_1(x)\phi_1(y)(x - y) dx dy = 0.$$

This cancellation is algebraic and does not require central symmetry of D . Using $F(0) = L^{-\kappa}|a|^{-\kappa}$ in (4.8) and integrating (4.11) gives

$$|t_{ij}(L) - L^{-\kappa}(M_{\mathbf{a}})_{ij}| \leq c_{d,s} C_{\kappa,R} m_1^2 L^{-\kappa-2} |a_i - a_j|^{-\kappa-2}.$$

The operator norm of a real symmetric matrix is at most its largest absolute row sum. Taking that row sum proves (4.10). $\blacksquare$

Theorem 4.6 (Finite multi-well effective Hamiltonian). Let D , μ_1 , g , ϕ_1 , and m_1 be as fixed at the beginning of this section, and let the multi-well geometry be as in Definition 4.1. If

$$L\delta_{\mathbf{a}} \geq 4R, \quad \gamma_L \leq \frac{g}{4}, \quad (4.12)$$

then, for $1 \leq k \leq N$,

$$|\lambda_k(\Omega_{\mathbf{a},L}) - (\mu_1 + L^{-\kappa}\theta_k)| \leq \varepsilon_L + \frac{2\gamma_L^2}{g}. \quad (4.13)$$

The cluster is separated from the rest of the spectrum by

$$\lambda_{N+1}(\Omega_{\mathbf{a},L}) \geq \mu_1 + g - \gamma_L \geq \mu_1 + \frac{3g}{4}. \quad (4.14)$$

Consequently, as $L \rightarrow \infty$, for every $1 \leq k \leq N$,

$$L^{d+2s}(\lambda_k(\Omega_{\mathbf{a},L}) - \mu_1) \longrightarrow \theta_k. \quad (4.15)$$

Proof. Lemma 4.2 identifies the full operator with $\mathcal{A}_0 + \mathcal{V}_L$. The lowest eigenvalue of $\mathcal{A}_0$ is μ_1 , with multiplicity N , its spectral projection is P_1 , and the rest of its spectrum lies in $[\mu_1 + g, \infty)$. Apply Lemma 4.3 with $A = \mathcal{A}_0$, $W = \mathcal{V}_L$, and $b = \|\mathcal{V}_L\| \leq \gamma_L$. If $\tau_1(L) \leq \dots \leq \tau_N(L)$ are the eigenvalues of T_L , this gives

$$-\frac{2\gamma_L^2}{g} \leq \lambda_k(\Omega_{\mathbf{a},L}) - (\mu_1 + \tau_k(L)) \leq 0. \quad (4.16)$$

The finite-dimensional min-max principle and Lemma 4.5 give

$$|\tau_k(L) - L^{-\kappa}\theta_k| \leq \varepsilon_L.$$

Combining these two estimates proves (4.13), and (4.14) follows from (4.7). Finally, $L^\kappa \varepsilon_L = O(L^{-2})$ and $L^\kappa \gamma_L^2 = O(L^{-\kappa})$; since $\kappa = d + 2s > 0$, multiplying (4.13) by L^κ proves (4.15). $\blacksquare$

Corollary 4.7 (*Recovery of the doublet*). Assume in addition that $D = -D$. For $N = 2$, $a_1 = -e/2$, and $a_2 = e/2$, the eigenvalues of $M_{\mathbf{a}}$ are $-c_{d,s}m_1^2$ and $c_{d,s}m_1^2$. The bound (4.13) then agrees with (3.4)–(3.5).

Proof. In this case

$$M_{\mathbf{a}} = \begin{pmatrix} 0 & -c_{d,s}m_1^2 \\ -c_{d,s}m_1^2 & 0 \end{pmatrix},$$

whose ordered eigenvalues are the two stated numbers. Moreover, $\Sigma_\kappa(\mathbf{a}) = \Sigma_{\kappa+2}(\mathbf{a}) = 1$, so $\gamma_L = \beta_L$. In the exact compression, central symmetry of D and evenness of ϕ_1 show, after the change of variables $y \mapsto -y$, that the translated two-well kernel integral equals I_L . Thus the compression errors, and hence the two displayed remainder bounds, coincide. $\blacksquare$

Corollary 4.8 (*Collective ground state*). For every multi-well configuration, $\theta_1 < 0$ is simple and has an eigenvector with strictly positive entries. Moreover, $\theta_N > 0$. Consequently, for all sufficiently large L , the lowest eigenvalue in the cluster is simple and lies below μ_1 , while at least one cluster eigenvalue lies above μ_1 .

Proof. The off-diagonal entries of $M_{\mathbf{a}}$ are strictly negative. Its Rayleigh quotient at $(1, \dots, 1)$ is negative, so $\theta_1 < 0$. If z is a minimizer of the Rayleigh quotient, replacing z by its componentwise absolute value cannot increase that quotient, and equality forces all nonzero components to have the same sign. The eigenvalue equation then shows that no component can vanish. Thus every eigenvector for θ_1 has one strict sign. Two linearly independent such eigenvectors would have a nonzero linear combination with a zero component, so θ_1 is simple. Finally, $\text{tr } M_{\mathbf{a}} = 0$ and $\theta_1 < 0$, hence $\theta_N > 0$. Put

$$r_L = \varepsilon_L + \frac{2\gamma_L^2}{g}.$$

Theorem 4.6 gives $L^\kappa r_L \rightarrow 0$. Since θ_1 is simple, $\theta_2 - \theta_1 > 0$. Choose L so large that

$$L^\kappa r_L < \min \left\{ \frac{\theta_2 - \theta_1}{2}, -\theta_1, \theta_N \right\}.$$

Then (4.13) gives $\lambda_2 - \lambda_1 > 0$, $\lambda_1 < \mu_1$, and $\lambda_N > \mu_1$, which proves all three assertions for the full operator. $\blacksquare$

Corollary 4.9 (*Regular-simplex cluster*). *Suppose the centers satisfy $|a_i - a_j| = r$ for all $i \neq j$, as for the vertices of a regular simplex. Such a configuration in $\mathbb{R}^d$ necessarily has $N \leq d + 1$. Put*

$$w = c_{d,s} m_1^2 r^{-\kappa}.$$

Then $\theta_1 = -(N-1)w$ and $\theta_2 = \dots = \theta_N = w$. Thus one collective level is lowered by $(N-1)wL^{-\kappa}$, while the remaining $N-1$ levels are raised by $wL^{-\kappa}$, up to the remainder in (4.13).

Proof. After translating a_N to the origin, the $N-1$ vectors $a_i - a_N$ have Gram matrix with diagonal entries r^2 and off-diagonal entries $r^2/2$. This matrix is positive definite, so $N-1 \leq d$. Here $M_{\mathbf{a}} = -w(J - I)$, where J is the all-ones matrix. The vector $(1, \dots, 1)$ has eigenvalue $-(N-1)w$, and its orthogonal complement has eigenvalue w . $\blacksquare$

5. A reproducible interval check

We finish with a conforming fixed-mesh Galerkin check of the two- and three-well limits. It is included as a reproducibility test of the formulas, not as an input to their proofs. Take $d = 1$, $0 < s < 1/2$, and partition each unit interval into cells of length h . Proposition 5.1 shows directly that cell indicators have finite form energy in this range, so they span a conforming finite-dimensional subspace of the form domain.

Proposition 5.1 (*Exact cell-integral matrix*). *Let $I_1, \dots, I_M$ be pairwise disjoint cells of length h , set $G = \bigcup_{\alpha=1}^M I_\alpha$, let x_α be their centers, put $r_{\alpha\beta} = |x_\alpha - x_\beta|$, and set $q = 1 - 2s$. In the $L^2(G)$ -orthonormal basis $e_\alpha = h^{-1/2} \mathbf{1}_{I_\alpha}$, the Galerkin matrix H_h of the zero-exterior*

form Π_G has entries

$$(H_h)_{\alpha\alpha} = \frac{c_{1,s}}{h} \frac{2h^q}{2sq}, \quad (5.1)$$

$$(H_h)_{\alpha\beta} = -\frac{c_{1,s}}{h} \frac{2r_{\alpha\beta}^q - (r_{\alpha\beta} + h)^q - (r_{\alpha\beta} - h)^q}{2sq}, \quad \alpha \neq \beta. \quad (5.2)$$

Proof. For two disjoint cells with center distance $r \geq h$, direct integration twice gives

$$\iint_{I_\alpha \times I_\beta} |x - y|^{-1-2s} dx dy = \frac{2r^q - (r + h)^q - (r - h)^q}{2sq}.$$

The off-diagonal form entry is minus $c_{1,s}$ times this integral. For one cell,

$$\int_{I_\alpha} \int_{\mathbb{R} \setminus I_\alpha} |x - y|^{-1-2s} dy dx = \frac{2h^q}{2sq}.$$

Dividing both expressions by h for the normalized indicators proves (5.1)–(5.2). $\blacksquare$

Corollary 5.2 (*Fixed-mesh cluster asymptotic*). Let $D = (-1/2, 1/2)$, and fix a uniform partition of D . Let $\mu_{1,h}$ and $\phi_{1,h}$ be the lowest eigenvalue and a positive, L^2 -normalized piecewise-constant eigenfunction of its Galerkin matrix. Put $m_{1,h} = \int_D \phi_{1,h}$. For the corresponding Galerkin matrix on $\bigcup_{j=1}^N (D + La_j)$, write its ordered eigenvalues as $\lambda_{k,h}(L)$. Then, as $L \rightarrow \infty$, for fixed distinct $a_1, \dots, a_N$,

$$\lambda_{k,h}(L) = \mu_{1,h} + L^{-1-2s} \theta_{k,h} + O_{h,a,s}(L^{-3-2s} + L^{-2-4s}), \quad 1 \leq k \leq N, \quad (5.3)$$

where $\theta_{1,h} \leq \dots \leq \theta_{N,h}$ are the eigenvalues of the matrix with zero diagonal and off-diagonal entries $-c_{1,s} m_{1,h}^2 |a_i - a_j|^{-1-2s}$.

Proof. All off-diagonal entries of the one-well matrix in (5.2) are negative. The Rayleigh-quotient argument in the proof of Corollary 4.8 therefore gives a simple lowest eigenvalue and a strictly positive eigenvector. On the direct sum of N fixed Galerkin spaces, the block expansion used in Lemma 4.2 is exact, its off-diagonal norm is $O_{h,a,s}(L^{-1-2s})$, and the complement of the N copied ground states is separated by the positive discrete one-well gap. The Taylor argument in Lemma 4.5, now in Euclidean operator norm, gives an $O_{h,a,s}(L^{-3-2s})$ compression error. Applying Lemma 4.3 to this finite matrix gives the additional $O_{h,a,s}(L^{-2-4s})$ term and proves (5.3). $\blacksquare$

For the computation we take $s = 1/4$, $D = (-1/2, 1/2)$, and the standard normalization

$$c_{1,s} = \frac{4^s s \Gamma(1/2 + s)}{\sqrt{\pi} \Gamma(1 - s)}.$$

Table 1 records the one-well quantities. The stabilization of $m_{1,h}$ is the relevant check for the interaction coefficient.

On the 180-cell mesh, the two-well component centers are $(-L/2, L/2)$, while the three-well centers are $(-L, 0, L)$. The corresponding fixed-mesh effective spectra are

$$\begin{aligned} (-L/2, L/2) : & \quad (-0.18521477, 0.18521477), \\ (-L, 0, L) : & \quad (-0.29671332, 0.06548331, 0.23123001). \end{aligned}$$

cells	$\mu_{1,h}$	$\mu_{2,h} - \mu_{1,h}$	$m_{1,h}$
60	1.3794549573	0.9105539931	0.9628386433
120	1.3753692130	0.8998883318	0.9633505837
180	1.3741292905	0.8969647744	0.9636021778

Table 1. One-well Galerkin data for $s = 1/4$.

The direct Galerkin eigenvalues are shown in Table 2; the last column is the largest absolute difference from the corresponding effective-matrix eigenvalue.

L	$L^{3/2}(\lambda_{1,h} - \mu_{1,h})$	$L^{3/2}(\lambda_{2,h} - \mu_{1,h})$	max. error
3	-0.190740	0.189832	5.52×10^{-3}
6	-0.186544	0.186335	1.33×10^{-3}
12	-0.185551	0.185486	3.36×10^{-4}
24	-0.185286	0.185295	8.04×10^{-5}

L	$L^{3/2}(\lambda_{1,h} - \mu_{1,h})$	$L^{3/2}(\lambda_{2,h} - \mu_{1,h})$	$L^{3/2}(\lambda_{3,h} - \mu_{1,h})$	max. error
3	-0.304984	0.065562	0.237544	8.27×10^{-3}
6	-0.298738	0.065554	0.232747	2.03×10^{-3}
12	-0.297232	0.065510	0.231597	5.19×10^{-4}
24	-0.296843	0.065504	0.231323	1.29×10^{-4}

Table 2. Scaled two-well (top) and three-well (bottom) cluster shifts.

The scaled errors decrease to zero in both configurations, as asserted at fixed mesh by Corollary 5.2. The calculation uses the exact entries (5.1)–(5.2); no singular quadrature or fitted coefficient enters the tables.

AI-use disclosure

The proof architecture and manuscript were developed with assistance from OpenAI’s GPT-5.6 Sol model (`gpt-5.6-sol`). The model served as an interactive research assistant for argument development and stress testing, citation and calculation audits, preparation of exact verification code, and revision. The author directed the research, selected and checked the final arguments, and takes responsibility for all mathematical claims, citations, computations, and wording.

Code availability

The script `tools/check_asymptotics.py` reconstructs every entry in Tables 1 and 2. Before solving the eigenproblems, it checks the assembled constant-function energy identity and prints progress after each separation. The pinned numerical dependencies are listed in `requirements-numeric.txt`; the reference output was replayed with Python 3.12.3, NumPy 1.26.4, and SciPy 1.11.4. A frozen copy of the scripts, pinned dependencies, release audit, and reference output is included in the submission source pack-

age; this is the version corresponding to the manuscript. The immutable repository tag for this version is `paper-2026-09-01`. The development repository is available at github.com/skypher/laplace-tunneling.

References

- [AFN20] Laura Abatangelo, Veronica Felli, and Benedetta Noris. On simple eigenvalues of the fractional Laplacian under removal of small fractional capacity sets. *Communications in Contemporary Mathematics*, 22(8):1950071, 2020.
[doi:10.1142/S0219199719500718](https://doi.org/10.1142/S0219199719500718).
- [BDVV23] Stefano Biagi, Serena Dipierro, Enrico Valdinoci, and Eugenio Vecchi. A Hong–Krahn–Szegő inequality for mixed local and nonlocal operators. *Mathematics in Engineering*, 5(1):1–25, 2023.
[doi:10.3934/mine.2023014](https://doi.org/10.3934/mine.2023014).
- [BP16] Lorenzo Brasco and Enea Parini. The second eigenvalue of the fractional p -Laplacian. *Advances in Calculus of Variations*, 9(4):323–355, 2016.
[doi:10.1515/acv-2015-0007](https://doi.org/10.1515/acv-2015-0007).
- [DFW21] Sidy Moctar Djitte, Mouhamed Moustapha Fall, and Tobias Weth. A fractional Hadamard formula and applications. *Calculus of Variations and Partial Differential Equations*, 60(6):231, 2021.
[doi:10.1007/s00526-021-02094-3](https://doi.org/10.1007/s00526-021-02094-3).
- [DNPV12] Eleonora Di Nezza, Giampiero Palatucci, and Enrico Valdinoci. Hitchhiker’s guide to the fractional Sobolev spaces. *Bulletin des Sciences Mathématiques*, 136(5):521–573, 2012.
[doi:10.1016/j.bulsci.2011.12.004](https://doi.org/10.1016/j.bulsci.2011.12.004).
- [DPLSV26] Serena Dipierro, Edoardo Proietti Lippi, Caterina Sportelli, and Enrico Valdinoci. Nonlocal eigenvalue problems and superposition operators, 2026.
[arXiv:2602.18035v1 \[math.AP\]](https://arxiv.org/abs/2602.18035v1), 20 February 2026.
- [Kwa12] Mateusz Kwaśnicki. Eigenvalues of the fractional Laplace operator in the interval. *Journal of Functional Analysis*, 262(5):2379–2402, 2012.
[doi:10.1016/j.jfa.2011.12.004](https://doi.org/10.1016/j.jfa.2011.12.004).
- [PS20] Enea Parini and Ariel Salort. Compactness and dichotomy in nonlocal shape optimization. *Mathematische Nachrichten*, 293(11):2208–2232, 2020.
[doi:10.1002/mana.201800234](https://doi.org/10.1002/mana.201800234).
- [Wu26] Zijian Wu. Equilibrium configurations and stability analysis of a discrete toy model arising from the minimization of eigenvalues of the fractional Laplacian. *Applied Mathematics and Computation*, 531:130195, 2026.
[doi:10.1016/j.amc.2026.130195](https://doi.org/10.1016/j.amc.2026.130195).
- [Zah26] Alvis Zahl. Minimizing eigenvalues of the fractional Laplacian. *Calculus of Variations and Partial Differential Equations*, 65(3):98, 2026.
[doi:10.1007/s00526-026-03270-z](https://doi.org/10.1007/s00526-026-03270-z).
- [Zha26] Cheng Zhang. Eigenvalue asymptotics and uniform eigenfunction bounds for the fractional Laplacian in the interval, 2026.
[arXiv:2608.23457v1 \[math.CA\]](https://arxiv.org/abs/2608.23457v1), 24 August 2026.