

Solvability of nonlocal Dirichlet problem for generalized Helmholtz equation in a unit ball

Turmetov, B. Kh and Karachik, V. V.

Published 2022 | COMPLEX VARIABLES AND ELLIPTIC EQUATIONS

0  
Times  
Cited

## Complex Variables And Elliptic Equations

Journal Impact Factor™

2020 Five Year

0.846

0.831

JCR Category	Category Rank	Category Quartile
MATHEMATICS in SCIE edition	195/330	Q3

Source: Journal Citation Reports™ 2020

Close

0  
Times  
Cited

2  
Times  
Cited

## Complex Variables and Elliptic Equations

Годы охвата Scopus: с 2007 по настоящий момент

Издатель: Taylor & Francis

ISSN: 1747-6933 E-ISSN: 1747-6941

Отрасль знаний: [Mathematics: Analysis](#) [Mathematics: Applied Mathematics](#) [Mathematics: Numerical Analysis](#) [Mathematics: Computational Mathematics](#)

Тип источника: Журнал

[Просмотреть все документы >](#)

[Настроить уведомление о документах](#)

[Сохранить в список источников](#) [Source Homepage](#)

CiteScore 2020

1.4

SJR 2020

0.457

SNIP 2020

0.995

CiteScore CiteScore рейтинг и тренды Содержание Scopus

### Улучшенная методика расчета CiteScore

Рейтинг CiteScore 2020 отражает количество цитирований в 2017-2020 гг. статей, обзоров, материалов конференций, глав книг и информационных документов, опубликованных в 2017-2020 гг., деленное на количество публикаций за 2017-2020 гг. [Подробнее >](#)

CiteScore 2020

1.4 =  $\frac{668 \text{ цитирований за 2017 - 2020 гг.}}{482 \text{ документов за 2017 - 2020 гг.}}$

Вычисление выполнено 05 May, 2021


CiteScoreTracker 2021

1.6 =  $\frac{761 \text{ цитирований на текущую дату}}{488 \text{ документов на текущую дату}}$

Последнее обновление 06 April, 2022 • Обновляется ежемесячно

### Рейтинг CiteScore 2020

Категория	Рейтинг	Процентиль
Mathematics Analysis	#86/164	47-й
Mathematics Applied Mathematics	#342/548	37-й



# Complex Variables and Elliptic Equations

## An International Journal

ISSN: (Print) (Online) Journal homepage: <https://www.tandfonline.com/loi/gcov20>

## Solvability of nonlocal Dirichlet problem for generalized Helmholtz equation in a unit ball

B. Kh. Turmetov & V. V. Karachik

To cite this article: B. Kh. Turmetov & V. V. Karachik (2022): Solvability of nonlocal Dirichlet problem for generalized Helmholtz equation in a unit ball, Complex Variables and Elliptic Equations, DOI: [10.1080/17476933.2022.2040021](https://doi.org/10.1080/17476933.2022.2040021)

To link to this article: <https://doi.org/10.1080/17476933.2022.2040021>



Published online: 28 Feb 2022.



Submit your article to this journal [↗](#)




View related articles [↗](#)



View Crossmark data [↗](#)



# Solvability of nonlocal Dirichlet problem for generalized Helmholtz equation in a unit ball

B. Kh. Turmetov <sup>a</sup> and V. V. Karachik <sup>b</sup>

<sup>a</sup>Department of Mathematics, Khoja Akhmet Yassawi International Kazakh-Turkish University, Turkistan, Kazakhstan; <sup>b</sup>Department of Mathematical Analysis, South Ural State University, Chelyabinsk, Russia

## ABSTRACT

In this paper, we study the solvability of a new class of nonlocal boundary value problems for the generalized Helmholtz equation in the unit ball. These problems are a generalization of the classical Dirichlet boundary value problem to the Helmholtz equation. For the considered problems, existence and uniqueness theorems are proved. Integral representation of solutions is established. Corresponding spectral questions are also investigated, namely, eigenfunctions and eigenvalues of the problem are found.

## ARTICLE HISTORY

Received 10 February 2021  
Accepted 5 February 2022

## COMMUNICATED BY

M. Ruzhansky

## KEYWORDS

Nonlocal boundary value problem; Helmholtz equation; orthogonal matrix; involution; Dirichlet problem; eigenfunction

## AMS SUBJECT CLASSIFICATIONS

34K06; 35J05; 35J25

## 1. Introduction

The concept of a nonlocal operator and the related concepts of a nonlocal differential equation and a nonlocal boundary value problem appeared in mathematics relatively recently. For example, in [1] loaded equations, equations containing fractional derivatives of the desired function, equations with deviating arguments, in other words, equations in which the unknown function and its derivatives enter, generally speaking, for different values of arguments are called nonlocal differential equations. Nonlocal boundary value problems for elliptic equations, in which boundary conditions are specified in the form of a connection between values of an unknown function and its derivatives at different points of the domain boundary, are called Bitsadze–Samarskii type problems [2]. Numerous applications of nonlocal boundary value problems for elliptic equations to problems in physics, technology, and other branches of science are described in detail in [3,4]. Solvability of nonlocal boundary value problems for elliptic equations is discussed in [5–8]. Boundary value problems for second- and fourth-order elliptic equations with involution, as special cases of nonlocal problems, are considered in [9–14]. Some nonlocal boundary value problems for the biharmonic equation are discussed in [15].

In this paper, we investigate the solvability conditions for one class of boundary value problems for a generalized Helmholtz equation in the unit ball with nonlocality both in the equation and in the boundary conditions.

Let  $\Omega = \{x \in \mathbb{R}^n : |x| < 1\}$ ,  $n \geq 2$ , be the unit ball,  $\partial\Omega$  be the unit sphere and  $S$  be an orthogonal matrix  $S \cdot S^T = I$ . Moreover, suppose that there exists  $m \in \mathbb{N}$  such that  $S^m = I$ . Note that if  $x \in \Omega$ , or  $x \in \partial\Omega$ , then for any  $k \in \mathbb{N}$  the following inclusions  $S^k x \in \Omega$ , or  $S^k x \in \partial\Omega$  hold. This is so since the transformation of the space  $\mathbb{R}^n$  with the matrix  $S$  preserves the norm  $|x|^2 = (x, x) = (S^T S x, x) = (Sx, Sx) = |Sx|^2$ .

Let  $a_0, a_1, \dots, a_{m-1}$  and  $b_0, b_1, \dots, b_{m-1}$  be some real numbers,  $f(x)$  and  $g(x)$  be functions defined in  $\Omega$  and on  $\partial\Omega$ , respectively. Consider the following nonlocal boundary value problem for the generalized Helmholtz equation.

**Problem:** Find a function  $u(x) \in C^2(\Omega) \cap C(\bar{\Omega})$  satisfying the following conditions

$$-\sum_{k=0}^{m-1} a_k \Delta u(S^k x) = \lambda u(x) + f(x), \quad x \in \Omega, \quad (1)$$

$$\sum_{k=0}^{m-1} b_k u(S^k x)|_{\partial\Omega} = g(x), \quad x \in \partial\Omega. \quad (2)$$

If  $a_0 = b_0 = 1$ ,  $a_k = b_k = 0$ ,  $k = 1, 2, \dots, m-1$ , then we have the classical Dirichlet boundary value problem for the Helmholtz equation. If  $\lambda = 0$  and  $b_0 = 1$ ,  $b_k = 0$ ,  $k = 1, 2, \dots, m-1$ , then this problem is investigated in [16], and if  $\lambda = 0$  and  $a_0 = 1$ ,  $a_k = 0$ ,  $k = 1, 2, \dots, m-1$ , it is considered in [17].

## 2. Auxiliary statements

To study the above posed problem, we need some auxiliary results. The following statement is proved in Lemma 3.1 from [16].

**Lemma 2.1:** The operator  $I_S u(x) = u(Sx)$  and the Laplace operator  $\Delta$  commute  $\Delta I_S u(x) = I_S \Delta u(x)$  on the functions  $u(x) \in C^2(\Omega)$ . The operator  $\Lambda = \sum_{i=1}^n x_i u_{x_i}(x)$  and the operator  $I_S$  also commute  $\Lambda I_S u(x) = I_S \Lambda u(x)$  on the functions  $u(x) \in C^1(\bar{\Omega})$ .

**Lemma 2.2:** Let  $\mu_0, \mu_1, \dots, \mu_{m-1}$  be different  $m$ th roots of unity, then

$$M^{-1} \equiv \begin{pmatrix} \mu_0^0 & \mu_0^1 & \dots & \mu_0^{m-1} \\ \mu_1^0 & \mu_1^1 & \dots & \mu_1^{m-1} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_{m-1}^0 & \mu_{m-1}^1 & \dots & \mu_{m-1}^{m-1} \end{pmatrix}^{-1} = \frac{1}{m} \overline{M}^T.$$

**Proof:** Let us find element  $e_{ij}$  from  $i$ th row and  $j$ th column in the product of the matrix  $M$  and the matrix from the right-hand side of the equality above. We obtain

$$e_{ij} = \frac{1}{m} \sum_{k=0}^{m-1} \mu_i^k \bar{\mu}_j^k = \frac{1}{m} \sum_{k=0}^{m-1} (\mu_i \bar{\mu}_j)^k = \begin{cases} 0, & i \neq j \\ 1, & i = j \end{cases} = \delta_{ij},$$

since  $\mu_i \bar{\mu}_j$  is also  $m$ th root of unity, not equal to 1, if  $i \neq j$  and  $\mu_i \bar{\mu}_i = |\mu_i|^2 = 1$ . Therefore, the product of these matrices is the identity matrix. This proves the lemma.  $\blacksquare$

A similar result is established during the proof of Lemma 2.3 from [16]. Consider the following matrix

$$A = \begin{pmatrix} a_0 & a_1 & \dots & a_{m-1} \\ a_{m-1} & a_0 & \dots & a_{m-2} \\ \vdots & \vdots & \ddots & \vdots \\ a_1 & a_2 & \dots & a_0 \end{pmatrix}, \quad (3)$$

which is formed based on coefficients  $a_0, a_1, \dots, a_{m-1}$  from (1).

**Lemma 2.3:** *Let  $\mu$  be a  $m$ th root of unity and*

$$\sigma_\mu(A) = \sum_{k=0}^{m-1} a_k \bar{\mu}^k, \quad (4)$$

*then for the matrix  $A$  from (3), we have the equality*

$$\begin{pmatrix} a_0 & a_1 & \dots & a_{m-1} \\ a_{m-1} & a_0 & \dots & a_{m-2} \\ \vdots & \vdots & \ddots & \vdots \\ a_1 & a_2 & \dots & a_0 \end{pmatrix} \begin{pmatrix} \mu^0 \\ \mu^1 \\ \vdots \\ \mu^{m-1} \end{pmatrix} = \sigma_{\bar{\mu}}(A) \begin{pmatrix} \mu^0 \\ \mu^1 \\ \vdots \\ \mu^{m-1} \end{pmatrix},$$

*i.e. the vector  $\mathbf{m}_\mu = (\mu^0, \mu^1, \dots, \mu^{m-1})^T$  is an eigenvector of the matrix  $A$ , and the number  $\sigma_{\bar{\mu}}(A)$  is an eigenvalue corresponding to it.*

*If  $\mu_0, \mu_1, \dots, \mu_{m-1}$  are different  $m$ th roots of unity, then eigenvectors  $\mathbf{m}_{\mu_0}, \mathbf{m}_{\mu_1}, \dots, \mathbf{m}_{\mu_{m-1}}$  are linear independent and*

$$\det A = \prod_{k=0}^{m-1} (\mathbf{a} \cdot \mathbf{m}_{\mu_k}),$$

*where  $\mathbf{a} = (a_0, a_1, \dots, a_{m-1})^T$ .*

**Proof:** It is not hard to see that the element of the  $i$ th row in the product of the matrix  $A$  and the vector  $\mathbf{m}_\mu$  is equal to

$$\begin{aligned} A_i^{\text{row}} \cdot \mathbf{m}_\mu &= \sum_{k=0}^{i-1} a_{m+k-i} \mu^k + \sum_{k=i}^{m-1} a_{k-i} \mu^k = \mu^i \sum_{k=0}^{i-1} a_{m+k-i} \mu^{k-i} + \mu^i \sum_{k=i}^{m-1} a_{k-i} \mu^{k-i} \\ &= \mu^i \left( \sum_{k'=m-i}^{m-1} a_{k'} \mu^{k'} + \sum_{k'=0}^{m-1-i} a_{k'} \mu^{k'} \right) = \mu^i \sum_{k'=0}^{m-1} a_{k'} \mu^{k'} = \sigma_{\bar{\mu}}(A) \mu^i. \end{aligned}$$

Here, in the first sum from the second line, the index is replaced by  $m+k-i = k'$ , and it is taken into account that  $\mu^{k'-m} = \mu^{k'}$ . In the second sum, the index is replaced by  $k-i = k'$ . Consequently,  $A \mathbf{m}_\mu = \sigma_{\bar{\mu}}(A) \mathbf{m}_\mu$ . The first lemma's statement is proved.

Due to Lemma 2.2, the matrix  $M = (\mathbf{m}_{\mu_0}, \dots, \mathbf{m}_{\mu_{m-1}})^T$  is invertible, and hence nonsingular, and therefore its row rank is equal to  $m$ , that is, rows are linearly independent. Further, it is known that  $\det A = \sigma_{\bar{\mu}_0}(A) \cdots \sigma_{\bar{\mu}_{m-1}}(A)$  and since

$$\sigma_{\bar{\mu}}(A) = \sum_{k=0}^{m-1} a_k \mu^k = \mathbf{a} \cdot \mathbf{m}_{\mu}, \quad (5)$$

we get

$$\det A = \sigma_{\bar{\mu}_0}(A) \cdots \sigma_{\bar{\mu}_{m-1}}(A) = \prod_{k=0}^{m-1} (\mathbf{a} \cdot \mathbf{m}_{\mu_k}).$$

The lemma is proved. ■

**Lemma 2.4:** *Let matrix  $A$  be constructed based on the numbers  $a_0, a_1, \dots, a_{m-1}$ , and matrix  $B$  be constructed based on the numbers  $b_0, b_1, \dots, b_{m-1}$  analogically to (3). Then the matrices  $A$  and  $B$  commute  $AB = BA$  and the matrix  $AB$  has the structure of matrices the  $A$  and  $B$ . If  $\mu_0, \mu_1, \dots, \mu_{m-1}$  are different  $m$ th roots of unity and  $\sigma_{\mu_k}(A) \neq 0$ ,  $k = 0, 1, \dots, m-1$ , then*

$$A^{-1} = \frac{1}{m} \left( \sum_{k=0}^{m-1} \frac{1}{\mu_k^{j-i} \sigma_{\bar{\mu}_k}(A)} \right)_{i,j=\overline{0,m-1}}. \quad (6)$$

**Proof:** Let us write the matrices  $A$  and  $B$  in the form

$$A = (a_{j-i})_{i,j=\overline{0,m-1}}, \quad B = (b_{j-i})_{i,j=\overline{0,m-1}},$$

where the subindex  $j-i$  is taken by  $\text{mod } m$ . Therefore,

$$\begin{aligned} (AB)_{i,j} &= \sum_{k=0}^{m-1} a_{k-i} b_{j-k} = \sum_{k'=-i}^{m-1-i} a_{k'} b_{j-i-k'} = \sum_{k'=-i}^{-1} a_{k'} b_{j-i-k'} + \sum_{k'=0}^{m-1-i} a_{k'} b_{j-i-k'} \\ &= \sum_{k'=m-i}^{m-1} a_{k'} b_{j-i-k'} + \sum_{k'=0}^{m-1-i} a_{k'} b_{j-i-k'} = \sum_{k=0}^{m-1} a_k b_{j-i-k}. \end{aligned}$$

If we denote  $c_i = \sum_{k=0}^{m-1} a_k b_{i-k}$ , then we get  $(AB)_{i,j} = c_{j-i}$  and hence the matrix  $AB$  has the structure of the matrices  $A$  and  $B$ . Since for  $i = 0, 1, \dots, m-1$

$$\begin{aligned} (AB)_{0,i} &= c_i = \sum_{k=0}^i a_k b_{i-k} + \sum_{k=i+1}^{m-1} a_k b_{i+m-k} = \sum_{k=0}^i b_k a_{i-k} \\ &+ \sum_{k=i+1}^{m-1} b_k a_{i+m-k} = \sum_{k=0}^{m-1} b_k a_{i-k} = (BA)_{0,i} \end{aligned}$$

and because elements of the matrices  $AB$  and  $BA$  are completely determined by their first row, we get  $AB = BA$ .

Let us multiply the matrix  $A$  by the matrix given by equality (6). It is easy to see that, according to the formula (5), similarly to the last equality from Lemma 2.2, the following equalities hold

$$\begin{aligned} (AA^{-1})_{ij} &= \frac{1}{m} \sum_{l=0}^{m-1} a_{l-i} \sum_{k=0}^{m-1} \frac{1}{\mu_k^{j-l} \sigma_{\mu_k}(A)} \\ &= \frac{1}{m} \sum_{k=0}^{m-1} \frac{\mu_k^{i-j}}{\sigma_{\mu_k}(A)} \sum_{l=0}^{m-1} \mu_k^{l-i} a_{l-i} = \frac{1}{m} \sum_{k=0}^{m-1} \mu_k^{i-j} = \delta_{ij}, \end{aligned}$$

where  $i, j = 0, \dots, m-1$ . This yields the lemma's statement  $AA^{-1} = I$ . ■

### 3. Solvability conditions of the problem

**Theorem 3.1:** *Let the function  $u(x)$  be a solution to the nonlocal Dirichlet problem (1) and (2) and let  $\mu$  be some  $m$ th root of unity, and the coefficients  $a_0, a_1, \dots, a_{m-1}$  of Equation (1) be such that  $\sigma_\mu(A) \neq 0$ , and the coefficients  $b_0, b_1, \dots, b_{m-1}$  of condition (2) be such that  $\sigma_\mu(B) \neq 0$ . Then the function*

$$v_\mu(x) = \sum_{k=0}^{m-1} \mu^k u(S^k x), \quad (7)$$

having the smoothness  $v_\mu(x) \in C^2(\Omega) \cap C(\bar{\Omega})$ , is a solution to the following Dirichlet problem for the Helmholtz equation

$$-\Delta v_\mu(x) = \lambda_\mu v_\mu(x) + f_\mu(x), \quad x \in \Omega, \quad (8)$$

$$v_\mu(x)|_{\partial\Omega} = g_\mu(x), \quad x \in \partial\Omega, \quad (9)$$

where

$$\lambda_\mu = \frac{\lambda}{\sigma_\mu(A)}, \quad f_\mu(x) = \frac{1}{\sigma_\mu(A)} \sum_{k=0}^{m-1} \mu^k f(S^k x), \quad g_\mu(s) = \frac{1}{\sigma_\mu(B)} \sum_{k=0}^{m-1} \mu^k g(S^k s). \quad (10)$$

**Proof:** Let a function  $u(x)$  be a solution to the problem (1) and (2). Obviously in this case  $v_\mu(x) \in C^2(\Omega) \cap C(\bar{\Omega})$ . Consider the operator  $I_S u(x) = u(Sx)$ . By Lemma 2.1, operators  $I_S$  and  $\Delta$  commute. Therefore, it is easy to see that in  $\Omega$  we have the equalities

$$\begin{aligned} -\Delta (a_0 u(S^0 x) + a_1 u(S^1 x) + \dots + a_{m-1} u(S^{m-1} x)) &= \lambda u(x) + f(x), \\ -\Delta (a_{m-1} u(S^0 x) + a_0 u(S^1 x) + \dots + a_{m-2} u(S^{m-1} x)) &= \lambda u(Sx) + f(Sx), \\ &\dots \\ -\Delta (a_1 u(S^0 x) + a_2 u(S^1 x) + \dots + a_0 u(S^{m-1} x)) &= \lambda u(S^{m-1} x) + f(S^{m-1} x), \end{aligned}$$

or in the matrix form

$$-A\Delta U(x) = \lambda U(x) + F(x), \quad x \in \Omega, \quad (11)$$

where

$$U(x) = \begin{pmatrix} u(S^0x) \\ u(S^1x) \\ \vdots \\ u(S^{m-1}x) \end{pmatrix}, \quad F(x) = \begin{pmatrix} f(S^0x) \\ f(S^1x) \\ \vdots \\ f(S^{m-1}x) \end{pmatrix}.$$

Let  $\mathbf{m}_\mu = (\mu^0, \mu^1, \dots, \mu^{m-1})^T$ . If multiply the vector equality (11) scalarly by the vector  $\mathbf{m}_\mu$ , then we obtain

$$-(\mathbf{m}_\mu, A \Delta U(x)) = \lambda(\mathbf{m}_\mu, U(x)) + (\mathbf{m}_\mu, F(x))$$

and hence

$$-(A^T \mathbf{m}_\mu, \Delta U(x)) = \lambda(\mathbf{m}_\mu, U(x)) + (\mathbf{m}_\mu, F(x)).$$

Let us write the matrix  $A$  in the form  $A = (a_{j-i})_{i,j=\overline{0,m-1}}$ , where, similarly to Lemma 2.4, the indices  $j-i$  are taken by  $\text{mod } m$ . Then the  $i$ th element of the vector  $A^T \mathbf{m}_\mu$ , due to equality (4)  $\sigma_\mu(A) = \sum_{k=0}^{m-1} a_k \bar{\mu}^k$ , has the form

$$(A^T \mathbf{m}_\mu)_i = \sum_{k=0}^{m-1} a_{i-k} \mu^k = \mu^i \sum_{k=0}^{m-1} a_{i-k} \mu^{k-i} = \mu^i \sum_{k=0}^{m-1} a_k \mu^{-k} = \mu^i \sigma_\mu(A) = \sigma_\mu(A) (\mathbf{m}_\mu)_i$$

and then

$$-\sigma_\mu(A) \Delta(\mathbf{m}_\mu, U(x)) = \lambda(\mathbf{m}_\mu, U(x)) + (\mathbf{m}_\mu, F(x)). \quad (12)$$

Therefore, if we use the notations of (7) and (10)

$$v_\mu(x) = (\mathbf{m}_\mu, U(x)), \quad f_\mu(x) = \frac{1}{\sigma_\mu(A)} (\mathbf{m}_\mu, F(x)),$$

we have

$$-\sigma_\mu(A) \Delta v_\mu(x) = \lambda_\mu v_\mu(x) + f_\mu(x).$$

Thus, if  $\sigma_\mu(A) \neq 0$  and  $\lambda_\mu = \frac{\lambda}{\sigma_\mu(A)}$ , then for the functions  $v_\mu(x)$  we obtain Equation (8).

Let us investigate the boundary condition (9). Obviously, similarly to (11), from the boundary condition of the original problem (2) we can obtain the equality

$$BU(x)|_{\partial\Omega} = G(x), \quad x \in \partial\Omega,$$

where  $G(x) = (g(S^0x), g(S^1x), \dots, g(S^{m-1}x))^T$ . Hence, multiplying the obtained vector equality scalarly by the vector  $\mathbf{m}_\mu$  and acting similarly to Equation (12), we obtain

$$(\mathbf{m}_\mu, BU(x))|_{\partial\Omega} = (B^T \mathbf{m}_\mu, U(x))|_{\partial\Omega} = \sigma_\mu(B) v_\mu(x)|_{\partial\Omega} = (\mathbf{m}_\mu, G(x)).$$

Since  $\sigma_\mu(B) \neq 0$ , then we have

$$v_\mu(x)|_{\partial\Omega} = g_\mu(x), \quad x \in \partial\Omega,$$

i.e. the boundary condition (9) is satisfied. The theorem is proved. ■



**Remark 3.1:** If a solution to Equation (1) exists, then the function  $v_\mu(x)$  from (7) satisfying Equation (8) must also satisfy the equality

$$v_\mu(Sx) = \sum_{k=0}^{m-1} \mu^k u(S^{k+1}x) = \frac{1}{\mu} \sum_{k=0}^{m-1} \mu^{k+1} u(S^{k+1}x) = \bar{\mu} v_\mu(x).$$

Consider the operator  $\Lambda u = \sum_{k=1}^n \xi_k u_{\xi_k}$ , which on  $\partial\Omega$  has the property

$$\frac{\partial u}{\partial \nu} = \Lambda_\xi u, \quad (13)$$

where  $\nu$  is the unit outward normal to  $\partial\Omega$ .

**Lemma 3.2:** Let the function  $v_\mu(x)$  be a solution of the problem (8) and (9), where the functions  $f_\mu(x)$  and  $g_\mu(x)$  satisfy the conditions

$$f_\mu(Sx) = \bar{\mu} f_\mu(x), \quad x \in \Omega, \quad g_\mu(Sx) = \bar{\mu} g_\mu(x), \quad x \in \partial\Omega, \quad (14)$$

and  $\lambda_\mu$  is not an eigenvalue of the problem, i.e. the problem (8) and (9) has a unique solution. Then for the function  $v_\mu(x)$  we have the equality

$$v_\mu(Sx) = \bar{\mu} v_\mu(x). \quad (15)$$

**Proof:** First, note that since  $S^T = S^{-1} = S^{m-1}$  and  $\mu^{m-1} = 1/\mu = \bar{\mu}$ , then the equalities (14) yield

$$f_\mu(S^T x) = \bar{\mu}^{m-1} f_\mu(x) = \mu f_\mu(x), \quad x \in \Omega, \quad g_\mu(S^T x) = \mu g_\mu(x), \quad x \in \partial\Omega \quad (16)$$

and vice versa.

Let  $G(x, \xi)$  be the Green's function of the Dirichlet problem for the Poisson equation in the domain  $\Omega$

$$\Delta u(x) = f_1(x), \quad x \in \Omega, \quad u(x)|_{\partial\Omega} = g_1(x), \quad x \in \partial\Omega.$$

It is well known that the solution to this problem can be written in the form [18]

$$u(x) = -\frac{1}{\omega_n} \int_{\Omega} G(x, \xi) f_1(\xi) \, d\xi - \frac{1}{\omega_n} \int_{\partial\Omega} \frac{\partial G(x, \xi)}{\partial \nu_\xi} g_1(\xi) \, ds_\xi,$$

where  $\omega_n$  is an area of the unit sphere in  $\mathbb{R}^n$ . Therefore, from Equation (8), taking into account the equality (13), we obtain

$$v_\mu(x) = \frac{1}{\omega_n} \int_{\Omega} G(x, \xi) (\lambda_\mu v_\mu(\xi) + f_\mu(\xi)) \, d\xi - \frac{1}{\omega_n} \int_{\partial\Omega} g_\mu(\xi) \Lambda_\xi G(x, \xi) \, ds_\xi. \quad (17)$$

Since the number  $\lambda_\mu$  is not an eigenvalue of the Dirichlet problem (8) and (9), the solution of the resulting integral equation is unique. Let us prove that the function  $\mu v_\mu(Sx)$  also satisfies the integral Equation (17), and hence  $\mu v_\mu(Sx) = v_\mu(x)$ .

Consider the function

$$F(x) = \mu v_\mu(Sx) - \frac{1}{\omega_n} \int_{\Omega} G(x, \xi) (\lambda_\mu \mu v_\mu(S\xi) + f_\mu(\xi)) d\xi \\ + \frac{1}{\omega_n} \int_{\partial\Omega} g_\mu(\xi) \Lambda_\xi G(x, \xi) ds_\xi$$

and put here  $x = S^T x$ . Then we have

$$F(S^T x) = \mu v_\mu(x) - \frac{1}{\omega_n} \int_{\Omega} G(S^T x, \xi) (\lambda_\mu \mu v_\mu(S\xi) + f_\mu(\xi)) d\xi \\ + \frac{1}{\omega_n} \int_{\Omega} \Lambda_\xi G(S^T x, \xi) g_\mu(\xi) ds_\xi.$$

Make the change of variables  $y = S\xi$  under the integral sign which gives  $\xi = S^T y$ . Then, due to commutability of the operators  $\Lambda_\xi$  and  $I_S$  (see Lemma 2.1) and the equality  $\int_{\partial\Omega} \varphi(Sx) ds_x = \int_{\partial\Omega} \varphi(x) ds_x$  (see [15, Lemma 5.1]), we obtain

$$F(S^T x) = \mu v_\mu(x) - \frac{1}{\omega_n} \int_{\Omega} G(S^T x, S^T y) (\mu \lambda_\mu v_\mu(y) + f_\mu(S^T y)) \left| \frac{\partial \xi}{\partial y} \right| dy \\ + \frac{1}{\omega_n} \int_{\partial\Omega} g_\mu(S^T \xi) \Lambda_\xi G(S^T x, S^T \xi) ds_\xi.$$

Now, if we take into account the equalities (16) and also that  $|\frac{\partial \xi}{\partial y}| = |S^T| = 1$ , we obtain

$$F(S^T x) = \mu v_\mu(x) - \mu \frac{1}{\omega_n} \int_{\Omega} G(S^T x, S^T y) (\lambda_\mu v_\mu(y) + f_\mu(y)) dy \\ + \mu \frac{1}{\omega_n} \int_{\partial\Omega} g_\mu(\xi) \Lambda_\xi G(S^T x, S^T \xi) ds_\xi.$$

It is known that for  $\Omega$

$$G(x, \xi) = E(x, \xi) - E\left(\frac{x}{|x|}, |x|\xi\right),$$

where

$$E(x, \xi) = \begin{cases} \frac{1}{n-2} |x - \xi|^{2-n}, & n > 2 \\ -\ln |x - \xi|, & n = 2 \end{cases}$$

is an elementary solution to the Laplace equation. Then it is easy to see that due to orthogonality of the matrix  $S$  we have

$$E(S^T x, S^T y) = \frac{1}{n-2} |S^T x - S^T y|^{2-n} = \frac{1}{n-2} |x - y|^{2-n} = E(x, y), \\ E\left(\frac{S^T x}{|S^T x|}, |S^T x| S^T y\right) = E\left(\frac{S^T x}{|x|}, |x| S^T y\right)$$

$$= \frac{1}{n-2} \left| S^T \left( \frac{x}{|x|} - |x|y \right) \right|^{2-n} = E \left( \frac{x}{|x|}, |x|y \right),$$

if  $n > 2$ . This yields  $G(S^T x, S^T y) = G(x, y)$ . Therefore, due to (17), we get

$$\begin{aligned} F(S^T x) &= \mu \left( v_\mu(x) - \frac{1}{\omega_n} \int_{\Omega} G(x, y) (\lambda_\mu v_\mu(y) + f_\mu(y)) dy \right. \\ &\quad \left. + \frac{1}{\omega_n} \int_{\partial\Omega} g_\mu(\xi) \Lambda_\xi G(x, \xi) ds_\xi \right) = 0 \end{aligned}$$

for  $x \in \bar{\Omega}$  and hence, due to bijectivity of the mapping  $S : \bar{\Omega} \rightarrow \bar{\Omega}$ , we have  $F(x) = 0$  for  $x \in \bar{\Omega}$ . Thus, the function  $\mu v_\mu(Sx)$  is also a solution to the integral equation (17), and therefore, as noted above,  $\mu v_\mu(Sx) = v_\mu(x)$ . This immediately implies (15).  $\blacksquare$

The converse statement to Theorem 3.1 is also true.

**Theorem 3.3:** Let functions  $v_{\mu_0}(x), v_{\mu_1}(x), \dots, v_{\mu_{m-1}}(x) \in C^2(\Omega) \cap C(\bar{\Omega})$ , where  $\mu_0, \mu_1, \dots, \mu_{m-1}$  are different  $m$ th roots of unity, be solutions to the Dirichlet problems (8) and (9) in which

$$\sigma_{\mu_k}(A) = \sum_{i=0}^{m-1} a_i \bar{\mu}_k^i \neq 0, \quad \sigma_{\mu_k}(B) = \sum_{i=0}^{m-1} b_i \bar{\mu}_k^i \neq 0,$$

functions  $f_{\mu_k}(x)$  and  $g_{\mu_k}(x)$  be defined through the functions  $f(x)$  and  $g(x)$  by the equalities similar to (10)

$$f_{\mu_k}(x) = \frac{1}{\sigma_{\mu_k}(A)} \sum_{i=0}^{m-1} \mu_k^i f(S^i x), \quad g_{\mu_k}(x) = \frac{1}{\sigma_{\mu_k}(B)} \sum_{i=0}^{m-1} \mu_k^i g(S^i x).$$

Let moreover, the numbers  $\lambda_{\mu_k} = \lambda / \sigma_{\mu_k}(A)$  for  $k = 0, 1, \dots, m-1$  be not eigenvalues corresponding to the problem (8) and (9), then the function

$$u(x) = \frac{1}{m} \sum_{k=0}^{m-1} v_{\mu_k}(x) \tag{18}$$

is a solution to the problem (1) and (2). Under the assumptions made, the solution to the problem (1) and (2) is unique.

**Proof:** Let the function  $u(x)$  be taken in the form of (18). It is obvious that  $u(x) \in C^2(\Omega) \cap C(\bar{\Omega})$ . Further, it is easy to see that the functions  $f_{\mu_k}(x)$  and  $g_{\mu_k}(x)$  satisfy the equalities (14). Indeed, if  $\mu = \mu_k$ , we have

$$f_\mu(Sx) = \frac{1}{\sigma_\mu(A)} \sum_{i=0}^{m-1} \mu^i f(S^{i+1}x) = \frac{1}{\mu \sigma_\mu(A)} \sum_{i=0}^{m-1} \mu^i f(S^i x) = \bar{\mu} f_\mu(x), \quad x \in \Omega, \tag{19}$$

and similarly

$$g_\mu(Sx) = \bar{\mu} g_\mu(x), \quad x \in \partial\Omega.$$

Since  $\lambda_{\mu_k} = \lambda/\sigma_{\mu_k}(A)$  for  $k = 0, 1, \dots, m-1$  are not eigenvalues of the corresponding problem (8) and (9), conditions of Lemma 3.2 hold. By this lemma the equality (15)  $v_\mu(Sx) = \bar{\mu}v_\mu(x)$  holds. Thus, for  $x \in \Omega$  we have

$$u(S^j x) = \frac{1}{m} \sum_{k=0}^{m-1} \bar{\mu}_k^j v_{\mu_k}(x) \quad (20)$$

and, moreover, due to (19) we get

$$\Delta v_\mu(S^j x) = \lambda_\mu v_\mu(S^j x) + f_\mu(S^j x) = \bar{\mu}^j (\lambda_\mu v_\mu(x) + f_\mu(x)).$$

By using the equalities (20) obtained above, due to Equation (8), we find

$$\begin{aligned} & -\Delta \left( \sum_{j=0}^{m-1} a_j u(S^j x) \right) \\ &= -\sum_{j=0}^{m-1} a_j \frac{1}{m} \sum_{k=0}^{m-1} \bar{\mu}_k^j \Delta v_{\mu_k}(x) \\ &= \frac{\lambda}{m} \sum_{k=0}^{m-1} v_{\mu_k}(x) \frac{1}{\sigma_{\mu_k}(A)} \sum_{j=0}^{m-1} a_j \bar{\mu}_k^j + \frac{1}{m} \sum_{j=0}^{m-1} a_j \sum_{k=0}^{m-1} \bar{\mu}_k^j f_{\mu_k}(x) \\ &= \frac{\lambda}{m} \sum_{k=0}^{m-1} v_{\mu_k}(x) + \frac{1}{m} \sum_{k=0}^{m-1} \sigma_{\mu_k}(A) f_{\mu_k}(x) = \lambda u(x) + \frac{1}{m} \sum_{i=0}^{m-1} f(S^i x) \sum_{k=0}^{m-1} \mu_k^i \\ &= \lambda u(x) + f(x), \end{aligned}$$

i.e. Equation (1) is satisfied in  $\Omega$ .

Let us check the boundary condition (2). Using the equalities (20) and (9), we can write

$$\begin{aligned} \left. \sum_{j=0}^{m-1} b_j u(S^j x) \right|_{\partial\Omega} &= \left. \sum_{j=0}^{m-1} b_j \frac{1}{m} \sum_{k=0}^{m-1} \bar{\mu}_k^j v_{\mu_k}(x) \right|_{\partial\Omega} = \frac{1}{m} \sum_{j=0}^{m-1} b_j \sum_{k=0}^{m-1} \bar{\mu}_k^j g_{\mu_k}(x) \\ &= \frac{1}{m} \sum_{k=0}^{m-1} g_{\mu_k}(x) \sum_{j=0}^{m-1} b_j \bar{\mu}_k^j = \frac{1}{m} \sum_{k=0}^{m-1} \sigma_{\mu_k}(B) g_{\mu_k}(x) \\ &= \frac{1}{m} \sum_{k=0}^{m-1} \sum_{i=0}^{m-1} \mu_k^i g(S^i x) = \sum_{i=0}^{m-1} g(S^i x) \frac{1}{m} \sum_{k=0}^{m-1} \mu_k^i = g(x). \end{aligned}$$

Thus, the function  $u(x)$  defined by the formula (18) is the solution to the problem (1) and (2). Now suppose that the solution to the problem (1) and (2) is not unique. Then  $u(x)$  – the difference of two solutions to this problem is a solution to the problem (1) and (2) for  $f(x) = 0$ ,  $g(x) = 0$  and hence  $f_{\mu_k}(x) = 0$ ,  $g_{\mu_k}(x) = 0$ . Since  $\lambda_{\mu_k}$  are not eigenvalues to the corresponding problem (8) and (9), then  $v_{\mu_k}(x) = 0$  and hence according to (18) we get  $u(x) = 0$ . This proves the uniqueness of the solution to the problem (1) and (2). The theorem is proved. ■

#### 4. Eigenfunctions and eigenvalues of the nonlocal problem

By virtue of Theorem 3.3 proved above, a solution to the homogeneous problem (1) and (2) can be represented through the solutions of the homogeneous problem (8) and (9), which have only a discrete spectrum. Therefore, it is natural to investigate the existence of eigenfunctions of the homogeneous problem (1) and (2)

$$-\sum_{k=0}^{m-1} a_k \Delta u(S^k x) = \lambda u(x), \quad x \in \Omega, \quad (21)$$

$$\sum_{k=0}^{m-1} b_k u(S^k x) = 0, \quad x \in \partial\Omega. \quad (22)$$

**Theorem 4.1:** *Let  $\mu_0, \mu_1, \dots, \mu_{m-1}$  be different  $m$ th roots of unity, numbers  $a_0, a_1, \dots, a_{m-1}$  and  $b_0, b_1, \dots, b_{m-1}$  be such that  $\sigma_{\mu_k}(A) \neq 0$  and  $\sigma_{\mu_k}(B) \neq 0$  for  $k = 0, 1, \dots, m-1$ , where numbers  $\sigma_{\mu_k}(A)$  and  $\sigma_{\mu_k}(B)$  are defined in (4). Then  $\lambda = \lambda^d \sigma_{\mu_k}(A)$ , where  $\lambda^d$  is an eigenvalue of the homogeneous problem (8) and (9), is an eigenvalue of the problem (21) and (22) and the system of eigenfunctions of the problem (21) and (22) is complete in  $L_2(\Omega)$ .*

**Proof:** Assume that a solution of the problem (21) and (22) exists for some  $\lambda$ . Apply Theorem 3.3 to the problem (21) and (22) for  $\mu = \mu_k$ ,  $k = 0, 1, \dots, m-1$ . In this case from the formula (10) it follows that  $f_{\mu_k}(x) = g_{\mu_k}(x) = 0$ ,  $\lambda_{\mu_k} = \lambda / \sigma_{\mu_k}(A)$ , and the problem (8) and (9) is reduced to the problem

$$-\Delta v(x) = \lambda v(x), \quad x \in \Omega, \quad v(x) = 0, \quad x \in \partial\Omega, \quad (23)$$

solution of which  $-v_\mu(x)$  exists and is given by the formula (7). If  $v_\mu(x) \neq 0$ , then  $\lambda_{\mu_k}$  must coincide with some eigenvalue of the Dirichlet problem (23), i.e.  $\lambda_{\mu_k} = \lambda^{(k)}$ .

Let  $\lambda^d$  be an eigenvalue of the problem (23), and  $v(x)$  be an eigenfunction of this problem corresponding to it. It is clear that  $\lambda^d > 0$ . Denote

$$v_k(x) = \frac{1}{m} \sum_{i=0}^{m-1} \mu_k^i v(S^i x). \quad (24)$$

It is easy to see that, due to Lemma 2.1 and the equality  $S(\partial\Omega) = \partial\Omega$ , the functions  $v_k(x)$  for  $k = 0, 1, \dots, m-1$  are also eigenfunctions of the problem (23)

$$\begin{aligned} -\Delta v_k(x) &= -\frac{1}{m} \sum_{i=0}^{m-1} \mu_k^i (\Delta v)(S^i x) = \lambda^d \frac{1}{m} \sum_{i=0}^{m-1} \mu_k^i v(S^i x) = \lambda^d v_k(x), \\ v_k(x)|_{\partial\Omega} &= \frac{1}{m} \sum_{i=0}^{m-1} \mu_k^i v(S^i x)|_{\partial\Omega} = 0. \end{aligned}$$

The following representation takes place

$$v(x) = \sum_{k=0}^{m-1} v_k(x). \quad (25)$$

Indeed,

$$\sum_{k=0}^{m-1} v_k(x) = \sum_{k=0}^{m-1} \frac{1}{m} \sum_{i=0}^{m-1} \mu_k^i v(S^i x) = \frac{1}{m} \sum_{i=0}^{m-1} v(S^i x) \frac{1}{m} \sum_{k=0}^{m-1} \mu_k^i = v(x),$$

since

$$\sum_{k=0}^{m-1} \mu_k^i = \begin{cases} m, & i = 0 \\ 0, & i = 1, 2, \dots, m-1 \end{cases}.$$

Note that the formula (24) can give null functions (see Example 4.2 below). The functions  $v_k(x)$  also have the property

$$v_k(Sx) = \frac{1}{m} \sum_{i=0}^{m-1} \mu_k^i v(S^{i+1}x) = \bar{\mu}_k \frac{1}{m} \sum_{i=0}^{m-1} \mu_k^{i+1} v(S^{i+1}x) = \bar{\mu}_k v_k(x).$$

Therefore, since  $v_k(x)$  is the eigenfunction of the problem (23), we have

$$\begin{aligned} -\Delta \left( \sum_{j=0}^{m-1} a_j v_k(S^j x) \right) &= - \sum_{j=0}^{m-1} a_j \bar{\mu}_k^j \Delta v_k(x) \\ &= \lambda^d v_k(x) \sum_{j=0}^{m-1} a_j \bar{\mu}_k^j = \lambda^d \sigma_{\mu_k}(A) v_k(x), \end{aligned}$$

i.e. the function  $v_k(x)$  satisfies Equation (21) at  $\lambda = \lambda^d \sigma_{\mu_k}(A)$ . Moreover,

$$\sum_{i=0}^{m-1} b_i v_k(S^i x)|_{\partial\Omega} = v_k(x)|_{\partial\Omega} \sum_{i=0}^{m-1} b_i \bar{\mu}_k^i = \sigma_{\mu_k}(B) v_k(x)|_{\partial\Omega} = 0$$

and, hence  $v_k(x)$  is the eigenfunction of the problem (21) and (22) for  $\lambda = \lambda^d \sigma_{\mu_k}(A)$ . It is known (see, e.g. [19]), that the problem (23) has a complete in  $L_2(\Omega)$  orthogonal system of eigenfunctions  $\{v^{(j)}(x), j \in \mathbb{N}\}$  corresponding to the eigenvalues  $\{\lambda^{(j)}(x), j \in \mathbb{N}\}$ . By virtue of the formula (25), each such a function is a linear combination of the eigenfunctions  $v_{k,j}(x)$  of the problem (21) and (22) for  $\lambda = \lambda^{(j)} \sigma_{\mu_k}(A)$

$$v^{(j)}(x) = \sum_{k=0}^{m-1} v_{k,j}(x).$$

Therefore, the system of eigenfunctions  $\{v_{k,j}(x)\}$  of the problem (21) and (22) is complete in  $L_2(\Omega)$ .

Note that the functions  $v_k(x)$  and  $v_s(x)$  in the expansion (25) are orthogonal if  $k \neq s$ . Indeed, if  $k \neq s$  we have

$$\int_{\Omega} v_k(x) \overline{v_s(x)} dx = \frac{1}{m^2} \int_{\Omega} \sum_{i,j=0}^{m-1} \mu_k^i \bar{\mu}_s^j v(S^i x) \overline{v(S^j x)} dx$$

$$\begin{aligned}
&= \frac{1}{m^2} \sum_{i,j=0}^{m-1} \mu_k^i \bar{\mu}_s^j \int_{\Omega} v(S^i x) \bar{v}(S^j x) dx \\
&= \sum_{l=0}^{m-1} \sum_{i-j=l \pmod{m}}^{m-1} \frac{\mu_k^i \bar{\mu}_s^j}{m^2} \int_{\Omega} v(S^i x) \overline{v(S^j x)} dx \\
&= \sum_{l=0}^{m-1} \frac{\mu_k^l}{m^2} \sum_{j=0}^{m-1} (\mu_k \bar{\mu}_s)^j \int_{\Omega} v(S^{l+j} x) \overline{v(S^j x)} dx \\
&= \frac{1}{m} \sum_{l=0}^{m-1} \mu_k^l \int_{\Omega} v(S^l y) \overline{v(y)} dy \frac{1}{m} \sum_{j=0}^{m-1} (\mu_k \bar{\mu}_s)^j = 0,
\end{aligned}$$

since  $\det |S| = 1$  and  $\mu_k \bar{\mu}_s \neq 1$  is also  $m$ th root of unity. The theorem is proved. ■

**Remark 4.1:** Since the orthogonal system of eigenfunctions of the problem (23) has the form (see, for example, [20,21])

$$v^{(\lambda_i, k, j)}(x) = g_{n+2k}(\lambda_i |x|^2) H_k^{(j)}(x) \equiv \frac{1}{|x|^{n/2-1}} J_{k+n/2-1}(\sqrt{\lambda_i} |x|) H_k^{(j)}\left(\frac{x}{|x|}\right), \quad j \in \mathbb{N}_0,$$

where  $J_\nu(x)$  is the Bessel function of the first kind,  $\sqrt{\lambda_i}$  is a root of Bessel function  $J_\nu(x)$ ,  $\{H_k^{(j)}(x)\}$  is a system of orthogonal on  $\partial\Omega$  homogeneous harmonic polynomials of order  $k$  (see, for example, [22]), and  $|x| = |Sx|$ , then the expansion (25) rather refers to the expansion of the homogeneous harmonic polynomial  $H_k(x)$ .

**Example 4.2:** Consider the case  $S = -I$  and  $a_0 \neq \pm a_1$ ,  $b_0 = 1$ ,  $b_1 = 0$ , i.e. consider the problem

$$a_0 \Delta u(x) + a_1 \Delta u(-x) = \lambda u(x), \quad x \in \Omega, \quad u(x)|_{\partial\Omega} = 0. \quad (26)$$

In this case  $m = 2$ ,  $\mu_0 = 1$ ,  $\mu_1 = -1$ , and by the formula (4) we get  $\sigma_{\mu_0}(A) = a_0 + a_1$ ,  $\sigma_{\mu_1}(A) = a_0 - a_1$ . Let  $v(x)$  be a solution to the problem (23)

$$\Delta v(x) + \mu u(x) = 0, \quad x \in \Omega, \quad v(x)|_{\partial\Omega} = 0. \quad (27)$$

Then  $\mu > 0$ , and the eigenfunction of this problem

$$v(x) = \frac{1}{|x|^{n/2-1}} J_{k+n/2-1}(\sqrt{\mu} |x|) H_k\left(\frac{x}{|x|}\right)$$

is either even or odd depending on the parity of  $k \in \mathbb{N}_0$  (see [13], Lemma 1)). Therefore, only one of the functions given by the equalities (24)

$$u_0(x) = \frac{1}{2} [v(x) + v(-x)], \quad u_1(x) = \frac{1}{2} [v(x) - v(-x)].$$

is nonzero. Theorem 4.1 implies, but it is easy to verify directly, that these functions are eigenfunctions of the problem (26). If  $H_k(x)$  is an even polynomial, then  $u_0(x) = v(x)$  is an

eigenfunction corresponding to the eigenvalue  $\lambda_0 = \mu\sigma_{\mu_0}(A) = \mu(a_0 + a_1)$  and  $u_1(x) = 0$ . If  $H_k(x)$  is an odd polynomial, then  $u_1(x) = v(x)$  is an eigenfunction corresponding to the eigenvalue  $\lambda_1 = \mu\sigma_{\mu_1}(A) = \mu(a_0 - a_1)$ . Thus, the eigenfunctions of the problem (26) coincide with the eigenfunctions of the problem (27), and therefore the system of eigenfunctions of the nonlocal problem (26) is orthogonal and complete in  $L_2(\Omega)$ .

**Remark 4.2:** Using the results of [23] and Remark 4.1, we can be convinced of the statement similar to Example 4.2: The eigenfunctions of the problem (21) and (22) coincide with the eigenfunctions of the problem (23), and therefore the system of eigenfunctions of the nonlocal problem (21) and (22) is a basis in  $L_2(\Omega)$ .

## Disclosure statement

No potential conflict of interest was reported by the author(s).

## Funding

The work was supported by a grant from the Ministry of Science and Education of the Republic of Kazakhstan (grant no. AP08855810).

## ORCID

B. Kh. Turmetov  <http://orcid.org/0000-0001-7735-6484>

V. V. Karachik  <http://orcid.org/0000-0002-3077-3595>

## References

- [1] Nahushev AM. Equations of mathematical biology. Moscow (M): Publishing house for foreign literature; 1995. (Russian)SEP.
- [2] Bitsadze AV, Samarskii AA. Some elementary generalizations of linear elliptic boundary value problems. Dokl Akad Nauk SSSR. 1996;185:739–740. (in Russian).
- [3] Skubachevskii AL. Nonclassical boundary value problems. I. J Math Sci. 2008;155:199–334.
- [4] Skubachevskii AL. Nonclassical boundary-value problems. II. J Math Sci. 2010;166:377–561.
- [5] Criado F, Criado FJ, Odishelidze N. On the solution of some non-local problems. Czech Math J. 2004;54:487–498.
- [6] Kadirkulov BJ, Kirane M. On solvability of a boundary value problem for the Poisson equation with a nonlocal boundary operator. Acta Math Sci. 2015;35:970–980.
- [7] Kirane M, Torebek BT. On a nonlocal problem for the Laplace equation in the unit ball with fractional boundary conditions. Math Methods Appl Sci. 2016;39:1121–1128.
- [8] Kishkis KY. On some nonlocal problem for harmonic functions in multiply connected domain. Differ Equ. 1987;23:174–177.
- [9] Przeworska-Rolewicz D. Some boundary value problems with transformed argument. Comment Math. 1974;17:451–457.
- [10] Karachik VV, Turmetov BKh. On solvability of some Neumann-type boundary value problems for biharmonic equation. Electron J Differential Equations. 2017;218(2017):1–17.
- [11] Sadybekov MA, Dukenbayeva AA. On boundary value problems of the Samarskii-Ionkin type for the Laplace operator in a ball. Complex Var Elliptic Equations. 2020;1–15. doi:10.1080/17476933.2020.1828377
- [12] Sadybekov MA, Dukenbayeva AA. Direct and inverse problems for the Poisson equation with equality of flows on a part of the boundary. Complex Var Elliptic Equations. 2019;64:777–791.
- [13] Sadybekov MA, Turmetov BKh. On analogues of periodic boundary value problems for the Laplace operator in ball. Eurasian Math J. 2012;3:143–146.



- [14] Sadybekov MA, Turmetov BKh. On an analog of periodic boundary value problems for the Poisson equation in the disk. *Differ Equ.* [2014](#);50:268–273.
- [15] Karachik VV, Turmetov BKh. On solvability of some nonlocal boundary value problems for biharmonic equation. *Math Slovaca.* [2020](#);70:329–342.
- [16] Karachik VV, Sarsenbi AM, Turmetov BKh. On the solvability of the main boundary value problems for a nonlocal Poisson equation. *Turkish J Math.* [2019](#);43:1604–1625.
- [17] Karachik VV, Turmetov BKh. Solvability of one nonlocal Dirichlet problem for the Poisson equation. *Novi Sad J Math.* [2020](#);50:67–88.
- [18] Evans LC. *Partial differential equations*. Vol. 19. Providence (RI): American Mathematical Society; [1998](#). (Graduate Studies in Mathematics).
- [19] Gilbarg D, Trudinger NS. *Elliptic partial differential equations of second order*. Berlin: Springer-Verlag; [1986](#).
- [20] Karachik VV. Normalized system of functions with respect to the Laplace operator and its applications. *J Math Anal Appl.* [2003](#);287:577–592.
- [21] Karachik VV, Antropova NA. On the solution of the inhomogeneous polyharmonic equation and the inhomogeneous Helmholtz equation. *Differ Equ.* [2010](#);46:387–399.
- [22] Karachik VV. On some special polynomials. *Proc Amer Math Soc.* [2004](#);132:1049–1058.
- [23] Turmetov B, Karachik V. On eigenfunctions and eigenvalues of a nonlocal Laplace operator with multiple involution. *Symmetry.* [2021](#);13(10):1781.