

# MCAC 201: Design and Analysis of Algorithms

Neelima Gupta

ngupta@cs.du.ac.in

April 17, 2023

# Asymptotic Notations 1

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

# Asymptotic Notations 1

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .

# Asymptotic Notations 1

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

# Asymptotic Notations 1

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

# Asymptotic Notations 1

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

- ▶ **Notation:**  $f(n) \leq g(n)$  asymptotically.

# Asymptotic Notations 1

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

- ▶ **Notation:**  $f(n) \leq g(n)$  asymptotically. Note: By asymptotically we mean within constant multiplicative factor and for large  $n$

# Asymptotic Notations 1

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

- ▶ **Notation:**  $f(n) \leq g(n)$  asymptotically. Note: By asymptotically we mean within constant multiplicative factor and for large  $n$
- ▶ **Examples:**

$$1. \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{7n^2 - 2n - 5}{5n^2 + 3n + 10} = \lim_{n \rightarrow \infty} \frac{7 - \frac{2}{n} - \frac{5}{n^2}}{5 + \frac{3}{n} + \frac{10}{n^2}} = \frac{7}{5}.$$



# Asymptotic Notations 1

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

- ▶ **Notation:**  $f(n) \leq g(n)$  asymptotically. Note: By asymptotically we mean within constant multiplicative factor and for large  $n$
- ▶ **Examples:**

1.  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{7n^2 - 2n - 5}{5n^2 + 3n + 10} = \lim_{n \rightarrow \infty} \frac{7 - \frac{2}{n} - \frac{5}{n^2}}{5 + \frac{3}{n} + \frac{10}{n^2}} = \frac{7}{5}.$
2.  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n+10}{n^2-100} = 0.$

# Asymptotic Notations 1

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = O(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \leq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = O(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 \leq c < \infty$$

$c$  is a non negative finite constant.  $c$  cannot be  $\infty$ .

- ▶ **Notation:**  $f(n) \leq g(n)$  asymptotically. Note: By asymptotically we mean within constant multiplicative factor and for large  $n$
- ▶ **Examples:**

1.  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{7n^2 - 2n - 5}{5n^2 + 3n + 10} = \lim_{n \rightarrow \infty} \frac{7 - \frac{2}{n} - \frac{5}{n^2}}{5 + \frac{3}{n} + \frac{10}{n^2}} = \frac{7}{5}.$
2.  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n+10}{n^2-100} = 0.$

## Asymptotic Notations 2

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .

## Asymptotic Notations 2

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

## Asymptotic Notations 2

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

- ▶ **Notation:**  $f(n) \geq g(n)$  asymptotically.

## Asymptotic Notations 2

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

- ▶ **Notation:**  $f(n) \geq g(n)$  asymptotically.
- ▶ **Examples:**

$$1. \lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{5n^2 + 3n + 10}{7n^2 - 2n - 5} = \lim_{n \rightarrow \infty} \frac{5 + \frac{3}{n} + \frac{10}{n^2}}{7 - \frac{2}{n} - \frac{5}{n^2}} = \frac{5}{7}.$$

# Asymptotic Notations 2

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

- ▶ **Notation:**  $f(n) \geq g(n)$  asymptotically.
- ▶ **Examples:**

1.  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{5n^2 + 3n + 10}{7n^2 - 2n - 5} = \lim_{n \rightarrow \infty} \frac{5 + \frac{3}{n} + \frac{10}{n^2}}{7 - \frac{2}{n} - \frac{5}{n^2}} = \frac{5}{7}.$
2.  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n^2 - 100}{n + 10} = \infty.$

# Asymptotic Notations 2

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \Omega(g(n))$  iff  $\exists$  constants  $c > 0, n_o > 0 : f(n) \geq cg(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \Omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c \leq \infty$$

$c$  is a (non zero) positive constant. It can be  $\infty$  but not 0.

- ▶ **Notation:**  $f(n) \geq g(n)$  asymptotically.
- ▶ **Examples:**

1.  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{5n^2 + 3n + 10}{7n^2 - 2n - 5} = \lim_{n \rightarrow \infty} \frac{5 + \frac{3}{n} + \frac{10}{n^2}}{7 - \frac{2}{n} - \frac{5}{n^2}} = \frac{5}{7}.$
2.  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n^2 - 100}{n + 10} = \infty.$



## Asymptotic Notations 3

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_o > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_o$ .

## Asymptotic Notations 3

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_o > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \theta(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c < \infty$$

$c$  is a (non-zero) positive finite constant.  $c$  is neither 0 nor  $\infty$ .

# Asymptotic Notations 3

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_o > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \theta(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c < \infty$$

$c$  is a (non-zero) positive finite constant.  $c$  is neither 0 nor  $\infty$ .

- ▶ **Notation:**  $f(n) = g(n)$  asymptotically.

# Asymptotic Notations 3

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \theta(g(n))$  iff  $\exists$  constants  $c_1, c_2 > 0, n_o > 0$  :  
 $c_1 g(n) \leq f(n) \leq c_2 g(n)$  for all  $n \geq n_o$ .
- ▶ **Equivalent Definition:**  $f(n) = \theta(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = c, 0 < c < \infty$$

$c$  is a (non-zero) positive finite constant.  $c$  is neither 0 nor  $\infty$ .

- ▶ **Notation:**  $f(n) = g(n)$  asymptotically.
- ▶ **Example:**

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{7n^2 - 2n - 5}{5n^2 + 3n + 10} = \lim_{n \rightarrow \infty} \frac{7 - \frac{2}{n} - \frac{5}{n^2}}{5 + \frac{3}{n} + \frac{10}{n^2}} = \frac{7}{5}.$$

## Asymptotic Notations 4

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = o(g(n))$  iff for every constant  $c > 0 \exists n_c > 0 :$   
 $f(n) < cg(n)$  for all  $n \geq n_c$ .

## Asymptotic Notations 4

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = o(g(n))$  iff for every constant  $c > 0 \exists n_c > 0 :$   
 $f(n) < cg(n)$  for all  $n \geq n_c$ .
- ▶ **Equivalent Definition:**  $f(n) = o(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

## Asymptotic Notations 4

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = o(g(n))$  iff for every constant  $c > 0 \exists n_c > 0 :$   
 $f(n) < cg(n)$  for all  $n \geq n_c$ .
- ▶ **Equivalent Definition:**  $f(n) = o(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

- ▶ **Notation:**  $f(n) < g(n)$  asymptotically.

## Asymptotic Notations 4

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = o(g(n))$  iff for every constant  $c > 0 \exists n_c > 0 :$   
 $f(n) < cg(n)$  for all  $n \geq n_c$ .
- ▶ **Equivalent Definition:**  $f(n) = o(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$$

- ▶ **Notation:**  $f(n) < g(n)$  asymptotically.
- ▶ **Example:**  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n+10}{n^2-100} = 0$ .



## Asymptotic Notations 5

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \omega(g(n))$  iff for every constant  $c > 0 \exists n_c > 0 :$   
 $f(n) > cg(n)$  for all  $n \geq n_c$

# Asymptotic Notations 5

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \omega(g(n))$  iff for every constant  $c > 0 \exists n_c > 0 :$   
 $f(n) > cg(n)$  for all  $n \geq n_c$
- ▶ **Equivalent Definition:**  $f(n) = \omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

# Asymptotic Notations 5

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \omega(g(n))$  iff for every constant  $c > 0 \exists n_c > 0 :$   
 $f(n) > cg(n)$  for all  $n \geq n_c$
- ▶ **Equivalent Definition:**  $f(n) = \omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

- ▶ **Notation:**  $f(n) > g(n)$  asymptotically.

# Asymptotic Notations 5

Consider a set  $\mathcal{F}$  of functions  $f : \mathbf{I}^+ \rightarrow \mathbf{R}$ . Let  $f, g \in \mathcal{F}$ .

- ▶  $f(n) = \omega(g(n))$  iff for every constant  $c > 0 \exists n_c > 0 :$   
 $f(n) > cg(n)$  for all  $n \geq n_c$
- ▶ **Equivalent Definition:**  $f(n) = \omega(g(n))$  iff

$$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty$$

- ▶ **Notation:**  $f(n) > g(n)$  asymptotically.
- ▶ **Example:**  $\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \lim_{n \rightarrow \infty} \frac{n^2 - 100}{n + 10} = \infty$ .

## Some easy to prove Observations

1.  $f(n) \leq g(n)$  iff either  $f(n) < g(n)$  or  $f(n) = g(n)$  is true.

## Some easy to prove Observations

1.  $f(n) \leq g(n)$  iff either  $f(n) < g(n)$  or  $f(n) = g(n)$  is true.
2.  $f(n) \leq g(n)$  iff  $g(n) \geq f(n)$ .

## Some easy to prove Observations

1.  $f(n) \leq g(n)$  iff either  $f(n) < g(n)$  or  $f(n) = g(n)$  is true.
2.  $f(n) \leq g(n)$  iff  $g(n) \geq f(n)$ .
3.  $f(n) = g(n)$  iff  $f(n) \leq g(n)$  and  $f(n) \geq g(n)$ .

Let's do it

1.  $f(n) = 6n^3 + 7n + 100$ ,  $g(n) = 8n^2 + 20$ .



Let's do it

1.  $f(n) = 6n^3 + 7n + 100$ ,  $g(n) = 8n^2 + 20$ .
2.  $f(n) = \log^2 n$ ,  $g(n) = (1/2)\text{sqrt}(n)$ .

## Let's do it

1.  $f(n) = 6n^3 + 7n + 100$ ,  $g(n) = 8n^2 + 20$ .
2.  $f(n) = \log^2 n$ ,  $g(n) = (1/2)\text{sqrt}(n)$ .
3.  $f(n) = 2^n$ ,  $g(n) = 3^n$ .

# Practice Questions

Use the equivalent definitions (limits) to prove the following:

1. Show that a polynomial of degree  $d$ , with positive leading coefficient is  $\Theta(n^d)$ .
2. For  $g(n) = f(n) + o(f(n))$ , show that  $g(n) = \Theta(f(n))$ .
3. Show that
  - a.  $\log^M n = o(n^\epsilon)$  where  $M$  and  $\epsilon$  are positive constants.
  - b.  $\log n = o(n)$ .
4.  $a^n = o(b^n)$  for all  $a < b$ .