

Chapter 5: Exploratory Data Analysis

Alex Chi


Exercises

5.1 (Exploring percentages of small classes).


The variable Pct.20 in the college dataset contains the percentage of “small classes” (defined as 20 or fewer students) in the National Universities.

- Construct a dotplot of the small-class percentages using the stripchart function. To see the density of points, it is helpful to use either the method=stack or method=jitter arguments. What is the shape of this data?
- There is a single school with an unusually large small-class percentage. Use the identify function to find the name of this unusual school.
- Find the median small-class percentage and draw a vertical line (using the abline function) on the dotplot at the location of the median.

```
college = read.table("Rx-data/college.txt",  
                     header=TRUE, sep="\t")  
stripchart(college$Pct.20, method="stack")
```



```
stripchart(college$Pct.20, method="jitter")  
abline(v=median(college$Pct.20, na.rm=TRUE))
```



```
college[order(college$Pct.20,decreasing=TRUE),][1:2,]
```


```
##           School Rank Tier Retention Grad.rate Pct.20 Pct.50 Full.time
## 210 Golden Gate   NA    4         NA        20      94        0        NA
## 151 New School   NA    3         81        61      88        1        37
##      Top.10 Accept.rate Alumni.giving
## 210      NA           100           NA
## 151      18           51            8
```

5.2 (Relationship between the percentages of small classes and large classes).


The variables Pct.20 and Pct.50 in the college dataset contain respectively the percentage of “small classes” (defined as 20 or fewer students) and the percentage of “large classes” (defined as 50 or more students) in the National Universities.

- Use the plot function to construct a scatterplot of Pct.20 (horizontal) against Pct.50 (vertical).
- Use the line function to find a resistant line to these data. Add this resistant line to the scatterplot constructed in part a.
- If 60% of the classes at a particular college have 20 or fewer students, use the fitted line to predict the percentage of classes that have 50 or more students.
- Construct a graph of the residuals (vertical) against Pct.20 (horizontal) and add a horizontal line at zero (using the abline function).
- Is there a distinctive pattern to the residuals? (Compare the sizes of the residuals for small Pct.20 and the sizes of the residuals for large Pct.50.)
- Use the identify function to identify the schools that have residuals that exceed 10 in absolute value. Interpret these large residuals in the context of the problem.

```
plot(Pct.50~Pct.20, college)
fit = lm(Pct.50 ~ Pct.20, college,
        na.action=na.exclude)
abline(fit)
```



```
plot(college$Pct.20, residuals(fit))
abline(h=0)
```



```
college[na.omit(abs(residuals(fit)) > 10), ]
```

##	School	Rank	Tier	Retention	Grad.rate	Pct.20	Pct.50
## 24	UCLA	24	1	97	89	54	20
## 37	Univ Calif San Diego	35	1	94	85	42	30
## 43	Univ Calif Davis	42	1	90	81	34	28
## 112	New Hamsire	110	2	87	73	44	16
## 139	Bowling Green	NA	3	76	58	41	6
## 190	South Florida	NA	3	83	48	27	14
## 191	St. Thomas	NA	3	88	72	38	2
## 213	Indiana State	NA	4	67	43	50	5

##	Full.time	Top.10	Accept.rate	Alumni.giving
## 24	90	97	23	14
## 37	93	100	42	7
## 43	94	98	53	11
## 112	84	24	65	10
## 139	94	12	87	9
## 190	97	25	46	15
## 191	73	20	81	15
## 213	87	10	66	13


5.3 (Relationship between acceptance rate and “top-ten” percent- age).

The variables Accept.rate and Top.10 in the college dataset contain respectively the acceptance rate and the percentage of incoming students in the top 10 percent of their high school class in the National Universities.


One would believe that these two variables are strongly associated, since, for example, “exclusive” colleges with small acceptance rates would be expected to have a large percentage of “top-ten” students.

- Explore the relationship between Accept.rate and Top.10. This exploration should include a graph and linear fit that describe the basic pattern in the relationship and a residual graph that shows how schools differ from the basic pattern.
- Schools are often classified into “elite” and “non-elite” colleges depending on the type of students they admit. Based on your work in part a, is there any evidence from Accept.rate and Top.10 that schools do indeed cluster into “elite” and “non-elite” groups? Explain.

```
plot(Accept.rate ~ Top.10, college)
fit = lm(Accept.rate ~ Top.10, college,
        na.action=na.exclude)
abline(fit)
```



```
plot(college$Top.10, residuals(fit))
abline(h=0)
```



5.4 (Exploring the pattern of college enrollment in the United States).

The U.S. National Center for Education Statistics lists the total enrollment at Institutions of Higher Education for years 1900-1985 at their website <http://nces.ed.gov>. Define the ordered pair (x,y), where y is the total enrollment in thousands in year x. Then we observe the data (1955, 2653), (1956, 2918), (1957, 3324), (1959, 3640), (1961, 4145), (1963, 4780), (1964, 5280), (1965, 5921), (1966, 6390), (1967, 6912), (1968,


7513), (1969, 8005), (1970, 8581).

- a. Enter this data into R.
- b. Use the `lm` function to fit a line to the pattern of enrollment growth in the period 1955 to 1970. By inspecting a graph of the residuals, decide if a line is a reasonable model of the change in enrollment.
- c. Transform the enrollment by a logarithm, and fit a line to the (year, log enrollment) data. Inspect the pattern of residuals and explain why a line is a better fit to the log enrollment data.
- d. By interpreting the fit to the log enrollment data, explain how the college enrollment is changing in this time period. How does this growth compare to the growth of the BGSU enrollment in Section 5?


```
year =      c(1955, 1956, 1957, 1959, 1961, 1963, 1964, 1965, 1966, 1967, 1968, 1969, 1970)
enrollment = c(2653, 2918, 3324, 3640, 4145, 4780, 5280, 5921, 6390, 6912, 7513, 8005, 8581)
data = data.frame(cbind(year, enrollment))
data
```

```
##   year enrollment
## 1  1955      2653
## 2  1956      2918
## 3  1957      3324
## 4  1959      3640
## 5  1961      4145
## 6  1963      4780
## 7  1964      5280
## 8  1965      5921
## 9  1966      6390
## 10 1967      6912
## 11 1968      7513
## 12 1969      8005
## 13 1970      8581
```


```
fit = lm(enrollment ~ year, data)
plot(enrollment ~ year, data)
abline(fit)
```




```
plot(data$year, residuals(fit))
abline(h=0)
```



```
data$log.enrollment = log(data$enrollment)
fit = lm(log.enrollment ~ year, data)
plot(log.enrollment ~ year, data)
abline(fit)
```



```
plot(data$year, residuals(fit))
abline(h = 0)
```



5.5 (Exploring percentages of full-time faculty).

The variable Full.time in the college dataset (see Example 5.3) contains the percentage of faculty who are hired full-time in the group of National Universities.


- Using the hist function, construct a histogram of the full-time percentages and comment on the

shape of the distribution.

- b. Use the froot and flog transformations to reexpress the full-time percentages. Construct histograms of the collection of froots and the collection of flogs. Is either transformation successful in making the full-time percentages approximately symmetric?
- c. For data that is approximately normally distributed, about 68% of the data fall within one standard deviation of the mean. Assuming you have found a transformation in part (b) that makes the full-time percentages approximately normal, find an interval that contains roughly 68% of the data on the new scale.


```
data = college$Full.time  
hist(data)
```

Histogram of data




```
froot = sqrt(data) - sqrt(100 - data)  
flog = log(data + 0.5) - log(100 - data + 0.5)  
hist(froot)
```

Histogram of froot



```
hist(flog)
```

Histogram of flog




5.6 (Exploring alumni giving rates).

The variable `Alumni.giving` contains the percentage of alumni from the college who make financial contributions.

- Construct a “stacked” dotplot of the alumni giving percentages using the `stripchart` function.
- Identify the names of the three schools with unusually large giving percentages.
- It can be difficult to summarize these giving percentages since the distribution is right-skewed. One can make the dataset more symmetric by applying either a square root transformation or a log transformation.

Apply both square root and log transformations. Which transformation makes the alumni giving rates approximately symmetric?

```
roots = sqrt(college$Alumni.giving)
logs = log(college$Alumni.giving)
stripchart(college$Alumni.giving, method="stack")
```




```
college[order(college$Alumni.giving,decreasing=TRUE),][1:3,]
```

```
##           School Rank Tier Retention Grad.rate Pct.20 Pct.50 Full.time
## 211 Idaho State   NA    4      56         27    NA    NA      NA
## 2   Princeton    1    1      98         96    75     9      92
## 11  Dartmouth    11    1      98         95    63     9      93
##           Top.10 Accept.rate Alumni.giving
## 211      NA              NA              67
## 2       97              10              61
## 11      90              13              51
```


```
hist(roots)
```

Histogram of roots



```
hist(logs)
```

Histogram of logs




5.7 (Exploring alumni giving rates (continued)).


In this exercise, we focus on the comparison of the alumni giving percentages between the four tiers of colleges.

- Using the `stripchart` function with the `stacked` option, construct parallel dotplots of alumni giving by tier.
- As one moves from Tier 4 to Tier 1, how does the average giving change?
- As one moves from Tier 4 to Tier 1, how does the spread of the giving rates change?
- We note from parts (b) and (c), that small giving rates tend to have small variation, and large giving rates tend to have large variation. One way of removing the dependence of average with spread is to apply a power transformation such as a square root or a log. Construct parallel stripcharts of the square roots of the giving rates, and parallel boxplots of the log giving rates.
- Looking at the two sets of parallel stripcharts in part (d), were the square root rates or the log rates successful in making the spreads approximately the same between groups?

```
stripchart(Alumni.giving ~ Tier, college, method="stack")
```



```
stripchart(sqrt(Alumni.giving) ~ Tier, college, method="stack")
```



```
boxplot(logs ~ Tier, college, method="stack")
```

