

Lecture 3 Random Sampling and Multivariate Normal Distribution

Shiwei Lan¹

¹School of Mathematical and Statistical Sciences
Arizona State University

STP533 Multivariate Analysis
Spring 2025

Overview

S.Lan

Radom Sampling

Radom Samples and
Expectation of Sample
Statistics

Generalized Variance
and Measurement of
Sample Variation

Multivariate Normal Distribution

Multivariate Normal
Density and Its
Properties

Parameter Estimation:
Maximum Likelihood
Estimation

The Sampling
Distribution of $\bar{\mathbf{X}}$ and \mathbf{S}

Large-Sample Behavior
of $\bar{\mathbf{X}}$ and \mathbf{S}

1 Radom Sampling

Radom Samples and Expectation of Sample Statistics

Generalized Variance and Measurement of Sample Variation

2 Multivariate Normal Distribution

Multivariate Normal Density and Its Properties

Parameter Estimation: Maximum Likelihood Estimation

The Sampling Distribution of $\bar{\mathbf{X}}$ and \mathbf{S}

Large-Sample Behavior of $\bar{\mathbf{X}}$ and \mathbf{S}

- Recall the data array \mathbf{X} is arranged as an $n \times p$ matrix

$$\mathbf{X} = \begin{bmatrix} x_{11} & x_{12} & \cdots & x_{1j} & \cdots & x_{1p} \\ x_{21} & x_{22} & \cdots & x_{2j} & \cdots & x_{2p} \\ \vdots & \vdots & & \vdots & & \vdots \\ x_{i1} & x_{i2} & \cdots & x_{ij} & \cdots & x_{ip} \\ \vdots & \vdots & & \vdots & & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{nj} & \cdots & x_{np} \end{bmatrix} = \begin{bmatrix} X_1^T \\ X_2^T \\ \vdots \\ X_n^T \end{bmatrix} = [X_1 \quad X_2 \quad \cdots \quad X_p]$$

- Each row $X_i^T = [X_{i1}, X_{i2}, \dots, X_{ip}]$ represents a *independent observation* from a joint distribution p -dimensional random vector.
- Each column $X_j = [X_{1j}, X_{2j}, \dots, X_{nj}]^T$ represents a *random sample* (collection of observations) of a random variable X_j .

- Random sample is often assumed to be a collection of *independently identically distributed (i.i.d.)* observations.
- Assume the p -dimensional distribution has a density function $f(\mathbf{x}) = f(x_1, x_2, \dots, x_p)$. We denote random sample $\{X_i\}_{i=1}^n \stackrel{iid}{\sim} f(\mathbf{x})$.
- For the joint distribution of all the samples, based on the iid assumption, we have

$$f(\mathbf{X}) = \prod_{i=1}^n f(\mathbf{x}_i).$$

- Note, in general $f(\mathbf{x}) \neq \prod_{j=1}^p f(x_j)$ where each $f(x_j)$ is the marginal density of random variable X_j .

- Now we assume a random sample $\{X_i\}_{i=1}^n \stackrel{iid}{\sim} f(\mathbf{x})$ from a joint distribution with mean $\boldsymbol{\mu} \in \mathbb{R}^p$ and covariance $\boldsymbol{\Sigma} \in \mathbb{R}^{p \times p}$.
- Previously we had sample mean $\bar{\mathbf{X}} = \frac{1}{n} \sum_{i=1}^n X_i$ and

$$\mathbb{E}[\bar{\mathbf{X}}] = \boldsymbol{\mu}, \quad \text{Cov}(\bar{\mathbf{X}}) = \frac{1}{n} \boldsymbol{\Sigma}$$

- Then we have for sample covariance $\mathbf{S}_n = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{\mathbf{X}})(X_i - \bar{\mathbf{X}})^T$

$$\mathbb{E}[\mathbf{S}_n] = \frac{n-1}{n} \boldsymbol{\Sigma}$$

- Therefore, we often consider the unbiased sample covariance matrix $\mathbf{S} = \frac{n}{n-1} \mathbf{S}_n$.


- For p -dimensional random sample $\mathbf{X}_{n \times p}$, the *generalized sample variance* is defined as the determinant of sample covariance \mathbf{S} :

$$\text{generalized sample variance} = |\mathbf{S}| = (n - 1)^p \text{vol}^2$$

where vol is the volume generated by p residual (deviation) vectors $\{\mathbf{x}_j - \bar{\mathbf{x}}_j\}_{j=1}^p$.

- It can be shown that $\text{vol}\{\mathbf{x} : (\mathbf{x} - \bar{\mathbf{x}})^T \mathbf{S}^{-1} (\mathbf{x} - \bar{\mathbf{x}}) \leq c^2\} = k_p |\mathbf{S}|^{\frac{1}{2}} c^p$.
- This quantity measures the variability of the random sample of size n .
- It can be used to detect *multi-collinearity*, i.e. X_1, X_2, \dots, X_p are linearly dependent when $|\mathbf{S}| = 0$.
- If $n \leq p$, then $|\mathbf{S}| = 0$ for all samples.

Consider the area generated within the plane by two deviation vectors $\mathbf{d}_1 = \mathbf{y}_1 - \bar{x}_1 \mathbf{1}$ and $\mathbf{d}_2 = \mathbf{y}_2 - \bar{x}_2 \mathbf{1}$. Let $L_{\mathbf{d}_1}$ be the length of \mathbf{d}_1 and $L_{\mathbf{d}_2}$ the length of \mathbf{d}_2 . By elementary geometry, we have the diagram



and the area of the trapezoid is $|L_{\mathbf{d}_1} \sin(\theta)| L_{\mathbf{d}_2}$. Since $\cos^2(\theta) + \sin^2(\theta) = 1$, we can express this area as

$$\text{Area} = L_{\mathbf{d}_1} L_{\mathbf{d}_2} \sqrt{1 - \cos^2(\theta)}$$

From (3-5) and (3-7),

$$L_{\mathbf{d}_1} = \sqrt{\sum_{j=1}^n (x_{j1} - \bar{x}_1)^2} = \sqrt{(n-1)s_{11}}$$

$$L_{\mathbf{d}_2} = \sqrt{\sum_{j=1}^n (x_{j2} - \bar{x}_2)^2} = \sqrt{(n-1)s_{22}}$$

and

$$\cos(\theta) = r_{12}$$

Therefore,

$$\text{Area} = (n-1) \sqrt{s_{11}} \sqrt{s_{22}} \sqrt{1 - r_{12}^2} = (n-1) \sqrt{s_{11}s_{22}(1 - r_{12}^2)} \quad (3-13)$$

Also,

$$\begin{aligned} |S| &= \left| \begin{bmatrix} s_{11} & s_{12} \\ s_{12} & s_{22} \end{bmatrix} \right| = \left| \begin{bmatrix} s_{11} & \sqrt{s_{11}} \sqrt{s_{22}} r_{12} \\ \sqrt{s_{11}} \sqrt{s_{22}} r_{12} & s_{22} \end{bmatrix} \right| \\ &= s_{11}s_{22} - s_{11}s_{22}r_{12}^2 = s_{11}s_{22}(1 - r_{12}^2) \end{aligned} \quad (3-14)$$

Overview

S.Lan

Random Sampling

Random Samples and
Expectation of Sample
Statistics

Generalized Variance
and Measurement of
Sample Variation

Multivariate Normal Distribution

Multivariate Normal
Density and Its
Properties

Parameter Estimation:
Maximum Likelihood
Estimation

The Sampling
Distribution of \bar{X} and S

Large-Sample Behavior
of \bar{X} and S

$$\mathbf{X} = \begin{bmatrix} 1 & 9 & 10 \\ 4 & 12 & 16 \\ 2 & 10 & 12 \\ 5 & 8 & 13 \\ 3 & 11 & 14 \end{bmatrix}$$

- Consider the *generalized sample variance of the standardized variables*

$$|\mathbf{R}| = (n - 1)^p \text{vol}^2$$

where vol is the volume generated by p standardized vectors $\left\{ \frac{\mathbf{x}_j - \bar{\mathbf{x}}_j}{\sqrt{s_{jj}}} \right\}_{j=1}^p$.

- What is the relationship between $|\mathbf{R}|$ and $|\mathbf{S}|$?
- Another generalization of variance is *total sample variance* defined as $\text{tr}(\mathbf{S})$.

- Recall we had the following matrix representation of sample statistics:

$$\bar{\mathbf{X}} = \frac{\mathbf{1}_n^T}{n} \mathbf{X}, \quad \mathbf{S} = \frac{1}{n-1} \mathbf{X}^T (\mathbf{I}_n - \mathbf{J}) \mathbf{X}, \quad \mathbf{J} = \frac{\mathbf{1}_n \mathbf{1}_n^T}{n}$$

- Now suppose we have two linear combinations $\mathbf{X}\mathbf{b}$ and $\mathbf{X}\mathbf{c}$. Then we have

$$\overline{\mathbf{X}\mathbf{b}} = \bar{\mathbf{X}}\mathbf{b}, \quad s_{\mathbf{X}\mathbf{b}, \mathbf{X}\mathbf{c}} = \mathbf{b}^T \mathbf{S} \mathbf{c}$$

- For example, $\mathbf{X} = \begin{bmatrix} 1 & 2 & 5 \\ 4 & 1 & 6 \\ 4 & 0 & 4 \end{bmatrix}$, $\mathbf{b} = [2, 2, -1]^T$ and $\mathbf{c} = [1, -1, 3]^T$.

Overview

S.Lan

Radom Sampling

Radom Samples and Expectation of Sample Statistics

Generalized Variance and Measurement of Sample Variation

Multivariate Normal Distribution

Multivariate Normal Density and Its Properties

Parameter Estimation: Maximum Likelihood Estimation

The Sampling Distribution of $\bar{\mathbf{X}}$ and \mathbf{S}

Large-Sample Behavior of $\bar{\mathbf{X}}$ and \mathbf{S}

1 Radom Sampling

Radom Samples and Expectation of Sample Statistics

Generalized Variance and Measurement of Sample Variation

2 Multivariate Normal Distribution

Multivariate Normal Density and Its Properties

Parameter Estimation: Maximum Likelihood Estimation

The Sampling Distribution of $\bar{\mathbf{X}}$ and \mathbf{S}

Large-Sample Behavior of $\bar{\mathbf{X}}$ and \mathbf{S}

- The read data are not exactly multivariate normal, but normal density can serve as a good approximation.
- The density of multivariate normal random vector $\mathbf{X} \sim N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ is

$$f(\mathbf{x}) = (2\pi)^{-\frac{p}{2}} |\boldsymbol{\Sigma}|^{-\frac{1}{2}} \exp\left\{-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x}-\boldsymbol{\mu})\right\}$$

where the covariance matrix $\boldsymbol{\Sigma}$ is PSD.




Figure 4.2 Two bivariate normal distributions. (a) $\sigma_{11} = \sigma_{22}$ and $\rho_{12} = 0$. (b) $\sigma_{11} = \sigma_{22}$ and $\rho_{12} = .75$.

- The contour of MVN density is determined by

$$(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) = c^2$$

- This is an ellipsoid centered at $\boldsymbol{\mu}$ and having axes $\pm \sqrt{\lambda_i} \mathbf{v}_i$ with eigen-pairs $\{\lambda_i, \mathbf{v}_i\}$ of $\boldsymbol{\Sigma}$.




Figure 4.3 A constant-density contour for a bivariate normal distribution with $\sigma_{11} = \sigma_{22}$ and $\sigma_{12} > 0$ (or $\rho_{12} > 0$).

- The linear combination of MVN is another MVN. Let $\mathbf{A} \in \mathbb{R}^{q \times p}$. Then

$$\mathbf{AX} \sim N_q(\mathbf{A}\boldsymbol{\mu}, \mathbf{A}\boldsymbol{\Sigma}\mathbf{A}^T)$$

- The marginal of MVN is also MVN. Consider $\mathbf{A} = [\mathbf{I}_q \quad \mathbf{0}]$.
- The conditional pdf of MVN is also MVN. Let $\mathbf{X} = \begin{bmatrix} \mathbf{X}_1 \\ \mathbf{X}_2 \end{bmatrix}$, $\boldsymbol{\mu} = \begin{bmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{bmatrix}$,

$$\boldsymbol{\Sigma} = \begin{bmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{21} & \boldsymbol{\Sigma}_{22} \end{bmatrix}. \text{ Then}$$

$$\mathbf{X}_1 | \mathbf{X}_2 = \mathbf{x}_2 \sim N_{p_1}(\boldsymbol{\mu}_1 - \boldsymbol{\Sigma}_{12}\boldsymbol{\Sigma}_{22}^{-1}(\mathbf{x}_2 - \boldsymbol{\mu}_2), \boldsymbol{\Sigma}_{11} - \boldsymbol{\Sigma}_{12}\boldsymbol{\Sigma}_{22}^{-1}\boldsymbol{\Sigma}_{21})$$

- Note $\mathbf{X}_1 \perp \mathbf{X}_2$ if and only if $\boldsymbol{\Sigma}_{12} = \mathbf{0}$. What is the caveat?
- What is the distribution of $(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})$?

- Consider the random sample $\mathbf{X}_{n \times p}$. The *likelihood* of the sample is the joint density

$$L_{\mathbf{X}}(\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \prod_{i=1}^n f(\mathbf{x}_i) = (2\pi)^{-\frac{np}{2}} |\boldsymbol{\Sigma}|^{-\frac{n}{2}} \exp \left\{ -\frac{1}{2} \sum_{i=1}^n (\mathbf{x}_i - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x}_i - \boldsymbol{\mu}) \right\}$$

- We notice the sum of quadratic form can be rewritten as

$$\sum_{i=1}^n (\mathbf{x}_i - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x}_i - \boldsymbol{\mu}) = \text{tr} \left[(\mathbf{X} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{X} - \boldsymbol{\mu}) \right]$$

- The maximum likelihood estimation (MLE) is to maximize the following log-likelihood with respect to $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$:

$$\ell_{\mathbf{X}}(\boldsymbol{\mu}, \boldsymbol{\Sigma}) = \log L_{\mathbf{X}}(\boldsymbol{\mu}, \boldsymbol{\Sigma}) = -\frac{n}{2} \log |\boldsymbol{\Sigma}| - \frac{1}{2} \text{tr} \left[(\mathbf{X} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{X} - \boldsymbol{\mu}) \right]$$

- Setting $\frac{\partial \ell}{\partial \boldsymbol{\mu}} = 0$ and $\frac{\partial \ell}{\partial \boldsymbol{\Sigma}} = 0$, we obtain the MLE for $\boldsymbol{\mu}, \boldsymbol{\Sigma}$ as

$$\hat{\boldsymbol{\mu}} = \bar{\mathbf{X}}, \quad \hat{\boldsymbol{\Sigma}} = \frac{n-1}{n} \mathbf{S}$$

- Note that $\bar{\mathbf{X}}$ and \mathbf{S} are also *sufficient statistics*.

Suppose $\mathbf{X}_i \stackrel{iid}{\sim} N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, then we have

- ① $\bar{\mathbf{X}} \sim N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma}/n)$.
- ② $(n-1)\mathbf{S} \sim W_{n-1}(\boldsymbol{\Sigma})$, Wishart distribution with degree of freedom $n-1$.
- ③ $\bar{\mathbf{X}} \perp \mathbf{S}$.

Definition (Wishart distribution)

A square matrix $\mathbf{A} \sim W_m(\boldsymbol{\Sigma})$ Wishart distribution with degree of freedom m if it can be expressed as $\mathbf{A} = \sum_{j=1}^m \mathbf{Z}_j \mathbf{Z}_j^T$, where $\mathbf{Z}_j \stackrel{iid}{\sim} N_p(\mathbf{0}, \boldsymbol{\Sigma})$. The density of \mathbf{A} is

$$f_m(\mathbf{A}|\boldsymbol{\Sigma}) = \frac{|\mathbf{A}|^{(m-p-1)/2} \exp\{-\text{tr}(\mathbf{A}\boldsymbol{\Sigma}^{-1})/2\}}{2^{pm/2} \pi^{p(p-1)/4} |\boldsymbol{\Sigma}|^{m/2} \prod_{i=1}^p \Gamma((m+1-i)/2)}$$

- If $\mathbf{A}_1 \sim W_{m_1}(\boldsymbol{\Sigma})$ and $\mathbf{A}_2 \sim W_{m_2}(\boldsymbol{\Sigma})$, then $\mathbf{A}_1 + \mathbf{A}_2 \sim W_{m_1+m_2}(\boldsymbol{\Sigma})$.
- If $\mathbf{A} \sim W_m(\boldsymbol{\Sigma})$, then $\mathbf{CAC}^T \sim W_m(\mathbf{C}\boldsymbol{\Sigma}\mathbf{C}^T)$.

Suppose $\mathbf{X}_i \stackrel{iid}{\sim} (\boldsymbol{\mu}, \boldsymbol{\Sigma})$, then we have

LLN $\bar{\mathbf{X}} \xrightarrow{P} \boldsymbol{\mu}$, i.e. for any $\epsilon > 0$, $P[|\bar{\mathbf{X}} - \boldsymbol{\mu}| > \epsilon] \rightarrow 0$ as $n \rightarrow \infty$.

CLT $\sqrt{n}(\bar{\mathbf{X}} - \boldsymbol{\mu}) \xrightarrow{L} N_p(\mathbf{0}, \boldsymbol{\Sigma})$, i.e $P[\sqrt{n}(\bar{\mathbf{X}} - \boldsymbol{\mu}) \leq \mathbf{x}] \rightarrow p_N(\mathbf{x}; \mathbf{0}, \boldsymbol{\Sigma})$ as $n \rightarrow \infty$.

- We also have $n(\bar{\mathbf{X}} - \boldsymbol{\mu})^T \mathbf{S}^{-1}(\bar{\mathbf{X}} - \boldsymbol{\mu}) \dot{\sim} \chi_p^2$.

- For univariate normality:

The steps leading to a $Q-Q$ plot are as follows:

1. Order the original observations to get $x_{(1)}, x_{(2)}, \dots, x_{(n)}$ and their corresponding probability values $(1 - \frac{1}{2})/n, (2 - \frac{1}{2})/n, \dots, (n - \frac{1}{2})/n$;
2. Calculate the standard normal quantiles $q_{(1)}, q_{(2)}, \dots, q_{(n)}$; and
3. Plot the pairs of observations $(q_{(1)}, x_{(1)}), (q_{(2)}, x_{(2)}), \dots, (q_{(n)}, x_{(n)})$, and examine the “straightness” of the outcome.

- For bivariate normality:

To construct the chi-square plot,

1. Order the squared distances in (4-32) from smallest to largest as $d_{(1)}^2 \leq d_{(2)}^2 \leq \dots \leq d_{(n)}^2$.
2. Graph the pairs $(q_{c,p}((j - \frac{1}{2})/n), d_{(j)}^2)$, where $q_{c,p}((j - \frac{1}{2})/n)$ is the $100(j - \frac{1}{2})/n$ quantile of the chi-square distribution with p degrees of freedom.