

**Example 5.4 (Simultaneous confidence intervals as shadows of the confidence ellipsoid)**

In Example 5.3, we obtained the 95% confidence ellipse for the means of the fourth roots of the door-closed and door-open microwave radiation measurements. The 95% simultaneous  $T^2$  intervals for the two component means are, from (5-24),

$$\begin{aligned} & \left( \bar{x}_1 - \sqrt{\frac{p(n-1)}{(n-p)} F_{p,n-p}(.05)} \sqrt{\frac{s_{11}}{n}}, \quad \bar{x}_1 + \sqrt{\frac{p(n-1)}{(n-p)} F_{p,n-p}(.05)} \sqrt{\frac{s_{11}}{n}} \right) \\ &= \left( .564 - \sqrt{\frac{2(41)}{40}} 3.23 \sqrt{\frac{.0144}{42}}, \quad .564 + \sqrt{\frac{2(41)}{40}} 3.23 \sqrt{\frac{.0144}{42}} \right) \text{ or } (.516, .612) \\ & \left( \bar{x}_2 - \sqrt{\frac{p(n-1)}{(n-p)} F_{p,n-p}(.05)} \sqrt{\frac{s_{22}}{n}}, \quad \bar{x}_2 + \sqrt{\frac{p(n-1)}{(n-p)} F_{p,n-p}(.05)} \sqrt{\frac{s_{22}}{n}} \right) \\ &= \left( .603 - \sqrt{\frac{2(41)}{40}} 3.23 \sqrt{\frac{.0146}{42}}, \quad .603 + \sqrt{\frac{2(41)}{40}} 3.23 \sqrt{\frac{.0146}{42}} \right) \text{ or } (.555, .651) \end{aligned}$$

In Figure 5.2, we have redrawn the 95% confidence ellipse from Example 5.3. The 95% simultaneous intervals are shown as shadows, or projections, of this ellipse on the axes of the component means. ■