

Lecture 7 Multivariate Linear Regression

Shiwei Lan¹

¹School of Mathematical and Statistical Sciences
Arizona State University

STP533 Multivariate Analysis
Spring 2025

Lecture 7

S.Lan

The Classical Linear Regression Model

The Classical Linear
Regression
Least Square Estimation
Inferences About the
Regression Model

Multivariate Multiple Regression

Multivariate Multiple
Regression
Least Square Estimation
Likelihood Ratio Tests
for Regression
Parameters
Predictions

1 The Classical Linear Regression Model

The Classical Linear Regression

Least Square Estimation

Inferences About the Regression Model

2 Multivariate Multiple Regression

Multivariate Multiple Regression

Least Square Estimation

Likelihood Ratio Tests for Regression Parameters

Predictions

Lecture 7

S.Lan

The Classical Linear Regression Model

The Classical Linear
Regression

Least Square Estimation

Inferences About the
Regression Model

Multivariate Multiple Regression

Multivariate Multiple
Regression

Least Square Estimation

Likelihood Ratio Tests
for Regression
Parameters

Predictions

- Regression analysis is the statistical methodology for predicting values of one or more *response* (dependent) variables from a collection of *predictor* (independent) variables.
- It can also be used for assessing the effects of the predictor variables on the responses.
- The name *regression*, dated back to 1885 by F. Galton.
- We first review the classical linear regression model with a single response. Then we generalize to linear model for several dependent variables.

- Suppose we have p predictor variables X_1, \dots, X_p and a response variable Y .
- For example, Y =current market value of a house, X_1 =square feet, X_2 =location, X_3 =appraised value of last year, and X_4 =quality of construction.
- A classical linear regression relates the average value of Y with a linear combination of X_i 's.

$$Y_i = \beta_0 + X_{i1}\beta_1 + \dots + X_{ip}\beta_p + \epsilon_i, \quad i = 1, \dots, n,$$

where we assume $\epsilon_i \stackrel{iid}{\sim} (0, \sigma^2)$.

- If we denote $\mathbf{Y} = [Y_1, \dots, Y_n]^T$, $\mathbf{X} = \begin{bmatrix} 1 & x_{11} & \dots & x_{1p} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & x_{n1} & \dots & x_{np} \end{bmatrix}$, and

$\boldsymbol{\beta} = [\beta_0, \beta_1, \dots, \beta_p]^T$, then we can rewrite

$$\mathbf{Y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}, \quad \boldsymbol{\epsilon} \sim (\mathbf{0}, \sigma^2 \mathbf{I}_n).$$

Example 7.2 (The design matrix for one-way ANOVA as a regression model)
Determine the design matrix if the linear regression model is applied to the one-way ANOVA situation in Example 6.6.

We create so-called *dummy* variables to handle the three population means: $\mu_1 = \mu + \tau_1$, $\mu_2 = \mu + \tau_2$, and $\mu_3 = \mu + \tau_3$. We set

$$z_1 = \begin{cases} 1 & \text{if the observation is} \\ & \text{from population 1} \\ 0 & \text{otherwise} \end{cases} \quad z_2 = \begin{cases} 1 & \text{if the observation is} \\ & \text{from population 2} \\ 0 & \text{otherwise} \end{cases}$$

$$z_3 = \begin{cases} 1 & \text{if the observation is} \\ & \text{from population 3} \\ 0 & \text{otherwise} \end{cases}$$

and $\beta_0 = \mu$, $\beta_1 = \tau_1$, $\beta_2 = \tau_2$, $\beta_3 = \tau_3$. Then

$$Y_j = \beta_0 + \beta_1 z_{j1} + \beta_2 z_{j2} + \beta_3 z_{j3} + \varepsilon_j, \quad j = 1, 2, \dots, 8$$

where we arrange the observations from the three populations in sequence. Thus, we obtain the observed response vector and design matrix


$$\mathbf{Y}_{(8 \times 1)} = \begin{bmatrix} 9 \\ 6 \\ 9 \\ 0 \\ 2 \\ 3 \\ 1 \\ 2 \end{bmatrix}; \quad \mathbf{Z}_{(8 \times 4)} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \end{bmatrix}$$

- The least square estimation (LSE) minimizes the sum of square $S(\beta) = \|\mathbf{Y} - \mathbf{X}\beta\|_2^2$ with respect to β .
- Let \mathbf{X} be full rank $p + 1 \leq n$. The LSE result of β is $\hat{\beta} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y}$.

- Let $\hat{\mathbf{y}} = \mathbf{X}\hat{\beta} = \mathbf{H}\mathbf{y}$ be the *fitted values* of \mathbf{y} , where $\mathbf{H} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$ is called *hat matrix*.
- The residual vector can now be written

$$\mathbf{e} = \mathbf{y} - \hat{\mathbf{y}} = (\mathbf{I} - \mathbf{H})\mathbf{y}$$

- The residual sum of squares becomes $S(\hat{\beta}) = \|\mathbf{e}\|_2^2 = \mathbf{y}^T (\mathbf{I} - \mathbf{H})\mathbf{y}$.



- Note $\mathbf{X} \perp \mathbf{e}$ and $\hat{\mathbf{y}} \perp \mathbf{e}$. Why?
- Then we have $\|\mathbf{y}\|^2 = \|\hat{\mathbf{y}}\|^2 + \|\mathbf{e}\|^2$.
- Further we have decomposition of the sum of squares about mean

$$\underbrace{\sum_{i=1}^n (y_i - \bar{y})^2}_{SST} = \underbrace{\sum_{i=1}^n (\hat{y}_i - \bar{y})^2}_{SSR} + \underbrace{\sum_{i=1}^n (y_i - \hat{y}_i)^2}_{SSE}$$

- The decomposition of the sum of squares can also be written as $\mathbf{y}^T(\mathbf{I} - \mathbf{J})\mathbf{y} = \mathbf{y}^T(\mathbf{H} - \mathbf{J})\mathbf{y} + \mathbf{y}^T(\mathbf{I} - \mathbf{H})\mathbf{y}$.

- We define the *coefficient of determination* as

$$R^2 = \frac{SSR}{SST} = 1 - \frac{SSE}{SST}$$

- This quantity measure the proportion of the total variation in y 's "explained" by the model with p predictors \mathbf{X} .
- If we plot $\hat{\mathbf{y}}$ against \mathbf{y} , what is the slope?

Lecture 7

S.Lan

The Classical
Linear
Regression
ModelThe Classical Linear
Regression

Least Square Estimation

Inferences About the
Regression ModelMultivariate
Multiple
RegressionMultivariate Multiple
Regression

Least Square Estimation

Likelihood Ratio Tests
for Regression
Parameters

Predictions

- We have the following property for LSE $\hat{\beta} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$

$$E[\hat{\beta}] = \beta, \quad \text{Cov}[\hat{\beta}] = \sigma^2 (\mathbf{X}^T \mathbf{X})^{-1}$$

- The residual vector $\mathbf{e} = \mathbf{y} - \hat{\mathbf{y}}$ has the following property

$$E[\mathbf{e}] = \mathbf{0}, \quad \text{Cov}[\mathbf{e}] = \sigma^2 [\mathbf{I} - \mathbf{H}]$$

Lecture 7

S.Lan

The Classical Linear Regression Model

The Classical Linear Regression

Least Square Estimation

Inferences About the Regression Model

Multivariate Multiple Regression

Multivariate Multiple Regression

Least Square Estimation

Likelihood Ratio Tests for Regression Parameters

Predictions

- Now we consider $\mathbf{Y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$ with $\boldsymbol{\epsilon} \sim N_n(\mathbf{0}, \sigma^2 \mathbf{I}_n)$.
- Then the maximum likelihood estimator (MLE) of $\boldsymbol{\beta}$ is the same as LSE $\hat{\boldsymbol{\beta}} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$. Moreover, we have

$$\hat{\boldsymbol{\beta}} \sim N(\boldsymbol{\beta}, \sigma^2 (\mathbf{X}^T \mathbf{X})^{-1})$$

- The residual $\mathbf{e} = \mathbf{y} - \hat{\mathbf{y}}$ is independent of $\hat{\boldsymbol{\beta}}$ and $SSE/n = \|\mathbf{e}\|^2/n$ is the MLE of σ^2 . Moreover,

$$\frac{\|\mathbf{e}\|^2}{\sigma^2} \sim \chi^2(n - p - 1).$$

- $MSE = \frac{SSE}{n-p-1} = \frac{\|\mathbf{e}\|^2}{n-p-1} =: s^2$ is an unbiased estimator of σ^2 .

Lecture 7

S.Lan

The Classical Linear Regression Model

The Classical Linear Regression

Least Square Estimation

Inferences About the Regression Model

Multivariate Multiple Regression

Multivariate Multiple Regression

Least Square Estimation

Likelihood Ratio Tests for Regression Parameters

Predictions

- $100(1 - \alpha)\%$ CR for β is determined by

$$(\beta - \hat{\beta})^T (\mathbf{X}^T \mathbf{X})^{-1} (\beta - \hat{\beta}) \leq (p + 1) s^2 F_{1-\alpha}(p + 1, n - p - 1).$$

- The $100(1 - \alpha)\%$ SCI for β_j 's are given by

$$\hat{\beta}_j \pm \sqrt{\widehat{\text{Var}}(\hat{\beta}_j)} \sqrt{(p + 1) F_{1-\alpha}(p + 1, n - p - 1)}, \quad j = 0, 1, \dots, p.$$

where $\widehat{\text{Var}}(\hat{\beta}_j)$ is the j -th diagonal element of $s^2 (\mathbf{X}^T \mathbf{X})^{-1}$.

- For each β_j , the $100(1 - \alpha)\%$ individual CI is

$$\hat{\beta}_j \pm t_{1-\alpha/2}(n - p - 1) \sqrt{\widehat{\text{Var}}(\hat{\beta}_j)}, \quad j = 0, 1, \dots, p.$$

Lecture 7

S.Lan

The Classical Linear Regression Model

The Classical Linear Regression

Least Square Estimation

Inferences About the Regression Model

Multivariate Multiple Regression

Multivariate Multiple Regression

Least Square Estimation

Likelihood Ratio Tests for Regression Parameters

Predictions

- Suppose you hypothesize that only the first $q \leq p$ predictors are significant in explaining the response variable.
- We want to test $H_0 : \beta_{q+1} = \beta_{q+2} = \cdots = \beta_p = 0$. Denote $\beta_2 = [\beta_{q+1}, \cdots, \beta_p]^T$.
- We divide $\mathbf{X} = [(\mathbf{X}_1)_{n \times (q+1)} | (\mathbf{X}_2)_{n \times (p-q)}]$ and $\beta = [\beta_1^T, \beta_2^T]^T$. Then

$$\mathbf{Y} = \mathbf{X}\beta + \epsilon = \mathbf{X}_1\beta_1 + \mathbf{X}_2\beta_2 + \epsilon$$

- The LRT rejects $H_0 : \beta_2 = \mathbf{0}$ if

$$\frac{(SSE(\mathbf{X}_1) - SSE(\mathbf{X})) / (p - q)}{s^2} > F_{1-\alpha}(p - q, n - p - 1).$$

Example 7.5 (Testing the importance of additional predictors using the extra sum-of-squares approach) Male and female patrons rated the service in three establishments (locations) of a large restaurant chain. The service ratings were converted into an index. Table 7.2 contains the data for $n = 18$ customers. Each data point in the table is categorized according to location (1, 2, or 3) and gender (male = 0 and female = 1). This categorization has the format of a two-way table with unequal numbers of observations per cell. For instance, the combination of location 1 and male has 5 responses, while the combination of location 2 and female has 2 responses. Introducing three dummy variables to account for location and two dummy variables to account for gender, we can develop a regression model linking the service index Y to location, gender, and their “interaction” using the design matrix

Table 7.2 Restaurant-Service Data		
Location	Gender	Service (Y)
1	0	15.2
1	0	21.2
1	0	27.3
1	0	21.2
1	0	21.2
1	1	36.4
1	1	92.4
2	0	27.3
2	0	15.2
2	0	9.1
2	0	18.2
2	0	50.0
2	1	44.0
2	1	63.6
3	0	15.2
3	0	30.3
3	1	36.4
3	1	40.9

Lecture 7

S.Lan

The Classical Linear Regression Model

The Classical Linear
Regression
Least Square Estimation
Inferences About the
Regression Model

Multivariate Multiple Regression

Multivariate Multiple
Regression
Least Square Estimation
Likelihood Ratio Tests
for Regression
Parameters
Predictions

1 The Classical Linear Regression Model

The Classical Linear Regression

Least Square Estimation

Inferences About the Regression Model

2 Multivariate Multiple Regression

Multivariate Multiple Regression

Least Square Estimation

Likelihood Ratio Tests for Regression Parameters

Predictions

Lecture 7

S.Lan

The Classical Linear Regression Model

The Classical Linear Regression
Least Square Estimation
Inferences About the Regression Model

Multivariate Multiple Regression

Multivariate Multiple Regression
Least Square Estimation
Likelihood Ratio Tests for Regression Parameters
Predictions

- Now we consider the problem of modeling the relationship between m responses Y_1, Y_2, \dots, Y_m and p predictor variables X_1, \dots, X_p :

$$Y_k = \beta_{0k} + \sum_{j=1}^p X_j \beta_{jk} + \epsilon_k, \quad k = 1, \dots, m$$

- Denote $\mathbf{Y} = [y_{ik}]_{n \times m}$, $\boldsymbol{\epsilon} = [\epsilon_{ik}]_{n \times m}$ and $\boldsymbol{\beta} = [\beta_{jk}]_{(p+1) \times m}$. The *multivariate linear regression model* is

$$\mathbf{Y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

where we assume $E[\epsilon_k] = \mathbf{0}$ and $\text{Cov}[\epsilon_k, \epsilon_{k'}] = \sigma_{kk'} \mathbf{I}_n$. Denote the inter-trial covariance as $\boldsymbol{\Sigma} = [\sigma_{kk'}]_{m \times m}$.

- Viewing the multivariate linear regression as m parallel classical regression, we get LSE for each

$$\hat{\beta}_k = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y}_k, \quad k = 1, \dots, m$$

- Denote $\hat{\beta} = [\hat{\beta}_1, \dots, \hat{\beta}_m]$. We have $\epsilon = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y}$.
- The predicted values and residuals now become

$$\hat{\mathbf{Y}} = \mathbf{X} \hat{\beta} = \mathbf{H} \mathbf{Y}, \quad \mathbf{E} = \mathbf{Y} - \hat{\mathbf{Y}}$$

- Then we have

$$\mathbf{X}^T \mathbf{E} = \mathbf{0}, \quad \hat{\mathbf{Y}}^T \mathbf{E} = \mathbf{0}$$

- Therefore the decomposition of sum of squares

$$\mathbf{Y}^T \mathbf{Y} = \hat{\mathbf{Y}}^T \hat{\mathbf{Y}} + \mathbf{E}^T \mathbf{E}$$

Example 7.8 (Fitting a multivariate straight-line regression model) To illustrate the calculations of $\hat{\beta}$, \hat{Y} , and $\hat{\epsilon}$, we fit a straight-line regression model (see Panel 7.2),

$$Y_{j1} = \beta_{01} + \beta_{11}z_{j1} + \epsilon_{j1}$$

$$Y_{j2} = \beta_{02} + \beta_{12}z_{j1} + \epsilon_{j2}, \quad j = 1, 2, \dots, 5$$

to two responses Y_1 and Y_2 using the data in Example 7.3. These data, augmented by observations on an additional response, are as follows:

z_1	0	1	2	3	4
y_1	1	4	3	8	9
y_2	-1	-1	2	3	2

The design matrix \mathbf{Z} remains unchanged from the single-response problem. We find that

$$\mathbf{Z}' = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 3 & 4 \end{bmatrix} \quad (\mathbf{Z}'\mathbf{Z})^{-1} = \begin{bmatrix} .6 & -.2 \\ -.2 & .1 \end{bmatrix}$$

- We have the following property for LSE $\hat{\beta} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y}$

$$E[\hat{\beta}] = \beta, \quad \text{Cov}[\hat{\beta}_k, \hat{\beta}_{k'}] = \sigma_{kk'} (\mathbf{X}^T \mathbf{X})^{-1}$$

- The residual $\mathbf{E} = \mathbf{Y} - \hat{\mathbf{Y}}$ has the following property

$$E[\mathbf{E}] = \mathbf{0}, \quad \text{Cov}[\mathbf{e}_k, \mathbf{e}_{k'}] = \sigma_{kk'} (n - p - 1), \quad E[\mathbf{E}^T \mathbf{E}] / (n - p - 1) = \mathbf{\Sigma}.$$

- Moreover, $\text{Cov}[\hat{\beta}_k, \mathbf{e}_{k'}] = \mathbf{0}$.

- So far we have not imposed any distribution assumption.
- For the following inference, we need the matrix valued normal distribution.

Definition

A random matrix $\mathbf{Y}_{n \times m}$ follows *matrix normal distribution*, $\mathcal{MN}(\boldsymbol{\mu}, \mathbf{C}_{n \times n}, \boldsymbol{\Sigma}_{m \times m})$, if $\text{vec}(\mathbf{Y}) \sim N_{nm}(\text{vec}(\boldsymbol{\mu}), \boldsymbol{\Sigma} \otimes \mathbf{C})$, where $\text{vec}(\mathbf{Y}) = [\mathbf{Y}_1^T, \mathbf{Y}_2^T, \dots, \mathbf{Y}_m^T]^T$.

- The density of $\mathbf{Y} \sim \mathcal{MN}(\boldsymbol{\mu}, \mathbf{C}_{n \times n}, \boldsymbol{\Sigma}_{m \times m})$ is

$$(2\pi)^{-mn/2} |\mathbf{C}|^{-m/2} |\boldsymbol{\Sigma}|^{-n/2} \exp \left\{ -\frac{1}{2} \text{tr}[\boldsymbol{\Sigma}^{-1}(\mathbf{Y} - \boldsymbol{\mu})^T \mathbf{C}^{-1}(\mathbf{Y} - \boldsymbol{\mu})] \right\}.$$

- In particular, we assume $\mathbf{C} = \mathbf{I}_n$. And $\boldsymbol{\Sigma}_{m \times m}$ is the inter-trial covariance.

- Now we consider the similar hypothesis test in the multivariate case

$$H_0 : \beta_2 = \mathbf{0}_{(p-q) \times m}.$$

- The LRT involves the extra sum of squares and cross products

$$(\mathbf{Y} - \mathbf{X}_1 \hat{\beta}_1)^T (\mathbf{Y} - \mathbf{X}_1 \hat{\beta}_1) - (\mathbf{Y} - \mathbf{X} \hat{\beta})^T (\mathbf{Y} - \mathbf{X} \hat{\beta}) = n(\hat{\Sigma}_1 - \hat{\Sigma})$$

- The LRT statistic is defined as

$$\Lambda = \frac{\max_{\beta_1, \Sigma} L(\beta_1, \Sigma)}{\max_{\beta, \Sigma} L(\beta, \Sigma)} = \frac{L(\hat{\beta}_1, \hat{\Sigma})}{L(\hat{\beta}, \hat{\Sigma})} = \left(\frac{|\hat{\Sigma}|}{|\hat{\Sigma}_1|} \right)^{n/2}$$

- The corresponding Wilk's lambda can be used in the following test statistic

$$- \left[n - p - 1 - \frac{1}{2}(m - p + q + 1) \right] \log \left(\frac{|\hat{\Sigma}|}{|\hat{\Sigma}_1|} \right) \sim \chi^2(m(p - q)).$$

Example 7.9 (Testing the importance of additional predictors with a multivariate response) The service in three locations of a large restaurant chain was rated according to two measures of quality by male and female patrons. The first service-quality index was introduced in Example 7.5. Suppose we consider a regression model that allows for the effects of location, gender, and the location-gender interaction on both service-quality indices. The design matrix (see Example 7.5) remains the same for the two-response situation. We shall illustrate the test of no location-gender interaction in either response using Result 7.11. A computer program provides

$$\begin{pmatrix} \text{residual sum of squares} \\ \text{and cross products} \end{pmatrix} = n\hat{\Sigma} = \begin{bmatrix} 2977.39 & 1021.72 \\ 1021.72 & 2050.95 \end{bmatrix}$$

$$\begin{pmatrix} \text{extra sum of squares} \\ \text{and cross products} \end{pmatrix} = n(\hat{\Sigma}_1 - \hat{\Sigma}) = \begin{bmatrix} 441.76 & 246.16 \\ 246.16 & 366.12 \end{bmatrix}.$$

- Recall we have $\mathbf{Y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\epsilon}$ with $\boldsymbol{\epsilon} \sim \mathcal{MN}(\mathbf{0}, \mathbf{I}_n, \boldsymbol{\Sigma})$.
- The task is to predict the mean response corresponding to \mathbf{x}_0 . Note

$$\hat{\boldsymbol{\beta}}^T \mathbf{x}_0 \sim N_m(\boldsymbol{\beta}^T \mathbf{x}_0, \mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0 \boldsymbol{\Sigma}), \quad \perp \quad n \hat{\boldsymbol{\Sigma}} \sim W_{n-p-1}(\boldsymbol{\Sigma}).$$

- Therefore we have the T^2 -statistic

$$T^2 = \left(\frac{\hat{\boldsymbol{\beta}}^T \mathbf{x}_0 - \boldsymbol{\beta}^T \mathbf{x}_0}{\sqrt{\mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0}} \right)^T \left(\frac{n}{n-p-1} \hat{\boldsymbol{\Sigma}} \right)^{-1} \left(\frac{\hat{\boldsymbol{\beta}}^T \mathbf{x}_0 - \boldsymbol{\beta}^T \mathbf{x}_0}{\sqrt{\mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0}} \right) \sim \frac{m(n-p-1)}{n-p-m} F(m, n-p-m)$$

- The $100(1 - \alpha)\%$ CR is determined by $T^2 \leq \frac{m(n-p-1)}{n-p-m} F_{1-\alpha}(m, n-p-m)$.
- The $100(1 - \alpha)\%$ SCI for $E[Y_j] = \mathbf{x}_0^T \boldsymbol{\beta}_j$'s are

$$\mathbf{x}_0^T \hat{\boldsymbol{\beta}}_j \pm \sqrt{\frac{m(n-p-1)}{n-p-m} F_{1-\alpha}(m, n-p-m)} \sqrt{\mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0 \left(\frac{n}{n-p-1} \hat{\sigma}_{jj} \right)}, \quad j = 1, \dots, m.$$

- To predict the new response corresponding to \mathbf{x}_0 , i.e. $\mathbf{Y}_0 = \beta^T \mathbf{x}_0 + \epsilon_0$, we note

$$\mathbf{Y}_0 - \hat{\beta}^T \mathbf{x}_0 = (\beta - \hat{\beta})^T \mathbf{x}_0 + \epsilon_0 \sim N_m(\mathbf{0}, (1 + \mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0) \Sigma).$$

- Therefore, the $100(1 - \alpha)\%$ CR for \mathbf{Y}_0 becomes

$$(\mathbf{Y}_0 - \hat{\beta}^T \mathbf{x}_0)^T \left(\frac{n}{n - p - 1} \hat{\Sigma} \right)^{-1} (\mathbf{Y}_0 - \hat{\beta}^T \mathbf{x}_0) \leq (1 + \mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0) \frac{m(n - p - 1)}{n - p - m} F_{1-\alpha}(m, n - p - m)$$

- The $100(1 - \alpha)\%$ SCI for Y_{0j} 's are

$$\mathbf{x}_0^T \hat{\beta}_j \pm \sqrt{\frac{m(n - p - 1)}{n - p - m} F_{1-\alpha}(m, n - p - m)} \sqrt{(1 + \mathbf{x}_0^T (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{x}_0) \left(\frac{n}{n - p - 1} \hat{\sigma}_{jj} \right)}, \quad j = 1, \dots, m.$$

Example 7.10 (Constructing a confidence ellipse and a prediction ellipse for bivariate responses) A second response variable was measured for the computer-requirement problem discussed in Example 7.6. Measurements on the response Y_2 , disk input/output capacity, corresponding to the z_1 and z_2 values in that example were

$$\mathbf{y}'_2 = [301.8, 396.1, 328.2, 307.4, 362.4, 369.5, 229.1]$$

Obtain the 95% confidence ellipse for $\boldsymbol{\beta}'\mathbf{z}_0$ and the 95% prediction ellipse for $\mathbf{Y}'_0 = [Y_{01}, Y_{02}]$ for a site with the configuration $\mathbf{z}'_0 = [1, 130, 7.5]$.

Computer calculations provide the fitted equation

$$\hat{y}_2 = 14.14 + 2.25z_1 + 5.67z_2$$

with $s = 1.812$. Thus, $\hat{\boldsymbol{\beta}}'_{(2)} = [14.14, 2.25, 5.67]$. From Example 7.6,

$$\hat{\boldsymbol{\beta}}'_{(1)} = [8.42, 1.08, 42], \quad \mathbf{z}'_0\hat{\boldsymbol{\beta}}_{(1)} = 151.97, \quad \text{and} \quad \mathbf{z}'_0(\mathbf{Z}'\mathbf{Z})^{-1}\mathbf{z}_0 = .34725$$

We find that

$$\mathbf{z}'_0\hat{\boldsymbol{\beta}}_{(2)} = 14.14 + 2.25(130) + 5.67(7.5) = 349.17$$

and

$$\begin{aligned} n\hat{\boldsymbol{\Sigma}} &= \begin{bmatrix} (\mathbf{y}_{(1)} - \mathbf{Z}\hat{\boldsymbol{\beta}}_{(1)})'(\mathbf{y}_{(1)} - \mathbf{Z}\hat{\boldsymbol{\beta}}_{(1)}) & (\mathbf{y}_{(1)} - \mathbf{Z}\hat{\boldsymbol{\beta}}_{(1)})'(\mathbf{y}_{(2)} - \mathbf{Z}\hat{\boldsymbol{\beta}}_{(2)}) \\ (\mathbf{y}_{(2)} - \mathbf{Z}\hat{\boldsymbol{\beta}}_{(2)})'(\mathbf{y}_{(1)} - \mathbf{Z}\hat{\boldsymbol{\beta}}_{(1)}) & (\mathbf{y}_{(2)} - \mathbf{Z}\hat{\boldsymbol{\beta}}_{(2)})'(\mathbf{y}_{(2)} - \mathbf{Z}\hat{\boldsymbol{\beta}}_{(2)}) \end{bmatrix} \\ &= \begin{bmatrix} 5.80 & 5.30 \\ 5.30 & 13.13 \end{bmatrix} \end{aligned}$$

Since

$$\hat{\boldsymbol{\beta}}'\mathbf{z}_0 = \begin{bmatrix} \hat{\boldsymbol{\beta}}'_{(1)} \\ \hat{\boldsymbol{\beta}}'_{(2)} \end{bmatrix} \mathbf{z}_0 = \begin{bmatrix} \mathbf{z}'_0\hat{\boldsymbol{\beta}}_{(1)} \\ \mathbf{z}'_0\hat{\boldsymbol{\beta}}_{(2)} \end{bmatrix} = \begin{bmatrix} 151.97 \\ 349.17 \end{bmatrix}$$