

Drug Overview

DESCRIPTION

Metformin hydrochloride is an oral antihyperglycemic drug used in the management of type 2 diabetes mellitus. Metformin hydrochloride (N,N-dimethylimidodicarbonimidic diamide hydrochloride) is not chemically or pharmacologically related to any other class of oral antihyperglycemic agents.

DOSAGE & ADMINISTRATION

Adults: Starting dose 500 mg twice daily or 850 mg once daily with meals. Doses should be increased in increments of 500 mg weekly or 850 mg every 2 weeks, up to a maximum of 2550 mg per day. Pediatrics (10-16 years): 500 mg twice daily with meals.

Metformin

metformin hydrochloride

Biguanide Antidiabetic Agent

Drug Overview			Clinical Information			Safety Information		
DESCRIPTION metformin hydrochloride is an oral antihyperglycemic drug used in the management of type 2 diabetes mellitus. Metformin hydrochloride (N,N-dimethylimidodicarbonimidic diamide hydrochloride) is not chemically or pharmacologically related to any other class of oral antihyperglycemic agents.			INDICATIONS & USAGE metformin HCl tablets are indicated as an adjunct to diet and exercise to improve glycemic control in adults and pediatric patients 10 years of age and older with type 2 diabetes mellitus.			WARNINGS & PRECAUTIONS Lactic Acidosis. Postmarketing cases of metformin-associated lactic acidosis have resulted in death, hypothermia, hypotension, and resistant bradyarrhythmias. Symptoms include malaise, myalgias, respiratory distress, somnolence, and abdominal pain. Discontinue if lactic acidosis suspected.		
DOSAGE & ADMINISTRATION Adults. Starting dose 500 mg twice daily or 850 mg once daily with meals. Doses should be increased in increments of 500 mg weekly or 850 mg every 2 weeks, up to a maximum of 2550 mg per day. Pediatrics (10-16 years): 500 mg twice daily with meals.			CONTRAINDICATIONS metformin is contraindicated in patients with: Severe renal impairment (eGFR < 30 mL/min/1.73 m2). Known hypersensitivity to metformin hydrochloride. Acute or chronic metabolic acidosis, including diabetic ketoacidosis.			ADVERSE REACTIONS The most common adverse reactions include diarrhea, nausea/vomiting, flatulence, asthenia, indigestion, abdominal discomfort, and headache. Decreased Vitamin B12 levels observed in approximately 7% of patients.		