

DESCRIPTION

1.1 DESCRIPTION Metformin hydrochloride tablets, USP contain the antihyperglycemic agent metformin, which is a biguanide, in the form of monohydrochloride. The chemical name of metformin hydrochloride is N,N-dimethylimidodicarbonimidic diamide hydrochloride. The structural formula is as shown below: Metformin hydrochloride, USP is a white to off-white crystalline compound with a molecular formula of C₄ H₁₁ N₅ • HCl and a molecular weight of 165.63. Metformin hydrochloride is freely soluble ...

DOSAGE & ADMINISTRATION

2.1 DOSAGE & ADMINISTRATION Adult Dosage for metformin hydrochloride tablets: Starting dose: 500 mg orally twice a day or 850 mg once a day, with meals (2.1) Increase the dose in increments of 500 mg weekly or 850 mg every 2 weeks, up to a maximum dose of 2550 mg per day, given in divided doses (2.1) Doses above 2000 mg may be better tolerated given 3 times a day with meals (2.1) Pediatric Dosage for metformin hydrochloride tablets: Starting dose: 500 mg orally twice a day, with meals (2.2 ...

Metformin

METFORMIN HYDROCHLORIDE

Pharmaceutical Agent

Drug Overview		Clinical Information	Safety Information
DESCRIPTION 1 DESCRIPTION Metformin hydrochloride tablets, USP contain the antihyperglycemic agent metformin, which is a biguanide, in the form of monohydrochloride. The chemical name of metformin hydrochloride is N,N-dimethylimidodicarbonimidic diamide hydrochloride. The structural formula is as shown below: Metformin hydrochloride, USP is a white to off-white crystalline compound with a molecular formula of C 4 H 11 N 5 • HCl and a molecular weight of 165.63. Metformin hydrochloride is freely soluble ... DOSAGE & ADMINISTRATION 2 DOSAGE & ADMINISTRATION Adult Dosage for metformin hydrochloride tablets: Starting dose: 500 mg orally twice a day or 850 mg once a day, with meals (2.1) Increase the dose in increments of 500 mg weekly or 850 mg every 2 weeks, up to a maximum dose of 2550 mg per day, given in divided doses (2.1) Doses above 2000 mg may be better tolerated given 3 times a day with meals (2.1) Pediatric Dosage for metformin hydrochloride tablets: Starting dose: 500 mg orally twice a day, with meals (2.2 ...		INDICATIONS & USAGE 1 INDICATIONS & USAGE Metformin hydrochloride tablets are indicated as an adjunct to diet and exercise to improve glycemic control in adults and pediatric patients 10 years of age and older with type 2 diabetes mellitus. Metformin hydrochloride tablets are biguanide indicated as an adjunct to diet and exercise to improve glycemic control in adults and pediatric patients 10 years of age and older with type 2 diabetes mellitus. (1) CONTRAINDICATIONS 4 CONTRAINDICATIONS Metformin hydrochloride tablets are contraindicated in patients with: Severe renal impairment (eGFR below 30 mL/min/1.73 m 2) [see Warnings and Precautions (5.1)]. Hypersensitivity to metformin. Acute or chronic metabolic acidosis, including diabetic ketoacidosis, with or without coma. Severe renal impairment (eGFR below 30 mL/min/1.73 m 2) (4 , 5.1) Hypersensitivity to ...	WARNINGS & PRECAUTIONS 3 WARNINGS AND PRECAUTIONS Lactic Acidosis. See boxed warning. (5.1) Vitamin B 12 Deficiency: Metformin may lower vitamin B12 levels. Measure hematological parameters annually and vitamin B12 at 2 to 3 year intervals and manage any abnormalities. (5.2) Hypoglycemia with Concomitant Use with Insulin and Insulin Secretagogues: Increased risk of hypoglycemia when used in combination with insulin and/or an insulin secretagogue. Lower dose of ... ADVERSE REACTIONS 6 ADVERSE REACTIONS The following adverse reactions are also discussed elsewhere in the labeling: Lactic Acidosis [see Boxed Warning and Warnings and Precautions (5.1)]. Vitamin B12 Deficiency [see Warnings and Precautions (5.2)]. Hypoglycemia [see Warnings and Precautions (5.3) For metformin hydrochloride tablets, the most common adverse reactions (>5.0%) are diarrhea, nausea/vomiting, ...