




1 speciesthindex: How scientifically popular is a species?


2 Jessica Tam¹, Malgorzata Lagisz¹, Shinichi Nakagawa¹, and Will
3 Cornwell¹

4 ¹ Evolution & Ecology Research Centre and School of Biological, Earth and Environmental
5 Sciences, University of New South Wales, Sydney, Australia

DOI: [10.21105/joss.03776](https://doi.org/10.21105/joss.03776)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: Hugo Gruson 

Reviewers:

- [@kamapu](#)
- [@tom-gu](#)
- [@DrMattG](#)

Submitted: 21 September 2021

Published: 10 November 2021

License

Authors of papers retain
copyright and release the work
under a Creative Commons
Attribution 4.0 International
License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

6 Summary

7 Conservation efforts rely heavily on the body of scientific research on the relevant species.
8 Reintroduction projects, pest eradication, breeding programs and habitat restoration, all re-
9 quire sound scientific knowledge for their design and implementations. Unfortunately, there
10 is an uneven spread of research among species, taxonomic bias ([Bonnet et al., 2002](#)), such
11 that certain species receive more research interest than the others ([Donaldson et al., 2017](#);
12 [Santos et al., 2020](#); [Seddon et al., 2005](#); [Titley et al., 2017](#); [Troudet et al., 2017](#)). Over time,
13 taxonomic bias in research has widened the gap between well- and poorly-studied species.
14 Quantifying taxonomic bias is the first step towards allowing the scientific community to fill
15 the gaps and to direct scientific efforts towards where they are most needed for conservation.

16 Over the last decade, the quantification of research biases have gained popularity, as shown
17 by the growing number of studies on taxonomic bias among species of plants and animals
18 ([Adamo et al., 2021](#); [Donaldson et al., 2017](#); [Ducatez, 2019](#); [Fleming & Bateman, 2016](#);
19 [McKenzie & Robertson, 2015](#); [Robertson & McKenzie, 2015](#); [Santos et al., 2020](#); [Schiesari
20 et al., 2007](#); [Tensen, 2018](#); [Trimble & van Aarde, 2010](#)). Most of these studies evaluated
21 small groups of species, no more than a few hundred. Additionally, the majority of the studies
22 used publication or citation counts to identify the bias. However, publication count can only
23 capture the total research productivity. While citation count can capture research interest, it
24 can be easily inflated by highly cited papers ([Hirsch, 2005](#)), without equally considering each
25 publication. Here, we propose and implement the use of Hirsch's *h*-index ([Hirsch, 2005](#)) as
26 a better alternative in quantifying research interest for taxonomic groups of species including
27 genus and species.

28 Statement of need

29 speciesthindex aims to make streamline the calculation of *h*-index in the context of measuring
30 popularity of species in the scientific literature. The *h*-index was introduced by Hirsch ([Hirsch,
31 2005](#)) to compare the influence of academics with a single number ([Hirsch & Buéla-Casal,
32 2014](#)). The *h*-index is calculated as the total number of publications (*n*) that have at least
33 been cited *n* times, after ranking the publications in a descending order by their number of
34 citations. The *h*-index is now also being used to measure the research interest and influence of
35 the publications of different academic disciplines ([Banks, 2006](#); [Harzing & Alakangas, 2016](#)),
36 journals ([Braun et al., 2006](#)), countries ([Csajbók et al., 2007](#)), species of animals ([Fleming &
37 Bateman, 2016](#); [McKenzie & Robertson, 2015](#); [Robertson & McKenzie, 2015](#)), plants ([Adamo
38 et al., 2021](#)), and pathogens ([Cox et al., 2016](#)). speciesthindex calculates the *h*-index of
39 different species and genera.

40 The package speciesthindex connects to the Scopus and Web of Science databases (in-
41 terdisciplinary broad-range databases of academic literature) and extracts citation records of

relevant publications, as identified via a search query. It returns bibliometric information including the publication title, number of citations, publication type, etc. The binomial or genus names of the species should be used in the search query, instead of their common names, since the latter are less specific and may refer to more than 1, and even unrelated species. An example of such is 'pig,' which can refer to the domestic pig (*Sus scrofa* - an ungulate), or the guinea pig (*Cavia porcellus* - a rodent). Although this package also connects to Bielefeld Academic Search Engine (BASE), article extraction is not available, hence *h*-index calculation is unavailable with BASE.

There are two types of functions that connect to the literature databases. The count functions, e.g. `CountSpT()` and `CountSpTAK()` for species, and `CountGenusT()` and `CountGenusTAK()` for genus, return the total publication count without extracting any citation records. Whereas the fetch functions, e.g. `FetchSpT()` and `FetchSpTAK()` for species, and `FetchSpT()` and `FetchSpTAK()` for genus, extract citation records for index calculations. These functions are further distinguished by their suffixes "T" and "TAK." "T" functions only find publications with the species' name in the publication title, whereas "TAK" functions query articles' title, abstract and keywords. Apart from the *h*-index, *specieshindex* can also compute other established influence indices, including the *h5* index, the *m*-index, and the *i10* index. The *h5* index is the *h*-index of articles published in the most recent 5 years (Crotty, 2017). The *m*-index is the *h*-index divided by the number of years since the first publication (Hirsch, 2005). The *i10* index is the total number of articles with 10 or more citations; it is currently used by Google Scholar (Jacso, 2012).

Implementation

The following packages are required for *specieshindex* to work.

```
# Installation from GitHub
install.packages("rscopus")
install.packages("wosr")
install.packages("rbace")
install.packages("taxize")
install.packages("XML")
install.packages("jsonlite")
install.packages("httr")
install.packages("dplyr")
install.packages("data.table")
install.packages("tidyr")
devtools::install_github("jessicatytam/specieshindex", force = TRUE, build_vignettes = TRUE)

# Load the library
library(specieshindex)
```

Connecting to Scopus

An API key from Scopus is required to extract citation records from their database legally. Here are the steps to obtain the key.

1. Go to <https://dev.elsevier.com/> and click on the button I want an API key.
2. Create an account and log in.
3. Go to the My API Key tab on top of the page and click Create API Key.
4. Read the legal documents and check the boxes.

87 Afterwards, run the following line of code to gain access to the Scopus database:

```
88 # Setup API key
89 apikey <- "your_api_key_from_scopus"
```

90 Connecting to Web of Science

91 You do not need a key to extract data from Web of Science if you are using this package from
92 your institution's location. To gain access to the Web of Science database, run the following
93 line of code:

```
94 # Setup session ID
95 sid <- auth(username = NULL, password = NULL)
```

96 Connecting to BASE

97 Having a whitelisted IP address is essential when gaining access to the BASE database, which
98 you can get on <https://www.base-search.net/about/en/contact.php>. A token or API key,
99 however, is not required. Only count functions, e.g. CountSpT() and CountSpTAK(), are
100 available as BASE does not return citation counts. Hence, index calculations will also be
101 unavailable using this database.

102 Example

103 The following example will demonstrate the calculation of various indices with citation records
104 from Scopus. The species' binomial name (e.g. *Sarcophilus harrisii*) or genus name (e.g. *Sar-*
105 *cophilus*) is required for the extraction:

```
106 # Extract citation data
107 Woylie <- FetchSpTAK(db = "scopus", genus = "Bettongia", species = "penicillata")
108 Quokka <- FetchSpTAK(db = "scopus", genus = "Setonix", species = "brachyurus")
109 Platypus <- FetchSpTAK(db = "scopus", genus = "Ornithorhynchus", species = "anatin")
110 Koala <- FetchSpTAK(db = "scopus", genus = "Phascolarctos", species = "cinereus")
```

111 These four datasets are readily available within the package. An efficient way to calculate all
112 of the indices at once is to use the function Allindices().

```
113 # Calculate indices
114 W <- Allindices(data = Woylie, genus = "Bettongia", species = "penicillata")
115 Q <- Allindices(data = Quokka, genus = "Setonix", species = "brachyurus")
116 P <- Allindices(data = Platypus, genus = "Ornithorhynchus", species = "anatinus")
117 K <- Allindices(data = Koala, genus = "Phascolarctos", species = "cinereus")
118
119 # Combine citation records into a single dataframe
120 CombineSp <- rbind(W, Q, P, K)
121 CombineSp
```

122 ##	genus_species	species	genus	publications	citations
123 ## 1	Bettongia_penicillata	penicillata	Bettongia	113	1903
124 ## 2	Setonix_brachyurus	brachyurus	Setonix	242	3427
125 ## 3	Ornithorhynchus_anatinus	anatinus	Ornithorhynchus	321	6365

```

126 ## 4   Phascolarctos_cinereus   cinereus   Phascolarctos           773      14291
127 ##      journals articles reviews years_publishing h      m i10 h5
128 ## 1         55         110         3         43 26 0.605  54  7
129 ## 2         107         237         5         66 29 0.439 121  4
130 ## 3         153         308        13         67 41 0.612 177  7
131 ## 4         227         744        29        139 53 0.381 427 14

```


132 Plots

133 This package has 2 built-in plotting functions. They are `plotAllindices()` and `plotPub()`.
 134 `plotAllindices()` plots the *h*-index, *m*-index, *i10* index, and *h5* index in the same plot.

```

135 # Plot the indices
136 plotAllindices(CombineSp)

```



137
 138 **Figure 1.** The *h*-index, *m*-index, *i10* index, and *h5* index of the Woylie (*Bettongia penicillata*),
 139 Platypus (*Ornithorhynchus anatinus*), Koala (*Phascolarctos cinereus*), and Quokka (*Setonix*
 140 *brachyurus*).


141 `plotPub()` plots the total publications per year for species, after extracting data on years and
 142 number of publications per year with the `getYear()` function.

```

143 # Extract year and frequency
144 extract_year_W <- getYear(data = Woylie, genus = "Bettongia", species = "penicilla
145 extract_year_Q <- getYear(data = Quokka, genus = "Setonix", species = "brachyurus"
146 extract_year_P <- getYear(data = Platypus, genus = "Ornithorhynchus", species = "a
147 extract_year_K <- getYear(data = Koala, genus = "Phascolarctos", species = "cinere
148
149 # Combine year and frequency into a single dataframe
150 Combine_pub <- rbind(extract_year_W, extract_year_Q, extract_year_P, extract_year_

```

```
151
152 # Plot the number of publications by year
153 plotPub(Combine_pub)
```



154
155 **Figure 2.** The total number of publications per year of the Woylie (*Bettongia penicillata*),
156 Platypus (*Ornithorhynchus anatinus*), Koala (*Phascolarctos cinereus*), and Quokka (*Setonix*
157 *brachyurus*).

158 Acknowledgements

159 I acknowledge the contributions of the authors of the dependence packages: `rscopus`, `wosr`,
160 `rbace`, `taxize`, `XML`, `jsonlite`, `httr`, `dplyr`, `data.table`, `tidyr`. `speciesindex` is
161 enabled by Scopus, Web of Science, and BASE (Pieper & Summann, 2006).

162 References

- 163 Adamo, M., Chialva, M., Calevo, J., Bertoni, F., Dixon, K., & Mammola, S. (2021). Plant sci-
164 entists' research attention is skewed towards colourful, conspicuous and broadly distributed
165 flowers. *Nature Plants*, 7. <https://doi.org/10.1038/s41477-021-00912-2>
- 166 Banks, M. G. (2006). An extension of the Hirsch index: Indexing scientific topics and com-
167 pounds. *Scientometrics*, 69(1), 161–168. <https://doi.org/10.1007/s11192-006-0146-5>
- 168 Bonnet, X., Shine, R., & Lourda, O. (2002). Taxonomic chauvinism. *Trends in Ecology &*
169 *Evolution*, 17(1), 1–3. [https://doi.org/10.1016/S0169-5347\(01\)02381-3](https://doi.org/10.1016/S0169-5347(01)02381-3)
- 170 Braun, T., Glänzel, W., & Schubert, A. (2006). A Hirsch-type index for journals. *Sciento-*
171 *metrics*, 69(1), 169–173. <https://doi.org/10.1007/s11192-006-0147-4>

- 172 Cox, R., McIntyre, K. M., Sanchez, J., Setzkorn, C., Baylis, M., & Revie, C. W. (2016).
173 Comparison of the h-Index Scores Among Pathogens Identified as Emerging Hazards in
174 North America. *Transboundary and Emerging Diseases*, 63(1), 79–91. <https://doi.org/10.1111/tbed.12221>
175
- 176 Crotty, D. (2017). Other metrics: Beyond the impact factor. *European Heart Journal*, 38(35),
177 2646–2647. <https://doi.org/10.1093/eurheartj/ehx446>
- 178 Csajbók, E., Berhidi, A., Vasas, L., & Schubert, A. (2007). Hirsch-index for countries based
179 on Essential Science Indicators data. *Scientometrics*, 73(1), 91–117. <https://doi.org/10.1007/s11192-007-1859-9>
180
- 181 Donaldson, M. R., Burnett, N. J., Braun, D. C., Suski, C. D., Hinch, S. G., Cooke, S. J., &
182 Kerr, J. T. (2017). Taxonomic bias and international biodiversity conservation research.
183 *FACETS*, 1(1), 105–113. <https://doi.org/10.1139/facets-2016-0011>
- 184 Ducatez, S. (2019). Which sharks attract research? Analyses of the distribution of research
185 effort in sharks reveal significant non-random knowledge biases. *Reviews in Fish Biology*
186 *and Fisheries*, 29. <https://doi.org/10.1007/s11160-019-09556-0>
- 187 Fleming, P. A., & Bateman, P. W. (2016). The good, the bad, and the ugly: Which Australian
188 terrestrial mammal species attract most research? *Mammal Review*, 46(4), 241–254.
189 <https://doi.org/10.1111/mam.12066>
- 190 Harzing, A. W., & Alakangas, S. (2016). Google Scholar, Scopus and the Web of Science: A
191 longitudinal and cross-disciplinary comparison. *Scientometrics*, 106(2), 787–804. <https://doi.org/10.1007/s11192-015-1798-9>
192
- 193 Hirsch, J. E. (2005). An index to quantify an individual's scientific research output. *Pro-*
194 *ceedings of the National Academy of Sciences of the United States of America*, 102(46),
195 16569–16572. <https://doi.org/10.1073/pnas.0507655102>
- 196 Hirsch, J. E., & Buela-Casal, G. (2014). The meaning of the h-index. *Inter-*
197 *national Journal of Clinical and Health Psychology*, 14(2), 161–164. [https://doi.org/10.1016/S1697-2600\(14\)70050-X](https://doi.org/10.1016/S1697-2600(14)70050-X)
198
- 199 Jacso, P. (2012). Google scholar author citation tracker: Is it too little, too late? *Online*
200 *Information Review*, 36, 126–141. <https://doi.org/10.1108/14684521211209581>
- 201 McKenzie, A. J., & Robertson, P. A. (2015). Which Species Are We Researching and Why?
202 A Case Study of the Ecology of British Breeding Birds. *PLOS ONE*, 10(7), e0131004.
203 <https://doi.org/10.1371/journal.pone.0131004>
- 204 Pieper, D., & Summann, F. (2006). Bielefeld academic search engine (BASE) an end-user
205 oriented institutional repository search service. *Library Hi Tech*, 24. <https://doi.org/10.1108/07378830610715473>
206
- 207 Robertson, P. A., & Mckenzie, A. J. (2015). The scientific profiles of terrestrial mammals
208 in Great Britain as measured by publication metrics. *Mammal Review*, 45(2), 128–132.
209 <https://doi.org/10.1111/mam.12038>
- 210 Santos, J. W. dos, Correia, R. A., Malhado, A. C., Campos-Silva, J., Teles, D., Jepson, P., &
211 Ladle, R. (2020). Drivers of taxonomic bias in conservation research: A global analysis of
212 terrestrial mammals. *Animal Conservation*. <https://doi.org/10.1111/acv.12586>
- 213 Schiesari, L., Britta, G., & Grillitsch, H. (2007). Biogeographic biases in research and their
214 consequences for linking amphibian declines to pollution. *Conservation Biology : The*
215 *Journal of the Society for Conservation Biology*, 21, 465–471. <https://doi.org/10.1111/j.1523-1739.2006.00616.x>
216
- 217 Seddon, P. J., Soorae, P. S., & Launay, F. (2005). Taxonomic bias in reintroduction projects.
218 *Animal Conservation*, 8(1), 51–58. <https://doi.org/10.1017/S1367943004001799>

- 219 Tensen, L. (2018). Biases in wildlife and conservation research, using felids and canids as
220 a case study. *Global Ecology and Conservation*, 15, e00423. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.gecco.2018.e00423)
221 [gecco.2018.e00423](https://doi.org/10.1016/j.gecco.2018.e00423)
- 222 Titley, M. A., Snaddon, J. L., & Turner, E. C. (2017). Scientific research on animal biodiversity
223 is systematically biased towards vertebrates and temperate regions. *PLOS ONE*, 12(12),
224 e0189577. <https://doi.org/10.1371/journal.pone.0189577>
- 225 Trimble, M., & van Aarde, R. (2010). Species inequality in scientific study. *Conservation*
226 *Biology : The Journal of the Society for Conservation Biology*, 24, 886–890. [https:](https://doi.org/10.1111/j.1523-1739.2010.01453.x)
227 [//doi.org/10.1111/j.1523-1739.2010.01453.x](https://doi.org/10.1111/j.1523-1739.2010.01453.x)
- 228 Troudet, J., Grandcolas, P., Blin, A., Vignes-Lebbe, R., & Legendre, F. (2017). Taxonomic
229 bias in biodiversity data and societal preferences. *Scientific Reports*, 7(1), 1–14. [https:](https://doi.org/10.1038/s41598-017-09084-6)
230 [//doi.org/10.1038/s41598-017-09084-6](https://doi.org/10.1038/s41598-017-09084-6)

DRAFT