




%PT_GSDesign: A SAS Macro for Group Sequential Designs with Time-to-event Data using the Concept of Proportional Time

Milind A. Phadnis^{*1} and Nadeesha Thewarapperuma¹

¹ University of Kansas Medical Center, Department of Biostatistics & Data Science

DOI: [10.21105/joss.03813](https://doi.org/10.21105/joss.03813)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Vincent Knight](#) 

Reviewers:

- [@uagla](#)
- [@lrackley](#)

Submitted: 04 March 2021

Published: 11 October 2021

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

Treatments that are found to be promising in a Phase II trial are studied more comprehensively in a Phase III trial where researchers aim to investigate the effectiveness and safety of the new treatment against the current standard-of-care. While traditional approaches require the calculation of a fixed sample size depending on the type I error, power and clinically important treatment effect, in the medical setting they suffer from the limitation that patients are continually being accrued into a study based on the accrual rate, the availability of qualified patients (based on inclusion/exclusion criteria) and the possibility of random dropouts among many factors. Thus, the primary outcome of interest is not available simultaneously on all patients and researchers may be interested to look at outcomes on the early enrollees and use that as a basis to decide whether the trial should be continued. Sequential testing in large-sized Phase III trials with interim points can be used to - {i} stop the trial early for overwhelming evidence of efficacy, {ii} stop the trial early for overwhelming evidence for futility, and {iii} continue the trial for lack of evidence of efficacy or futility.

A Group Sequential Design (GSD) formalizes the concept by providing a statistical framework under which either of the three decisions can be taken after looking at interim results. Ethical, financial and administrative requirements often guide the statistical designs of GSDs (see [Enas et al. \(1989\)](#); [Jennison & Turnbull \(1990\)](#); [Ellenberg et al. \(2002\)](#)). Such GSDs have been well developed for continuous and binary outcomes and have a long history starting with quality control applications ([Wald \(1947\)](#)) and progressing to the medical setting ([Armitage \(1960\)](#)). Vast literature is available on this topic in many books ([Whitehead \(1997\)](#); [Jennison & Turnbull \(2000\)](#); [Proschan et al. \(2006\)](#); [Dmitrienko et al. \(2005\)](#); [Wassmer & Brannath \(2016\)](#)) and overview articles ([Whitehead \(1999\)](#); [Todd \(2007\)](#); [Mazumdar & Bang \(2008\)](#)). When dealing with time-to-event outcome, a repeated significance testing approach incorporating a family of designs ([Pocock \(1977\)](#); [O'Brien & Fleming \(1979\)](#); [Wang & Tsatis \(1987\)](#)) can be combined with the error spending method ([Lan & DeMets \(1983\)](#)) to implement a GSD using a log-rank test or by using the proportional hazards (PH) assumption. Popular statistical software such as GPower, PASS, and nQuery often implement GSDs for time-to-event outcome using the weighted and unweighted versions of the log-rank test either explicitly assuming exponentially distributed survival times or with the PH assumption and are able to incorporate complexities of survival outcomes such as random dropouts, prespecified accrual and follow-up times, varying accrual patterns, equal/unequal spaced interim testing points (looks), efficacy-only designs, efficacy and futility designs, binding and non-binding futility rules, and many other flexible features specific to time-to-event outcomes.

^{*}Corresponding author

Statement of need

When the underlying assumptions that drive the analytical and simulation-based approaches using the framework of the log-rank test are not valid, hardly any alternate methods are available in literature or in standard statistical software. Recent developments in this field have considered relaxing the PH assumption in favor of a 'proportionality of time (PT)' assumption leading to development of GSDs in the context of an accelerated failure time (AFT) model (Phadnis & Mayo (2020)). The authors have described various scenarios in the biomedical setting where their approach could be advantageous compared to the standard methods with the help of real-life examples. Their proposed GSD method provides an alternate approach when the PH assumption is not appropriate and allows various hazard shapes (increasing/decreasing monotonically over time, bathtub shaped, arc-shaped) using the generalized gamma distribution. The purpose of this paper is to present a fully functional SAS macro that can be used to implement their GSD method (SAS Institute Inc (2010)). The SAS macro incorporates multitude of design features specific to a two-arm GSD for time-to-event outcomes and includes validation for any parameters defined by the user, as well as suggestions for correcting erroneous input.

Acknowledgements

The High performance computing capabilities, which were used to conduct some of the analyses described in this paper, were supported in part by the National Cancer Institute (NCI) Cancer Center Support Grant P30 CA168524; the Kansas IDeA Network of Biomedical Research Excellence Bioinformatics Core, supported by the National Institute of General Medical Science award P20 GM103418; and the Kansas Institute for Precision Medicine COBRE, supported by the National Institute of General Medical Science award P20 GM130423.

References

- Armitage, P. (1960). *Sequential medical trials* (1st ed.). Thomas.
- Dmitrienko, A., Molenberghs, G., Chuang-Stein, C., & Offen, W. W. (2005). *Analysis of clinical trials using SAS: A practical guide*. SAS Institute.
- Ellenberg, S., Fleming, T., & DeMets, D. (2002). *Data monitoring committees in clinical trials: A practical perspective*. John Wiley & Sons.
- Enas, G., Dornseif, B., Sampson, C., Rochhol, F., & Wu, J. (1989). Monitoring versus interim analysis of clinical trials: Perspective from the pharmaceutical industry. *Controlled Clinical Trials*, 10, 57–70. [https://doi.org/10.1016/0197-2456\(89\)90018-4](https://doi.org/10.1016/0197-2456(89)90018-4)
- Jennison, C., & Turnbull, B. (1990). Statistical approaches to interim monitoring of medical trials: A review and commentary. *Statistical Science*, 5, 299–317. <https://doi.org/10.1214/ss/1177012099>
- Jennison, C., & Turnbull, B. (2000). *Group sequential methods with applications to clinical trials*. Chapman & Hall/CRC. <https://doi.org/10.1201/9780367805326>
- Lan, K., & DeMets, D. (1983). Discrete sequential boundaries for clinical trials. *Biometrika*, 70, 659–663. <https://doi.org/10.2307/2336502>
- Mazumdar, M., & Bang, H. (2008). *Sequential and group sequential designs in clinical trials: Guidelines for practitioners; handbook of statistics 27; series title: Epidemiology and medical statistics*. Elsevier.

- 82 O'Brien, P., & Fleming, T. (1979). A multiple testing procedure for clinical trials. *Biometrics*,
83 35, 549–556. <https://doi.org/10.2307/2530245>
- 84 Phadnis, M., & Mayo, M. (2020). Group sequential design for time-to-event data using the
85 concept of proportional time. *Statistical Methods in Medical Research*, 29, 1867–1890.
86 <https://doi.org/10.1177/0962280219876313>
- 87 Pocock, S. (1977). Group sequential methods in the design and analysis of clinical trials.
88 *Biometrika*, 64, 191–199. <https://doi.org/10.2307/2335684>
- 89 Proschan, M., Lan, K., & Wittes, J. (2006). *Statistical monitoring of clinical trial*. Springer.
- 90 SAS Institute Inc. (2010). *SAS software, version 9.4* (Version 0.20.2) [Computer software].
91 https://www.sas.com/en_us/software/sas9.html
- 92 Todd, S. (2007). A 25-year review of sequential methodology in clinical trials. *Statistics in*
93 *Medicine*, 26, 237–252. <https://doi.org/10.1002/sim.2763>
- 94 Wald, A. (1947). *Sequential analysis*. John Wiley & Sons.
- 95 Wang, S., & Tsiatis, A. (1987). Approximately optimal one-parameter boundaries for group
96 sequential trials. *Biometrics*, 43, 193–199. <https://doi.org/10.2307/2531959>
- 97 Wassmer, G., & Brannath, W. (2016). *Group sequential and confirmatory adaptive designs*
98 *in clinical trials*. Springer.
- 99 Whitehead, J. (1997). *The design and analysis of sequential clinical trials. 2nd edition*. John
100 Wiley & Sons.
- 101 Whitehead, J. (1999). A unified theory for sequential clinical trials. *Statistics in Medicine*, 18,
102 2271–2286. [https://doi.org/10.1002/\(sici\)1097-0258\(19990915/30\)18:17/18%3C2271::](https://doi.org/10.1002/(sici)1097-0258(19990915/30)18:17/18%3C2271::aid-sim254%3E3.0.co;2-z)
103 [aid-sim254%3E3.0.co;2-z](https://doi.org/10.1002/(sici)1097-0258(19990915/30)18:17/18%3C2271::aid-sim254%3E3.0.co;2-z)