FTL FOR OQO PROBLEMS

- One popular instantiation of the online learning problem is the problem of *online quadratic optimization* (OQO).
- In its most general form, the loss function is thereby defined as

$$\ell(a_t, z_t) = \frac{1}{2}||a_t - z_t||_2^2,$$

where $\mathcal{A}, \mathcal{Z} \subset \mathbb{R}^d$.

- **Proposition:** Using FTL on any online quadratic optimization problem with $\mathcal{A} = \mathbb{R}^d$ and $V = \sup_{z \in \mathcal{Z}} ||z||_2$, leads to a regret of

$$R_T^{\mathrm{FTL}} \leq 4V^2((\log(T) + 1)).$$

FTL FOR OQO PROBLEMS: ANALYSIS

- **Proof:**
  - In the following, we denote $a_1^{\mathrm{FTL}}, a_2^{\mathrm{FTL}}, \ldots$ simply by $a_1, a_2, \ldots$ $a_1, a_2, \ldots$

FTL FOR OQO PROBLEMS: ANALYSIS

- **Proof:**
  - In the following, we denote $a_1^{\mathrm{FTL}}, a_2^{\mathrm{FTL}}, \ldots$ simply by $a_1, a_2, \ldots$

$a_1, a_2, \ldots$

Reminder (Useful Lemma):

$$R_T^{\mathrm{FTL}} \leq \sum_{t=1}^{T} L(a_t^{\mathrm{FTL}}, z_t) - \sum_{t=1}^{T} L(a_{t+1}^{\mathrm{FTL}}, z_t)$$

FTL FOR OQO PROBLEMS: ANALYSIS

- **Proof:**
  - In the following, we denote $a_1^{\mathrm{FTL}}, a_2^{\mathrm{FTL}}, \ldots$ simply by $a_1, a_2, \ldots$

Reminder (Useful Lemma):

$$R_T^{\mathrm{FTL}} \leq \sum_{t=1}^{T} L(a_t^{\mathrm{FTL}}, z_t) - \sum_{t=1}^{T} L(a_{t+1}^{\mathrm{FTL}}, z_t)$$

  - Using this lemma, we just have to show that

$$\sum_{t=1}^{T} (L(a_t, z_t) - L(a_{t+1}, z_t)) \leq 4L^2 \cdot (\log(T) + 1) . \qquad (1)$$

FTL FOR OQO PROBLEMS: ANALYSIS

- **Proof:**
  - In the following, we denote $a_1^{\mathrm{FTL}}, a_2^{\mathrm{FTL}}, \ldots$ simply by $a_1, a_2, \ldots$

Reminder (Useful Lemma):

$$R_T^{\mathrm{FTL}} \leq \sum_{t=1}^{T} L(a_t^{\mathrm{FTL}}, z_t) - \sum_{t=1}^{T} L(a_{t+1}^{\mathrm{FTL}}, z_t)$$

  - Using this lemma, we just have to show that

$$\sum_{t=1}^{T} (L(a_t, z_t) - L(a_{t+1}, z_t)) \leq 4L^2 \cdot (\log(T) + 1) . \quad (1)$$

  - So, we will prove (1). For this purpose, we compute the explicit form of the actions of FTL for this type of online learning problem.

FTL FOR OQO PROBLEMS: ANALYSIS

- Claim: It holds that $a_t = \frac{1}{t-1} \cdot \sum_{s=1}^{t-1} z_s$, if $(a, z) \mapsto \frac{1}{2}\frac{1}{2}\|a - z\|_2^2$.

FTL FOR OQO PROBLEMS: ANALYSIS

- Claim: It holds that $a_t = \frac{1}{t-1} \cdot \sum_{s=1}^{t-1} z_s$, if $\ell(a, z) = \frac{1}{2}\|a - z\|_2^2$.
  - Recall that

$$a_t^{\mathrm{FTL}} = \arg\min_{a \in \mathcal{A}} \sum_{s=1}^{t-1} \ell(a, z_s) = \arg\min_{a \in \mathcal{A}} \sum_{s=1}^{t-1} \frac{1}{2}\|a - z_s\|_2^2 .$$

FTL FOR OQO PROBLEMS: ANALYSIS

- Claim: It holds that $a_t = \frac{1}{t-1} \cdot \sum_{s=1}^{t-1} z_s$, if $\ell(a, z) = \frac{1}{2}\|a - z\|_2^2$.
  - Recall that

$$a_t^{\mathrm{FTL}} = \arg\min_{a \in \mathcal{A}} \sum_{s=1}^{t-1} \ell(a, z_s) = \arg\min_{a \in \mathcal{A}} \sum_{s=1}^{t-1} \frac{1}{2}\|a - z_s\|_2^2.$$

  - So, we have to find the minimizer of the function

$$f(a) := \sum_{s=1}^{t-1} \frac{1}{2}\|a - z_s\|_2^2 = \sum_{s=1}^{t-1} \frac{1}{2}(a - z_s)^\top (a - z_s).$$

FTL FOR OQO PROBLEMS: ANALYSIS

- Claim: It holds that $a_t = \frac{1}{t-1} \cdot \sum_{s=1}^{t-1} z_s$, if $\ell(a, z) = \frac{1}{2}\|a - z\|_2^2$.
  - Recall that

$$a_t^{\mathrm{FTL}} = \arg\min_{a \in \mathcal{A}} \sum_{s=1}^{t-1} \ell(a, z_s) = \arg\min_{a \in \mathcal{A}} \sum_{s=1}^{t-1} \frac{1}{2}\|a - z_s\|_2^2.$$

  - So, we have to find the minimizer of the function

$$f(a) := \sum_{s=1}^{t-1} \frac{1}{2}\|a - z_s\|_2^2 = \sum_{s=1}^{t-1} \frac{1}{2}(a - z_s)^\top (a - z_s).$$

  - Compute $\nabla f(a) = \sum_{s=1}^{t-1} a - z_s = (t-1)a - \sum_{s=1}^{t-1} z_s$, which we set to zero and solve with respect to a to obtain the claim.

    (f is convex, so that this leads indeed to a minimizer.)

FTL FOR OQO PROBLEMS: ANALYSIS

- Hence, a_t is the empirical average of $z_1, \ldots, z_{t-1}$ and we can provide the following incremental update formula for its computation

$$a_{t+1} = \frac{1}{t} \cdot \sum_{s=1}^{t} z_s = \frac{1}{t}\left(z_t + \sum_{s=1}^{t-1} z_s\right)$$
$$= \frac{1}{t}(z_t + (t-1)a_t) = \frac{1}{t}z_t + \left(1 - \frac{1}{t}\right) a_t.$$

- From the last display we derive that

$$a_{t+1} - z_t = \left(1 - \frac{1}{t}\right) \cdot a_t + \frac{1}{t}z_t - z_t = \left(1 - \frac{1}{t}\right) \cdot (a_t - z_t).$$

- Claim:

$$L(a_t, z_t) - L(a_{t+1}, z_t) \leq \frac{1}{t} \|a_t - z_t\|_2^2. \qquad (2)$$

FTL FOR OQO PROBLEMS: ANALYSIS

Reminder: $a_{t+1} - z_t = \left(1 - \frac{1}{t}\right) \cdot (a_t - z_t).$

- Indeed, this can be seen as follows

$$
\begin{aligned}
L(a_t, z_t) - L(a_{t+1}, z_t) &= \frac{1}{2}\|a_t - z_t\|_2^2 - \frac{1}{2}\|a_{t+1} - z_t\|_2^2 \\
&= \frac{1}{2}\left(\|a_t - z_t\|_2^2 - \|a_{t+1} - z_t\|_2^2\right) \\
&= \frac{1}{2}\left(\|a_t - z_t\|_2^2 - \left\|\left(1 - \frac{1}{t}\right) \cdot (a_t - z_t)\right\|_2^2\right).
\end{aligned}
$$

FTL FOR OQO PROBLEMS: ANALYSIS

Reminder: $a_{t+1} - z_t = \left(1 - \frac{1}{t}\right) \cdot (a_t - z_t).$

- Indeed, this can be seen as follows

$$
\begin{aligned}
L(a_t, z_t) - L(a_{t+1}, z_t) &= \frac{1}{2}\|a_t - z_t\|_2^2 - \frac{1}{2}\|a_{t+1} - z_t\|_2^2 \\
&= \frac{1}{2}\left(\|a_t - z_t\|_2^2 - \|a_{t+1} - z_t\|_2^2\right) \\
&= \frac{1}{2}\left(\|a_t - z_t\|_2^2 - \left\|\left(1 - \frac{1}{t}\right) \cdot (a_t - z_t)\right\|_2^2\right).
\end{aligned}
$$

- And from this,

$$
\begin{aligned}
L(a_t, z_t) - L(a_{t+1}, z_t) &= \frac{1}{2}\left(\|a_t - z_t\|_2^2 - \left(1 - \frac{1}{t}\right)^2 \cdot \|a_t - z_t\|_2^2\right) \\
&= \frac{1}{2}\left(1 - \left(1 - \frac{1}{t}\right)^2\right) \|a_t - z_t\|_2^2 \\
&= \left(\frac{1}{t} - \frac{1}{2t^2}\right) \|a_t - z_t\|_2^2 \\
&\leq \frac{1}{t} \|a_t - z_t\|_2^2.
\end{aligned}
$$

FTL FOR OQO PROBLEMS: ANALYSIS

Reminder: $L(a_t, z_t) - L(a_{t+1}, z_t) \leq \frac{1}{t} \|a_t - z_t\|_2^2$. (2)

- Since by assumption $L = \sup_{z \in \mathcal{Z}} \|z\|_2$ and a_t is the empirical average of $z_1, \ldots, z_{t-1}$, we have that $\|a_t\|_2 \leq L$.

FTL FOR OQO PROBLEMS: ANALYSIS

Reminder: $L(a_t, z_t) - L(a_{t+1}, z_t) \leq \frac{1}{t} \|a_t - z_t\|_2^2$. (2)

- Since by assumption $L = \sup_{z \in \mathcal{Z}} \|z\|_2$ and a_t is the empirical average of $z_1, \ldots, z_{t-1}$, we have that $\|a_t\|_2 \leq L$.
- Now the triangle inequality states that for any two vectors $x, y \in \mathbb{R}^d$ it holds that

$$\|x + y\|_2 \leq \|x\|_2 + \|y\|_2,$$

so that

$$\|a_t - z_t\|_2 \leq \|a_t\|_2 + \|z_t\|_2 \leq 2L. \quad (3)$$

FTL FOR OQO PROBLEMS: ANALYSIS

Reminder: $L(a_t, z_t) - L(a_{t+1}, z_t) \leq \frac{1}{t} \|a_t - z_t\|_2^2$. (2)

- Since by assumption $L = \sup_{z \in \mathcal{Z}} \|z\|_2$ and a_t is the empirical average of $z_1, \ldots, z_{t-1}$, we have that $\|a_t\|_2 \leq L$.
- Now the triangle inequality states that for any two vectors $x, y \in \mathbb{R}^d$ it holds that

$$\|x + y\|_2 \leq \|x\|_2 + \|y\|_2,$$

so that

$$\|a_t - z_t\|_2 \leq \|a_t\|_2 + \|z_t\|_2 \leq 2L. \tag{3}$$

- Summing over all t in (2) and using (3) we arrive at

$$\sum_{t=1}^{T} (L(a_t, z_t) - L(a_{t+1}, z_t)) \leq \sum_{t=1}^{T} \left(\frac{1}{t} \|a_t - z_t\|_2^2 \right) \leq \sum_{t=1}^{T} \frac{1}{t} (2L)^2$$

$$= 4L^2 \cdot \sum_{t=1}^{T} \frac{1}{t}.$$

FTL FOR OQO PROBLEMS: ANALYSIS

Reminder: $\sum_{t=1}^{T} \left(\ell(a_t, z_t) - \ell(a_{t+1}, z_t) \right) \leq 4L^2 \cdot \sum_{t=1}^{T} \frac{1}{t}$

FTL FOR OQO PROBLEMS: ANALYSIS

Reminder: $\sum_{t=1}^{T} \left((a_t, z_t) - (a_{t+1}, z_t) \right) \leq 4L^2 \cdot \sum_{t=1}^{T} \frac{1}{t}$

- Now, it holds that $\sum_{t=1}^{T} \frac{1}{t} \leq \log(T) + 1$, so that we obtain

$$\sum_{t=1}^{T} \left((a_t, z_t) - (a_{t+1}, z_t) \right) \leq 4L^2 \cdot \sum_{t=1}^{T} \frac{1}{t} \leq 4L^2 \cdot (\log(T) + 1),$$

which is what we wanted to prove. □