Introduction to Machine Learning

ML-Basics
Optimization



Learning goals

- Understand how the risk function is optimized to learn the optimal parameters of a model
- Understand the idea of gradient descent as a basic risk optimizer

LEARNING AS PARAMETER OPTIMIZATION

- We can operationalize the search for a model f that matches training data best, by looking for its parametrization $\boldsymbol{\theta} \in \Theta$ with lowest empirical risk $\mathcal{R}_{\text{emp}}(\boldsymbol{\theta})$.
- Therefore, we usually traverse the error surface downwards; often by local search from a starting point to its minimum.



LOCAL MINIMA



If $\mathcal{R}_{\text{emp}}$ is continuous in $\boldsymbol{\theta}$ we can define a **local minimum** $\hat{\boldsymbol{\theta}}$:

$$\exists \epsilon > 0 \; \forall \boldsymbol{\theta} \text{ with } \left\| \hat{\boldsymbol{\theta}} - \boldsymbol{\theta} \right\| < \epsilon : \quad \mathcal{R}_{\text{emp}}(\hat{\boldsymbol{\theta}}) \leq \mathcal{R}_{\text{emp}}(\boldsymbol{\theta}).$$

Clearly every global minimum is also a local minimum. Finding a local minimum is easier than finding a global minimum.



LOCAL MINIMA AND STATIONARY POINTS

If $\mathcal{R}_{\text{emp}}$ is continuously differentiable in $\boldsymbol{\theta}$ then a **sufficient condition** for a local minimum is that $\hat{\boldsymbol{\theta}}$ is **stationary** with 0 gradient, so no local improvement is possible:

$$\frac{\partial}{\partial \boldsymbol{\theta}} \mathcal{R}_{\text{emp}}(\hat{\boldsymbol{\theta}}) = 0$$

and the Hessian $\frac{\partial^2}{\partial \boldsymbol{\theta}^2} \mathcal{R}_{\text{emp}}(\hat{\boldsymbol{\theta}})$ is positive definite. While the neg. gradient points into the direction of fastest local decrease, the Hessian measures local curvature of $\mathcal{R}_{\text{emp}}$.



$\frac{\partial}{\partial \boldsymbol{\theta}} \mathcal{R}_{\text{emp}}(\boldsymbol{\theta})$ const. pos. def. Hessian const. neg. def. Hessian

GRADIENT DESCENT

The idea of GD is to iteratively go from the current candidate $\boldsymbol{\theta}^{[t]}$ in the direction of the negative gradient, i.e., the direction of the steepest descent, with learning rate α to the next $\boldsymbol{\theta}^{[t+1]}$:

$$\boldsymbol{\theta}^{[t+1]} = \boldsymbol{\theta}^{[t]} - \alpha \frac{\partial}{\partial \boldsymbol{\theta}} \mathcal{R}_{\text{emp}}(\boldsymbol{\theta}^{[t]}).$$



We choose a random start $\boldsymbol{\theta}^{[0]}$ with risk $\mathcal{R}_{\text{emp}}(\boldsymbol{\theta}^{[0]}) = 76.25$.

GRADIENT DESCENT - EXAMPLE



Now we follow in the direction of the negative gradient at $\boldsymbol{\theta}^{[0]}$.



We arrive at $\boldsymbol{\theta}^{[1]}$ with risk $\mathcal{R}_{\text{emp}}(\boldsymbol{\theta}^{[1]}) \approx 42.73$.
We improved:
$\mathcal{R}_{\text{emp}}(\boldsymbol{\theta}^{[1]}) < \mathcal{R}_{\text{emp}}(\boldsymbol{\theta}^{[0]})$.

GRADIENT DESCENT - EXAMPLE



Again we follow in the direction of the negative gradient, but now at $\boldsymbol{\theta}^{[1]}$.



Now $\boldsymbol{\theta}^{[2]}$ has risk $\mathcal{R}_{\text{emp}}(\boldsymbol{\theta}^{[2]}) \approx 25.08$.

GRADIENT DESCENT - EXAMPLE



We iterate this until some form of convergence or termination.



We arrive close to a stationary $\hat{\boldsymbol{\theta}}$ which is hopefully at least a local minimum.

GRADIENT DESCENT - LEARNING RATE

- The negative gradient is a direction that looks locally promising to reduce $\mathcal{R}_{\text{emp}}$.
- Hence it weights components higher in which $\mathcal{R}_{\text{emp}}$ decreases more.
- However, the length of $-\frac{\partial}{\partial \boldsymbol{\theta}} \mathcal{R}_{\text{emp}}$ measures only the local decrease rate, i.e., there are no guarantees that we will not go "too far".
- We use a learning rate α to scale the step length in each iteration. Too much can lead to overstepping and no convergence, too low leads to slow convergence.
- Usually, a simple constant rate or rate-decrease mechanisms to enforce local convergence are used



good convergence for α_1



poor convergence for $\alpha_2 (< \alpha_1)$



no convergence for $\alpha_3 (> \alpha_1)$

FURTHER TOPICS

- Few models, e.g. linear regression, can be optimized analytically.
- GD is a so-called first-order method. Second-order methods (like Newton-Raphson) use the Hessian to refine the search direction for faster convergence.
- There exist many improvements of GD (momentum, ADAM), e.g., to smartly control the learn rate, to escape saddle points, to mimic second order behavior without computing the Hessian.
- If the gradient of is not computed on the complete data, but instead on small, random batches, this is **stochastic gradient descent** (SGD). For large-scale problems, this is usually more efficient.