OUT-OF-BAG VS IN-BAG OBSERVATIONS

ID	Color	Form	Length	Origin	Banana
1	yellow	oblong	14	imported	yes
2	brown	oblong	10	imported	yes
3	red	round	16	domestic	no

OOB

IB

predict

Bootstrapping to train tree 1

ID	Color	Form	Length	Origin	Banana
1	yellow	oblong	14	imported	yes
3	red	round	16	domestic	no
3	red	round	16	domestic	no

Tree 1

- IB observations for m-th bootstrap:
  $\mathrm{IB}^{[m]} = \{i \in \{1, \ldots, n\} \mid \left(\mathbf{x}^{(i)}, y^{(i)}\right) \in \mathcal{D}^{[m]}\}$
- OOB observations for m-th bootstrap:
  $\mathrm{OOB}^{[m]} = \{i \in \{1, \ldots, n\} \mid \left(\mathbf{x}^{(i)}, y^{(i)}\right) \notin \mathcal{D}^{[m]}\}$
- Nr. of trees where i-th observation is OOB:
  $S_{\mathrm{OOB}}^{(i)} = \sum_{m=1}^{M} \mathbb{I}(i \in \mathrm{OOB}^{[m]}).$

OUT-OF-BAG ERROR ESTIMATE

Predict i-th observation with all trees $\hat{b}^{[m]}$ for which it is OOB:

ID	Color	Form	Length	Origin	Banana	OOB trees
1	yellow	oblong	14	imported	yes	{2}
2	brown	oblong	10	imported	yes	{1, 3, 4}
3	red	round	16	domestic	no	{2, 4}



OOB prediction $\hat{\pi}^{(2)}_{\text{OOB}} = 2/3$. Evaluating all OOB predictions with some loss function L or set-based metric ρ estimates the GE.

As we do not violate the **untouched test set principle**, $\widehat{\text{GE}}$ is not *optimistically* biased.

OUT-OF-BAG ERROR PSEUDO CODE

Out-Of-Bag error estimation

1: **Input:** $\mathrm{OOB}^{[m]}, \hat{b}^{[m]} \forall m \in \{1, \ldots, M\}$
2: **for** $i = 1 \rightarrow n$ **do**
3: Compute the ensemble OOB prediction for observation i, e.g., for regression:

$$\hat{f}^{(i)}_{\mathrm{OOB}} = \frac{1}{S_{\mathrm{OOB}^{(i)}}} \sum_{m=1}^{M} \mathbb{I}(i \in \mathrm{OOB}^{[m]}) \cdot \hat{f}^{[m]}(\mathbf{x}^{(i)})$$

4: **end for**
5: Average losses over all observations:

$$\widehat{\mathrm{GE}}_{\mathrm{OOB}} = \frac{1}{n} \sum_{i=1}^{n} L(y^{(i)}, \hat{f}^{(i)}_{\mathrm{OOB}})$$

USING THE OUT-OF-BAG ERROR ESTIMATE

- Gives us a (proper) estimator of GE, computable during training
- Can even compute this for all smaller ensemble sizes (after we fitted M models)



OOB ERROR: COMPARABILITY, BEST PRACTICE

OOB Size: The probability that an observation is out-of-bag (OOB) is:

$$\mathbb{P}\left(i \in \mathrm{OOB}^{[m]}\right) = \left(1 - \frac{1}{n}\right)^n \overset{n\to\infty}{\longrightarrow} \frac{1}{e} \approx 0.37$$

$\Rightarrow$ similar to holdout or 3-fold CV (1/3 validation, 2/3 training)

Comparability Issues:

- **OOB error** rather unique to RFs / bagging
- To compare models, we often still use CV, etc., to be consistent

Use the OOB Error for:

- Get first impression of RF performance
- Select ensemble size
- Efficiently evaluate different RF hyperparameter configurations