

The solutions for the implementation exercises for this sheet can be found in the files *hw_sol_3-1_PDP_ICE_Interactions.R* and *hw_sol_3-4_ME.R* on Moodle.

Solution 1:

a) Derivation of PD function for $S = \{1\}$ (with $C = \{2\}$) given

$$\hat{f}(\mathbf{x}) = \hat{f}(x_1, x_2) = \hat{\beta}_1 x_1 + \hat{\beta}_2 x_2 + \hat{\beta}_3 x_1 x_2 + \hat{\beta}_0$$

$$\begin{aligned} f_{1,PD}(x_1) &= \mathbb{E}_{x_2} \left(\hat{f}(x_1, x_2) \right) = \int_{-\infty}^{\infty} \left(\hat{\beta}_1 x_1 + \hat{\beta}_2 x_2 + \hat{\beta}_3 x_1 x_2 + \hat{\beta}_0 \right) d\mathbb{P}(x_2) \\ &= \hat{\beta}_1 x_1 + \int_{-\infty}^{\infty} \hat{\beta}_2 x_2 + \hat{\beta}_3 x_1 x_2 d\mathbb{P}(x_2) + \hat{\beta}_0 \\ &= \hat{\beta}_1 x_1 + \int_{-\infty}^{\infty} (\hat{\beta}_2 + \hat{\beta}_3 x_1) x_2 d\mathbb{P}(x_2) + \hat{\beta}_0 \\ &= \hat{\beta}_1 x_1 + (\hat{\beta}_2 + \hat{\beta}_3 x_1) \cdot \int_{-\infty}^{\infty} x_2 d\mathbb{P}(x_2) + \hat{\beta}_0 \\ &= \hat{\beta}_1 x_1 + (\hat{\beta}_2 + \hat{\beta}_3 x_1) \cdot \mathbb{E}_{x_2}(x_2) + \hat{\beta}_0 \end{aligned}$$

b) PD function for $\hat{\beta}_0 = 0$, $\hat{\beta}_1 = -8$, $\hat{\beta}_2 = 0.2$, $\hat{\beta}_3 = 16$, $X_1 \sim \text{Unif}(-1, 1)$ and $X_2 \sim B(1, 0.5)$.


$$\begin{aligned} f_{1,PD}(x_1) &= \hat{\beta}_1 x_1 + (\hat{\beta}_2 + \hat{\beta}_3 x_1) \cdot \mathbb{E}_{x_2}(x_2) + \hat{\beta}_0 = -8x_1 + (0.2 + 16x_1) \cdot \mathbb{E}_{x_2}(x_2) + 0 \\ &= -8x_1 + (0.2 + 16x_1) \cdot 0.5 \\ &= -8x_1 + 0.1 + 8x_1 \\ &= 0.1 \end{aligned}$$

c) ICE functions for group $X_2 = 1$ and for group $X_2 = 0$:

$$f_1(x_1) = \begin{cases} -8x_1 + (0.2 + 16x_1) \cdot 1 = 8x_1 + 0.2 & x_2 = 1 \\ -8x_1 + (0.2 + 16x_1) \cdot 0 = -8x_1 & x_2 = 0 \end{cases}$$

The light green dots correspond to group $X_2 = 1$, the light blue dots to group $X_2 = 0$.

d) The example illustrates that by the averaging of ICE curves for a PD plot we might obfuscate heterogeneous effects and interactions. Although the ICE curves here show a strong effect of X_1 on Y , the effect is not apparent in the PDP. Therefore, it is highly recommended to plot PD plots and ICE curves together.



e) , f) For the last two parts of the exercise, please find the solution in the file *hw_sol_3_1_PDP_ICE_Interactions.R*.

Solution 2:

The fitted model is

$$\hat{f}(x_1, x_2) = \theta_0 + \theta_1 x_1^2 + \theta_2 x_2^2 + \theta_{1,2} x_1 x_2, \quad (\theta_0, \theta_1, \theta_2, \theta_{1,2}) = (0, 1, 0.5, 2).$$

a) Derivative marginal effect (dME)

$$\text{dME}_1(x_1, x_2) = \frac{\partial \hat{f}}{\partial x_1} = 2\theta_1 x_1 + \theta_{1,2} x_2 = 2x_1 + 2x_2.$$

Discussion. Because the model is quadratic in x_1 , the derivative is a *linear* function of both x_1 and x_2 . If the surface were exactly linear, this rate would coincide with any finite difference.

b) Forward marginal effect (fME) with step $h_1 > 0$

$$\text{fME}_1(x_1, x_2; h_1) = \hat{f}(x_1 + h_1, x_2) - \hat{f}(x_1, x_2).$$

Expand $\hat{f}(x_1 + h_1, x_2)$:

$$\begin{aligned} \hat{f}(x_1 + h_1, x_2) &= \theta_1 (x_1 + h_1)^2 + \theta_2 x_2^2 + \theta_{1,2} (x_1 + h_1) x_2 \\ &= \theta_1 [x_1^2 + 2x_1 h_1 + h_1^2] + \theta_2 x_2^2 + \theta_{1,2} x_1 x_2 + \theta_{1,2} h_1 x_2. \end{aligned}$$

Subtracting $\hat{f}(x_1, x_2)$ gives

$$\boxed{\text{fME}_1(x_1, x_2; h_1) = 2\theta_1 x_1 h_1 + \theta_1 h_1^2 + \theta_{1,2} x_2 h_1} = 2x_1 h_1 + h_1^2 + 2x_2 h_1.$$

Discussion. The extra term $\theta_1 h_1^2$ captures curvature; it vanishes faster than linearly when $h_1 \rightarrow 0$, at which point fME converges to dME.

c) Numerical evaluation at $(x_1, x_2, h_1) = (1, 2, 1)$

$$\text{dME}_1(1, 2) = 2(1) + 2(2) = 6,$$

$$\text{fME}_1(1, 2; 1) = 2 \cdot 1 \cdot 1 + 1^2 + 2 \cdot 2 \cdot 1 = 2 + 1 + 4 = 7.$$

Why different? The derivative 6 is the *instantaneous* slope. The finite step of 1 exposes curvature; the quadratic term makes the actual change go up to 7. Hence, using dME for a sizeable perturbation would misestimate (over- or under-predict) the effect.

d) Non-Linearity Measure (NLM) (R^2 of the linear secant along the path)

Hints.

- Compute $\text{SSR} = \sum (\hat{f} - g)^2$ and $\text{SST} = \sum (\hat{f} - \bar{f})^2$ from the table.
- Then evaluate $\text{NLM} = 1 - \text{SSR}/\text{SST}$.
- Comment: Is the resulting NLM close enough to 1 to accept the secant as a faithful local explanation?

Step 1: Path points. For $T = 10$ equidistant $t_i \in [0, 1]$:

$$t_i = \frac{i}{9+1}, \quad x_1^{(i)} = 1 + t_i h_1 = 1 + 0.5 t_i, \quad x_2^{(i)} = 2.$$

Step 2: Model and secant values.

$$f_i = \hat{f}(x_1^{(i)}, 2), \quad g_i = \hat{f}(1, 2) + t_i \cdot \text{fME}_1(1, 2; 0.5), \quad \hat{f}(1, 2) = 7, \text{ fME} = 3.25.$$

Step 3: Compute R^2 .

$$\text{NLM} = 1 - \frac{\sum_{i=1}^9 (f_i - g_i)^2}{\sum_{i=1}^9 (f_i - \bar{f})^2}, \quad \bar{f} = \frac{1}{9} \sum f_i.$$

Numerically

$$\boxed{\text{NLM} \approx 0.9967}.$$

Interpretation. Along the half-unit move in x_1 , the quadratic surface is *almost* linear (NLM very close to 1). If we doubled h_1 , the numerator would grow faster than the denominator, lowering NLM and signaling stronger curvature.

e) See the implementation in *hw_sol3_4_ME.R*.