



# Interpretable Machine Learning

## Intro to IML

## Correlation and Dependencies




### Learning goals

- Pearson correlation
- Coefficient of determination  $R^2$
- Mutual information
- Correlation vs. dependence

# PEARSON'S CORRELATION COEFFICIENT $\rho$


**Correlation** often refers to Pearson's correlation (measures only **linear** relationship)

$$\rho(X_1, X_2) = \frac{\sum_{i=1}^n (x_1^{(i)} - \bar{x}_1) \cdot (x_2^{(i)} - \bar{x}_2)}{\sqrt{\sum_{i=1}^n (x_1^{(i)} - \bar{x}_1)^2 \sum_{i=1}^n (x_2^{(i)} - \bar{x}_2)^2}} \in [-1, 1]$$




Geometric interpretation of  $\rho$ :

- Numerator is sum of rectangle's area with width  $x_1^{(i)} - \bar{x}_1$  and height  $x_2^{(i)} - \bar{x}_2$
- Areas enter numerator with positive (+) or negative (-) sign, depending on position
- Denominator scales the sum into the range  $[-1, 1]$




# PEARSON'S CORRELATION COEFFICIENT $\rho$

**Correlation** often refers to Pearson's correlation (measures only **linear** relationship)



$$\rho(X_1, X_2) = \frac{\sum_{i=1}^n (x_1^{(i)} - \bar{x}_1) \cdot (x_2^{(i)} - \bar{x}_2)}{\sqrt{\sum_{i=1}^n (x_1^{(i)} - \bar{x}_1)^2 \sum_{i=1}^n (x_2^{(i)} - \bar{x}_2)^2}} \in [-1, 1]$$




Geometric interpretation of  $\rho$ :

- Numerator is sum of rectangle's area with width  $x_1^{(i)} - \bar{x}_1$  and height  $x_2^{(i)} - \bar{x}_2$
- Areas enter numerator with positive (+) or negative (-) sign, depending on position
- Denominator scales the sum into the range  $[-1, 1]$

- $\rho > 0$  if **positive areas** dominate **negative areas**  
 $\rightsquigarrow X_1, X_2$  positive correlated
- $\rho < 0$  if **negative areas** dominate **positive areas**  
 $\rightsquigarrow X_1, X_2$  negative correlated
- $\rho = 0$  if area of rectangles cancels out  $\rightsquigarrow X_1, X_2$  linearly uncorrelated

# COEFFICIENT OF DETERMINATION $R^2$

Another method to evaluate **linear dependency** between features is  $R^2$




- Fit a linear model:

$$\hat{x}_2 = \hat{f}_{LM}(x_1) = \theta_0 + \theta_1 x_1$$


↪ Slope  $\theta_1 = 0 \Rightarrow$  no dependence

↪ Large slope  $\Rightarrow$  strong dependence



# COEFFICIENT OF DETERMINATION $R^2$

Another method to evaluate **linear dependency** between features is  $R^2$



- Fit a linear model:

$$\hat{x}_2 = \hat{f}_{LM}(x_1) = \theta_0 + \theta_1 x_1$$


↪ Slope  $\theta_1 = 0 \Rightarrow$  no dependence

↪ Large slope  $\Rightarrow$  strong dependence

- Exact  $\theta_1$  score problematic



↪ Re-scaling of  $x_1$  or  $x_2$  changes  $\theta_1$

↪ °F  $\rightarrow$  °C  $\Rightarrow \theta_1 = 78 \rightarrow \theta_1^* = 141$



# COEFFICIENT OF DETERMINATION $R^2$

Another method to evaluate **linear dependency** between features is  $R^2$



- Fit a linear model:

$$\hat{x}_2 = \hat{f}_{LM}(x_1) = \theta_0 + \theta_1 x_1$$

↪ Slope  $\theta_1 = 0 \Rightarrow$  no dependence

↪ Large slope  $\Rightarrow$  strong dependence

- Exact  $\theta_1$  score problematic

↪ Re-scaling of  $x_1$  or  $x_2$  changes  $\theta_1$

- Set  $SSE_{LM}$  in relation to  $SSE$  of a constant model  $\hat{f}_c = \bar{x}_2$

$$SSE_{LM} = \sum_{i=1}^n (x_2^{(i)} - \hat{f}_{LM}(x_1^{(i)}))^2$$

$$SSE_c = \sum_{i=1}^n (x_2^{(i)} - \bar{x}_2)^2$$

$\Rightarrow$  Measure of fitting quality of LM:  $R^2 = 1 - \frac{SSE_{LM}}{SSE_c} \in [0, 1]$

$\Rightarrow \rho(X_1, X_2) = R$


# JOINT, MARGINAL AND CONDITIONAL DISTRIBUTION

For two discrete random variables  $X_1, X_2$ :

## Joint distribution

$$p_{X_1, X_2}(x_1, x_2) = \mathbb{P}(X_1 = x_1, X_2 = x_2)$$

$p_{X_1, X_2}$	$\mathbb{P}(X_2 = 0)$	$\mathbb{P}(X_2 = 1)$	$p_{X_1}$
$\mathbb{P}(X_1 = 0)$	0.2	0.3	0.5
$\mathbb{P}(X_1 = 1)$	0.1	0.4	0.5
$p_{X_2}$	0.3	0.7	1



# JOINT, MARGINAL AND CONDITIONAL DISTRIBUTION

For two discrete random variables  $X_1, X_2$ :

## Joint distribution

$$p_{X_1, X_2}(x_1, x_2) = \mathbb{P}(X_1 = x_1, X_2 = x_2)$$


$p_{X_1, X_2}$	$\mathbb{P}(X_2 = 0)$	$\mathbb{P}(X_2 = 1)$	$p_{X_1}$
$\mathbb{P}(X_1 = 0)$	0.2	0.3	0.5
$\mathbb{P}(X_1 = 1)$	0.1	0.4	0.5
$p_{X_2}$	0.3	0.7	1

## Marginal distribution

$$p_{X_1}(x_1) = \mathbb{P}(X_1 = x_1) = \sum_{x_2 \in \mathcal{X}_2} p(x_1, x_2)$$

$p_{X_1, X_2}$	$\mathbb{P}(X_2 = 0)$	$\mathbb{P}(X_2 = 1)$	$p_{X_1}$
$\mathbb{P}(X_1 = 0)$	0.2	0.3	0.5
$\mathbb{P}(X_1 = 1)$	0.1	0.4	0.5
$p_{X_2}$	0.3	0.7	1

↪ In continuous case with integrals





# JOINT, MARGINAL AND CONDITIONAL DISTRIBUTION

For two discrete random variables  $X_1, X_2$ :

## Joint distribution

$$p_{X_1, X_2}(x_1, x_2) = \mathbb{P}(X_1 = x_1, X_2 = x_2)$$

$p_{X_1, X_2}$	$\mathbb{P}(X_2 = 0)$	$\mathbb{P}(X_2 = 1)$	$p_{X_1}$
$\mathbb{P}(X_1 = 0)$	0.2	0.3	0.5
$\mathbb{P}(X_1 = 1)$	0.1	0.4	0.5
$p_{X_2}$	0.3	0.7	1

## Marginal distribution

$$p_{X_1}(x_1) = \mathbb{P}(X_1 = x_1) = \sum_{x_2 \in \mathcal{X}_2} p(x_1, x_2)$$


$p_{X_1, X_2}$	$\mathbb{P}(X_2 = 0)$	$\mathbb{P}(X_2 = 1)$	$p_{X_1}$
$\mathbb{P}(X_1 = 0)$	0.2	0.3	0.5
$\mathbb{P}(X_1 = 1)$	0.1	0.4	0.5
$p_{X_2}$	0.3	0.7	1

↪ In continuous case with integrals

## Conditional distribution

$$\begin{aligned} p_{X_1|X_2}(x_1|x_2) &= \mathbb{P}(X_1 = x_1 | X_2 = x_2) \\ &= \frac{p_{X_1, X_2}(x_1, x_2)}{p_{X_2}(x_2)} \end{aligned}$$

	$x_2 = 0$	$x_2 = 1$
$\mathbb{P}(X_1 = 0   X_2 = x_2)$	0.67	0.43
$\mathbb{P}(X_1 = 1   X_2 = x_2)$	0.33	0.57
$\sum$	1	1




# DEPENDENCE

**Dependence:** Describes general dependence structure (e.g., non-lin. relationships)

- Definition:  $X_j, X_k$  independent  $\Leftrightarrow$  joint distribution is product of marginals:

$$\mathbb{P}(X_j, X_k) = \mathbb{P}(X_j) \cdot \mathbb{P}(X_k)$$



# DEPENDENCE


**Dependence:** Describes general dependence structure (e.g., non-lin. relationships)

- Definition:  $X_j, X_k$  independent  $\Leftrightarrow$  joint distribution is product of marginals:

$$\mathbb{P}(X_j, X_k) = \mathbb{P}(X_j) \cdot \mathbb{P}(X_k)$$

- Equivalent definition (knowing  $X_k$  gives no info about  $X_j$  and vice versa):

$$\mathbb{P}(X_j|X_k) = \mathbb{P}(X_j) \text{ and } \mathbb{P}(X_k|X_j) = \mathbb{P}(X_k) \text{ (follows from cond. probability)}$$



# DEPENDENCE

**Dependence:** Describes general dependence structure (e.g., non-lin. relationships)

- Definition:  $X_j, X_k$  independent  $\Leftrightarrow$  joint distribution is product of marginals:

$$\mathbb{P}(X_j, X_k) = \mathbb{P}(X_j) \cdot \mathbb{P}(X_k)$$


- Equivalent definition (knowing  $X_k$  gives no info about  $X_j$  and vice versa):

$$\mathbb{P}(X_j|X_k) = \mathbb{P}(X_j) \text{ and } \mathbb{P}(X_k|X_j) = \mathbb{P}(X_k) \text{ (follows from cond. probability)}$$

- Measuring complex dependencies is difficult but different measures exist

Examples

- $\rightsquigarrow$  Spearman correlation (measures monotonic dependencies via ranks)
- $\rightsquigarrow$  Information-theoretical measures like mutual information
- $\rightsquigarrow$  Kernel-based measures like Hilbert-Schmidt Independence Criterion (HSIC)



# DEPENDENCE

**Dependence:** Describes general dependence structure (e.g., non-lin. relationships)

- Definition:  $X_j, X_k$  independent  $\Leftrightarrow$  joint distribution is product of marginals:

$$\mathbb{P}(X_j, X_k) = \mathbb{P}(X_j) \cdot \mathbb{P}(X_k)$$

- Equivalent definition (knowing  $X_k$  gives no info about  $X_j$  and vice versa):

$$\mathbb{P}(X_j|X_k) = \mathbb{P}(X_j) \text{ and } \mathbb{P}(X_k|X_j) = \mathbb{P}(X_k) \text{ (follows from cond. probability)}$$


- Measuring complex dependencies is difficult but different measures exist  
Examples

~> Spearman correlation (measures monotonic dependencies via ranks)

~> Information-theoretical measures like mutual information

~> Kernel-based measures like Hilbert-Schmidt Independence Criterion (HSIC)

- **N.B.:**  $X_j, X_k$  indep.  $\Rightarrow \rho(X_j, X_k) = 0$  **but**  $\rho(X_j, X_k) = 0 \nRightarrow X_j, X_k$  indep.  
Equivalency holds if distribution is jointly normal




# MUTUAL INFORMATION

- MI describes expected amount of information shared by two RVs:

$$MI(X_1, X_2) = \mathbb{E}_{p(x_1, x_2)} \left[ \log \left( \frac{p(x_1, x_2)}{p(x_1)p(x_2)} \right) \right]$$


- MI measures amount of "dependence" between features by looking how different the joint distribution is from pure indep.  $p(x_1, x_2) = p(x_1)p(x_2)$   
 $\rightsquigarrow MI(X_1, X_2) = \mathbb{E}_{p(x_1, x_2)} \left[ \log \left( \frac{p(x_1, x_2)}{p(x_1)p(x_2)} \right) \right] = \mathbb{E}_{p(x_1, x_2)} [\log(1)] = 0$   
 $\rightsquigarrow MI(X_j, X_k) = 0$  if and only if the features are independent
- Unlike (Pearson) correlation, MI is also defined for categorical features



# MUTUAL INFORMATION: EXAMPLE

For two discrete RV  $X_1$  and  $Y$ :

$$MI(X_1; Y) = \mathbb{E}_{p(x_1, y)} \left[ \log \left( \frac{p(x_1, y)}{p(x_1)p(y)} \right) \right] = \sum_{x_1 \in \mathcal{X}_1} \sum_{y \in \mathcal{Y}} p(x_1, y) \log \left( \frac{p(x_1, y)}{p(x_1)p(y)} \right)$$




$X_1$	...	$Y$
yes	...	yes
yes	...	no
no	...	yes
no	...	no

	$\mathbb{P}(X_1 = \text{yes})$	$\mathbb{P}(X_1 = \text{no})$	$p_Y$
$\mathbb{P}(Y = \text{yes})$	0.25	0.25	0.5
$\mathbb{P}(Y = \text{no})$	0.25	0.25	0.5
$p_{X_1}$	0.5	0.5	1

# MUTUAL INFORMATION: EXAMPLE

For two discrete RV  $X_1$  and  $Y$ :

$$MI(X_1; Y) = \mathbb{E}_{p(x_1, y)} \left[ \log \left( \frac{p(x_1, y)}{p(x_1)p(y)} \right) \right] = \sum_{x_1 \in \mathcal{X}_1} \sum_{y \in \mathcal{Y}} p(x_1, y) \log \left( \frac{p(x_1, y)}{p(x_1)p(y)} \right)$$



$X_1$	...	$Y$
yes	...	yes
yes	...	no
no	...	yes
no	...	no


	$\mathbb{P}(X_1 = \text{yes})$	$\mathbb{P}(X_1 = \text{no})$	$p_Y$
$\mathbb{P}(Y = \text{yes})$	0.25	0.25	0.5
$\mathbb{P}(Y = \text{no})$	0.25	0.25	0.5
$p_{X_1}$	0.5	0.5	1

$$\begin{aligned} MI(X_1; Y) &= 0.25 \log \left( \frac{0.25}{0.5 \cdot 0.5} \right) + 0.25 \log \left( \frac{0.25}{0.5 \cdot 0.5} \right) \\ &\quad + 0.25 \log \left( \frac{0.25}{0.5 \cdot 0.5} \right) + 0.25 \log \left( \frac{0.25}{0.5 \cdot 0.5} \right) \\ &= 0.25 \log \left( \frac{0.25}{0.25} \right) \cdot 4 \\ &= 0.25 \log(1) \cdot 4 = 0 \end{aligned}$$



# DEPENDENCE AND INDEPENDENCE


## Example:



Conditional distributions at different vertical and horizontal slices (after normalizing area to 1) match their marginal distributions



$$\Rightarrow P(X_1|X_2) = P(X_1)$$

$$P(X_2|X_1) = P(X_2)$$



# DEPENDENCE AND INDEPENDENCE

## Example:




Conditional distributions at different vertical and horizontal slices (after normalizing area to 1) match their marginal distributions

Conditional distributions do not match their marginal distributions

$$\Rightarrow P(X_1|X_2) = P(X_1)$$

$$P(X_2|X_1) = P(X_2)$$




# CORRELATION VS. DEPENDENCE


Illustration of bivariate normal distribution with different correlations

$X_1, X_2 \sim N(0, 1)$



$\rho(X_1, X_2) = 0$   
(independent)



$\rho(X_1, X_2) = 0.8$





$\rho(X_1, X_2) = -0.8$



# CORRELATION VS. DEPENDENCE

Illustration of bivariate normal distribution with different correlations

$X_1, X_2 \sim N(0, 1)$



Examples with Pearson's corr.  $\rho \approx 0$  but non-linear dependencies ( $MI \neq 0$ ):

$\rho(X_1, X_2) = 0$ ,  $MI(X_1, X_2) = 0.52$        $\rho(X_1, X_2) = 0.01$ ,  $MI(X_1, X_2) = 0.37$        $\rho(X_1, X_2) = -0.06$ ,  $MI(X_1, X_2) = 0.61$

