

Interpretable Machine Learning




Regional Effects
REPID



Learning goals


- Difference between feature effects and feature interactions
- REPID

WHY REGIONAL EXPLANATIONS?




Problem: PD & ICE plots can be confounded by feature interactions.
Solution: Group homogeneous ICE curves in such a way that reduces the presence of individual interaction effects within a group
$\rightsquigarrow$ Regional effect plots (REPs).


Global Effect



Regional Effect (1)




Regional Effect (2)



- Splitting by the "workingday" revealed 2 different patterns that we're clashed together in the initial plot
- $\cup_i(\texttt{regional_explanation}_i) = \texttt{global_explanation}$
- *Fidelity*$(\texttt{regional_explanation}_i) >$ *Fidelity*$(\texttt{global_explanation})$


ICE CURVE: LOCAL FEATURE EFFECTS



Question: How do feature changes affect the prediction for **one obs.**?
Idea: Split $\boldsymbol{x} = (x_j, \boldsymbol{x}_{-j})$ into x_j (feat of interest) and $\boldsymbol{x}_{-j}$ (remaining feats)

- Replace observed values x_j with grid values $\tilde{x}_j$ while keeping $\boldsymbol{x}_{-j}$ fixed
- Visualize function $\hat{f}(\tilde{x}_j, \boldsymbol{x}_{-j})$ for varying $\tilde{x}_j$ (ICE)


ICE CURVE: LOCAL FEATURE EFFECTS




Question: How do feature changes affect the prediction for **one obs.**?
Idea: Split $\boldsymbol{x} = (x_j, \boldsymbol{x}_{-j})$ into x_j (feat of interest) and $\boldsymbol{x}_{-j}$ (remaining feats)


- Replace observed values x_j with grid values $\tilde{x}_j$ while keeping $\boldsymbol{x}_{-j}$ fixed
- Visualize function $\hat{f}(\tilde{x}_j, \boldsymbol{x}_{-j})$ for varying $\tilde{x}_j$ (ICE)

Example: SVM prediction surface (left), select obs. and visualize changes in prediction for varying x_2 while keeping x_1 fixed $\Rightarrow$ **local interpretation**





PD PLOT - GLOBAL FEATURE EFFECTS




Question: How do changes of feat values affect model prediction **on avg.**?


- **PD function**: Integrate out effect of X_{-j} to obtain marginal effect of x_j

$$f_j^{PD}(\tilde{x}_j) = \mathbb{E}_{X_{-j}}[\hat{f}(\tilde{x}_j, X_{-j})] = \int \hat{f}(\tilde{x}_j, X_{-j}) d\mathbb{P}(X_{-j})$$


- **Estimate (MC integration):** Average ICE curves at grid points $\tilde{x}_j$

$$\hat{f}_j^{PD}(\tilde{x}_j) = \frac{1}{n} \sum_{i=1}^{n} \hat{f}(\tilde{x}_j, \boldsymbol{x}_{-j}^{(i)})$$





FEATURE INTERACTIONS



Hooker (2004, 2007): Functional ANOVA decomp. of a function

$$\hat{f}(\mathbf{x}) = g_0 + \underbrace{\sum_{j=1}^{p} g_j(x_j)}_{\text{main effect}} + \underbrace{\sum_{j \neq k} g_{j,k}(x_j, x_k)}_{\text{two-way interaction effect}} + \cdots + \underbrace{g_{1,2,\ldots,p}(\mathbf{x})}_{\text{p-way interaction effect}}$$


Friedman and Popescu (2008):

$\Rightarrow$ If x_j and $\boldsymbol{x}_{-j}$ don't interact, we can decomp. $f(\boldsymbol{x}) = g_j(x_j) + g_{-j}(\boldsymbol{x}_{-j})$


$\Rightarrow$ If x_j and x_k don't interact, decomposition: $f(\boldsymbol{x}) = g_{-j}(\boldsymbol{x}_{-j}) + g_{-k}(\boldsymbol{x}_{-k})$

Example: Not additively separable:

$f(\boldsymbol{x}) = x_1 + x_2 + x_1 \cdot x_2 \neq g(x_1) + g(x_2)$



FEATURE INTERACTIONS



Hooker (2004, 2007): Functional ANOVA decomp. of a function

$$\hat{f}(\mathbf{x}) = g_0 + \underbrace{\sum_{j=1}^{p} g_j(x_j)}_{\text{main effect}} + \underbrace{\sum_{j \neq k} g_{j,k}(x_j, x_k)}_{\text{two-way interaction effect}} + \cdots + \underbrace{g_{1,2,\ldots,p}(\mathbf{x})}_{\text{p-way interaction effect}}$$


Friedman and Popescu (2008):

$\Rightarrow$ If x_j and $\boldsymbol{x}_{-j}$ don't interact, we can decomp. $f(\boldsymbol{x}) = g_j(x_j) + g_{-j}(\boldsymbol{x}_{-j})$

$\Rightarrow$ If x_j and x_k don't interact, decomposition: $f(\boldsymbol{x}) = g_{-j}(\boldsymbol{x}_{-j}) + g_{-k}(\boldsymbol{x}_{-k})$


Example: Separable:

$f(\boldsymbol{x}) = x_1 + x_2 + \log(x_1 \cdot x_2) = (x_1 + \log(x_1)) + (x_2 + \log(x_2)) = g_1(x_1) + g_2(x_2)$




REPID: REGIONAL EFFECT PLOTS

▸ "Herbinger et al." 2022




Recall: Different shapes of ICE curves ⇒ interactions (ignore vertical shifts)
⇒ Focus on shape differences of **mean-centered ICE curves**.
Mean-centered ICE curve for obs. $\boldsymbol{x}$ evaluated at m grid points $\tilde{x}_j^{(1)}, \ldots, \tilde{x}_j^{(m)}$ is:

$$\hat{f}^c(\tilde{x}_j, \boldsymbol{x}_{-j}) = \hat{f}(\tilde{x}_j, \boldsymbol{x}_{-j}) - \frac{1}{m} \sum_{k=1}^{m} \hat{f}(\tilde{x}_j^{(k)}, \boldsymbol{x}_{-j})$$



REGIONAL EFFECTS - SYNTHETIC EXAMPLE




Example: $X_1, X_2, X_6 \sim \mathcal{U}(-1, 1)$, $X_3, X_4, X_5 \sim \mathcal{B}(n, 0.5)$ (all iid)

$\rightsquigarrow$ Ground truth: $f(X) = 0.2X_1 \; -8X_2 + 8X_2 \mathbb{1}_{(X_1>0)} + 16X_2 \mathbb{1}_{(X_3=0)} \; + \epsilon$


$\rightsquigarrow$ Model: Random forest

Problem:

- PD curve of X_2 is misleading due to interactions $\rightsquigarrow$ ICE
- ICE curves do not identify the interacting features



REGIONAL EFFECTS - SYNTHETIC EXAMPLE



Example: $X_1, X_2, X_6 \sim \mathcal{U}(-1, 1)$, $X_3, X_4, X_5 \sim \mathcal{B}(n, 0.5)$ (all iid)


$\rightsquigarrow$ Ground truth: $f(X) = 0.2X_1 - 8X_2 + 8X_2 \mathbb{1}_{(X_1>0)} + 16X_2 \mathbb{1}_{(X_3=0)} + \epsilon$

$\rightsquigarrow$ Model: Random forest


Problem:

- PD curve of X_2 is misleading due to interactions $\rightsquigarrow$ ICE
- ICE curves do not identify the interacting features

Idea: Find regions with similar ICE curves and aggregate them to regional effects



REGIONAL EFFECTS - SYNTHETIC EXAMPLE



Example: $X_1, X_2, X_6 \sim \mathcal{U}(-1, 1)$, $X_3, X_4, X_5 \sim \mathcal{B}(n, 0.5)$ (all iid)

$\rightsquigarrow$ Ground truth: $f(X) = 0.2X_1 - 8X_2 + 8X_2 \mathbb{1}_{(X_1>0)} + 16X_2 \mathbb{1}_{(X_3=0)} + \epsilon$


$\rightsquigarrow$ Model: Random forest

Problem:

- PD curve of X_2 is misleading due to interactions $\rightsquigarrow$ ICE
- ICE curves do not identify the interacting features

Idea: Find regions with similar ICE curves and aggregate them to regional effects

Regional effect (blue curves) $\hat{=}$ Estimate PD curve in each region




$\hat{f}_2^{PD}(X_2) \approx 8X_2 \quad \approx 16X_2 \quad \approx -8X_2 \quad \approx 0$


$\Rightarrow$ Additive decomposition of global feat effect

REGIONAL EFFECTS - DETAILS

Question: How to split curves into regions? Define risk as L2 loss of mean-centered ICE curves:

$$\mathcal{R}_j(\mathcal{N}) = \sum_{\boldsymbol{x} \in \mathcal{N}} \sum_{k=1}^{m} \underbrace{\left(\hat{f}^c(\tilde{x}_j^{(k)}, \boldsymbol{x}_{-j}) - \hat{f}_{j|\mathcal{N}}^{PD,c}(\tilde{x}_j^{(k)})\right)^2}_{d_k}$$





with the avg. feature effect in region $\mathcal{N} \subseteq \mathcal{X}$:

$$\hat{f}_{j|\mathcal{N}}^{PD,c}(\tilde{x}_j) = \frac{1}{|\mathcal{N}|} \sum_{\boldsymbol{x} \in \mathcal{N}} \hat{f}^c(\tilde{x}_j, \boldsymbol{x}_{-j})$$

$\rightsquigarrow$ Measures interaction-related heterogeneity (variance) of ICE curves in $\mathcal{N}$


$\rightsquigarrow$ Recursive partitioning (CART): Find best feat-split combo that solves


$$\arg\min_{z,t} \; \mathcal{R}_j(\mathcal{N}_{left}) + \mathcal{R}_j(\mathcal{N}_{right})$$

- $\mathcal{N}_{left} = \{\boldsymbol{x} \in \mathcal{N} | x_z \leq t\}$
- $\mathcal{N}_{right} = \{\boldsymbol{x} \in \mathcal{N} | x_z > t\}$
- Split point t for feature x_z, $z \in -j$

Intuition: Is another feature x_z responsible for the heterogeneity (measured by $\mathcal{R}_j$)?


REGIONAL EFFECT PLOTS - REAL EXAMPLE






- Identify feature with highly heterogeneous local effects
  $\rightsquigarrow$ `hour`: Most important; highly heterogeneous feat (highest variance)
- Find regions in feature space where this heterogeneity is minimal
  $\rightsquigarrow$ Partition feature space using CART to minimize variance of mean-centered ICE curves within each region


REGIONAL EFFECT PLOTS - REAL EXAMPLE




Regional effects of `hour`




workingday=False




workingday=True




temp=cold



temp=hot



temp=cold



temp=hot

