

Interpretable Machine Learning




Intro to IML
Correlation and Dependencies



Learning goals


- Pearson correlation
- Coefficient of determination R^2
- Mutual information
- Correlation vs. dependence

PEARSON'S CORRELATION COEFFICIENT ρ



Correlation often refers to Pearson's correlation (captures only **linear relationship**)


$$\rho(X_1, X_2) = \frac{\sum_{i=1}^{n}(x_1^{(i)} - \bar{x}_1)(x_2^{(i)} - \bar{x}_2)}{\sqrt{\sum_{i=1}^{n}(x_1^{(i)} - \bar{x}_1)^2 \sum_{i=1}^{n}(x_2^{(i)} - \bar{x}_2)^2}} \in [-1, 1]$$



Geometric interpretation of ρ:


- Numerator is sum of rectangle's area with width $x_1^{(i)} - \bar{x}_1$ and height $x_2^{(i)} - \bar{x}_2$
- Areas enter numerator with positive (**+**) or negative (**-**) sign, depending on position
- Denominator scales the sum into the range $[-1, 1]$

PEARSON'S CORRELATION COEFFICIENT ρ



Correlation often refers to Pearson's correlation (captures only **linear relationship**)

$$\rho(X_1, X_2) = \frac{\sum_{i=1}^{n}(x_1^{(i)} - \bar{x}_1)(x_2^{(i)} - \bar{x}_2)}{\sqrt{\sum_{i=1}^{n}(x_1^{(i)} - \bar{x}_1)^2 \sum_{i=1}^{n}(x_2^{(i)} - \bar{x}_2)^2}} \in [-1, 1]$$




Geometric interpretation of ρ:

- Numerator is sum of rectangle's area with width $x_1^{(i)} - \bar{x}_1$ and height $x_2^{(i)} - \bar{x}_2$
- Areas enter numerator with positive (**+**) or negative (**-**) sign, depending on position
- Denominator scales the sum into the range $[-1, 1]$

- $\rho > 0$ if positive areas dominate negative areas $\rightsquigarrow$ X_1, X_2 positive correlated
- $\rho < 0$ if negative areas dominate positive areas $\rightsquigarrow$ X_1, X_2 negative correlated
- $\rho = 0$ if area of rectangles cancels out$\rightsquigarrow$ X_1, X_2 linearly uncorrelated

COEFFICIENT OF DETERMINATION R^2


Another method to evaluate **linear dependency** between features is R^2



- Fit a linear model:
  $\hat{x}_2 = \hat{f}_{LM}(x_1) = \theta_0 + \theta_1 x_1$


$\rightsquigarrow$ Slope $\theta_1 = 0 \Rightarrow$ no dependence

$\rightsquigarrow$ Large slope $\Rightarrow$ strong dependence



COEFFICIENT OF DETERMINATION R^2

Another method to evaluate **linear dependency** between features is R^2



- Fit a linear model:
  $\hat{x}_2 = \hat{f}_{LM}(x_1) = \theta_0 + \theta_1 x_1$


$\rightsquigarrow$ Slope $\theta_1 = 0 \Rightarrow$ no dependence

$\rightsquigarrow$ Large slope $\Rightarrow$ strong dependence

- Exact θ_1 score problematic


$\rightsquigarrow$ Re-scaling of x_1 or x_2 changes θ_1

$\rightsquigarrow$ °F $\rightarrow$ °C $\Rightarrow \theta_1 = 78 \rightarrow \theta_1^* = 141$



COEFFICIENT OF DETERMINATION R^2

Another method to evaluate **linear dependency** between features is R^2



- Fit a linear model:
  $\hat{x}_2 = \hat{f}_{LM}(x_1) = \theta_0 + \theta_1 x_1$

$\rightsquigarrow$ Slope $\theta_1 = 0 \Rightarrow$ no dependence

$\rightsquigarrow$ Large slope $\Rightarrow$ strong dependence

- Exact θ_1 score problematic

$\rightsquigarrow$ Re-scaling of x_1 or x_2 changes θ_1

- Set SSE_{LM} in relation to SSE of a constant model $\hat{f}_c = \bar{x}_2$


$$SSE_{LM} = \sum_{i=1}^{n} (x_2^{(i)} - \hat{f}_{LM}(x_1^{(i)}))^2$$

$$SSE_c = \sum_{i=1}^{n} (x_2^{(i)} - \bar{x}_2)^2$$

$\Rightarrow$ Measure of fitting quality of LM: $R^2 = 1 - \frac{SSE_{LM}}{SSE_c} \in [0, 1]$

$\Rightarrow \rho(X_1, X_2) = R$

JOINT, MARGINAL AND CONDITIONAL DISTRIBUTION




For two discrete random variables X_1, X_2:

Joint distribution

$$p_{X_1,X_2}(x_1, x_2) = \mathbb{P}(X_1 = x_1, X_2 = x_2)$$

p_{X_1,X_2}	$\mathbb{P}(X_2 = 0)$	$\mathbb{P}(X_2 = 1)$	p_{X_1}
$\mathbb{P}(X_1 = 0)$	0.2	0.3	0.5
$\mathbb{P}(X_1 = 1)$	0.1	0.4	0.5
p_{X_2}	0.3	0.7	1

JOINT, MARGINAL AND CONDITIONAL DISTRIBUTION



For two discrete random variables X_1, X_2:

Joint distribution

$$p_{X_1,X_2}(x_1, x_2) = \mathbb{P}(X_1 = x_1, X_2 = x_2)$$

p_{X_1,X_2}	$\mathbb{P}(X_2 = 0)$	$\mathbb{P}(X_2 = 1)$	p_{X_1}
$\mathbb{P}(X_1 = 0)$	0.2	0.3	0.5
$\mathbb{P}(X_1 = 1)$	0.1	0.4	0.5
p_{X_2}	0.3	0.7	1


Marginal distribution

$$p_{X_1}(x_1) = \mathbb{P}(X_1 = x_1) = \sum_{x_2 \in \mathcal{X}_2} p(x_1, x_2)$$

p_{X_1,X_2}	$\mathbb{P}(X_2 = 0)$	$\mathbb{P}(X_2 = 1)$	p_{X_1}
$\mathbb{P}(X_1 = 0)$	0.2	0.3	0.5
$\mathbb{P}(X_1 = 1)$	0.1	0.4	0.5
p_{X_2}	0.3	0.7	1

$\rightsquigarrow$ In continuous case with integrals

JOINT, MARGINAL AND CONDITIONAL DISTRIBUTION



For two discrete random variables X_1, X_2:

Joint distribution

$$p_{X_1,X_2}(x_1, x_2) = \mathbb{P}(X_1 = x_1, X_2 = x_2)$$

p_{X_1,X_2}	$\mathbb{P}(X_2 = 0)$	$\mathbb{P}(X_2 = 1)$	p_{X_1}
$\mathbb{P}(X_1 = 0)$	0.2	0.3	0.5
$\mathbb{P}(X_1 = 1)$	0.1	0.4	0.5
p_{X_2}	0.3	0.7	1

Marginal distribution

$$p_{X_1}(x_1) = \mathbb{P}(X_1 = x_1) = \sum_{x_2 \in \mathcal{X}_2} p(x_1, x_2)$$

p_{X_1,X_2}	$\mathbb{P}(X_2 = 0)$	$\mathbb{P}(X_2 = 1)$	p_{X_1}
$\mathbb{P}(X_1 = 0)$	0.2	0.3	0.5
$\mathbb{P}(X_1 = 1)$	0.1	0.4	0.5
p_{X_2}	0.3	0.7	1

$\rightsquigarrow$ In continuous case with integrals


Conditional distribution

$$p_{X_1|X_2}(x_1 \mid x_2) = \mathbb{P}(X_1 = x_1 \mid X_2 = x_2) = \frac{p_{X_1,X_2}(x_1, x_2)}{p_{X_2}(x_2)}$$

	$x_2 = 0$	$x_2 = 1$
$\mathbb{P}(X_1 = 0 \mid X_2 = x_2)$	0.67	0.43
$\mathbb{P}(X_1 = 1 \mid X_2 = x_2)$	0.33	0.57
$\sum$	1	1

DEPENDENCE


Dependence: Describes general dependence structure (e.g., non-lin. relationships)



- Definition: X_j, X_k independent $\Leftrightarrow$ joint distribution is product of marginals:

$$\mathbb{P}(X_j, X_k) = \mathbb{P}(X_j) \cdot \mathbb{P}(X_k)$$

DEPENDENCE



Dependence: Describes general dependence structure (e.g., non-lin. relationships)


- Definition: X_j, X_k independent $\Leftrightarrow$ joint distribution is product of marginals:

$$\mathbb{P}(X_j, X_k) = \mathbb{P}(X_j) \cdot \mathbb{P}(X_k)$$

- Equivalent definition (knowing X_k gives no info about X_j and vice versa):

$$\mathbb{P}(X_j|X_k) = \mathbb{P}(X_j) \text{ and } \mathbb{P}(X_k|X_j) = \mathbb{P}(X_k) \text{ (follows from cond. probability)}$$

DEPENDENCE



Dependence: Describes general dependence structure (e.g., non-lin. relationships)

- Definition: X_j, X_k independent $\Leftrightarrow$ joint distribution is product of marginals:


$$\mathbb{P}(X_j, X_k) = \mathbb{P}(X_j) \cdot \mathbb{P}(X_k)$$

- Equivalent definition (knowing X_k gives no info about X_j and vice versa):

$$\mathbb{P}(X_j|X_k) = \mathbb{P}(X_j) \text{ and } \mathbb{P}(X_k|X_j) = \mathbb{P}(X_k) \text{ (follows from cond. probability)}$$

- Measuring complex dependencies is difficult but different measures exist
  Examples
  $\rightsquigarrow$ Spearman correlation (measures monotonic dependencies via ranks)
  $\rightsquigarrow$ Information-theoretical measures like mutual information
  $\rightsquigarrow$ Kernel-based measures like Hilbert-Schmidt Independence Criterion (HSIC)

DEPENDENCE



Dependence: Describes general dependence structure (e.g., non-lin. relationships)

- Definition: X_j, X_k independent $\Leftrightarrow$ joint distribution is product of marginals:


$$\mathbb{P}(X_j, X_k) = \mathbb{P}(X_j) \cdot \mathbb{P}(X_k)$$

- Equivalent definition (knowing X_k gives no info about X_j and vice versa):

$$\mathbb{P}(X_j|X_k) = \mathbb{P}(X_j) \text{ and } \mathbb{P}(X_k|X_j) = \mathbb{P}(X_k) \text{ (follows from cond. probability)}$$

- Measuring complex dependencies is difficult but different measures exist
  Examples
  $\rightsquigarrow$ Spearman correlation (measures monotonic dependencies via ranks)
  $\rightsquigarrow$ Information-theoretical measures like mutual information
  $\rightsquigarrow$ Kernel-based measures like Hilbert-Schmidt Independence Criterion (HSIC)
- **N.B.:** X_j, X_k indep. $\Rightarrow \rho(X_j, X_k) = 0$ **but** $\rho(X_j, X_k) = 0 \not\Rightarrow X_j$, X_k indep.
  Equivalency holds if distribution is jointly normal

MUTUAL INFORMATION




- MI describes expected amount of information shared by two RVs:

$$MI(X_1, X_2) = \mathbb{E}_{p(x_1,x_2)}\left[log\left(\frac{p(x_1,x_2)}{p(x_1)p(x_2)}\right)\right]$$

- MI measures amount of "dependence" between features by looking how different the joint distribution is from pure indep. $p(x_1, x_2) = p(x_1)p(x_2)$
  $\rightsquigarrow MI(X_1, X_2) = \mathbb{E}_{p(x_1,x_2)}\left[log\left(\frac{p(x_1,x_2)}{p(x_1,x_2)}\right)\right] = \mathbb{E}_{p(x_1,x_2)}\left[log(1)\right] = 0$
  $\rightsquigarrow MI(X_j, X_k) = 0$ if and only if the features are independent
- Unlike (Pearson) correlation, MI is also defined for categorical features

MUTUAL INFORMATION: EXAMPLE




For two discrete RV X_1 and Y:

$$MI(X_1; Y) = \mathbb{E}_{p(x_1,y)}\left[log\left(\frac{p(x_1, y)}{p(x_1)p(y)}\right)\right] = \sum_{x_1 \in \mathcal{X}_1} \sum_{y \in \mathcal{Y}} p(x_1, y) log\left(\frac{p(x_1, y)}{p(x_1)p(y)}\right)$$

$\mathbf{X}_1$	...	$\mathbf{Y}$
yes	...	yes
yes	...	no
no	...	yes
no	...	no

	$\mathbb{P}(X_1 = \text{yes})$	$\mathbb{P}(X_1 = \text{no})$	p_Y
$\mathbb{P}(Y = \text{yes})$	0.25	0.25	0.5
$\mathbb{P}(Y = \text{no})$	0.25	0.25	0.5
p_{X_1}	0.5	0.5	1

MUTUAL INFORMATION: EXAMPLE



For two discrete RV X_1 and Y:


$$MI(X_1; Y) = \mathbb{E}_{p(x_1,y)}\left[log\left(\frac{p(x_1,y)}{p(x_1)p(y)}\right)\right] = \sum_{x_1 \in \mathcal{X}_1}\sum_{y \in \mathcal{Y}} p(x_1,y) log\left(\frac{p(x_1,y)}{p(x_1)p(y)}\right)$$

X_1	...	Y
yes	...	yes
yes	...	no
no	...	yes
no	...	no


	$\mathbb{P}(X_1 = \text{yes})$	$\mathbb{P}(X_1 = \text{no})$	p_Y
$\mathbb{P}(Y = \text{yes})$	0.25	0.25	0.5
$\mathbb{P}(Y = \text{no})$	0.25	0.25	0.5
p_{X_1}	0.5	0.5	1

$$\begin{aligned} MI(X_1; Y) &= 0.25\, log\left(\frac{0.25}{0.5 \cdot 0.5}\right) + 0.25\, log\left(\frac{0.25}{0.5 \cdot 0.5}\right) \\ &\quad + 0.25\, log\left(\frac{0.25}{0.5 \cdot 0.5}\right) + 0.25\, log\left(\frac{0.25}{0.5 \cdot 0.5}\right) \\ &= 0.25\, log\left(\frac{0.25}{0.25}\right) \cdot 4 \\ &= 0.25\, log\,(1) \cdot 4 = 0 \end{aligned}$$

DEPENDENCE AND INDEPENDENCE




Example:




Conditional distributions at different vertical and horizontal slices (after normalizing area to 1) match their marginal distributions

$$\Rightarrow \mathbb{P}(X_1|X_2) = \mathbb{P}(X_1)$$
$$\mathbb{P}(X_2|X_1) = \mathbb{P}(X_2)$$

DEPENDENCE AND INDEPENDENCE




Example:



Conditional distributions at different vertical and horizontal slices (after normalizing area to 1) match their marginal distributions

$$\Rightarrow \mathbb{P}(X_1|X_2) = \mathbb{P}(X_1)$$
$$\mathbb{P}(X_2|X_1) = \mathbb{P}(X_2)$$



Conditional distributions do not match their marginal distributions

CORRELATION VS. DEPENDENCE





Illustration of bivariate normal distribution with different correlations
$X_1, X_2 \sim N(0, 1)$

$\rho(X_1, X_2) = 0$
(independent)

$\rho(X_1, X_2) = 0.8$

$\rho(X_1, X_2) = -0.8$



CORRELATION VS. DEPENDENCE





Illustration of bivariate normal distribution with different correlations
$X_1, X_2 \sim N(0, 1)$

$\rho(X_1, X_2) = 0$ (independent)

$\rho(X_1, X_2) = 0.8$

$\rho(X_1, X_2) = -0.8$



Examples with Pearson's corr. $\rho \approx 0$ but non-linear dependencies (MI $\neq 0$):

$\rho(X_1, X_2) = 0$, $MI(X_1, X_2) = 0.52$

$\rho(X_1, X_2) = 0.01$, $MI(X_1, X_2) = 0.37$

$\rho(X_1, X_2) = -0.06$, $MI(X_1, X_2) = 0.61$

