


Interpretable Machine Learning

Functional Decompositions: Further Methods



Learning goals

- Limitations of classical fANOVA
- Alternatives: Generalized fANOVA and ALE
- Advantages and relevance of functional decompositions


LIMITATIONS OF CLASSICAL FANOVA

- Standard fANOVA builds on PD-functions
- *Remember:* Problems of PDPs for **correlated / dependent features**
- Here: Dependent features \implies Standard fANOVA does NOT fulfill vanishing conditions



LIMITATIONS OF CLASSICAL FANOVA

- Standard fANOVA builds on PD-functions
- *Remember:* Problems of PDPs for **correlated / dependent features**
- Here: Dependent features \implies Standard fANOVA does NOT fulfill vanishing conditions



Example

Assume dependency $2x_1^2 = x_2$ and

$$\hat{f}(x_1, x_2, x_3) = -2x_1 - 2\sin(x_3) + |x_1|x_2 + 0.5x_2x_3 + 1.$$

\leadsto Following two decompositions would both “make sense”:

$$\hat{f}(x_1, x_2, x_3) = \underbrace{1}_{g_0} + \underbrace{(-2x_1)}_{g_1(x_1)} + \underbrace{(-2\sin(x_3))}_{g_3(x_3)} + \underbrace{|x_1|x_2}_{g_{1,2}(x_1, x_2)} + \underbrace{0.5x_2x_3}_{g_{2,3}(x_2, x_3)}$$

$$\hat{f}(x_1, x_2, x_3) = \underbrace{1}_{g_0} + \underbrace{(-2x_1 + 2|x_1|^3)}_{g_1(x_1)} + \underbrace{(-2\sin(x_3))}_{g_3(x_3)} + \underbrace{x_1^2x_3}_{g_{2,3}(x_1, x_3)}$$

LIMITATIONS OF CLASSICAL FANOVA

- Standard fANOVA builds on PD-functions
- *Remember:* Problems of PDPs for **correlated / dependent features**
- Here: Dependent features \implies Standard fANOVA does NOT fulfill vanishing conditions



Example

Assume dependency $2x_1^2 = x_2$ and

$$\hat{f}(x_1, x_2, x_3) = -2x_1 - 2\sin(x_3) + |x_1|x_2 + 0.5x_2x_3 + 1.$$

\rightsquigarrow Following two decompositions would both “make sense”:

$$\hat{f}(x_1, x_2, x_3) = \underbrace{1}_{g_0} + \underbrace{(-2x_1)}_{g_1(x_1)} + \underbrace{(-2\sin(x_3))}_{g_3(x_3)} + \underbrace{|x_1|x_2}_{g_{1,2}(x_1, x_2)} + \underbrace{0.5x_2x_3}_{g_{2,3}(x_2, x_3)}$$

$$\hat{f}(x_1, x_2, x_3) = \underbrace{1}_{g_0} + \underbrace{(-2x_1 + 2|x_1|^3)}_{g_1(x_1)} + \underbrace{(-2\sin(x_3))}_{g_3(x_3)} + \underbrace{x_1^2x_3}_{g_{2,3}(x_1, x_3)}$$

\rightarrow Extreme example, but again: Problem of definition

ALTERNATIVE: GENERALIZED FUNCTIONAL ANOVA


- Algorithm proposed by ▶ Hooker (2007)
- Generalizes standard fANOVA to situations with dependent features



ALTERNATIVE: GENERALIZED FUNCTIONAL ANOVA

- Algorithm proposed by ▶ Hooker (2007)
- Generalizes standard fANOVA to situations with dependent features
- Showed: Generalized fANOVA is solution to so-called “relaxed vanishing conditions”
(i.e., weaker form of vanishing condition)
- “Relaxed vanishing conditions” do not imply orthogonality, but “hierarchical orthogonality”:

$$\mathbb{E}_{\mathbf{x}}[g_V(\mathbf{x}_V)g_S(\mathbf{x}_S)] = 0 \quad \forall V \subsetneq S$$



ALTERNATIVE: GENERALIZED FUNCTIONAL ANOVA


- Algorithm proposed by ▶ Hooker (2007)
- Generalizes standard fANOVA to situations with dependent features
- Showed: Generalized fANOVA is solution to so-called “relaxed vanishing conditions”
(i.e., weaker form of vanishing condition)
- “Relaxed vanishing conditions” do not imply orthogonality, but “hierarchical orthogonality”:

$$\mathbb{E}_{\mathbf{x}}[g_V(\mathbf{x}_V)g_S(\mathbf{x}_S)] = 0 \quad \forall V \subsetneq S$$

\rightsquigarrow Only components are orthogonal where $g_V(\mathbf{x}_V)$ is “lower in hierarchy” than $g_S(\mathbf{x}_S)$

\implies Generalized fANOVA provides functional decomposition for arbitrary settings

- **Advantage:** Also provides a variance decomposition



ALTERNATIVE: GENERALIZED FUNCTIONAL ANOVA


- Algorithm proposed by ▶ Hooker (2007)
- Generalizes standard fANOVA to situations with dependent features
- Showed: Generalized fANOVA is solution to so-called “relaxed vanishing conditions”
(i.e., weaker form of vanishing condition)
- “Relaxed vanishing conditions” do not imply orthogonality, but “hierarchical orthogonality”:

$$\mathbb{E}_{\mathbf{x}}[g_V(\mathbf{x}_V)g_S(\mathbf{x}_S)] = 0 \quad \forall V \subsetneq S$$


↪ Only components are orthogonal where $g_V(\mathbf{x}_V)$ is “lower in hierarchy” than $g_S(\mathbf{x}_S)$

⇒ Generalized fANOVA provides functional decomposition for arbitrary settings



- **Advantage:** Also provides a variance decomposition
- **Problems:**
 - Difficult to estimate, involves manual choice of a “weight function”
 - Computationally very costly




REVISITING ALE PLOTS



$$\hat{f}_{S,ALE}(x) = \sum_{k=1}^{k_S(x)} \frac{1}{n_S(k)} \sum_{i: x_S^{(i)} \in [z_{k-1,S}, z_{k,S}]} \left[\hat{f}(z_{k,S}, \mathbf{x}_{-S}^{(i)}) - \hat{f}(z_{k-1,S}, \mathbf{x}_{-S}^{(i)}) \right]$$




Second order ALE



ALE DECOMPOSITION

- One can define ALE plots for arbitrary many variables (similar to PDPs vs. PD-functions)

→ Gives full functional decomposition of ALE plots




ALE DECOMPOSITION

- One can define ALE plots for arbitrary many variables (similar to PDPs vs. PD-functions)

→ Gives full functional decomposition of ALE plots

- **Advantages:** Handle dependencies well + computationally fast
- Constraints / orthogonality properties more complicated

⇒ ALE decomposition theoretically more involved, but good alternative in practice



CONCLUSION: HOW USEFUL ARE FUNCTIONAL DECOMPOSITIONS?

- If computed, offer a lot of insight into a model or function, i.p. high-dimensional
- Complete analysis of all interactions



CONCLUSION: HOW USEFUL ARE FUNCTIONAL DECOMPOSITIONS?

- If computed, offer a lot of insight into a model or function, i.p. high-dimensional
- Complete analysis of all interactions
- Very important theoretical concept:
 - Theoretical framework for general definition of interactions (H-statistic)
 - Theoretical background for many IML methods: GAMs and EBMs, ICE, PDPs and PD-functions, ALE plots, Shapley values, Feature importance methods (see later)




CONCLUSION: HOW USEFUL ARE FUNCTIONAL DECOMPOSITIONS?

- If computed, offer a lot of insight into a model or function, i.p. high-dimensional


→ Complete analysis of all interactions

- Very important theoretical concept:
 - Theoretical framework for general definition of interactions (H-statistic)
 - Theoretical background for many IML methods: GAMs and EBMs, ICE, PDPs and PD-functions, ALE plots, Shapley values, Feature importance methods (see later)
- In practice often infeasible (2^p components for p features)

⇒ Often only sparse decompositions feasible (E.g. EBMs)



CONCLUSION: HOW USEFUL ARE FUNCTIONAL DECOMPOSITIONS?




- If computed, offer a lot of insight into a model or function, i.p. high-dimensional

→ Complete analysis of all interactions

- Very important theoretical concept:
 - Theoretical framework for general definition of interactions (H-statistic)
 - Theoretical background for many IML methods: GAMs and EBMs, ICE, PDPs and PD-functions, ALE plots, Shapley values, Feature importance methods (see later)
 - In practice often infeasible (2^p components for p features)
- ⇒ Often only sparse decompositions feasible (E.g. EBMs)
- All single methods have disadvantages:
 - Standard fANOVA: Only independent features + compute intensive
 - Generalized fANOVA: Even more computational intensive, evtl. infeasible
 - ALE: No variance decomposition

CONCLUSION: HOW USEFUL ARE FUNCTIONAL DECOMPOSITIONS?



- If computed, offer a lot of insight into a model or function, i.p. high-dimensional

→ Complete analysis of all interactions

- Very important theoretical concept:
 - Theoretical framework for general definition of interactions (H-statistic)
 - Theoretical background for many IML methods: GAMs and EBM, ICE, PDPs and PD-functions, ALE plots, Shapley values, Feature importance methods (see later)
- In practice often infeasible (2^p components for p features)

⇒ Often only sparse decompositions feasible (E.g. EBM)

- All single methods have disadvantages:
 - Standard fANOVA: Only independent features + compute intensive
 - Generalized fANOVA: Even more computational intensive, evtl. infeasible
 - ALE: No variance decomposition

Overall: Very important concept and theoretical background, explains idea behind many other methods