

# Interpretable Machine Learning

## Feature Importances 1

## Conditional Feature Importance (CFI)






Figure: Bike Sharing Dataset

### Learning goals

- Extrapolation and Conditional Sampling
- Conditional Feature Importance (CFI)
- Interpretation of CFI and difference to PFI


# CFI MOTIVATION

- **PFI Idea:** Replace feature(s)  $X_S$  with perturbed  $\tilde{X}_S$  to preserve marginal distrib.  $\mathbb{P}(X_S)$  so that  $\tilde{X}_S \perp\!\!\!\perp Y$  (indep.), e.g., by random permutations




# CFI MOTIVATION

- **PFI Idea:** Replace feature(s)  $X_S$  with perturbed  $\tilde{X}_S$  to preserve marginal distrib.  $\mathbb{P}(X_S)$  so that  $\tilde{X}_S \perp\!\!\!\perp Y$  (indep.), e.g., by random permutations
- **Problem:** Breaks not only association between  $X_S$  and  $Y$  (what we want) but also between  $X_S, X_{-S} \Rightarrow \mathbb{P}(X_S, X_{-S}) \neq \mathbb{P}(\tilde{X}_S, X_{-S})$  (extrapolation)




# CFI MOTIVATION

- **PFI Idea:** Replace feature(s)  $X_S$  with perturbed  $\tilde{X}_S$  to preserve marginal distrib.  $\mathbb{P}(X_S)$  so that  $\tilde{X}_S \perp\!\!\!\perp Y$  (indep.), e.g., by random permutations
- **Problem:** Breaks not only association between  $X_S$  and  $Y$  (what we want) but also between  $X_S, X_{-S} \Rightarrow \mathbb{P}(X_S, X_{-S}) \neq \mathbb{P}(\tilde{X}_S, X_{-S})$  (extrapolation)
- **CFI Idea:** Replace feature(s)  $X_S$  with perturbed  $\tilde{X}_S$  to preserve joint distrib. so that  $\mathbb{P}(X_S, X_{-S}) = \mathbb{P}(\tilde{X}_S, X_{-S})$  (no extrapolation) while still  $\tilde{X}_S \perp\!\!\!\perp Y$




# CFI MOTIVATION

- **PFI Idea:** Replace feature(s)  $X_S$  with perturbed  $\tilde{X}_S$  to preserve marginal distrib.  $\mathbb{P}(X_S)$  so that  $\tilde{X}_S \perp\!\!\!\perp Y$  (indep.), e.g., by random permutations
- **Problem:** Breaks not only association between  $X_S$  and  $Y$  (what we want) but also between  $X_S, X_{-S} \Rightarrow \mathbb{P}(X_S, X_{-S}) \neq \mathbb{P}(\tilde{X}_S, X_{-S})$  (extrapolation)
- **CFI Idea:** Replace feature(s)  $X_S$  with perturbed  $\tilde{X}_S$  to preserve joint distrib. so that  $\mathbb{P}(X_S, X_{-S}) = \mathbb{P}(\tilde{X}_S, X_{-S})$  (no extrapolation) while still  $\tilde{X}_S \perp\!\!\!\perp Y$



**Example:** Conditional permutation scheme

**Black dots:**  $X_2 \sim \mathcal{U}(0, 1)$  and  $X_1 \sim \mathcal{N}(0, 1)$  (if  $X_2 < 0.5$ ) or  $\mathcal{N}(4, 4)$  (if  $X_2 \geq 0.5$ )



**Left:** For  $X_2 < 0.5$ , permuting  $X_1$  (crosses) preserves marginal (but not joint) distrib.

$\rightsquigarrow$  Bottom: Marginal density of  $X_1$

**Right:** Permuting  $X_1$  within subgroups  $X_2 < 0.5$  &  $X_2 \geq 0.5$  reduces extrapolation



$\rightsquigarrow$  Bottom:  $X_1$ -density cond. on groups

► Molnar 2020

# RECALL: EXTRAPOLATION IN PFI

**Recall:** Let  $y = x_3 + \epsilon_y$ , with  $\epsilon_y \sim \mathcal{N}(0, 0.1)$ .

- $x_1 := \epsilon_1$ ,  $x_2 := x_1 + \epsilon_2$ ; highly correlated ( $\epsilon_1 \sim \mathcal{N}(0, 1)$ ,  $\epsilon_2 \sim \mathcal{N}(0, 0.01)$ )
- $x_3 := \epsilon_3$ ,  $x_4 := \epsilon_4$ , with  $\epsilon_3, \epsilon_4 \sim \mathcal{N}(0, 1)$ ; all noise terms  $\epsilon_j$  are indep.
- Fitting a linear model yields  $\hat{f}(\mathbf{x}) \approx 0.3x_1 - 0.3x_2 + x_3$



Hexbin plot of  $(x_1, x_2)$  before (left) and after (center) permuting  $x_1$ ;  
PFI scores (right).

- $\Rightarrow x_1, x_2$  cancel in  $\hat{f}$  and should be irrelevant
- $\Rightarrow$  But PFI evaluates model on unrealistic inputs (caused by permutation)
  - $\leadsto PFI > 0$  for  $x_1, x_2$  due to extrapolation
  - $\leadsto x_1, x_2$  are misleadingly considered relevant

CFI for  $X_S$  using test data  $\mathcal{D}$ :

- Measure the error **with unperturbed features**  $x_S$ .
- Measure the error **with perturbed feature values**  $x_S \sim \mathbb{P}(X_S|X_{-S})$
- Repeat perturbing  $X_S$  (e.g.,  $m$  times) and avg. difference of both errors:

$$\widehat{CFI}_S = \frac{1}{m} \sum_{k=1}^m \mathcal{R}_{\text{emp}}(\hat{f}, \mathcal{DS} - S_{(k)}) - \mathcal{R}_{\text{emp}}(\hat{f}, \mathcal{D})$$


Here,  $\mathcal{DS} - S$  denotes data, where  $x_S$  values are conditionally resampled given  $x_{-S}$ .

**Illustrative example:** Conditional permutation when  $X_{-S}$  is categorical:

Original Data		
ID	$X_{-S}$	$X_S$
1	A	3.1
2	A	2.7
3	A	3.4
4	B	6.0
5	B	5.4
6	B	6.2

Permuted Conditionally on $X_{-S}$		
ID	$X_{-S}$	$X_S$
1	A	2.7
2	A	3.1
3	A	3.4
4	B	6.2
5	B	6.0
6	B	5.4


Here,  $X_S$  is permuted *within* each group of  $X_{-S}$  to preserve  $\mathbb{P}(X_S, X_{-S})$ .



# IMPLICATIONS OF CFI

► K\_NIG\_ET\_2020

**Interpretation:** Due to the conditional sampling w.r.t. all other features, CFI quantifies a feature's unique contribution to the model performance.






# IMPLICATIONS OF CFI

► K\_NIG\_ET\_2020

**Interpretation:** Due to the conditional sampling w.r.t. all other features, CFI quantifies a feature's unique contribution to the model performance.

## Entanglement with data:


- If feat  $x_S$  does not contrib. unique information about  $y$ , i.e.,  $x_S \perp\!\!\!\perp y | x_{-S}$   
 $\Rightarrow \text{CFI} = 0$
- Why? Under the conditional indep.  $\mathbb{P}(X_S, X_{-S}, Y) = \mathbb{P}(X_S, X_{-S}, Y)$   
 $\leadsto$  no prediction-relevant information is destroyed by permutation of  $x_S$  conditional on  $x_{-S}$



# IMPLICATIONS OF CFI

► K\_NIG\_ET\_2020

**Interpretation:** Due to the conditional sampling w.r.t. all other features, CFI quantifies a feature's unique contribution to the model performance.



## Entanglement with data:

- If feat  $x_S$  does not contrib. unique information about  $y$ , i.e.,  $x_S \perp\!\!\!\perp y | x_{-S}$   
 $\Rightarrow \text{CFI} = 0$
- Why? Under the conditional indep.  $\mathbb{P}(X_S, X_{-S}, Y) = \mathbb{P}(X_S, X_{-S}, Y)$   
 $\rightsquigarrow$  no prediction-relevant information is destroyed by permutation of  $x_S$  conditional on  $x_{-S}$


## Entanglement with model:

- If the model does not use a feature  $\Rightarrow \text{CFI} = 0$
- Why? Then the prediction is not affected by any perturbation of the feat  
 $\rightsquigarrow$  model performance does not change after conditional permutation

# IMPLICATIONS OF CFI

Can we gain insight into whether ...


- ❶ the feature  $x_j$  is causal for the prediction?
  - $CFI_j \neq 0 \Rightarrow$  model relies on  $x_j$   
(converse does not hold, see next slide)



# IMPLICATIONS OF CFI

Can we gain insight into whether ...


- ❶ the feature  $x_j$  is causal for the prediction?
  - $CFI_j \neq 0 \Rightarrow$  model relies on  $x_j$   
(converse does not hold, see next slide)
- ❷ the variable  $x_j$  contains prediction-relevant information?
  - If  $x_j \not\perp\!\!\!\perp y$  but  $x_j \perp\!\!\!\perp y | x_{-j}$  (e.g.,  $x_j$  and  $x_{-j}$  share information)  
 $\Rightarrow CFI_j = 0$
  - $x_j$  is not exploited by model (regardless of its usefulness for  $y$ )  
 $\Rightarrow CFI_j = 0$



# IMPLICATIONS OF CFI

Can we gain insight into whether ...



- ❶ the feature  $x_j$  is causal for the prediction?
  - $CFI_j \neq 0 \Rightarrow$  model relies on  $x_j$   
(converse does not hold, see next slide)
- ❷ the variable  $x_j$  contains prediction-relevant information?
  - If  $x_j \not\perp\!\!\!\perp y$  but  $x_j \perp\!\!\!\perp y|x_{-j}$  (e.g.,  $x_j$  and  $x_{-j}$  share information)  
 $\Rightarrow CFI_j = 0$
  - $x_j$  is not exploited by model (regardless of its usefulness for  $y$ )  
 $\Rightarrow CFI_j = 0$
- ❸ Does the model need access to  $x_j$  to achieve its prediction performance?
  - $CFI_j \neq 0 \Rightarrow x_j$  contributes unique information (meaning  $x_j \not\perp\!\!\!\perp y|x_{-j}$ )
  - Only uncovers the relationships that were exploited by the model



# EXTRAPOLATION: COMPARE PFI AND CFI

**Recall:** Let  $y = x_3 + \epsilon_y$ , with  $\epsilon_y \sim \mathcal{N}(0, 0.1)$ .

- $x_1 := \epsilon_1$ ,  $x_2 := x_1 + \epsilon_2$ ; highly correlated ( $\epsilon_1 \sim \mathcal{N}(0, 1)$ ,  $\epsilon_2 \sim \mathcal{N}(0, 0.01)$ )
- $x_3 := \epsilon_3$ ,  $x_4 := \epsilon_4$ , with  $\epsilon_3, \epsilon_4 \sim \mathcal{N}(0, 1)$ ; all noise terms  $\epsilon_j$  are indep.
- Fitting a linear model yields  $\hat{f}(\mathbf{x}) \approx 0.3x_1 - 0.3x_2 + x_3$



**Figure:** Density plot for  $x_1, x_2$  before permuting  $x_1$  (left). PFI and CFI (right).

- $x_1$  and  $x_2$  cancel in  $\hat{f}(\mathbf{x})$  and should be irrelevant for the prediction
- PFI evaluates model on unrealistic obs.  
     $\leadsto x_1, x_2$  appear relevant (PFI > 0)
- CFI evaluates model on realistic obs. (due to conditional sampling)  
     $\leadsto x_1, x_2$  appear irrelevant (CFI = 0)