

Interpretable Machine Learning

Permutation Feature Importance (PFI)





Figure: Bike Sharing Dataset

Learning goals

- Understand how PFI is computed
- Understanding strengths and weaknesses



Interpretable Machine Learning

Feature Importances 1

Permutation Feature Importance (PFI)





Figure: Bike Sharing Dataset


Learning goals

- Understand how PFI is computed
- Understanding strengths and weaknesses




MOTIVATION FOR PFI

- **Goal:** Assess how important feature(s) X_S are for predictive performance of a **fixed trained model** \hat{f} on a given dataset \mathcal{D}
- **Idea:** Estimate change in model performance when X_S is "made uninformative"




MOTIVATION FOR PFI

- **Goal:** Assess how important feature(s) X_S are for predictive performance of a **fixed trained model** \hat{f} on a given dataset \mathcal{D}
- **Idea:** Estimate performance change when X_S is "made uninformative"




MOTIVATION FOR PFI

- **Goal:** Assess how important feature(s) X_S are for predictive performance of a **fixed trained model** \hat{f} on a given dataset \mathcal{D}
- **Idea:** Estimate change in model performance when X_S is "made uninformative"
- **Question:** Can we make X_S uninformative by removing it from the model?
~~ No, \hat{f} was trained with X_S and retraining without X_S gives a different model



MOTIVATION FOR PFI

- **Goal:** Assess how important feature(s) X_S are for predictive performance of a **fixed trained model** \hat{f} on a given dataset \mathcal{D}
- **Idea:** Estimate performance change when X_S is "made uninformative"
- **Question:** Can we make X_S uninformative by removing it from model?
~~ No, \hat{f} was trained with X_S ; retraining without X_S gives a different model




MOTIVATION FOR PFI

- **Goal:** Assess how important feature(s) X_S are for predictive performance of a **fixed trained model** \hat{f} on a given dataset \mathcal{D}
- **Idea:** Estimate change in model performance when X_S is "made uninformative"
- **Question:** Can we make X_S uninformative by removing it from the model?
~~ No, \hat{f} was trained with X_S and retraining without X_S gives a different model
- **Solution:** Simulate feature removal by replacing X_S with a perturbed version \tilde{X}_S that is independent of (X_{-S}, Y) but preserves distribution $\mathbb{P}(X_S)$
~~ Compare **baseline predictions** $\hat{f}(X)$ with **perturbed predictions** $\hat{f}(\tilde{X}_S, X_{-S})$

$$\text{PFI}_S := \underbrace{\mathbb{E}\left[L(\hat{f}(\tilde{X}_S, X_{-S}), Y)\right]}_{\text{risk after "destroying" } X_S} - \underbrace{\mathbb{E}\left[L(\hat{f}(X), Y)\right]}_{\text{baseline risk}},$$

- **How to perturb X_S ?**

- Add random noise: distorts $\mathbb{P}(X_S)$ (not used)
- Permutation: preserves marginal $\mathbb{P}(X_S)$, breaks dependence with Y (used)




MOTIVATION FOR PFI

- **Goal:** Assess how important feature(s) X_S are for predictive performance of a **fixed trained model** \hat{f} on a given dataset \mathcal{D}
- **Idea:** Estimate performance change when X_S is "made uninformative"
- **Question:** Can we make X_S uninformative by removing it from model?
~~ No, \hat{f} was trained with X_S ; retraining without X_S gives a different model
- **Solution:** Simulate feature removal by replacing X_S with a perturbed version \tilde{X}_S that is independent of (X_{-S}, Y) but preserves distrib. $\mathbb{P}(X_S)$
~~ Compare **baseline predictions** $\hat{f}(X)$ with **perturbed predictions** $\hat{f}(\tilde{X}_S, X_{-S})$

$$\text{PFI}_S := \underbrace{\mathbb{E}\left[L(\hat{f}(\tilde{X}_S, X_{-S}), Y)\right]}_{\text{risk after "destroying" } X_S} - \underbrace{\mathbb{E}\left[L(\hat{f}(X), Y)\right]}_{\text{baseline risk}},$$

- **How to perturb X_S ?**

- Add random noise: distorts $\mathbb{P}(X_S)$ (not used)
- Permutation: preserves marginal $\mathbb{P}(X_S)$, breaks dependence with Y (used)



PERMUTATION FEATURE IMPORTANCE (PFI)

▶ Breiman (2001)

Sample estimator (using independent test set $\mathcal{D} = \{(\mathbf{x}^{(i)}, y^{(i)})\}_{i=1}^n$)

- Measure the error **with feat. values x_S** and **with permuted feat. values \tilde{x}_S**
- Repeat permutation (e.g., m times) and average difference of both errors:


$$\widehat{PFI}_S = \frac{1}{m} \sum_{k=1}^m [\mathcal{R}_{\text{emp}}(\hat{f}, \tilde{\mathcal{D}}_{(k)}^S) - \mathcal{R}_{\text{emp}}(\hat{f}, \mathcal{D})]$$

- $\tilde{\mathcal{D}}_S^{(k)}$: dataset where column(s) x_S are **permuted** once (in repetition k)
- $\mathcal{R}_{\text{emp}}(\hat{f}, \mathcal{D}) = \frac{1}{n} \sum_{(x,y) \in \mathcal{D}} L(\hat{f}(x), y)$: Measures performance of \hat{f} using \mathcal{D}
- Average over m permutations to reduce Monte-Carlo variance

Example of permuting feature x_S with $S = \{1\}$ and $m = 6$ permutations:

\mathcal{D}	$\tilde{\mathcal{D}}_{(1)}^S$	$\tilde{\mathcal{D}}_{(2)}^S$	$\tilde{\mathcal{D}}_{(3)}^S$	$\tilde{\mathcal{D}}_{(4)}^S$	$\tilde{\mathcal{D}}_{(5)}^S$	$\tilde{\mathcal{D}}_{(6)}^S$
$\begin{array}{ c c c }\hline x_1 & x_2 & x_3 \\ \hline 1 & 4 & 7 \\ \hline 2 & 5 & 8 \\ \hline 3 & 6 & 9 \\ \hline\end{array}$	$\begin{array}{ c c c }\hline x_S & x_2 & x_3 \\ \hline 1 & 4 & 7 \\ \hline 2 & 5 & 8 \\ \hline 3 & 6 & 9 \\ \hline\end{array}$	$\begin{array}{ c c c }\hline x_S & x_2 & x_3 \\ \hline 2 & 4 & 7 \\ \hline 1 & 5 & 8 \\ \hline 3 & 6 & 9 \\ \hline\end{array}$	$\begin{array}{ c c c }\hline x_S & x_2 & x_3 \\ \hline 2 & 4 & 7 \\ \hline 3 & 5 & 8 \\ \hline 1 & 6 & 9 \\ \hline\end{array}$	$\begin{array}{ c c c }\hline x_S & x_2 & x_3 \\ \hline 1 & 4 & 7 \\ \hline 3 & 5 & 8 \\ \hline 2 & 6 & 9 \\ \hline\end{array}$	$\begin{array}{ c c c }\hline x_S & x_2 & x_3 \\ \hline 3 & 4 & 7 \\ \hline 2 & 5 & 8 \\ \hline 1 & 6 & 9 \\ \hline\end{array}$	$\begin{array}{ c c c }\hline x_S & x_2 & x_3 \\ \hline 3 & 4 & 7 \\ \hline 2 & 5 & 8 \\ \hline 1 & 6 & 9 \\ \hline\end{array}$
\Rightarrow						

Note: S refers to a subset of features, here $|S| = 1$ to measure impact of permuting x_1 on performance



PERMUTATION FEATURE IMPORTANCE (PFI)

▶ BREIMAN_2001

Sample estimator (using independent test set $\mathcal{D} = \{(\mathbf{x}^{(i)}, y^{(i)})\}_{i=1}^n$)

- Measure error **with feat. values x_S** and **with permuted feat. values \tilde{x}_S**
- Repeat permutation (e.g., m times) and average difference of both errors:


$$\widehat{PFI}_S = \frac{1}{m} \sum_{k=1}^m [\mathcal{R}_{\text{emp}}(\hat{f}, \tilde{\mathcal{D}}_{S(k)}^S) - \mathcal{R}_{\text{emp}}(\hat{f}, \mathcal{D})]$$

- $\tilde{\mathcal{D}}_S^{(k)}$: dataset with column(s) x_S are **permuted** once (in repetition k)
- $\mathcal{R}_{\text{emp}}(\hat{f}, \mathcal{D}) = \frac{1}{n} \sum_{(x,y) \in \mathcal{D}} L(\hat{f}(x), y)$: Measures performance of \hat{f} using \mathcal{D}
- Average over m permutations to reduce Monte-Carlo variance

Example of permuting feature x_S with $S = \{1\}$ and $m = 6$ permutations:

\mathcal{D}	$\tilde{\mathcal{D}}_{(1)}^S$	$\tilde{\mathcal{D}}_{(2)}^S$	$\tilde{\mathcal{D}}_{(3)}^S$	$\tilde{\mathcal{D}}_{(4)}^S$	$\tilde{\mathcal{D}}_{(5)}^S$	$\tilde{\mathcal{D}}_{(6)}^S$
$\begin{array}{ c c c }\hline x_1 & x_2 & x_3 \\ \hline 1 & 4 & 7 \\ \hline 2 & 5 & 8 \\ \hline 3 & 6 & 9 \\ \hline\end{array}$	$\begin{array}{ c c c }\hline x_S & x_2 & x_3 \\ \hline 1 & 4 & 7 \\ \hline 2 & 5 & 8 \\ \hline 3 & 6 & 9 \\ \hline\end{array}$	$\begin{array}{ c c c }\hline x_S & x_2 & x_3 \\ \hline 2 & 4 & 7 \\ \hline 1 & 5 & 8 \\ \hline 3 & 6 & 9 \\ \hline\end{array}$	$\begin{array}{ c c c }\hline x_S & x_2 & x_3 \\ \hline 2 & 4 & 7 \\ \hline 3 & 5 & 8 \\ \hline 1 & 6 & 9 \\ \hline\end{array}$	$\begin{array}{ c c c }\hline x_S & x_2 & x_3 \\ \hline 1 & 4 & 7 \\ \hline 3 & 5 & 8 \\ \hline 2 & 6 & 9 \\ \hline\end{array}$	$\begin{array}{ c c c }\hline x_S & x_2 & x_3 \\ \hline 3 & 4 & 7 \\ \hline 2 & 5 & 8 \\ \hline 1 & 6 & 9 \\ \hline\end{array}$	$\begin{array}{ c c c }\hline x_S & x_2 & x_3 \\ \hline 3 & 4 & 7 \\ \hline 2 & 5 & 8 \\ \hline 1 & 6 & 9 \\ \hline\end{array}$
\Rightarrow						

Note: S refers to a subset of features, here $|S| = 1$ to measure impact of permuting x_1 on performance




PERMUTATION FEATURE IMPORTANCE

$\tilde{\mathcal{D}}_{(k)}^S$

i	x_1	x_2	x_3
1	2	4	7
:	1	5	8
n	3	6	9

\mathcal{D}

i	x_1	x_2	x_3
1	1	4	7
:	2	5	8
n	3	6	9




PERMUTATION FEATURE IMPORTANCE

$\mathcal{D}_{(k)}$

i	x_1	x_2	x_3
1	2	4	7
:	1	5	8
n	3	6	9

\mathcal{D}


i	x_1	x_2	x_3
1	1	4	7
:	2	5	8
n	3	6	9




1. **Perturbation:** Sample feature values from the distribution of x_S ($P(X_S)$).
 - ⇒ Randomly permute feature x_S
 - ⇒ Replace x_S with permuted feature \tilde{x}_S and create data $\tilde{\mathcal{D}}^S$ containing \tilde{x}_S

1. **Perturbation:** Sample feature values from the distribution of x_S ($P(X_S)$).
 - ⇒ Randomly permute feature x_S
 - ⇒ Replace x_S with permuted feat. x_S and create data \mathcal{D}_S containing x_S


PERMUTATION FEATURE IMPORTANCE




1. **Perturbation:** Sample feature values from the distribution of x_s ($P(X_s)$).
⇒ Randomly permute feature x_s
⇒ Replace x_s with permuted feature \tilde{x}_s and create data $\tilde{\mathcal{D}}^s$ containing \tilde{x}_s
2. **Prediction:** Make predictions for both data, i.e., \mathcal{D} and $\tilde{\mathcal{D}}^s$




PERMUTATION FEATURE IMPORTANCE



1. **Perturbation:** Sample feature values from the distribution of x_s ($P(X_s)$).
⇒ Randomly permute feature x_s
⇒ Replace x_s with permuted feat. x_s and create data $\tilde{\mathcal{D}}_s$ containing x_s
2. **Prediction:** Make predictions for both data, i.e., \mathcal{D} and $\tilde{\mathcal{D}}_s$




PERMUTATION FEATURE IMPORTANCE




3. Aggregation:

- Compute the loss for each observation in both data sets




PERMUTATION FEATURE IMPORTANCE



3. Aggregation:

- Compute the loss for each observation in both data sets




PERMUTATION FEATURE IMPORTANCE

$\tilde{D}_{(k)}^S$

i	x_1	x_2	x_3	D	x_1	x_2	x_3	ΔL
1	2	4	7		1	4	7	0.65
:	1	5	8		2	5	8	0.15
n	3	6	9		3	6	9	0

$$L(\hat{f}, y)$$

0.9	-	0.25
0.5	-	0.35
0.1	-	0.1



3. Aggregation:

- Compute the loss for each observation in both data sets
- Take the difference of both losses ΔL for each observation

PERMUTATION FEATURE IMPORTANCE

$D_{S(k)}$


i	x_1	x_2	x_3	D	x_1	x_2	x_3	ΔL
1	2	4	7		1	4	7	0.65
:	1	5	8		2	5	8	0.15
n	3	6	9		3	6	9	0

$$L(\hat{f}, y)$$

0.9	-	0.25
0.5	-	0.35
0.1	-	0.1

3. Aggregation:

- Compute the loss for each observation in both data sets
- Take the difference of both losses ΔL for each observation



PERMUTATION FEATURE IMPORTANCE


$$\mathcal{R}_{\text{emp}}(\hat{f}, \tilde{\mathcal{D}}_{(k)}^S) - \mathcal{R}_{\text{emp}}(\hat{f}, \mathcal{D})$$

i	\mathbf{x}_S	\mathbf{x}_2	\mathbf{x}_3
1	2	4	7
:	1	5	8
n	3	6	9

i	\mathbf{x}_1	\mathbf{x}_2	\mathbf{x}_3
1	1	4	7
:	2	5	8
n	3	6	9

	ΔL
1	0.65
2	0.15
n	0

$$= 0.267$$



PERMUTATION FEATURE IMPORTANCE


$$\mathcal{R}_{\text{emp}}(\hat{f}, \mathcal{DS}_{(k)}) - \mathcal{R}_{\text{emp}}(\hat{f}, \mathcal{D})$$

i	\mathbf{x}_S	\mathbf{x}_2	\mathbf{x}_3
1	2	4	7
:	1	5	8
n	3	6	9

i	\mathbf{x}_1	\mathbf{x}_2	\mathbf{x}_3
1	1	4	7
:	2	5	8
n	3	6	9

	ΔL
1	0.65
2	0.15
n	0

$$= 0.267$$



3. Aggregation:

- Compute the loss for each observation in both data sets
- Take the difference of both losses ΔL for each observation
- Average this change in loss across all observations

Note: Same as computing \mathcal{R}_{emp} on both data sets and taking difference

3. Aggregation:

- Compute the loss for each observation in both data sets
- Take the difference of both losses ΔL for each observation
- Average this change in loss across all observations

Note: Same as computing \mathcal{R}_{emp} on both data sets and taking difference

PERMUTATION FEATURE IMPORTANCE

$$\mathcal{R}_{\text{emp}}(\hat{f}, \tilde{\mathcal{D}}_{(k)}^S) - \mathcal{R}_{\text{emp}}(\hat{f}, \mathcal{D})$$

i	\mathbf{x}_S	\mathbf{x}_2	\mathbf{x}_3	\mathbf{x}_1	\mathbf{x}_2	\mathbf{x}_3
1	2	4	7	1	4	7
\vdots	1	5	8	2	5	8
n	3	6	9	3	6	9

i	ΔL
1	0.65
\vdots	0.15
n	0

$$= 0.267$$


$$\widehat{PFI}_S = \frac{1}{2} (0.267 + 0.4)$$

i	ΔL
1	0.85
\vdots	0
n	0.35

$$= 0.4$$

3. Aggregation:

- Compute the loss for each observation in both data sets
- Take the difference of both losses ΔL for each observation
- Average this change in loss across all observations
- Repeat perturbation and average over multiple repetitions



PERMUTATION FEATURE IMPORTANCE

$$\mathcal{R}_{\text{emp}}(\hat{f}, \tilde{\mathcal{D}}_{(k)}^S) - \mathcal{R}_{\text{emp}}(\hat{f}, \mathcal{D})$$

i	\mathbf{x}_S	\mathbf{x}_2	\mathbf{x}_3	\mathbf{x}_1	\mathbf{x}_2	\mathbf{x}_3
1	2	4	7	1	4	7
\vdots	1	5	8	2	5	8
n	3	6	9	3	6	9

i	ΔL
1	0.65
\vdots	0.15
n	0

$$= 0.267$$


$$\widehat{PFI}_S = \frac{1}{2} (0.267 + 0.4)$$

i	ΔL
1	0.85
\vdots	0
n	0.35


$$= 0.4$$

3. Aggregation:

- Compute the loss for each observation in both data sets
- Take the difference of both losses ΔL for each observation
- Average this change in loss across all observations
- Repeat perturbation and average over multiple repetitions




EXAMPLE: BIKE SHARING DATASET




Interpretation:

- 'yr' and 'temp' are most important features using mean absolute error (MAE)
- Destroying information about 'yr' by permuting it increases MAE of model by 816
- Error bars show 5% and 95% quantiles over multiple permutations




EXAMPLE: BIKE SHARING DATASET




Interpretation:

- 'yr' and 'temp' are most important feats using mean absolute error (MAE)
- Destroying info. about 'yr' by permuting it increases MAE of model by 816
- Error bars show 5% and 95% quantiles over multiple permutations




COMMENTS ON PFI

- Interpretation: Increase in error when feature's information is destroyed




COMMENTS ON PFI

- Interpretation: Increase in error when feature's information is destroyed




COMMENTS ON PFI

- Interpretation: Increase in error when feature's information is destroyed
- Results can be unreliable due to random permutations
⇒ Solution: Average results over multiple repetitions




COMMENTS ON PFI

- Interpretation: Increase in error when feature's information is destroyed
- Results can be unreliable due to random permutations
⇒ Solution: Average results over multiple repetitions




COMMENTS ON PFI

- Interpretation: Increase in error when feature's information is destroyed
- Results can be unreliable due to random permutations
⇒ Solution: Average results over multiple repetitions
- Permuting features despite correlation/dependence with other features can lead to unrealistic combinations of feature values ↪ Extrapolation issue




COMMENTS ON PFI

- Interpretation: Increase in error when feature's information is destroyed
- Results can be unreliable due to random permutations
⇒ Solution: Average results over multiple repetitions
- Permuting features despite correlation/dependence with other features can lead to unrealistic combinations of feature values
↪ Extrapolation issue




COMMENTS ON PFI

- Interpretation: Increase in error when feature's information is destroyed
- Results can be unreliable due to random permutations
⇒ Solution: Average results over multiple repetitions
- Permuting features despite correlation/dependence with other features can lead to unrealistic combinations of feature values ↪ Extrapolation issue
- PFI automatically includes importance of interaction effects with other features
⇒ Permuting x_j also destroys interactions with permuted feature
⇒ PFI score contains importance of all interactions with permuted feature




COMMENTS ON PFI

- Interpretation: Increase in error when feature's information is destroyed
- Results can be unreliable due to random permutations
⇒ Solution: Average results over multiple repetitions
- Permuting features despite correlation/dependence with other features can lead to unrealistic combinations of feature values
↪ Extrapolation issue
- PFI automatically includes importance of interaction effects with other features
⇒ Permuting x_j also destroys interactions with permuted feature
⇒ PFI score contains importance of all interactions with permuted feature




COMMENTS ON PFI

- Interpretation: Increase in error when feature's information is destroyed
- Results can be unreliable due to random permutations
⇒ Solution: Average results over multiple repetitions
- Permuting features despite correlation/dependence with other features can lead to unrealistic combinations of feature values ↪ Extrapolation issue
- PFI automatically includes importance of interaction effects with other features
⇒ Permuting x_j also destroys interactions with permuted feature
⇒ PFI score contains importance of all interactions with permuted feature
- Interpretation of PFI depends on whether training or test data is used



COMMENTS ON PFI


- Interpretation: Increase in error when feature's information is destroyed
- Results can be unreliable due to random permutations
⇒ Solution: Average results over multiple repetitions
- Permuting features despite correlation/dependence with other features can lead to unrealistic combinations of feature values
↪ Extrapolation issue
- PFI automatically includes importance of interaction effects with other features
⇒ Permuting x_j also destroys interactions with permuted feature
⇒ PFI score contains importance of all interactions with permuted feature
- Interpretation of PFI depends on whether training or test data is used



COMMENTS ON PFI - EXTRAPOLATION

Example: Let $y = x_3 + \epsilon_y$, with $\epsilon_y \sim \mathcal{N}(0, 0.1)$.


- $x_1 := \epsilon_1$, $x_2 := x_1 + \epsilon_2$ are highly correlated ($\epsilon_1 \sim \mathcal{N}(0, 1)$, $\epsilon_2 \sim \mathcal{N}(0, 0.01)$)
- $x_3 := \epsilon_3$, $x_4 := \epsilon_4$, with $\epsilon_3, \epsilon_4 \sim \mathcal{N}(0, 1)$ and all noise terms ϵ_j are independent
- Fitting a linear model yields $\hat{f}(\mathbf{x}) \approx 0.3x_1 - 0.3x_2 + x_3$



COMMENTS ON PFI - EXTRAPOLATION

Example: Let $y = x_3 + \epsilon_y$, with $\epsilon_y \sim \mathcal{N}(0, 0.1)$.


- $x_1 := \epsilon_1$, $x_2 := x_1 + \epsilon_2$; highly correlated ($\epsilon_1 \sim \mathcal{N}(0, 1)$, $\epsilon_2 \sim \mathcal{N}(0, 0.01)$)
- $x_3 := \epsilon_3$, $x_4 := \epsilon_4$, with $\epsilon_3, \epsilon_4 \sim \mathcal{N}(0, 1)$; all noise terms ϵ_j are indep.
- Fitting a linear model yields $\hat{f}(\mathbf{x}) \approx 0.3x_1 - 0.3x_2 + x_3$




COMMENTS ON PFI - EXTRAPOLATION

Example: Let $y = x_3 + \epsilon_y$, with $\epsilon_y \sim \mathcal{N}(0, 0.1)$.

- $x_1 := \epsilon_1$, $x_2 := x_1 + \epsilon_2$ are highly correlated ($\epsilon_1 \sim \mathcal{N}(0, 1)$, $\epsilon_2 \sim \mathcal{N}(0, 0.01)$)
- $x_3 := \epsilon_3$, $x_4 := \epsilon_4$, with $\epsilon_3, \epsilon_4 \sim \mathcal{N}(0, 1)$ and all noise terms ϵ_j are independent
- Fitting a linear model yields $\hat{f}(\mathbf{x}) \approx 0.3x_1 - 0.3x_2 + x_3$




Hexbin plot of (x_1, x_2) before (left) and after (center) permuting x_1 ; PFI scores (right).




COMMENTS ON PFI - EXTRAPOLATION

Example: Let $y = x_3 + \epsilon_y$, with $\epsilon_y \sim \mathcal{N}(0, 0.1)$.

- $X_1 := \epsilon_1$, $X_2 := X_1 + \epsilon_2$; highly correlated ($\epsilon_1 \sim \mathcal{N}(0, 1)$, $\epsilon_2 \sim \mathcal{N}(0, 0.01)$)
- $X_3 := \epsilon_3$, $X_4 := \epsilon_4$, with $\epsilon_3, \epsilon_4 \sim \mathcal{N}(0, 1)$; all noise terms ϵ_j are indep.
- Fitting a linear model yields $\hat{f}(\mathbf{x}) \approx 0.3x_1 - 0.3x_2 + x_3$




Hexbin plot of (x_1, x_2) before (left) and after (center) permuting x_1 ; PFI scores (right).



COMMENTS ON PFI - EXTRAPOLATION


Example: Let $y = x_3 + \epsilon_y$, with $\epsilon_y \sim \mathcal{N}(0, 0.1)$.

- $x_1 := \epsilon_1, x_2 := x_1 + \epsilon_2$ are highly correlated ($\epsilon_1 \sim \mathcal{N}(0, 1), \epsilon_2 \sim \mathcal{N}(0, 0.01)$)
- $x_3 := \epsilon_3, x_4 := \epsilon_4$, with $\epsilon_3, \epsilon_4 \sim \mathcal{N}(0, 1)$ and all noise terms ϵ_j are independent
- Fitting a linear model yields $\hat{f}(\mathbf{x}) \approx 0.3x_1 - 0.3x_2 + x_3$



Hexbin plot of (x_1, x_2) before (left) and after (center) permuting x_1 ; PFI scores (right).


- ⇒ x_1, x_2 cancel in \hat{f} since $x_1 \approx x_2$, hence $0.3x_1 - 0.3x_2 \approx 0 \rightsquigarrow$ should be irrelevant
- ⇒ Permuting x_1 breaks joint structure \rightsquigarrow unrealistic inputs
- ⇒ $PFI > 0$ due to extrapolation (PFI evaluates model on unrealistic inputs)
 $\rightsquigarrow x_1, x_2$ are misleadingly considered relevant



COMMENTS ON PFI - EXTRAPOLATION



Example: Let $y = x_3 + \epsilon_y$, with $\epsilon_y \sim \mathcal{N}(0, 0.1)$.

- $x_1 := \epsilon_1, x_2 := x_1 + \epsilon_2$; highly correlated ($\epsilon_1 \sim \mathcal{N}(0, 1), \epsilon_2 \sim \mathcal{N}(0, 0.01)$)
- $x_3 := \epsilon_3, x_4 := \epsilon_4$, with $\epsilon_3, \epsilon_4 \sim \mathcal{N}(0, 1)$; all noise terms ϵ_j are indep.
- Fitting a linear model yields $\hat{f}(\mathbf{x}) \approx 0.3x_1 - 0.3x_2 + x_3$



Hexbin plot of (x_1, x_2) before (left) and after (center) permuting x_1 ; PFI scores (right).

- ⇒ x_1, x_2 cancel in \hat{f} since $x_1 \approx x_2$, hence $0.3x_1 - 0.3x_2 \approx 0$
 \rightsquigarrow should be irrelevant
- ⇒ Permuting x_1 breaks joint structure \rightsquigarrow unrealistic inputs
- ⇒ $PFI > 0$ due to extrapolation (PFI evaluates model on unrealistic inputs)
 $\rightsquigarrow x_1, x_2$ are misleadingly considered relevant




COMMENTS ON PFI - INTERACTIONS

Example: Let x_1, \dots, x_4 be independently and uniformly sampled from $\{-1, 1\}$ and

$$y := x_1 x_2 + x_3 + \epsilon_Y \text{ with } \epsilon_Y \sim N(0, 1)$$

Fitting a LM yields $\hat{f}(x) \approx x_1 x_2 + x_3$.




COMMENTS ON PFI - INTERACTIONS

Example: Let x_1, \dots, x_4 be independently and uniformly sampled from $\{-1, 1\}$ and

$$y := x_1 x_2 + x_3 + \epsilon_Y \text{ with } \epsilon_Y \sim N(0, 1)$$

Fitting a LM yields $\hat{f}(x) \approx x_1 x_2 + x_3$.



COMMENTS ON PFI - INTERACTIONS



Example: Let x_1, \dots, x_4 be independently and uniformly sampled from $\{-1, 1\}$ and

$$y := x_1 x_2 + x_3 + \epsilon_Y \text{ with } \epsilon_Y \sim N(0, 1)$$

Fitting a LM yields $\hat{f}(x) \approx x_1 x_2 + x_3$.

Although x_3 alone contributes as much to the prediction as x_1 and x_2 jointly, all three are considered equally relevant.

⇒ PFI does not fairly attribute the performance to the individual features.



COMMENTS ON PFI - INTERACTIONS



Example: Let x_1, \dots, x_4 be independently and uniformly sampled from $\{-1, 1\}$ and

$$y := x_1 x_2 + x_3 + \epsilon_Y \text{ with } \epsilon_Y \sim N(0, 1)$$

Fitting a LM yields $\hat{f}(x) \approx x_1 x_2 + x_3$.

Although x_3 alone contributes as much to the prediction as x_1 and x_2 jointly, all three are considered equally relevant.


⇒ PFI does not fairly attribute the performance to the individual features.



COMMENTS ON PFI - TRAIN VS. TEST DATA


Example:

- x_1, \dots, x_{20}, y are independently sampled from $\mathcal{U}(-10, 10)$
- Train set: $n = 50$ (intentionally small) and large test set
- Model: xgboost with default settings (overfits strongly)



Observation:


- PFI on train data highlights features that the model overfitted to.
- PFI on test data detects no relevant features.



COMMENTS ON PFI - TRAIN VS. TEST DATA


Example:

- x_1, \dots, x_{20}, y are independently sampled from $\mathcal{U}(-10, 10)$
- Train set: $n = 50$ (intentionally small) and large test set
- Model: xgboost with default settings (overfits strongly)



Observation:


- PFI on train data highlights features that the model overfitted to.
- PFI on test data detects no relevant features.



COMMENTS ON PFI - TRAIN VS. TEST DATA

Example:


- x_1, \dots, x_{20}, y are independently sampled from $\mathcal{U}(-10, 10)$
- Train set: $n = 50$ (intentionally small) and large test set
- Model: xgboost with default settings (overfits strongly)



Observation:

- PFI on train data highlights features that the model overfitted to.
- PFI on test data detects no relevant features.


Why? $PFI \neq 0$ if permuting a feature breaks a dependency the model relies on.
Model overfits due to spurious feature-target dependencies in train that vanish on test.
⇒ To identify features that help the model to generalize, compute PFI on test data.



COMMENTS ON PFI - TRAIN VS. TEST DATA

Example:


- x_1, \dots, x_{20}, y are independently sampled from $\mathcal{U}(-10, 10)$
- Train set: $n = 50$ (intentionally small) and large test set
- Model: xgboost with default settings (overfits strongly)



Observation:

- PFI on train data highlights features that the model overfitted to.
- PFI on test data detects no relevant features.

Why? $PFI \neq 0$ if permuting a feature breaks a dependency the model relies on. Model overfits due to spurious feature-target dependencies in train that vanish on test.
⇒ To find features that help the model to generalize, compute PFI on test data.




IMPLICATIONS OF PFI

Can we get insight into whether the ...

- ➊ feature x_j is causal for the prediction?

- $PFI_j \neq 0 \Rightarrow$ model relies on x_j
- As the train vs. test data example shows, the converse does not hold




IMPLICATIONS OF PFI

Can we get insight into whether the ...

- ➊ feature x_j is causal for the prediction?

- $PFI_j \neq 0 \Rightarrow$ model relies on x_j
- As the train vs. test data example shows, the converse does not hold



IMPLICATIONS OF PFI


Can we get insight into whether the ...

- ➊ feature x_j is causal for the prediction?

- $PFI_j \neq 0 \Rightarrow$ model relies on x_j
 - As the train vs. test data example shows, the converse does not hold

- ➋ feature x_j contains prediction-relevant information?

- $PFI_j \neq 0 \Rightarrow x_j$ is dependent on y , x_{-j} , or both (due to extrapolation)
 - x_j is not exploited by model (regardless of whether it is useful for y or not)
 $\Rightarrow PFI_j = 0$



IMPLICATIONS OF PFI


Can we get insight into whether the ...

- ➊ feature x_j is causal for the prediction?

- $PFI_j \neq 0 \Rightarrow$ model relies on x_j
 - As the train vs. test data example shows, the converse does not hold

- ➋ feature x_j contains prediction-relevant information?


- $PFI_j \neq 0 \Rightarrow x_j$ is dependent on y , x_{-j} , or both (due to extrapolation)
 - x_j is not exploited by model (regardless of its usefulness for y)
 $\Rightarrow PFI_j = 0$



IMPLICATIONS OF PFI

Can we get insight into whether the ...

- ➊ feature x_j is causal for the prediction?
 - $PFI_j \neq 0 \Rightarrow$ model relies on x_j
 - As the train vs. test data example shows, the converse does not hold
- ➋ feature x_j contains prediction-relevant information?
 - $PFI_j \neq 0 \Rightarrow x_j$ is dependent on y , x_{-j} , or both (due to extrapolation)
 - x_j is not exploited by model (regardless of whether it is useful for y or not)
 $\Rightarrow PFI_j = 0$
- ➌ model requires access to x_j to achieve its prediction performance?
 - As the extrapolation example demonstrates, such insight is not possible



IMPLICATIONS OF PFI

Can we get insight into whether the ...

- ➊ feature x_j is causal for the prediction?
 - $PFI_j \neq 0 \Rightarrow$ model relies on x_j
 - As the train vs. test data example shows, the converse does not hold
- ➋ feature x_j contains prediction-relevant information?
 - $PFI_j \neq 0 \Rightarrow x_j$ is dependent on y , x_{-j} , or both (due to extrapolation)
 - x_j is not exploited by model (regardless of its usefulness for y)
 $\Rightarrow PFI_j = 0$
- ➌ model requires access to x_j to achieve its prediction performance?
 - As shown by the extrapolation example, such insight is not possible

