

Mathematical Concepts 1

**Exercise 1: Gradient**

Consider the bivariate function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}, (x_1, x_2) \mapsto x_1^2 + 0.5x_2^2 + x_1x_2$ .

- (a) Show that  $f$  is smooth (as defined in the lecture).
- (b) Find the direction of greatest increase of  $f$  at  $\mathbf{x} = (1, 1)$ .
- (c) Find the direction of greatest decrease of  $f$  at  $\mathbf{x} = (1, 1)$ .
- (d) Find a direction in which  $f$  does not instantly change at  $\mathbf{x} = (1, 1)$ .
- (e) Assume there exists a differentiable parametrization of a curve  $\tilde{\mathbf{x}} : \mathbb{R} \rightarrow \mathbb{R}^2, t \mapsto \tilde{\mathbf{x}}(t)$  such that  $\forall t \in \mathbb{R} : f(\tilde{\mathbf{x}}(t)) = f(1, 1)$ . Show that at each point of the curve  $\tilde{\mathbf{x}}$  the tangent line  $\frac{\partial \tilde{\mathbf{x}}}{\partial t}$  is perpendicular to the gradient  $\nabla f(\tilde{\mathbf{x}})$ .
- (f) Interpret (d), (e) geometrically

**Exercise 2: Convexity**

Consider two convex functions  $f, g : \mathbb{R} \rightarrow \mathbb{R}$ .

- (a) Show that  $f + g : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto f(x) + g(x)$  is convex.
- (b) Now, assume that  $g$  is additionally non-decreasing, i.e.,  $g(y) \geq g(x) \forall x \in \mathbb{R}, \forall y \in \mathbb{R}$  with  $y > x$ . Show that  $g \circ f$  is convex.

**Exercise 3: Taylor polynomials**

Consider the bivariate function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}, (x_1, x_2) \mapsto \exp(\pi \cdot x_1) - \sin(\pi \cdot x_2) + \pi \cdot x_1 \cdot x_2$

- (a) Compute the gradient of  $f$  for an arbitrary  $\mathbf{x}$ .
- (b) Compute the Hessian of  $f$  for an arbitrary  $\mathbf{x}$ .
- (c) State the first order taylor polynomial  $T_{1,\mathbf{a}}(\mathbf{x})$  expanded around the point  $\mathbf{a} = (0, 1)$ .
- (d) State the second order taylor polynomial  $T_{2,\mathbf{a}}(\mathbf{x})$  expanded around the point  $\mathbf{a} = (0, 1)$ .
- (e) Determine if  $T_{2,\mathbf{a}}$  is a convex function.