Optimization in Machine Learning

Mathematical Concepts
Conditions for optimality



Learning goals

- Local and global optima
- First & second order conditions

DEFINITION LOCAL AND GLOBAL MINIMUM

Given $\mathcal{S} \subseteq \mathbb{R}^d$, $f : \mathcal{S} \to \mathbb{R}$:

- f has **global minimum** in $\mathbf{x}^* \in \mathcal{S}$, if $f(\mathbf{x}^*) \leq f(\mathbf{x})$ for all $\mathbf{x} \in \mathcal{S}$
- f has a **local minimum** in $\mathbf{x}^* \in \mathcal{S}$, if $\epsilon > 0$ exists s.t. $f(\mathbf{x}^*) \leq f(\mathbf{x})$ for all $\mathbf{x} \in B_\epsilon(\mathbf{x}^*)$ ("ϵ"-ball around $\mathbf{x}^*$).





Source (**left**): `https://en.wikipedia.org/wiki/Maxima_and_minima`.

Source (**right**): `https://wngaw.github.io/linear-regression/`.

EXISTENCE OF OPTIMA



We regard the two main cases of $f : \mathcal{S} \to \mathbb{R}$:

- f **continuous**: If $\mathcal{S}$ is **compact**, f attains a minimum and a maximum (extreme value theorem).
- f **discontinuous**: **No general** statement possible about existence of optima.

Example: $\mathcal{S} = [0, 1]$ compact, f discontinuous with

$$f(x) = \begin{cases} 1/x & \text{if } x > 0, \\ 0 & \text{if } x = 0. \end{cases}$$

FIRST ORDER CONDITION FOR OPTIMALITY

Observation: At an interior local optimum of $f \in \mathcal{C}^1$, first order Taylor approximation is flat, i.e., first order derivatives are zero.

This condition is therefore **necessary** and called **first order**.



Strictly convex functions (**left:** univariate, **right:** multivariate) with unique local minimum, which is the global one. Tangent (hyperplane) is perfectly flat at the optimum. (Source: Watt, *Machine Learning Refined*, 2020)

SECOND ORDER CONDITION FOR OPTIMALITY

Second order condition: Hessian of $f \in \mathcal{C}^2$ at stationary point $\mathbf{x}^* \in \mathcal{S}$ is positive or negative definite:

$$H(\mathbf{x}^*) \succ 0 \text{ or } H(\mathbf{x}^*) \prec 0$$



Interpretation: Curvature of f at local optimum is either positive in all directions or negative in all directions.

The second order condition is **sufficient** for a stationary point.

Proof: Later.

CONDITIONS FOR OPTIMALITY AND CONVEXITY

Let $f : \mathcal{S} \to \mathbb{R}$ be **convex**. Then:

- Any local minimum is **also global** minimum
- If f **strictly convex**, f has **at most one** local minimum which would also be unique global minimum on $\mathcal{S}$



Three quadratic forms. **Left:** $H(\mathbf{x}^*)$ has two positive eigenvalues. **Middle:** $H(\mathbf{x}^*)$ has positive and negative eigenvalue. **Right:** $H(\mathbf{x}^*)$ has positive and a zero eigenvalue.