



Nelder-Mead method



- General idea
- Reflection, expansion, contraction
- Advantages & disadvantages
- Examples

- General idea
- Reflection, expansion, contraction
- Advantages & disadvantages
- Examples

NELDER-MEAD METHOD

- Derivative-free method \Rightarrow heuristic
- Generalization of bisection in d -dimensional space
- Based on d -simplex, defined by $d + 1$ points:
 - $d = 1$ interval
 - $d = 2$ triangle
 - $d = 3$ tetrahedron
 - ...



NELDER-MEAD METHOD / 2



A version of the **Nelder-Mead** method:

Initialization: Choose $d + 1$ random, affinely independent points \mathbf{v}_i (\mathbf{v}_i are vertices: corner points of the simplex/polytope).

❶ **Order:** Order points according to ascending function values

$$f(\mathbf{v}_1) \leq f(\mathbf{v}_2) \leq \dots \leq f(\mathbf{v}_d) \leq f(\mathbf{v}_{d+1}).$$




with \mathbf{v}_1 best point, \mathbf{v}_{d+1} worst point.



NELDER-MEAD METHOD / 3

② Compute **centroid** without worst point

$$\bar{\mathbf{v}} = \frac{1}{d} \sum_{i=1}^d \mathbf{v}_i.$$




NELDER-MEAD METHOD / 4

③ **Reflection:** Compute reflection point

$$\mathbf{v}_r = \bar{\mathbf{v}} + \rho(\bar{\mathbf{v}} - \mathbf{v}_{d+1}),$$

with $\rho > 0$. Compute $f(\mathbf{v}_r)$.



Note: Default value for reflection coefficient: $\rho = 1$



NELDER-MEAD METHOD / 5

Distinguish three cases:

- **Case 1:** $f(\mathbf{v}_1) \leq f(\mathbf{v}_r) < f(\mathbf{v}_d)$



\Rightarrow Accept \mathbf{v}_r and discard \mathbf{v}_{d+1}

- **Case 2:** $f(\mathbf{v}_r) < f(\mathbf{v}_1)$

⇒ **Expansion:**

$$\mathbf{v}_e = \bar{\mathbf{v}} + \chi(\mathbf{v}_r - \bar{\mathbf{v}}), \quad \chi > 1.$$

We discard \mathbf{v}_{d+1} and accept the better of \mathbf{v}_r and \mathbf{v}_e .



Note: Default value for expansion coefficient: $\chi = 2$

NELDER-MEAD

Advantages:

- No gradients needed
- Robust, often works well for non-differentiable functions.


Drawbacks:

- Relatively slow (not applicable in high dimensions)
- Not each step improves solution, only mean of corner values is reduced.
- No guarantee for convergence to local optimum / stationary point.

Visualization:



<http://www.benfrederickson.com/numerical-optimization/>

Note: Nelder-Mead is default method of R function `optim()`. If gradient is available and cheap, L-BFGS is preferred.





NELDER-MEAD VISUALIZATION IN 2D

$$\min_{\mathbf{x}} f(x_1, x_2) = x_1^2 + x_2^2 + x_1 \cdot \sin x_2 + x_2 \cdot \sin x_1$$





NELDER-MEAD VISUALIZATION IN 2D

$$\min_{\mathbf{x}} f(x_1, x_2) = x_1^2 + x_2^2 + x_1 \cdot \sin x_2 + x_2 \cdot \sin x_1$$





NELDER-MEAD VISUALIZATION IN 2D

$$\min_{\mathbf{x}} f(x_1, x_2) = x_1^2 + x_2^2 + x_1 \cdot \sin x_2 + x_2 \cdot \sin x_1$$




NELDER-MEAD VISUALIZATION IN 2D

$$\min_{\mathbf{x}} f(x_1, x_2) = x_1^2 + x_2^2 + x_1 \cdot \sin x_2 + x_2 \cdot \sin x_1$$



NELDER-MEAD VS. GD / 2



Even for only 5 cities, Nelder-Mead (bottom) performs poorly. However, gradient descent (top) still works.