


# Introduction to Machine Learning

## Regularization

## Perspectives on Ridge Regression (Deep-Dive)



### Learning goals

- Interpretation of  $L2$  regularization as row-augmentation
- Interpretation of  $L2$  regularization as minimizing risk under feature noise



## L2 AND ROW-AUGMENTATION I

We can also recover the ridge estimator by performing least-squares on a **row-augmented** data set: Let  $\tilde{\mathbf{X}} := \begin{pmatrix} \mathbf{X} \\ \sqrt{\lambda} \mathbf{I}_p \end{pmatrix}$  and  $\tilde{\mathbf{y}} := \begin{pmatrix} \mathbf{y} \\ \mathbf{0}_p \end{pmatrix}$ .

With the augmented data, the unreg. least-squares solution  $\tilde{\boldsymbol{\theta}}$  is:

$$\begin{aligned}\tilde{\boldsymbol{\theta}} &= \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^{n+p} \left( y^{(i)} - \boldsymbol{\theta}^T \mathbf{x}^{(i)} \right)^2 \\ &= \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^n \left( y^{(i)} - \boldsymbol{\theta}^T \mathbf{x}^{(i)} \right)^2 + \sum_{j=1}^p \left( 0 - \sqrt{\lambda} \theta_j \right)^2 \\ &= \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^n \left( y^{(i)} - \boldsymbol{\theta}^T \mathbf{x}^{(i)} \right)^2 + \lambda \|\boldsymbol{\theta}\|_2^2\end{aligned}$$

$\implies \hat{\boldsymbol{\theta}}_{\text{ridge}}$  is the least-squares solution  $\tilde{\boldsymbol{\theta}}$  but using  $\tilde{\mathbf{X}}, \tilde{\mathbf{y}}$  instead of  $\mathbf{X}, \mathbf{y}$ !

This is a sometimes useful “recasting” or “rewriting” for ridge.



# L2 AND NOISY FEATURES I

Now consider perturbed features  $\tilde{\mathbf{x}}^{(i)} := \mathbf{x}^{(i)} + \boldsymbol{\delta}^{(i)}$  where  $\boldsymbol{\delta}^{(i)} \stackrel{iid}{\sim} (\mathbf{0}, \lambda \mathbf{I}_p)$ .

We assume no specific distribution. Now minimize risk with L2 loss, we define it slightly different than usual, as here our data  $\mathbf{x}^{(i)}$ ,  $y^{(i)}$  are fixed, but we integrate over the random permutations  $\boldsymbol{\delta}$ :



$$\mathcal{R}(\boldsymbol{\theta}) := \mathbb{E}_{\boldsymbol{\delta}} \left[ \sum_{i=1}^n (y^{(i)} - \boldsymbol{\theta}^\top \tilde{\mathbf{x}}^{(i)})^2 \right] = \mathbb{E}_{\boldsymbol{\delta}} \left[ \sum_{i=1}^n (y^{(i)} - \boldsymbol{\theta}^\top (\mathbf{x}^{(i)} + \boldsymbol{\delta}^{(i)}))^2 \right] \quad \Big| \text{ expand}$$

$$\mathcal{R}(\boldsymbol{\theta}) = \mathbb{E}_{\boldsymbol{\delta}} \left[ \sum_{i=1}^n ((y^{(i)} - \boldsymbol{\theta}^\top \mathbf{x}^{(i)})^2 - 2\boldsymbol{\theta}^\top \boldsymbol{\delta}^{(i)}(y^{(i)} - \boldsymbol{\theta}^\top \mathbf{x}^{(i)}) + \boldsymbol{\theta}^\top \boldsymbol{\delta}^{(i)} \boldsymbol{\delta}^{(i)\top} \boldsymbol{\theta}) \right]$$

By linearity of expectation,  $\mathbb{E}_{\boldsymbol{\delta}}[\boldsymbol{\delta}^{(i)}] = \mathbf{0}_p$  and  $\mathbb{E}_{\boldsymbol{\delta}}[\boldsymbol{\delta}^{(i)} \boldsymbol{\delta}^{(i)\top}] = \lambda \mathbf{I}_p$ , this is

$$\begin{aligned} \mathcal{R}(\boldsymbol{\theta}) &= \sum_{i=1}^n ((y^{(i)} - \boldsymbol{\theta}^\top \mathbf{x}^{(i)})^2 - 2\boldsymbol{\theta}^\top \mathbb{E}_{\boldsymbol{\delta}}[\boldsymbol{\delta}^{(i)}](y^{(i)} - \boldsymbol{\theta}^\top \mathbf{x}^{(i)}) + \boldsymbol{\theta}^\top \mathbb{E}_{\boldsymbol{\delta}}[\boldsymbol{\delta}^{(i)} \boldsymbol{\delta}^{(i)\top}] \boldsymbol{\theta}) \\ &= \sum_{i=1}^n (y^{(i)} - \boldsymbol{\theta}^\top \mathbf{x}^{(i)})^2 + \lambda \|\boldsymbol{\theta}\|_2^2 \end{aligned}$$

$\implies$  Ridge regression on unperturbed features  $\mathbf{x}^{(i)}$  turns out to be the same as minimizing squared loss averaged over feature noise distribution!