



Supervised Learning

Refreshing Mathematical Tools



Learning goals


- Refresher on the basics of probability theory

PROBABILITY SPACE

Probability space. A probability space is a triple $(\Omega, \mathcal{F}, \mathbb{P})$ which is modeling a specific random experiment/process. The components are

- a sample space Ω , which is the set of all possible outcomes of the random process modeled by the probability space.
- a σ -algebra \mathcal{F} , which is a family of sets representing the allowable events of the random process modeled by the probability space. In particular, each set in \mathcal{F} is a subset of the sample space Ω .
- a probability measure \mathbb{P} , which assigns each allowable event a probability value, i.e., $\mathbb{P} : \mathcal{F} \rightarrow [0, 1]$. It satisfies the following axioms of probability:
 - *Completeness* — $\mathbb{P}(\Omega) = 1$,
 - *σ -additivity* — For any finite or countably infinite sequence of mutually disjoint events E_1, E_2, \dots , it holds that


$$\mathbb{P}(\cup_{i \geq 1} E_i) = \sum_{i \geq 1} \mathbb{P}(E_i).$$



RUNNING EXERCISE

An urn contains five blue, three green, and one red ball. Two balls are randomly selected (without replacement).

What is the sample space of this experiment?



What is the probability of each point in the sample space?

INDEPENDENCE


Independence of events:

- Two events $A, B \in \mathcal{F}$ are *independent* iff $\mathbb{P}(A \cap B) = \mathbb{P}(A)\mathbb{P}(B)$.
- Events $E_1, \dots, E_n \in \mathcal{F}$ are called *pairwise independent* iff any pair E_i, E_j with $i \neq j$ is independent.
- Events $E_1, \dots, E_n \in \mathcal{F}$ are called *mutually independent* iff for any subset $I \subset \{1, \dots, n\}$ it holds that $\mathbb{P}(\bigcap_{i \in I} E_i) = \prod_{i \in I} \mathbb{P}(E_i)$.

Note that pairwise independence does not imply mutual independence!

Example: One urn with four balls with the labels 110, 101, 011, 000. We select one ball randomly.


- For $i = 1, 2, 3$ let
 $E_i := \{\text{Selected ball has zero at the } i\text{-th position}\}.$
- $\mathbb{P}(E_i) = 1/2, i = 1, 2, 3$ and
 $\mathbb{P}(E_1 \cap E_2) = \mathbb{P}(E_1 \cap E_3) = \mathbb{P}(E_2 \cap E_3) = 1/4.$
- But $\mathbb{P}(E_1 \cap E_2 \cap E_3) = 1/4 \neq 1/8 = \mathbb{P}(E_1)\mathbb{P}(E_2)\mathbb{P}(E_3).$



RUNNING EXERCISE

An urn contains five blue, three green, and one red ball. Two balls are randomly selected (without replacement).

Consider the event $A = \{\text{First ball is red}\}$ and the event $B = \{\text{Second ball is red}\}$. Are these events independent?




Are the events independent if we put a ball back into the urn after each selection?

RANDOM VARIABLES

Random variables. A random variable X on a sample space Ω is a real-valued function on Ω , that is $X : \Omega \rightarrow \mathbb{R}$. The following observations can be made:

- A random variable defines a probability space $(\Omega_X, \mathcal{F}_X, \mathbb{P}_X)$ with $\Omega_X = \text{Im}(X)$ and $\mathbb{P}_X(A) = \mathbb{P}(\{\omega \in \Omega \mid X(\omega) \in A\})$ for any $A \subset \Omega_X$. Usually, one writes just $\mathbb{P}(X \in A)$ to denote the latter term, which is the *probability distribution of X* .
(Technical remark: \mathcal{F}_X is usually the Borel- σ -algebra on \mathbb{R} .)
- In practical applications oftentimes the original probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is not the interesting object, but rather the induced probability space by X . One is rather interested in the probability of the outcome of the random variable:
 - *Coin tossing* — Let X be the random variable counting the number of tails after 100 flips.
 - *Financial market* — Let X_t be the price of some asset in a future time t .



DISCRETE RANDOM VARIABLES

If the image Ω_X of X is discrete (e.g., finite or countably infinite), then X is called a *discrete RV*.

For a discrete RV X , the function

$$p: \Omega_X \rightarrow [0, 1], x \mapsto \mathbb{P}(X \in \{x\}) = \mathbb{P}(X = x)$$

is called a probability function or probability mass function of X .

Obviously, $p(x) \geq 0$ and $\sum_{x \in \Omega_X} p(x) = 1$.

Examples:

- *Bernoulli distribution*: For a binary RV with $\Omega_X = \{0, 1\}$, $X \sim \text{Ber}(\theta)$ if $p(1) = \theta$ and $p(0) = 1 - \theta$.
- *Binomial distribution*: $X \sim \text{Bin}(n, \theta)$ if

$$p(k) = \begin{cases} \binom{n}{k} \theta^k (1 - \theta)^{n-k} & \text{if } k \in \{0, \dots, n\} \\ 0 & \text{otherwise} \end{cases}.$$




CONTINUOUS RANDOM VARIABLES

X is a *continuous RV* if Ω_X is non-discrete and there exists a function $p : \mathbb{R} \rightarrow \mathbb{R}$ such that

- $p(x) \geq 0$ for all $x \in \mathbb{R}$
- $\int_{-\infty}^{\infty} p(x) dx = 1$,
- for all $a \leq b$ it holds that $\mathbb{P}(a \leq X \leq b) = \int_a^b p(x) dx$.

The function p is called the probability density function (PDF) of X .



Examples.

- *Uniform distribution*: $X \sim U(a, b)$ if

$$p(x) = \begin{cases} 1/(b-a) & \text{if } a \leq x \leq b \\ 0 & \text{otherwise} \end{cases}.$$

- *Normal/Gaussian distribution*: $X \sim \mathcal{N}(\mu, \sigma)$ if

$$p(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right).$$

CUMULATIVE DISTRIBUTION FUNCTION


The cumulative distribution function (CDF) of a random variable X is the function

$$F_X : \mathbb{R} \rightarrow [0, 1], x \mapsto \mathbb{P}(X \leq x).$$

A CDF fully characterizes a RV: If $F_X(x) = F_Y(x)$ for all $x \in \mathbb{R}$, then X and Y are identically distributed.

If X is


- discrete with probability mass function p , then for all $x \in \mathbb{R}$,
$$F_X(x) = \sum_{y \in \Omega_X \cap (-\infty, x]} p(y).$$
- continuous with probability density function p , then
$$F_X(x) = \int_{-\infty}^x p(t) dt \text{ for all } x \in \mathbb{R}, \text{ and } p(x) = F'_X(x) \text{ whenever } F_X \text{ is differentiable at } x.$$



RUNNING EXERCISE

An urn contains five blue, three green, and one red ball. Two balls are randomly selected (without replacement).

Let X be the number of green balls selected. What are the possible values of X ?



What is the cumulative distribution function of X ?

EXPECTED VALUE/EXPECTATION

Expectation is the most basic characteristic of a random variable. Let X be a random variable, then the expectation of X , denoted by $\mathbb{E}(X)$, is

$$\mathbb{E}(X) = \int x dF(x) = \begin{cases} \sum_{x \in \Omega_X} x p(x) & \text{if } X \text{ is discrete} \\ \int_{\Omega_X} x p(x) dx & \text{if } X \text{ is continuous} \end{cases}$$

provided the sum resp. the integral is well-defined and exists.

Some important properties of the expected value are:


- Linearity — For any constants $c_1, c_2 \in \mathbb{R}$ and any pair of random variables X and Y it holds that

$$\mathbb{E}(c_1 X + c_2 Y) = c_1 \mathbb{E}(X) + c_2 \mathbb{E}(Y).$$

- Transformations — If $f : \mathbb{R} \rightarrow \mathbb{R}$ is a (measurable) function, then the expectation of $f(X)$ is

$$\mathbb{E}(f(X)) = \int f(x) dF(x) = \begin{cases} \sum_{x \in \Omega_X} f(x) p(x) & \text{if } X \text{ is discrete} \\ \int_{\Omega_X} f(x) p(x) dx & \text{if } X \text{ is continuous} \end{cases}$$


(provided the sum resp. integral exists.)



RUNNING EXERCISE

An urn contains five blue, three green, and one red ball. Two balls are randomly selected (without replacement).

Let X be the number of green balls selected. What is the expected value of X ?



VARIANCE AND COVARIANCE

The variance of a RV X is defined as follows:


$$\text{Var}(X) = \mathbb{E} [(X - \mathbb{E}(X))^2] = \int_{\Omega_X} (x - \mathbb{E}(X))^2 dF(x),$$

provided the integral on the right-hand side exists.

The *standard deviation* is defined by $\sqrt{\text{Var}(X)}$.

The *covariance* between RVs X and Y is

$$\text{Cov}(X, Y) = \mathbb{E} [(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])]$$



MULTIVARIATE RANDOM VARIABLES

RVs X_1, \dots, X_n over the same probability space can be combined into a *random vector* $\mathbf{X} = (X_1, \dots, X_n)$.

Their joint distribution is specified by the joint mass/density function $p_{\mathbf{X}}$, such that for any measurable set A it holds that


$$\mathbb{P}(\mathbf{X} \in A) = \begin{cases} \sum_{(x_1, \dots, x_n) \in A} p_{\mathbf{X}}(x_1, \dots, x_n) & \text{if } X_1, \dots, X_n \text{ are discrete} \\ \int_A p_{\mathbf{X}}(x_1, \dots, x_n) dx_1 \dots dx_n & \text{if } X_1, \dots, X_n \text{ are continuous} \end{cases}$$

The marginal distribution p_1 of X_1 is given by

$$p_1(x_1) = \begin{cases} \sum_{(x_2, \dots, x_n) \in \Omega_{X_2} \times \dots \times \Omega_{X_n}} p_{\mathbf{X}}(x_1, \dots, x_n) & \text{if discrete} \\ \int_{\Omega_{X_2} \times \dots \times \Omega_{X_n}} p_{\mathbf{X}}(x_1, \dots, x_n) dx_2 \dots dx_n & \text{if continuous} \end{cases}$$

In the same way, the marginal distributions of X_2, \dots, X_n are defined.


The same type of projection (summation/integration over all remaining variables) is used to define marginal distributions on subsets of variables $(X_{i_1}, \dots, X_{i_k})$ with $\{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}$.



RUNNING EXERCISE

An urn contains five blue, three green, and one red ball. Two balls are randomly selected (without replacement).


Let X_1 be the number of green balls selected and X_2 the number of blue balls selected. What is the joint mass function of (X_1, X_2) ?



INDEPENDENCE OF RANDOM VARIABLES

- Two discrete random variables X and Y are independent iff for any $x \in \Omega_X, y \in \Omega_Y$ it holds that $p_{XY}(x, y) = p_X(x)p_Y(y)$.
- Random variables X_1, \dots, X_n are pairwise independent iff for any pair i, j and any $x_i \in \Omega_{X_i}, x_j \in \Omega_{X_j}$ it holds that $p_{X_i X_j}(x_i, x_j) = p_{X_i}(x_i)p_{X_j}(x_j)$.
- Random variables X_1, \dots, X_n are mutually independent iff for any subset $I \subset \{1, \dots, n\}$ it holds that the joint probability mass/density function of $(X_i)_{i \in I}$ is given by $\prod_{i \in I} p_{X_i}(x_i)$ for any $x_i \in \Omega_{X_i}, i \in I$.

Similar to independence of events, pairwise independence of random variables does not imply their mutual independence. If we say that random variables are independent without further specifications we are referring to mutual independence.



CONDITIONAL DISTRIBUTIONS

If (X, Y) have a joint distribution with mass function $p_{X,Y}$, then the *conditional probability mass function* for X given Y is defined by

$$p_{X|Y}(x | y) = \frac{p(X = x, Y = y)}{p(Y = y)} = \frac{p_{X,Y}(x, y)}{p_Y(y)}$$

provided $p(Y = y) > 0$.


Likewise, in the continuous case, the *conditional probability density function* is given by

$$p_{X|Y}(x | y) = \frac{p_{X,Y}(x, y)}{p_Y(y)}$$

provided $p_Y(y) > 0$. Then,

$$p(X \in A | Y = y) = \int_A p_{X|Y}(x | y) dx .$$

The soundness of this definition is less obvious than in the discrete case, due to conditioning on an event of probability 0 here.



CONDITIONAL EXPECTED VALUE/EXPECTATION


Let X and Y be random variables, then the conditional expectation of X given $Y = y$, denoted by $\mathbb{E}(X|Y = y)$, is given by

$$\mathbb{E}(X|Y = y) = \begin{cases} \sum_{x \in \Omega_X} x p_{X|Y}(x | y) & \text{discrete case} \\ \int_{\Omega_X} x p_{X|Y}(x | y) dx & \text{continuous case} \end{cases}$$

Interpretation: The expected value of X under the condition that $Y = y$ holds.

Note that $\mathbb{E}(X|Y = y)$ induces a mapping from Ω_Y to \mathbb{R} according to $y \mapsto \mathbb{E}(X|Y = y)$. This function is called the *conditional expectation of X given Y* and simply denoted by $\mathbb{E}(X|Y)$. Note that $\mathbb{E}(X|Y)$ is a random variable!

$\mathbb{E}(X|Y)$ can be also interpreted as a prediction of X under the information encoded by the random variable Y .



CONDITIONAL VARIANCE

Let X and Y be random variables, then the conditional variance of X given $Y = y$, denoted by $\text{Var}(X|Y = y)$, is given by

$$\text{Var}(X|Y = y) = \mathbb{E} [(X - \mathbb{E}[X | Y = y])^2 | Y = y]$$

Interpretation: Variance of the prediction $\mathbb{E}[X | Y]$ for X .

Note that $\text{Var}(X|Y = y)$ induces a mapping from Ω_Y to \mathbb{R} according to $y \mapsto \text{Var}(X|Y = y)$. This function is called the *conditional variance of X given Y* and simply denoted by $\text{Var}(X|Y)$. Note that $\text{Var}(X|Y)$ is a random variable!

An important property is the law of total variance:

$$\text{Var}(X) = \mathbb{E}(\text{Var}(X | Y)) + \text{Var}(\mathbb{E}(X | Y)).$$

