
Introduction to Machine Learning

Linear Support Vector Machines SVMs and Empirical Risk Minimization





Learning goals

- Know why the SVM problem can be understood as (regularized) empirical risk minimization problem
- Know that the corresponding loss is the hinge loss

REGULARIZED EMPIRICAL RISK MINIMIZATION

- We motivated SVMs from a geometrical point of view: The margin is a distance to be maximized.
- This is not really true anymore under margin violations: The slack variables are not really distances. Instead, $\gamma \cdot \zeta^{(i)}$ is the distance by which an observation violates the margin.
- This already indicates that transferring the geometric intuition from hard-margin SVMs to the soft-margin case has its limits.
- There is an alternative approach to understanding soft-margin SVMs: They are regularized empirical risk minimizers.



SOFT-MARGIN SVM WITH ERM AND HINGE LOSS

We derived this QP for the soft-margin SVM:

$$\begin{split} & \min_{\boldsymbol{\theta}, \boldsymbol{\theta}_0, \boldsymbol{\zeta}^{(i)}} & \frac{1}{2} \|\boldsymbol{\theta}\|^2 + C \sum_{i=1}^n \boldsymbol{\zeta}^{(i)} \\ & \text{s.t.} & \boldsymbol{y}^{(i)} \left(\left\langle \boldsymbol{\theta}, \boldsymbol{x}^{(i)} \right\rangle + \boldsymbol{\theta}_0 \right) \geq 1 - \boldsymbol{\zeta}^{(i)} & \forall \, i \in \{1, \dots, n\}, \\ & \text{and} & \boldsymbol{\zeta}^{(i)} \geq 0 & \forall \, i \in \{1, \dots, n\}. \end{split}$$



In the optimum, the inequalities will hold with equality (as we minimize the slacks), so $\zeta^{(i)} = 1 - y^{(i)} f(\mathbf{x}^{(i)})$, but the lowest value $\zeta^{(i)}$ can take is 0 (we do no get a bonus for points beyond the margin on the correct side). So we can rewrite the above:

$$\frac{1}{2}\|\theta\|^2 + C\sum_{i=1}^n L\left(y^{(i)}, f\left(\mathbf{x}^{(i)}\right)\right); \ L(y, f(\mathbf{x})) = \begin{cases} 1 - yf & \text{if } yf \leq 1\\ 0 & \text{if } yf > 1 \end{cases}$$

We can also write $L(y, f(\mathbf{x})) = \max(1 - yf, 0)$.

OTHER LOSSES

SVMs can easily be generalized by changing the loss function.

- Squared hinge loss / Least Squares SVM: $L(y, f(\mathbf{x})) = \max(0, (1 yf)^2)$
- Huber loss (smoothed hinge loss)
- Bernoulli/Log loss. This is L2-regularized logistic regression!
- NB: These other losses usually do not generate sparse solutions in terms of data weights and hence have no "support vectors".

