# Introduction to Machine Learning

# Information Theory
# Kullback-Leibler Divergence




**Learning goals**

- Know the KL divergence as distance between distributions
- Understand KL as expected log-difference
- Understand how KL can be used as loss
- Understand that KL is equivalent to the expected likelihood ratio

# KULLBACK-LEIBLER DIVERGENCE



We now want to establish a measure of distance between (discrete or continuous) distributions with the same support for $X \sim p(X)$:

$$D_{KL}(p\|q) = \mathbb{E}_{X\sim p}\left[\log \frac{p(X)}{q(X)}\right] = \sum_{x\in\mathcal{X}} p(x)\cdot \log \frac{p(x)}{q(x)},$$

or:

$$D_{KL}(p\|q) = \mathbb{E}_{X\sim p}\left[\log \frac{p(X)}{q(X)}\right] = \int_{x\in\mathcal{X}} p(x)\cdot \log \frac{p(x)}{q(x)}\mathrm{d}x.$$

In the above definition, we use the conventions that $0\log(0/0) = 0$, $0\log(0/q) = 0$ and $p\log(p/0) = \infty$ (based on continuity arguments where $p \to 0$). Thus, if there is any realization $x \in \mathcal{X}$ such that $p(x) > 0$ and $q(x) = 0$, then $D_{KL}(p\|q) = \infty$.

# KL-DIVERGENCE EXAMPLE

KL divergence between $p(x) = N(0,1)$ and $q(x) = LP(0,1.5)$ given by

$$D_{KL}(p\|q) = \int_{x\in\mathcal{X}} p(x) \cdot \log \frac{p(x)}{q(x)}.$$



# KL-DIVERGENCE EXAMPLE

KL divergence between $p(x) = LP(0, 1.5)$ and $q(x) = N(0, 1)$ is different since KL not symmetric



# KL-DIVERGENCE EXAMPLE

KL divergence of $p(x) = N(0, 1)$ and $q(x) = LP(0, \sigma)$ for varying $\sigma$



# INFORMATION INEQUALITY



$D_{KL}(p\|q) \geq 0$ holds always true for any pair of distributions, and holds with equality if and only if $p = q$.

We use Jensen's inequality. Let $A$ be the support of $p$:

$$
\begin{aligned}
-D_{KL}(p\|q) &= -\sum_{x \in A} p(x) \log \frac{p(x)}{q(x)} \\
&= \sum_{x \in A} p(x) \log \frac{q(x)}{p(x)} \\
&\leq \log \sum_{x \in A} p(x) \frac{q(x)}{p(x)} \\
&\leq \log \sum_{x \in \mathcal{X}} q(x) = log(1) = 0
\end{aligned}
$$

As log is strictly concave, Jensen also tells us that equality can only happen if $q(x)/p(x)$ is constant everywhere. That implies $p = q$.

# KL AS LOG-DIFFERENCE



Suppose that data is being generated from an unknown distribution $p(x)$ and we model $p(x)$ using an approximating distribution $q(x)$.

First, we could simply see KL as the expected log-difference between $p(x)$ and $q(x)$:

$$D_{KL}(p\|q) = \mathbb{E}_{X\sim p}[\log(p(X)) - \log(q(X))].$$

This is why we integrate out with respect to the data distribution $p$. A "good" approximation $q(x)$ should minimize the difference to $p(x)$.



# KL IN FITTING

In machine learning, KL divergence is commonly used to quantify how different one distribution is from another.
Because KL quantifies the difference between distributions, it can be used as a loss function between distributions.

In our example, we investigated the KL between $p = N(0, 1)$ and $q = LP(0, \sigma)$. Now, we identify an optimal $\sigma$ which minimizes the KL.



# KL AS LIKELIHOOD RATIO



- Let us assume we have some data and want to figure out whether $p(x)$ or $q(x)$ matches it better.
- How do we usually do that in stats? Likelihood ratio!

$$LR = \prod_i \frac{p(\mathbf{x}^{(i)})}{q(\mathbf{x}^{(i)})} \qquad LLR = \sum_i \log \frac{p(\mathbf{x}^{(i)})}{q(\mathbf{x}^{(i)})}$$

If for $\mathbf{x}^{(i)}$ we have $p(\mathbf{x}^{(i)})/q(\mathbf{x}^{(i)}) > 1$, then $p$ seems better, for $p(\mathbf{x}^{(i)})/q(\mathbf{x}^{(i)}) < 1$ $q$ seems better.

- Now assume that the data is generated by $p$. Can also ask:
- "How to quantify how much better does $p$ fit than $q$, on average?"

$$\mathbb{E}_p \left[ \log \frac{p(X)}{q(X)} \right]$$

That expected LLR is really KL!