# CURSE OF DIMENSIONALITY: EXAMPLE / 2



Based on the frequency of exclamation marks, we train a very simple classifier (a decision stump with split point $\mathbf{x} \equiv 0.25$):

- We divide the input space into 2 equally sized regions.
- In the second region $[0.25, 0.5]$, 7 out of 10 are spam.
- Given that at least 0.25% of all letters are exclamation marks, an email is spam with a probability of $\frac{7}{10} = 0.7$.

# CURSE OF DIMENSIONALITY: EXAMPLE / 3

Let us feed more information into our classifier. We include a feature that contains the length of the longest sequence of capital letters.




- In the 1D case we had 20 observations across 2 regions.
- In the 1D case we had 20 observations across 2 regions.
- The same number is now spread across 4 regions.
- The same number is now spread across 4 regions.

# CURSE OF DIMENSIONALITY: EXAMPLE / 4

Let us further increase the dimensionality to 3 by using the frequency of the word “your” in an email.



# CURSE OF DIMENSIONALITY: EXAMPLE / 5

- When adding a third dimension, the same number of observations is spread across 8 regions.
- In 4 dimensions the data points are spread across 16 cells, in 5 dimensions across 32 cells and so on ...
- As dimensionality increases, the data become **sparse**; some of the cells become empty.
- There might be too few data in each of the blocks to understand the distribution of the data and to model it.




Bishop, Pattern Recognition and Machine Learning, 2006

# THE HIGH-DIMENSIONAL CUBE / 2




$$a^p = \frac{1}{10} \Leftrightarrow a = \frac{1}{\sqrt[p]{10}}$$

- So: covering 10% of total volume in a cell requires cells with almost 50% of the entire range in 3 dimensions, 80% in 10 dimensions.

# THE HIGH-DIMENSIONAL SPHERE / 2

Consider a 20-dimensional sphere. Nearly all of the volume lies in its outer shell of thickness 0.2:



# HYPHERSPHERE WITHIN HYPERCUBE

Consider a p-dimensional hypersphere of radius $r$ and volume $S_p(r)$ inscribed in a p-dimensional hypercube with sides of length $2r$ and volume $C_p(r)$. Then it holds that

$$\lim_{p\to\infty} \frac{S_p(r)}{C_p(r)} = \lim_{p\to\infty} \frac{\left(\frac{\pi^{\frac{p}{2}}}{\Gamma(\frac{p}{2}+1)}\right) r^p}{(2r)^p} = \lim_{p\to\infty} \frac{\pi^{\frac{p}{2}}}{2^p\Gamma(\frac{p}{2}+1)} = 0,$$

i.e., as the dimensionality increases, most of the volume of the hypercube can be found in its corners.




Mohammed J. Zaki, Wagner Meira, Jr., Data Mining and Analysis: Fundamental Concepts and Algorithms, 2014

# HYPHERSPHERE WITHIN HYPERCUBE / 2

Consider a 10-dimensional sphere inscribed in a 10-dimensional cube.
Nearly all of the volume lies in the corners of the cube:




Note: For $r > 0$, the volume fraction $\frac{S_p(r)}{C_p(r)}$ is independent of $r$.
Note: For $r > 0$, the volume fraction $\frac{S_p(r)}{C_p(r)}$ is independent of $r$.

# GAUSSIANS IN HIGH DIMENSIONS

A further manifestation of the **curse of dimensionality** appears if we consider a standard Gaussian $N_p(\mathbf{0}, \boldsymbol{I}_p)$ in $p$ dimensions.

- After transforming from Cartesian to polar coordinates and integrating out the directional variables, we obtain an expression for the density $p(r)$ as a function of the radius $r$ (i.e., the point's distance from the origin), s.t.

$$p(r) = \frac{S_p(r) r^{p-1}}{(2\pi\sigma^2)^{p/2}} \exp\left(-\frac{r^2}{2\sigma^2}\right),$$

  where $S_p(r)$ is the surface area of the $p$-dimensional unit hypersphere.

- Thus $p(r)\delta r$ is the approximate probability mass inside a thin shell of thickness $\delta r$ located at radius $r$.

# GAUSSIANS IN HIGH DIMENSIONS / 2

- To verify this functional relationship empirically, we draw $10^4$ points from the p-dimensional standard normal distribution and plot $p(r)$ over the histogram of the points' distances to the origin:




- We can see that for large $p$ the probability mass of the Gaussian is concentrated in a fairly thin "shell" rather far away from the origin. This may seem counterintuitive, but:

# GAUSSIANS IN HIGH DIMENSIONS / 3

- For the probability mass of a hyperspherical shell it follows that

$$\int_{r-\frac{\delta r}{2}}^{r+\frac{\delta r}{2}} p(\bar{r})d\bar{r} = \int_{r-\frac{\delta r}{2} \leq ||\mathbf{x}||_2 \leq r+\frac{\delta r}{2}} f_p(\bar{\mathbf{x}})d\bar{\mathbf{x}},$$

where $f_p(\mathbf{x})$ is the density of the $p$-dimensional standard normal distribution and $p(r)$ the associated radial density.




Example: 2D normal distribution

- While $f_p$ becomes smaller with increasing $r$, the region of the integral -the hyperspherical shell- becomes bigger.

# INTERMEDIATE REMARKS

However, we can find effective techniques applicable to high-dimensional spaces if we exploit these properties of real data:

- Often the data is restricted to a manifold of a lower dimension. (Or at least the directions in the feature space over which significant changes in the target variables occur may be confined.)
- At least locally small changes in the input variables usually will result in small changes in the target variables.