MAXIMUM LIKELIHOOD

Let's consider regression from a maximum likelihood perspective. Assume:

$$y \mid \mathbf{x} \sim p(y \mid \mathbf{x}, \boldsymbol{\theta})$$

Common case: true underlying relationship f_{true} with additive noise:

$$y = f_{\text{true}}(\mathbf{x}) + \epsilon$$

where f_{true} has params $\boldsymbol{\theta}$ and ϵ a RV that follows some distribution $\mathbb{P}_\epsilon$, with $\mathbb{E}[\epsilon] = 0$. Also, assume $\epsilon \perp\!\!\!\perp \mathbf{x}$.

GAUSSIAN ERRORS - L2-LOSS

Assume $y = f_{\text{true}}(\mathbf{x}) + \epsilon$ with additive Gaussian errors, i.e. $\epsilon^{(i)} \sim \mathcal{N}(0, \sigma^2)$. Then

- Then, given i.i.d data $\mathcal{D} = \left(\left(\mathbf{x}^{(1)}, y^{(1)}\right), \ldots, \left(\mathbf{x}^{(n)}, y^{(n)}\right)\right)$ from P_{xy} the maximum-likelihood principle is to maximize the **likelihood**

$$y \mid \mathbf{x} \sim \mathcal{N}(f_{\text{true}}(\mathbf{x}), \sigma^2)$$

The likelihood is then

$$\mathcal{L}(\theta) = \prod_{i=1}^{n} p\left(y^{(i)} \mid \mathbf{x}^{(i)}, \theta\right)$$

$$\mathcal{L}(\theta) \propto \prod_{i=1}^{n} p\left(y^{(i)} \mid f\left(\mathbf{x}^{(i)} \mid \theta\right), \sigma^2\right)$$

or equivalently to minimize the **negative log-likelihood**

$$\propto \prod_{i=1}^{n} \exp\left(-\frac{1}{2\sigma^2}\left(y^{(i)} - f\left(\mathbf{x}^{(i)} \mid \theta\right)\right)^2\right)$$

$$-\ell(\theta) = -\sum_{i=1}^{n} \log p\left(y^{(i)} \mid \mathbf{x}^{(i)}, \theta\right)$$

DISTRIBUTIONS AND LOSSES

From an ML perspective we assume our hypothesis space corresponds to the space of the (parameterized) f.

Supervised learning: Likelihood function

$$\mathcal{L}(y, f(\mathbf{x} \mid \theta)) := -\log p(y \mid \mathbf{x}, \theta)$$

Then, maximum likelihood = ERM

$$\mathcal{R}_{\text{emp}}(\theta) = \sum_{i=1}^{n} L\left(y^{(i)}, f\left(\mathbf{x}^{(i)} \mid \theta\right)\right)$$

- For every error distribution $\mathbb{P}_\epsilon$ we can derive an equivalent loss function, which leads to the same point estimator for the parameter vector $\boldsymbol{\theta}$ as maximum likelihood. Formally,
  - $\hat{\theta} \in \arg\max_{\boldsymbol{\theta}} \mathcal{L}(\boldsymbol{\theta}) \implies \hat{\theta} \in \arg\min_{\boldsymbol{\theta}} -\log(\mathcal{L}(\boldsymbol{\theta}))$
- **But**: The other way around does not always work: We cannot derive a corresponding pdf or error distribution for every loss function – the Hinge loss is one prominent example, for which some probabilistic interpretation is still possible however, see Sollich 1999.
- NB: When we are only interested in the minimizer, we can ignore multiplicative or additive constants.
- We use $\propto$ as "proportional up to multiplicative and additive constants"

GAUSSIAN ERRORS - L2-LOSS

Assume $y = f_{\text{true}}(\mathbf{x}) + \epsilon$ with additive Gaussian errors, i.e. $\epsilon^{(i)} \sim \mathcal{N}(0, \sigma^2)$. Then

$$y \mid \mathbf{x} \sim N\left(f_{\text{true}}(\mathbf{x}), \sigma^2\right)$$

The likelihood is then

$$\begin{aligned}
\mathcal{L}(\theta) &= \prod_{i=1}^{n} p\left(y^{(i)} \,\middle|\, f\left(\mathbf{x}^{(i)} \mid \theta\right), \sigma^2\right) \\
&\propto \prod_{i=1}^{n} \exp\left(-\frac{1}{2\sigma^2}\left(y^{(i)} - f\left(\mathbf{x}^{(i)} \mid \theta\right)\right)^2\right)
\end{aligned}$$

GAUSSIAN ERRORS - L2-LOSS

Easy to see: minimizing neg. log-likelihood with Gaussian errors is the same as ERM with $L2$-loss:

$$
\begin{aligned}
-\ell(\boldsymbol{\theta}) &= -\log\left(\mathcal{L}(\boldsymbol{\theta})\right) \\
&\propto -\log\left(\prod_{i=1}^{n} \exp\left(-\frac{1}{2\sigma^2}\left(y^{(i)} - f\left(\mathbf{x}^{(i)} \mid \boldsymbol{\theta}\right)\right)^2\right)\right) \\
&\propto \sum_{i=1}^{n}\left(y^{(i)} - f\left(\mathbf{x}^{(i)} \mid \boldsymbol{\theta}\right)\right)^2
\end{aligned}
$$

GAUSSIAN ERRORS - L2-LOSS

- We simulate data $y \mid \mathbf{x} \sim \mathcal{N}\left(f_{\text{true}}(\mathbf{x}), 1\right)$ with $f_{\text{true}} = 0.2 \cdot \mathbf{x}$
- Let's plot empirical errors as histogram, after fitting our model with $L2$-loss
- Q-Q-plot compares empirical residuals vs. theoretical quantiles of Gaussian





DISTRIBUTIONS AND LOSSES

- For every error distribution $\mathbb{P}_\epsilon$ we can derive an equivalent loss function, which leads to the same point estimator for the parameter vector θ as maximum-likelihood. Formally,
  - $\hat{\theta} \in \arg\max_\theta \mathcal{L}(\theta) \implies \hat{\theta} \in \arg\min_\theta -\log(\mathcal{L}(\theta))$
- **But**: The other way around does not always work: We cannot derive a corresponding pdf or error distribution for every loss function – the Hinge loss is one prominent example, for which some probabilistic interpretation is still possible however, see Sollich 1999.

DISTRIBUTIONS AND LOSSES

When does the reverse direction hold?

- If we can write the loss as $L(y, f(\mathbf{x})) = L(y - f(\mathbf{x})) = L(r)$ for $r \in \mathbb{R}$, then minimizing $L(y - f(\mathbf{x}))$ is equivalent to maximizing a conditional log-likelihood $\log(p(y - f(\mathbf{x}|\theta))$ if
  - $\log(p(r))$ is affine trafo of L (undoing the $\propto$):

$$\log(p(r)) = a - bL(r), \;\; a \in \mathbb{R}, b > 0$$

  - p is a pdf (non-negative and integrates to one)

Thus, a loss L corresponds to MLE under *some* distribution if there exist $a \in \mathbb{R}$, $b > 0$ such that

$$\int_{\mathbb{R}} \exp(a - bL(r))\mathrm{d}r = 1$$