# Introduction to Machine Learning

## Boosting
## Gradient Boosting: Concept




**Learning goals**

- Understand idea of forward stagewise modelling
- Understand fitting process of gradient boosting for regression problems

# FORWARD STAGEWISE ADDITIVE MODELING



Assume a regression problem for now (as this is simpler to explain); and assume a space of base learners $\mathcal{B}$.

We want to learn an additive model:

$$f(\mathbf{x}) = \sum_{m=1}^{M} \alpha^{[m]} b(\mathbf{x}, \boldsymbol{\theta}^{[m]}).$$

Hence, we minimize the empirical risk:

$$\mathcal{R}_{\text{emp}}(f) = \sum_{i=1}^{n} L\left(y^{(i)}, f\left(\mathbf{x}^{(i)}\right)\right) = \sum_{i=1}^{n} L\left(y^{(i)}, \sum_{m=1}^{M} \alpha^{[m]} b(\mathbf{x}^{(i)}, \boldsymbol{\theta}^{[m]})\right)$$

# GRADIENT BOOSTING

The algorithm we just introduced is not really an algorithm, but rather an abstract principle. We need to find the new additive component $b\left(\mathbf{x}, \boldsymbol{\theta}^{[m]}\right)$ and its weight coefficient $\alpha^{[m]}$ in each iteration $m$. This can be done by gradient descent, but in function space.

**Thought experiment:** Consider a completely non-parametric model $f$ whose predictions we can arbitrarily define on every point of the training data $\mathbf{x}^{(i)}$. So we basically specify $f$ as a discrete, finite vector.

$$\left(f\left(\mathbf{x}^{(1)}\right), \ldots, f\left(\mathbf{x}^{(n)}\right)\right)^\top$$

This implies $n$ parameters $f\left(\mathbf{x}^{(i)}\right)$ (and the model would provide no generalization...).
Furthermore, we assume our loss function $L(\cdot)$ to be differentiable.



# GRADIENT BOOSTING

**Aim:** Define a movement in function space so we can push our current function towards the data points.

**Given:** Regression problem with one feature $x$ and target variable $y$.

**Initialization:** Set all parameters to the optimal constant value (e.g., the mean of $y$ for $L2$).



# PSEUDO RESIDUALS



How do we have to distort this function to move it towards the observations and drive loss down?

We minimize the risk of such a model with gradient descent (yes, this makes no sense, suspend all doubts for a few seconds).
So, we calculate the gradient at a point of the parameter space, that is, the derivative w.r.t. each component of the parameter vector (which is 0 for all terms with $i \neq j$):

$$\tilde{r}^{(i)} = -\frac{\partial \mathcal{R}_{\text{emp}}}{\partial f\left(\mathbf{x}^{(i)}\right)} = -\frac{\partial \sum_j L(y^{(j)}, f(\mathbf{x}^{(j)}))}{\partial f\left(\mathbf{x}^{(i)}\right)} = -\frac{\partial L\left(y^{(i)}, f\left(\mathbf{x}^{(i)}\right)\right)}{\partial f\left(\mathbf{x}^{(i)}\right)}.$$

**Reminder:** The pseudo-residuals $\tilde{r}\left(f\right)$ match the usual residuals for the squared loss:

$$-\frac{\partial L\left(y, f(\mathbf{x})\right)}{\partial f(\mathbf{x})} = -\frac{\partial 0.5(y - f(\mathbf{x}))^2}{\partial f(\mathbf{x})}$$
$$= y - f(\mathbf{x})$$



# BOOSTING AS GRADIENT DESCENT



Combining this with the iterative additive procedure of "forward stagewise modeling", we are at the spot $f^{[m-1]}$ during minimization. At this point, we now calculate the direction of the negative gradient or also called pseudo-residuals $\tilde{r}^{[m](i)}$:

$$\tilde{r}^{[m](i)} = -\left[\frac{\partial L\left(y^{(i)}, f\left(\mathbf{x}^{(i)}\right)\right)}{\partial f(\mathbf{x}^{(i)})}\right]_{f=f^{[m-1]}}$$

The gradient descent update for each vector component of $f$ is:

$$f^{[m]}(\mathbf{x}^{(i)}) = f^{[m-1]}(\mathbf{x}^{(i)}) - \alpha \frac{\partial L\left(y^{(i)}, f\left(\mathbf{x}^{(i)}\right)\right)}{\partial f^{[m-1]}(\mathbf{x}^{(i)})}.$$

This tells us how we could "nudge" our whole function $f$ in the direction of the data to reduce its empirical risk.

# GRADIENT BOOSTING

**Iteration 1:**

Let's move our function $f\left(\mathbf{x}^{(i)}\right)$ a fraction towards the pseudo-residuals with a learning rate of $\alpha = 0.6$.



# GRADIENT BOOSTING ALGORITHM



**Algorithm** Gradient Boosting Algorithm.

Initialize $\hat{f}^{[0]}(\mathbf{x}) = \arg\min_{\theta_0 \in \mathbb{R}} \sum_{i=1}^{n} L(y^{(i)}, \theta_0)$
**for** $m = 1 \rightarrow M$ **do**
    For all $i$: $\tilde{r}^{[m](i)} = -\left[\frac{\partial L(y,f)}{\partial f}\right]_{f=\hat{f}^{[m-1]}(\mathbf{x}^{(i)}), y=y^{(i)}}$
    Fit a regression base learner to the vector of pseudo-residuals $\tilde{r}^{[m]}$:
    $\hat{\boldsymbol{\theta}}^{[m]} = \arg\min_{\boldsymbol{\theta}} \sum_{i=1}^{n} (\tilde{r}^{[m](i)} - b(\mathbf{x}^{(i)}, \boldsymbol{\theta}))^2$
    Set $\alpha^{[m]}$ to $\alpha$ being a small constant value or via line search
    Update $\hat{f}^{[m]}(\mathbf{x}) = \hat{f}^{[m-1]}(\mathbf{x}) + \alpha^{[m]} b(\mathbf{x}, \hat{\boldsymbol{\theta}}^{[m]})$
**end for**
Output $\hat{f}(\mathbf{x}) = \hat{f}^{[M]}(\mathbf{x})$

Note that we also initialize the model in a loss-optimal manner.

# LINE SEARCH



The learning rate in gradient boosting influences how fast the algorithm converges. Although a small constant learning rate is commonly used in practice, it can also be replaced by a line search.

Line search is an iterative approach to find a local minimum. In the case of setting the learning rate, the following one-dimensional optimization problem has to be solved:

$$\hat{\alpha}^{[m]} = \arg\min_{\alpha} \sum_{i=1}^{n} L(y^{(i)}, f^{[m-1]}(\mathbf{x}) + \alpha b(\mathbf{x}, \hat{\boldsymbol{\theta}}^{[m]}))$$

Optionally, an (inexact) backtracking line search can be used to find the $\alpha^{[m]}$ that minimizes the above equation.