



Introduction to Machine Learning

Linear Support Vector Machines

Support Vector Machine Training




Learning goals

- Know that the SVM problem is not differentiable
- Know how to optimize the SVM problem in the primal via subgradient descent
- Know how to optimize SVM in the dual formulation via pairwise coordinate ascent

SUPPORT VECTOR MACHINE TRAINING

- Until now, we have ignored the issue of solving the various convex optimization problems.
- The first question is whether we should solve the **primal** or the **dual problem**.
- In the literature SVMs are usually trained in the dual.
- However, SVMs can be trained both in the primal and the dual – each approach has its advantages and disadvantages.
- It is not easy to create an efficient SVM solver, and often specialized approaches have been developed, we only cover basic ideas here.



TRAINING SVM IN THE PRIMAL

Unconstrained formulation of soft-margin SVM:


$$\min_{\theta, \theta_0} \quad \frac{\lambda}{2} \|\theta\|^2 + \sum_{i=1}^n L(y^{(i)}, f(\mathbf{x}^{(i)} | \theta))$$

where $L(y, f(\mathbf{x})) = \max(0, 1 - yf)$ and $f(\mathbf{x} | \theta) = \theta^T \mathbf{x} + \theta_0$.
(We inconsequentially changed the regularization constant.)

We cannot directly use GD, as the above is not differentiable.

Solutions:


- 1 Use smoothed loss (squared hinge, huber), then do GD.
NB: Will not create a sparse SVM if we do not add extra tricks.
- 2 Use **subgradient** methods.
- 3 Do stochastic subgradient descent.
Pegasos: Primal Estimated sub-GrAdient SOLver for SVM.



TRAINING SVM IN THE DUAL

The dual problem of the soft-margin SVM is

$$\begin{aligned} \max_{\alpha} \quad & \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \alpha_i \alpha_j y^{(i)} y^{(j)} \langle \mathbf{x}^{(i)}, \mathbf{x}^{(j)} \rangle \\ \text{s.t.} \quad & 0 \leq \alpha_i \leq C \quad \sum_{i=1}^n \alpha_i y^{(i)} = 0 \end{aligned}$$




We could solve this problem using coordinate ascent. That means we optimize w.r.t. α_1 , for example, while holding $\alpha_2, \dots, \alpha_n$ fixed.

But: We cannot make any progress since α_1 is determined by $\sum_{i=1}^n \alpha_i y^{(i)} = 0$!

TRAINING SVM IN THE DUAL

$$\begin{aligned} \max_{\alpha} \quad & \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \alpha_i \alpha_j y^{(i)} y^{(j)} \langle \mathbf{x}^{(i)}, \mathbf{x}^{(j)} \rangle \\ \text{s.t.} \quad & 0 \leq \alpha_i \leq C \quad \sum_{i=1}^n \alpha_i y^{(i)} = 0 \end{aligned}$$



We move on the linear constraint until the pair-optimum or the boundary (here: $C = 1$).

