



Introduction to Machine Learning

Nonlinear Support Vector Machines The Gaussian RBF Kernel





Learning goals

- Know the Gaussian (RBF) kernel
- Understand that all data sets are separable with this kernel
- Understand the effect of the kernel hyperparameter σ

RBF KERNEL

The “radial” **Gaussian kernel** is defined as

$$k(\mathbf{x}, \tilde{\mathbf{x}}) = \exp\left(-\frac{\|\mathbf{x} - \tilde{\mathbf{x}}\|^2}{2\sigma^2}\right) \quad \text{or} \quad k(\mathbf{x}, \tilde{\mathbf{x}}) = \exp(-\gamma\|\mathbf{x} - \tilde{\mathbf{x}}\|^2)$$



A straightforward extension is


$$k(\mathbf{x}, \tilde{\mathbf{x}}) = \exp\left(-(\mathbf{x} - \tilde{\mathbf{x}})^T C (\mathbf{x} - \tilde{\mathbf{x}})\right)$$

for a symmetric, positive definite matrix C .


WEIGHTED MIXTURE OF GAUSSIANS

Via the RKHS / basis function intuition we can understand the effect of the RBF kernel much better as a local model.

$$f(\mathbf{x}) = \sum_{i=1}^n \alpha_i y^{(i)} k(\mathbf{x}^{(i)}, \mathbf{x}) + \theta_0$$




All support vectors are assigned RBF "bumps", these are weighted with the dual variables / Lagrange multipliers α_i and labels $y^{(i)}$. We then "mix" these bumps together to form the decision score function. Which becomes a bumpy surface.




WEIGHTED MIXTURE OF GAUSSIANS

Via the RKHS / basis function intuition we can understand the effect of the RBF kernel much better as a local model.

$$f(\mathbf{x}) = \sum_{i=1}^n \alpha_i y^{(i)} k(\mathbf{x}^{(i)}, \mathbf{x}) + \theta_0$$




All support vectors are assigned RBF "bumps", these are weighted with the dual variables / Lagrange multipliers α_i and labels $y^{(i)}$. We then "mix" these bumps together to form the decision score function. Which becomes a bumpy surface.




WEIGHTED MIXTURE OF GAUSSIANS

Via the RKHS / basis function intuition we can understand the effect of the RBF kernel much better as a local model.

$$f(\mathbf{x}) = \sum_{i=1}^n \alpha_i y^{(i)} k(\mathbf{x}^{(i)}, \mathbf{x}) + \theta_0$$




All support vectors are assigned RBF "bumps", these are weighted with the dual variables / Lagrange multipliers α_i and labels $y^{(i)}$. We then "mix" these bumps together to form the decision score function. Which becomes a bumpy surface.




WEIGHTED MIXTURE OF GAUSSIANS

Via the RKHS / basis function intuition we can understand the effect of the RBF kernel much better as a local model.

$$f(\mathbf{x}) = \sum_{i=1}^n \alpha_i y^{(i)} k(\mathbf{x}^{(i)}, \mathbf{x}) + \theta_0$$




All support vectors are assigned RBF "bumps", these are weighted with the dual variables / Lagrange multipliers α_i and labels $y^{(i)}$. We then "mix" these bumps together to form the decision score function. Which becomes a bumpy surface.




WEIGHTED MIXTURE OF GAUSSIANS

Via the RKHS / basis function intuition we can understand the effect of the RBF kernel much better as a local model.

$$f(\mathbf{x}) = \sum_{i=1}^n \alpha_i y^{(i)} k(\mathbf{x}^{(i)}, \mathbf{x}) + \theta_0$$




All support vectors are assigned RBF "bumps", these are weighted with the dual variables / Lagrange multipliers α_i and labels $y^{(i)}$. We then "mix" these bumps together to form the decision score function. Which becomes a bumpy surface.




WEIGHTED MIXTURE OF GAUSSIANS

Via the RKHS / basis function intuition we can understand the effect of the RBF kernel much better as a local model.

$$f(\mathbf{x}) = \sum_{i=1}^n \alpha_i y^{(i)} k(\mathbf{x}^{(i)}, \mathbf{x}) + \theta_0$$




All support vectors are assigned RBF "bumps", these are weighted with the dual variables / Lagrange multipliers α_i and labels $y^{(i)}$. We then "mix" these bumps together to form the decision score function. Which becomes a bumpy surface.




WEIGHTED MIXTURE OF GAUSSIANS

Via the RKHS / basis function intuition we can understand the effect of the RBF kernel much better as a local model.

$$f(\mathbf{x}) = \sum_{i=1}^n \alpha_i y^{(i)} k(\mathbf{x}^{(i)}, \mathbf{x}) + \theta_0$$




All support vectors are assigned RBF "bumps", these are weighted with the dual variables / Lagrange multipliers α_i and labels $y^{(i)}$. We then "mix" these bumps together to form the decision score function. Which becomes a bumpy surface.




RBF KERNEL WIDTH

A large σ (or a small γ) will make the decision boundary very smooth and in the limit almost linear.



svm: kernel=radial; cost=1; gamma=0.01
Train: mmce=0.0800000; CV: mmce.test.mean=0.1100000



svm: kernel=radial; cost=1; gamma=0.08
Train: mmce=0.4766667; CV: mmce.test.mean=0.4900000

